1

National Library
of Canada

Bibliothèque nationale
du Canada

Canadian Theses Service

Services des thèses canadiennes

Ottawa, Canada
K1A 0N4

CANADIAN THESES

THÈSES CANADIENNES

NOTICE

The quality of this microfiche is heavily dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to ensure the highest quality of reproduction possible.

If pages are missing, contact the university which granted the degree.

Some pages may have indistinct print especially if the original pages were typed with a poor typewriter ribbon or if the university sent us an inferior photocopy.

Previously copyrighted materials (journal articles, published tests, etc.) are not filmed.

Reproduction in full or in part of this film is governed by the Canadian Copyright Act, R.S.C. 1970, c. C-30. Please read the authorization forms which accompany this thesis.

**THIS DISSERTATION
HAS BEEN MICROFILMED
EXACTLY AS RECEIVED**

AVIS

La qualité de cette microfiche dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

S'il manque des pages, veuillez communiquer avec l'université qui a conféré le grade.

La qualité d'impression de certaines pages peut laisser à désirer, surtout si les pages originales ont été dactylographiées à l'aide d'un ruban usé ou si l'université nous a fait parvenir une photocopie de qualité inférieure.

Les documents qui font déjà l'objet d'un droit d'auteur (articles de revue, examens publiés, etc.) ne sont pas microfilmés.

La reproduction, même partielle, de ce microfilm est soumise à la Loi canadienne sur le droit d'auteur, SRC 1970, c. C-30. Veuillez prendre connaissance des formules d'autorisation qui accompagnent cette thèse.

**LA THÈSE A ÉTÉ
MICROFILMÉE TELLE QUE
NOUS L'AVONS REÇUE**

National Library
of CanadaBibliothèque nationale
du Canada

Canadian Theses Division

Division des thèses canadiennes

Ottawa, Canada
K1A 0N4**PERMISSION TO MICROFILM — AUTORISATION DE MICROFILMER**

• Please print or type — Écrire en lettres moulées ou dactylographier

Full Name of Author — Nom complet de l'auteur

HAYES, A. CATHERINE

Date of Birth — Date de naissance

26/01/59

Country of Birth — Lieu de naissance

CANADA

Permanent Address — Résidence fixe

39 IMPERIAL AVE., OTTAWA, ONT. K1S 3C9

Title of Thesis — Titre de la thèse

"A STUDY OF THE SYSTEM CARBON DIOXIDE-
WATER - t-BUTANOL"

University — Université

CARLETON UNIVERSITY

Degree for which thesis was presented — Grade pour lequel cette thèse fut présentée

M.Sc.

Year this degree conferred — Année d'obtention de ce grade

1984

Name of Supervisor — Nom du directeur de thèse

DR. PEETER KRUIS

Permission is hereby granted to the NATIONAL LIBRARY OF
CANADA to microfilm this thesis and to lend or sell copies of
the film.The author reserves other publication rights, and neither the
thesis nor extensive extracts from it may be printed or other-
wise reproduced without the author's written permission.L'autorisation est, par la présente, accordée à la BIBLIOTHÈ-
QUE NATIONALE DU CANADA de microfilmer cette thèse et de
prêter ou de vendre des exemplaires du film.L'auteur se réserve les autres droits de publication; ni la thèse
ni de longs extraits de celle-ci ne doivent être imprimés ou
autrement reproduits sans l'autorisation écrite de l'auteur.

Date

Sept. 28/84

Signature

Catherine Hayes.

LIST OF SYMBOLS

TBA	tertiary butanol
W	water
1	solvent
2	solute
X	mole fraction
R	gas constant
k	Boltzmann's constant
T	temperature
F	Faraday's constant
N	Avagadro's number
P	total pressure
P_i	partial pressure
P°	vapour pressure, pure substance
P°	vapour pressure, Henry's law reference state
a	activity
γ	Raoult's law activity coefficient
μ	chemical potential
ΔG_M^E	excess free energy of mixing
ΔH_M^E	enthalpy of mixing
ΔS_M^E	excess entropy of mixing
k_H	Henry's law constant
ΔG_{sol}°	free energy of solution
ΔH_{sol}°	enthalpy of solution
ΔS_{sol}°	entropy of solution
ΔG_{sol}^E	excess free energy of solution
ΔG_{exc}	excess free energy of solution, SPT

LIST OF FIGURES

1. (a) Excess Enthalpy of Mixing in Alcohol-Water Mixtures
(b) Excess Mixing Curves for t-Butanol-Water Mixtures p. 10
2. Excess Molar Volumes of Various Nonelectrolytes in Aqueous Solution p. 12
3. Solubility of Various Gases in Alcohol-Water Mixtures p. 19
4. The Refractive Index of t-Butanol-Water Mixtures p. 28
5. Gas Solubility Apparatus p. 30
6. Vapour Pressures of t-Butanol-Water Mixtures at 10.00°C p. 36
7. Activity Coefficients of t-Butanol and Water p. 41
8. Excess Free Energy of Mixing for t-Butanol-Water Mixtures p. 43
9. Henry's Law Plots for Carbon Dioxide Solubility in t-Butanol-Water Mixtures at 10.00°C p. 48
10. Henry's Law Constant, k_H , as a function of t-Butanol Concentration at 25., 10., and 1.°C p. 52
11. ΔG_{sol}^* , ΔH_{sol}^* , and ΔS_{sol}^* for Carbon Dioxide in t-Butanol-Water Mixtures at 10.00°C p. 55
12. pH of t-Butanol-Water and t-Butanol-Water-CO₂ Mixtures at 25.°C p. 57
13. Titration of Aqueous Buffers with t-Butanol p. 60
14. pK_1' of Carbonic Acid in t-Butanol-Water Mixtures p. 66
15. Properties of t-Butanol-Water Mixtures p. 68
16. Rayleigh Scattering Intensity due to Concentration Fluctuations in t-Butanol-Water Mixtures p. 71
17. pH versus $1/\epsilon$ in t-Butanol-Water Mixtures p. 79
18. k_H as a Function of Mole Fraction and Volume Fraction of t-Butanol p. 84
19. ΔG_{sol}^E for CO₂ Solubility in t-Butanol-Water Mixtures at 25., 10. and 1.°C p. 87
20. (a) ΔG_{exc} and (b) ΔG_{exc}^* calculated from Scaled Particle Theory p. 101

The undersigned hereby recommends to the
Department of Chemistry acceptance of this report,
submitted by A. Catherine Hayes in partial
fulfillment of the requirements for the Degree of
Masters of Science.

Arthur Krum

Supervisor

Opewich

Chairman

ABSTRACT

The Henry's law constant for the solubility of carbon dioxide has been determined in tertiary butanol-water mixtures over the temperature range 1.-25.°C. The solubility exhibits a sharp, temperature-dependent minimum in water-rich solutions. In order to identify effects due to t-butanol-water interactions, activity coefficients for the t-butanol-water system were calculated from vapour pressure measurements. The activity coefficient of t-butanol in aqueous solution exhibits a temperature-dependent maximum in very dilute solutions which corresponds to the minimum solubility of carbon dioxide.

A model was developed in which the solubility of carbon dioxide is related to structural effects in the alcohol-water mixtures arising from hydrophobic effects. The application of the scaled particle theory indicated volume effects are also a governing factor in the gas solubility. The pH of t-butanol-water and t-butanol-water-CO₂ mixtures also reveals anomalous solvation effects.

ACKNOWLEDGEMENTS

I would like to thank Dr. Kruus and Mr. T. Patrabo for many helpful discussions and advice. I would also like to express my gratitude to those friends who helped me over the rough spots along the way.

TABLE OF CONTENTS

Statement of Approval.....	1
Abstract.....	ii
Acknowledgements.....	iii
Table of Contents.....	iv
List of Figures.....	vi
List of Tables.....	vii
List of Symbols.....	viii
 1. INTRODUCTION.....	 1
1.1 The Solubility of Nonelectrolytes in Water - Hydrophobic Hydration.....	 2
1.2 Aqueous Solutions of Alcohols.....	9
1.3 Solubility of Gases in Aqueous Alcohol Solutions.....	16
1.4 Aqueous Solutions of Carbon Dioxide.....	24
2. EXPERIMENTAL.....	27
3. RESULTS	
3.1 Vapour Pressures of t-Butanol-Water Mixtures.....	34
3.2 Solubility of Carbon Dioxide in t-Butanol-Water Mixtures.....	45
3.3 pH Measurements.....	56
4. DISCUSSION	
4.1 Component Activities in t-Butanol- Water Mixtures.....	67
4.1.1 Light-Scattering in t-Butanol- Water Mixtures.....	69
4.1.2 Hydrophobic Effects.....	72
4.1.3 Proton Activity in t-Butanol- Water Mixtures.....	75
4.2 Solubility of Carbon Dioxide in t-Butanol Water Mixtures.....	81

4.2.1	Solubility of Carbon Dioxide in Water.....	81
4.2.2	Thermodynamics of Gas Solubility in Alcohol-Water Mixtures.....	83
4.2.3	Activity Coefficient of Carbon Dioxide in t-Butanol-Water Mixtures.....	88
4.2.4	Models of Solutions of Gases in Alcohol-Water Mixtures.....	92
4.2.5	A Model for Carbon Dioxide Solubility in t-Butanol-Water Mixtures.....	94
4.2.6	Scaled Particle Theory Description of Carbon Dioxide Solubility.....	96
4.2.7	Dissociation of Carbonic Acid in t-Butanol-Water Mixtures.....	103
5.	CONCLUSIONS.....	104
6.	REFERENCES.....	105
	APPENDIX I: SPT Computer Program.....	111
	APPENDIX II: C-13 n.m.r. Experiments.....	112

LIST OF FIGURES

1. (a) Excess Enthalpy of Mixing in Alcohol-Water Mixtures (b) Excess Mixing Curves for t-Butanol-Water Mixtures	p. 10
2. Excess Molar Volumes of Various Nonelectrolytes in Aqueous Solution	p. 12
3. Solubility of Various Gases in Alcohol-Water Mixtures	p. 19
4. The Refractive Index of t-Butanol-Water Mixtures	p. 28
5. Gas Solubility Apparatus	p. 30
6. Vapour Pressures of t-Butanol-Water Mixtures at 10.00°C	p. 36
7. Activity Coefficients of t-Butanol and Water	p. 41
8. Excess Free Energy of Mixing for t-Butanol-Water Mixtures	p. 43
9. Henry's Law Plots for Carbon Dioxide Solubility in t-Butanol-Water Mixtures at 10.00°C	p. 48
10. Henry's Law Constant, k_H , as a function of t-Butanol Concentration at 25., 10., and 1.°C	p. 52
11. ΔG_{sol}^* , ΔH_{sol}^* , and ΔS_{sol}^* for Carbon Dioxide in t-Butanol-Water Mixtures at 10.00°C	p. 55
12. pH of t-Butanol-Water and t-Butanol-Water-CO ₂ Mixtures at 25.°C	p. 57
13. Titration of Aqueous Buffers with t-Butanol	p. 60
14. pK_1' of Carbonic Acid in t-Butanol-Water Mixtures	p. 66
15. Properties of t-Butanol-Water Mixtures	p. 68
16. Rayleigh Scattering Intensity due to Concentration Fluctuations in t-Butanol-Water Mixtures	p. 71
17. pH versus $1/\epsilon$ in t-Butanol-Water Mixtures	p. 79
18. k_H as a Function of Mole Fraction and Volume Fraction of t-Butanol	p. 84
19. ΔG_{sol}^E for CO ₂ Solubility in t-Butanol-Water Mixtures at 25., 10. and 1.°C	p. 87
20. (a) ΔG_{exc} and (b) ΔG_{exc}^* calculated from Scaled Particle Theory	p. 101

LIST OF TABLES

1. Experimentally Measured Vapour Pressures of t-Butanol-Water Mixtures at 10. and 1.°C	p. 35
2. Partial Pressures, Activity Coefficients and ΔG_M^E for t-Butanol-Water Mixtures	p. 38
3. Solubility of CO ₂ in Water at 25.°C	p. 46
4. Solubility of CO ₂ in t-Butanol-Water Mixtures	p. 49
5. pH of t-Butanol-Water and t-Butanol-Water-CO ₂ Mixtures at 25.°C	p. 58
6. pH of t-Butanol-Water Mixtures predicted from K_w	p. 62
7. pK_1' of Carbonic Acid in t-Butanol-Water Mixtures	p. 65
8. Solubility of Various Gases in Water	p. 82
9. Activity Coefficient of CO ₂ in t-Butanol-Water Mixtures	p. 91
10. Molecular Diameters for SPT Calculations	p. 99

LIST OF SYMBOLS

TBA	tertiary butanol
W	water
1	solvent
2	solute
X	mole fraction
R	gas constant
k	Boltzmann's constant
T	temperature
F	Faraday's constant
N	Avagadro's number
P	total pressure
P_i	partial pressure
P°	vapour pressure, pure substance
P°	vapour pressure, Henry's law reference state
a	activity
γ	Raoult's law activity coefficient
μ	chemical potential
ΔG_M^E	excess free energy of mixing
ΔH_M^E	enthalpy of mixing
ΔS_M^E	excess entropy of mixing
k_H	Henry's law constant
ΔG_{sol}°	free energy of solution
ΔH_{sol}°	enthalpy of solution
ΔS_{sol}°	entropy of solution
ΔG_{sol}^E	excess free energy of solution
ΔG_{exc}	excess free energy of solution, SPT

G_c	free energy of cavity formation
G_i	free energy of interaction
V	molar volume
$\bar{V}_1$	partial molar volume of 1
$\Phi_{V,2}$	apparent molal volume of 2
d	density
ϵ	static dielectric constant
I	ionic strength
z	ionic charge
e	electronic charge
E_j	liquid junction potential
K_w	dissociation constant of water
D	hard sphere diameter
α	polarizability
λ	wavelength
R_{id}	Rayleigh scattering intensity of an ideal solution
R_c	Rayleigh scattering intensity due to concentration fluctuations
S_o	volume of gas dissolved per kg. of liquid at STP

1. INTRODUCTION

Carbon dioxide plays an important role in both natural and industrial processes. Supercritical carbon dioxide has found wide-scale use for extractions in the food and mining industries, and in tertiary oil recovery. Although many bulk properties of carbon dioxide solutions have been characterized, little is known of the fundamental interactions of carbon dioxide with nonelectrolyte solutes in aqueous solution. The study of simple carbon dioxide-nonelectrolyte-water systems may provide models which can be applied to more complex systems.

Tertiary butanol was chosen as a cosolute in this study for several reasons: (i) it is miscible with water in all proportions, (ii) many properties of t-butanol-water solutions have been investigated, and (iii) the peculiar features of alcohol-water solutions are particularly well-developed in aqueous t-butanol due to its bulky alkyl group.

a.

b.

translational and/or rotational degrees of freedom by the solute, this in itself cannot explain the observed heat capacity or volume effects.

The most popular explanation of the observed thermodynamic properties is that based on the idea advanced by Frank and Evans (7) that the introduction of a solute causes an increase in the structure of the neighbouring water molecules. It is this ordering phenomenon in the solvent molecules that is commonly referred to as hydrophobic hydration. The thermodynamic evidence for hydrophobic hydration has been observed in aqueous solutions of rare gases, apolar nonelectrolytes and monofunctional nonelectrolytes containing a hydrophobic residue e.g. alcohols, amines, ethers, and ketones.

Many models for aqueous solutions of nonelectrolytes start with a mixture model for pure water. In this model it is proposed that two types of distinguishable molecular species are in equilibrium. One water species is highly structured, consisting of hydrogen-bonded lattice networks of low density. It can be designated as "bulky water", $(H_2O)_b$. The second water species contains a higher proportion of non-bonded protons and is considered to be "dense water", $(H_2O)_d$. The equilibrium which exists in the pure liquid:

is presumably shifted from the right to the left by the introduction of a hydrophobic solute.

Specific models which have been proposed deal with the specific natures of $(H_2O)_b$ and $(H_2O)_d$. Frank and Evans (7) coined the much abused term "icebergs" to describe the structuring of water about an apolar solute. In their own words, the word iceberg represents "... a microscopic region, either of pure water or surrounding a solute molecule or ion, in which water molecules are tied together in some sort of quasi-solid structure. It is not implied that the structure is exactly ice-like..." The existence of clathrate hydrates (8) for rare gases and hydrocarbons was cited as supporting evidence for the formation of long-lived water structures about the solute particle.

Nemethy and Scheraga (9) developed a statistical thermodynamic treatment for pure water based on the Frank and Wen (10) "flickering cluster" description of water. Frank and Wen postulated that the formation of hydrogen bonds is a co-operative process so that short-lived "clusters" of highly hydrogen-bonded regions are formed. These clusters are surrounded by monomeric water molecules and their formation is governed by local energy fluctuations. No assumption is made as to the arrangement of the molecules in the clusters.

The introduction of a hydrocarbon molecule was postulated (11) to energetically favour tetrahedrally-bonded water molecules due to attractive van der Waal's forces between this water species and the solute. The energy of a water molecule with three or less hydrogen bonds is increased by the presence of the solute molecule since it

displaces a water nearest neighbour. This means a strong water-water dipole interaction is replaced by a weaker water-solute interaction. This change in the energy levels of the hydrogen-bonded species causes a shift towards the tetrahedrally-coordinated water species and the formation of a partial "ice-cage" around the solute. The extent of this cage depends on the size of the solute and the system temperature.

The thermodynamic functions calculated for the solution of various gases agreed with the experimental data to varying degrees. There has been some criticism of the mathematical procedures employed. A glaring conceptual fault is that a tetrahedrally hydrogen-bonded water molecule cannot have a solute molecule as its nearest neighbour.

At about the same time Frank and Quist (12) extended the Pauling model (13) of water to aqueous methane solutions. According to the Pauling model, liquid water has a structure, or range of structures similar to those of the gas hydrates, which consists of a framework of ordered water containing voids which are occupied by non-bonding water molecules. The solute molecules compete for occupation of these interstitial sites. Frank and Franks (14) also proposed that $(H_2O)_b$ resembles a clathrate framework, and that $(H_2O)_d$ is a quasi-lattice liquid which mixes ideally with $(H_2O)_b$. The solute is partitioned between these two water species. The authors concluded that the large negative entropy of solution is not due to structure promotion but rather the result of the entry of a fraction of the solute

molecules into the clathrate-like environment.

In the model developed by Mikhailov (15,16), $(H_2O)_b$ resembles the lattice structure of ice and $(H_2O)_d$ water molecules occupy interstitial sites of the lattice. A solute molecule can occupy one or more cavities by the expulsion of interstitial water molecules. The water which is expelled from the voids can enter into the lattice or it can form structureless regions outside the lattice network. This third water species is assumed to mix regularly with the solute. The system can be described by the following equilibria:

The values of the equilibrium constants K , K_α and K_β can be obtained from the thermodynamic properties of a given system. The model has been successfully applied to the water-ethanol system over the entire composition range.

Glew (17) examined the concentration dependence of several properties of aqueous solutions of t-butanol, tetrahydrofuran, dioxane and acetone. The molar ratio of water to solute molecules at the maximum or minimum points in these properties was similar to the clathrate hydrate formula for a given solute. Based on this observation, he proposed that pure water is an ice-like lattice formed by bent hydrogen bonds. In dilute solutions the solute is

dissolved interstitially and is encaged by stable hydrogen-bonded water shells composed of approximately 28 water molecules. At higher concentrations, the solute breaks down the water lattice. This model provided a qualitative explanation of the molar volumes, ultrasound absorption, proton chemical shift and compressibility experimentally measured in these solutions.

Several computer simulations of very dilute solutions of nonelectrolytes have been reported (18) utilizing either molecular dynamics (19) or Monte Carlo (20,21,22) methods. Of most relevance to the present study is the Monte Carlo calculation on the hydration of t-butanol in an infinitely dilute solution (215 water molecules, one alcohol molecule) by Nakanishi et al. (22). The results of these calculations show that the potential energy and structure of water tend to be stabilized by the introduction of the alcohol molecule. The formation of two strong hydrogen bonds between t-butanol and water favours the formation of a bulky and stable hydration shell. The density diagrams for water clearly illustrated the highly localized water structure surrounding both the hydrophilic and hydrophobic portions of the alcohol. The influence of the t-butanol molecule on water structure extended at least within a sphere of 10 Å diameter.

The evidence to date indicates that the effect of the addition of a hydrophobic molecule is to increase water structure in the immediate vicinity of the solute molecule. This increased structure may take the form of a partial or

complete lattice of hydrogen bonded water molecules which encages the solute. The formation of a water shell is analagous to the formation of clathrate hydrates, in which the water cage is stabilized by dispersive interactions with the guest molecule. There is a solute concentration of maximum stabilization beyond which further addition of solute causes the breakdown of the water lattice structures.

It must be pointed out that there cannot be a complete understanding of the phenomenon of hydrophobic hydration until the structure and properties of pure water are understood.

1.2 Aqueous Solutions of Alcohols

Because of their sparing solubility in water, aqueous solutions of nonpolar solutes can only be studied over a very limited concentration range. In order to study the concentration dependence of the hydrophobic effect, many studies have been carried out on aqueous solutions of nonelectrolytes containing one polar functional group capable of hydrogen bonding to water, e.g. alcohols, ethers, amines and ketones. The alcohols have been the most studied of this group of compounds, and many of the properties of their aqueous solutions have been precisely measured.

The peculiar features of aqueous alcohol mixtures have been discussed by many reviewers (1-3,23-26). The concentration range of most interest is the water-rich region where many properties exhibit extrema. In the brief discussion which follows, emphasis will be placed on studies of the t-butanol-water system.

1.2.1 Thermodynamic Properties

The enthalpy of mixing curves for the alcohols (Figure 1a) show a concentration dependence which indicates two or more competing processes are occurring which involve hydrogen bonding between water-water, alcohol-water, and alcohol-alcohol. The excess free energy of mixing is positive over all the concentration range, so that the sign and the magnitude of the excess entropy of mixing is the controlling

FIGURE 1

- (a) Excess Enthalpy of Mixing in Alcohol-Water Mixtures
(after reference 24)
- (b) Excess Mixing Curves for t-Butanol-Water Mixtures
(after reference 23)

a.

b.

factor in the observed excess free energy. The enthalpy of mixing becomes less negative as the size of the alkyl group increases, but $T\Delta S_M^E$ does not show great sensitivity to size or shape of the alcohol. The large negative excess entropy of mixing is also observed in the solution of hydrocarbons and nonpolar gases in water, which suggests that the alcohols behave very much like soluble hydrocarbons.

At the alcohol-rich end of the concentration scale, depolymerization of alcohol aggregates occurs, thus in the higher molecular weight alcohols ΔH_M^E becomes positive. This is illustrated in the excess mixing curves for t-butanol-water (Figure 1b).

The excess enthalpy of mixing of the alcohols in water also exhibits a strong temperature dependence which indicates significant heat capacity effects. The excess heat capacity change for the alcohol, $\Delta C_{p,2}^E$, is large and positive over the entire concentration range. De Visser et al. (27) have determined the heat capacities of the t-butanol-water system in the temperature range 25-55°C. The excess molar heat capacity quickly reaches a maximum positive value at about $X_{TBA} = 0.06$. The excess apparent molal heat capacities of t-butanol exhibit a significant temperature-dependent hump at low alcohol concentrations. The heat capacity properties of the system are indicative of significant structural effects.

The volumetric properties of several alcohol-water systems also show unexpected anomalies. In Figure 2 the excess partial molar volume isotherms for several alcohols

FIGURE 2

Excess Partial Molar Volumes of Various Nonelectrolytes
in Aqueous Solution (25°C)
(after reference 3)

are compared with those of hydrophilic solutes. While in all cases $\bar{V}_2^E$ is negative, the alcohol curves exhibit minima at low concentrations. The depth of this minimum increases and moves to lower mole fraction of alcohol as the size of the alcohol alkyl group increases.

The volume properties of the t-butanol-water system (27,28) have been thoroughly investigated. The sharp minimum in the apparent molal of volume t-butanol in very dilute solutions has been attributed to the maximum structuring effect of the alcohol on neighbouring water molecules. The addition of further alcohol causes the rapid decrease in water structure and $\Phi_{V,2}$ sharply increases. The excess molar volume of water exhibits a small positive maximum in very dilute t-butanol mixtures, then quickly becomes negative.

1.2.2 Continuum Properties.

The ultrasonic absorption and relaxation in aqueous alcohol systems have been extensively studied by Blandamer (29). The ultrasonic absorption of t-butanol-water mixtures at 25° C (30,31) changes very little in the concentration range $X_{TBA}=0$ to 0.04, however over the concentration range $X_{TBA}=0.04$ to 0.08, the absorption increases dramatically. These observations have been explained in terms of a clathrate shell model (17). The sharp increase in the sound absorption is due to the rapid breakdown of the water shells.

The results of a light-scattering study in the mixtures of t-butanol and water have also been interpreted in terms

of a clathrate hydrate model (32). The authors developed an associated complex model related to the concentration fluctuations present in the solutions, and concluded that in dilute solutions the alcohol molecule is encaged by a 21-member water clathrate shell. As the alcohol concentration increases the shells are shared between solute molecules, and finally broken down when there are no longer sufficient water molecules to maintain stable structures.

Dielectric relaxation measurements on alcohol-water systems (33) provide information on the orientational motions of water in solution. In dilute alcohol solutions, the dielectric relaxation rate of water is slowed down. The shift in the dielectric relaxation frequency of water resulting from the addition of alcohol could be correlated with the corresponding entropy of transfer. The experimental results were analyzed in terms of a model in which water molecules are exchanged between the bulk solvent and the hydration layer around the alcohol. The rotational motion of water in the hydration layer was two or three times slower than in the bulk solvent.

1.2.3 Spectroscopic Studies

There have been several infra-red spectroscopic studies on alcohol-water systems (17,34-36) in which the effect of the solute on the infra-red spectrum of water is observed. Changes in the water spectrum have been associated with increased hydrogen-bonding in dilute solutions. The spectra

water. The results of these calculations indicated that the magnitude of the free energy of solution is very sensitive to the dimensions chosen for the solute and solvent molecules.

The SPT has been applied to the solution of non-polar gases in aqueous alcohol solutions by Lucas and co-workers (61,62) with some success. The thermodynamic functions for the transfer of the $(\text{Bu})_4\text{N}^+$ ion from water to aqueous mixed solvents have also been calculated using the SPT (63).

1.3 The Solubility of Non-Polar Gases in Aqueous Nonelectrolyte Mixtures.

1.3.1 Thermodynamic Description of Gas Solubility

Henry's law states that for dilute solutions of a non-dissociating solute at constant temperature the fugacity of the solute is proportional to its mole fraction:

$$f_2 = k_H X_2 \quad 1.3$$

where k_H is the Henry's law constant. As the solution becomes more dilute this law becomes more accurate, i.e.

$$\lim_{X_2 \rightarrow 0} f_2 / X_2 = k_H \quad 1.4$$

If it is assumed that the partial pressure of the solute, P_2 , behaves ideally at low pressures (up to 1 atm) then k_H can be determined from slope of P_2 versus X_2 which is linear if Henry's law is obeyed. The relationship between the molar free energy of solution, ΔG_{sol}° , and k_H is given by:

$$\Delta G_{sol}^\circ = RT \ln (k_H / \text{atm}) \quad 1.5$$

This quantity refers to the transfer of one mole of the solute from the gas phase at 1 atm partial pressure to the liquid. The molar enthalpy and entropy of solution can then be determined according to the following relationships:

$$\Delta H_{sol}^\circ = \partial(\Delta G_{sol}^\circ / T) / \partial(1/T) \quad 1.6$$

$$\Delta S_{sol}^\circ = (\Delta H_{sol}^\circ - \Delta G_{sol}^\circ) / T \quad 1.7$$

1.3.2 Experimental Studies

Battino and Clever (45) have provided a very good general review of the solubility of gases in liquids. They discuss some of the various experimental methods available for the determination of gas solubilities. The best example of current "state-of-the-art" experimental work is that of Battino and colleagues (46) who are systematically obtaining very high precision values for the solubility of several gases in water.

Ben-Naim has measured the solubility of argon in water-ethanol (47), water-p-dioxane (48) and water-methanol (49) mixtures in the range 4 - 30°C. In both water-alcohol systems the solubility, given in terms of the Ostwald absorption coefficient, exhibits a maximum closely followed by a minimum as a function of alcohol content. These extrema, which occur in dilute solution, are more pronounced at lower temperatures. As the temperature is raised the curves become smoother so that at higher temperatures the solubility increases quite smoothly. The enthalpy and entropy of solution, both of which are negative in the water-alcohol mixtures, show a steep increase in the range $X_{ROH} = 0$ to 0.25 beyond which the curves flatten out. Similar curves were found for the p-dioxane-water system; however, the solubility of argon exhibited only a minimum at about mole fraction 0.1 of p-dioxane.

By invoking the two-structure model for liquid water, Ben-Naim interpreted the shape of the solubility curves as

being due to shifts in the equilibria between the structured, icelike form of water and the monomer form. In the region $X < 0.03$ both the gas and the alcohol stabilize the structure of water by shifting the equilibrium towards the structured form of water. In the range $0.03 < X < 0.2$ the gas molecules continue to stabilize the water structure but the alcohol destabilizes it. When $X > 0.2$ both solutes have a destabilizing effect, i.e. the equilibrium is shifted towards the monomer form of water. Thus the effect of the alcohol is to reduce the size of structured clusters of water so that at $X = 0.2$ only very small clusters exist. In the p-dioxane-water system, the addition of small amounts of p-dioxane has a destabilizing effect so that no maximum in solubility is observed.

Cargill has made extensive studies of the solubility of argon (50), oxygen (51), helium and hydrogen (52) in ethanol-water mixtures and t-butanol-water mixtures in the range of 4-62°C. The solubility curves obtained for several of these studies are illustrated in Figure 3. They are similar to those observed by Ben-Naim in water-alcohol systems. The maxima and minima in solubility were most pronounced in the t-butanol-water system. From his observation of the dependence on alcohol concentration of the extrema in the solubility curves, Cargill concluded that there was some correlation between the size of the alkyl group and its stabilizing effect on water structure.

By assuming that the inflection points in the enthalpy and entropy of solution curves correspond to the total

FIGURE 3

Solubility of Various Gases in Alcohol-Water Mixtures

—— t-butanol-water mixtures
----- ethanol-water mixtures

CO₂ data: reference 53

Ar data: reference 50

He data: reference 52

destruction of the structured form of water by the added alcohol, Cargill was able to estimate the relative amounts of the structured form of water as a function of temperature.

The solubility of each gas in the water-alcohol mixtures was related to its size. Cargill concluded that the smallest gas molecule, He, best stabilized large clusters of water and the largest gas molecule studied, O_2 , best stabilized small clusters of water. The relative amounts as a function of temperature of large, medium and small clusters were estimated from the enthalpy of solution of helium, hydrogen and oxygen respectively in water. The number of small clusters increased with increasing temperature at the expense of the large clusters. This indicates a continuous dynamic equilibrium between clusters of different sizes exists, as postulated by Frank and Wen in their "flickering cluster" model of water. Within this model, Cargill described a possible mechanism of structure stabilization in which the thermal disruption of clusters is minimized by the cushioning effect of inert solute molecules.

Cargill has also measured the solubility of carbon dioxide in water-ethanol mixtures (53). No corrections were applied for the non-ideality of CO_2 in the gas phase or for the formation of bicarbonate. The dependence of the solubility of CO_2 on alcohol concentration was similar to that of other gases, however, the overall solubility of CO_2 is much higher than that of previous gases studied.

1.3.3 Theoretical Studies - Scaled Particle Theory

In the scaled particle theory (SPT) advanced by Reiss and colleagues (54) equations were derived for the reversible work required to introduce a hard sphere into a fluid of spherical particles. Based on these equations, a method was developed by Pierotti (55,56) for predicting the solubility and thermodynamic functions for solution of simple non-polar gases. The introduction of the gas molecule into the solvent is considered to consist of two steps:

1. The creation of a cavity^o in the solvent which has the same dimensions as the spherical solute particle;
2. The introduction of the solute particle into the cavity accompanied by interactions between the solute and solvent molecule.

The free energy of solution, $\Delta G_{\text{sol}}^{\circ} = RT \ln k_H$, is given by the following expression:

$$\Delta G_{\text{sol}}^{\circ} = G_c + G_i + RT \ln(RT/V) \quad 1.8$$

where G_c and G_i are the free energy of cavity formation (step 1) and the free energy of interaction (step 2) respectively; and V is the molar volume of the solvent.

G_c is a function of solvent density, temperature, pressure and the hard sphere diameters of the solute and solvent molecules. G_i includes terms for dispersion,

inductive and repulsive interactions. The dispersion and repulsive interactions can be described by the Lennard-Jones (6-12) potential energy function; the inductive term is the product of the solute polarizability and the dipole moment of the solvent.

In the SPT, the solvent molecules are assumed to be evenly distributed about the solute particle. However, some structural features of the solvent are implicitly taken into account since the experimental solvent density and molar volume is included in the calculations.

The calculations made by Pierotti for the solubility of nonpolar gases in water agreed well with experimental values. Lucas has discussed the influence of a hard-sphere solute on water structure from the point of view of the SPT (57,58). He concludes that a hard-sphere solute enhances water structure below 4 °C but lessens it at higher temperatures, so that the experimental entropies of solution, heat capacity changes and solute molal volume cannot be taken as evidence for water structure promotion at temperatures higher than 4 °C.

Stillinger (59) improved upon Pierotti's analysis for gas solution in water by incorporating the measured surface tensions and radial-distribution functions for pure water. It was pointed out that there must be an orientational effect on the water molecules forming the inner surface of the cavity. Lucas (60) made use of the Stillinger modification to the SPT to calculate gas solubilities in

water. The results of these calculations indicated that the magnitude of the free energy of solution is very sensitive to the dimensions chosen for the solute and solvent molecules.

The SPT has been applied to the solution of non-polar gases in aqueous alcohol solutions by Lucas and co-workers (61,62) with some success. The thermodynamic functions for the transfer of the $(\text{Bu})_4\text{N}^+$ ion from water to aqueous mixed solvents have also been calculated using the SPT (63).

1.4 Aqueous Solutions of Carbon Dioxide

In aqueous solutions of carbon dioxide the following equilibria exist (64):

Because it is a nonpolar gas, carbon dioxide is only sparingly soluble in water. However, it is more soluble than might be predicted on the basis of its molecular diameter and polarizability due to the formation of ionic species in aqueous solution (6).

The hydration of CO_2 , reaction 1.10, is a relatively slow reaction. The rate constant governing the forward reaction, k_{CO_2} , is on the order of 0.03 sec^{-1} (65); the rate constant for the dehydration reaction, $k_{\text{H}_2\text{CO}_3}$, is about 20 sec^{-1} (65) which indicates this process occurs very slowly. The ratio of $\text{CO}_2(\text{aq})$ to $\text{H}_2\text{CO}_3(\text{aq})$ is about 600:1 (65), so often the quantity $[\text{H}_2\text{CO}_3^*(\text{aq})]$ is defined (64):

$$\begin{aligned} [\text{H}_2\text{CO}_3^*(\text{aq})] &= [\text{CO}_2(\text{aq})] + [\text{H}_2\text{CO}_3(\text{aq})] \\ &\approx [\text{CO}_2(\text{aq})] \end{aligned} \quad 1.14$$

The ionization equilibria occur very rapidly. The formation of carbonate ions (reaction 1.12) is only

significant at pH values greater than 8 (64). The acid constant $K(\text{CO}_2)$ can be defined as (66):

$$K(\text{CO}_2) = \frac{[\text{HCO}_3^-(\text{aq})][\text{H}^+(\text{aq})]}{[\text{H}_2\text{CO}_3^*(\text{aq})]} \quad 1.15$$

and its value is usually given as the first dissociation constant of carbonic acid, 4.45×10^{-7} at 25°C . The small value of $K(\text{CO}_2)$ indicates that only very small concentration of the ionic species are formed in carbon dioxide solutions.

Reaction 1.11 and 1.12, which involve proton transfers, have rapid relaxation rates with relaxation times of the order of 10^{-9} sec (67). Thus C-13 n.m.r. experiments cannot distinguish $\text{CO}_3^{2-}(\text{aq})$ and $\text{HCO}_3^-(\text{aq})$ species. It is uncertain whether $\text{H}_2\text{CO}_3(\text{aq})$ is distinguishable; possibly because of its very low concentration no peak attributable to this species has been detected (67,68).

Separate peaks attributable to $\text{CO}_2(\text{aq})$, $\text{HCO}_3^-(\text{aq})$ and $\text{CO}_3^{2-}(\text{aq})$ have been observed in the Raman spectra of carbon dioxide solutions (69,70). The similarity of the Raman spectra of $\text{CO}_2(\text{g})$ and $\text{CO}_2(\text{aq})$ implies that the linear structure of the gas is unaltered when it is dissolved in water.

In a Raman study of the clathrate hydrate and aqueous solutions of carbon dioxide (71), the band positions of CO_2 in aqueous solution did not change appreciably up to 1.76 kbar gas pressure and were very near to those observed in the Raman spectrum of the hydrate. These observations seem to indicate that the arrangement of water molecules around a

CO_2 molecule in aqueous solution is similar to the clathrate hydrate.

The partial molar volume of CO_2 (aq) at infinite dilution in water has recently been determined from high precision density measurements (72). The value of V_2° (CO_2 , aq) was determined to be $32.8 \text{ cm}^3 \text{ mol}^{-1} \pm 1.2 \text{ cm}^3 \text{ mol}^{-1}$ by extrapolation of measurements in solutions up to 0.1 molal in carbon dioxide, where Henry's law is still obeyed. This value is lower than the molar volume of liquid CO_2 at its boiling point, which is about $40 \text{ cm}^3 \text{ mol}^{-1}$ (6).

2. EXPERIMENTAL

All water-alcohol solutions were prepared from filtered Millipore deionized water and freshly distilled Fisher Certified t-butanol. Matheson Research Grade carbon dioxide (99.995% CO_2) was used for all solubility measurements. Matheson Research Grade helium (99.9999% He) was used to calibrate the solubility apparatus.

Refractive index measurements were made with an Abbe refractometer (Officine Galileo) thermostatted at $25.00 \pm 0.05^\circ\text{C}$. Using solutions prepared from t-butanol dried over 4A molecular sieves, a calibration curve was constructed and fitted to a sixth-order polynomial (Figure 4). The compositions of all water-t-butanol solutions prepared were then calculated from refractive index measurements.

Measurements of pH were made with a combination glass/Ag-AgCl electrode (Fisher 13-639-92) and a digital pH meter (Fisher Model 620). The water-t-butanol solution was first degassed under vacuum by several freeze-pump-thaw cycles. The solution was then quickly transferred to the measuring flask and nitrogen gas was bubbled continuously through the sample. This bubbling action also provided sufficient stirring of the solution. The measuring flask was placed in a thermostatted bath at $25.00 \pm 0.05^\circ\text{C}$. The typical time for equilibration of the pH reading in samples with a nitrogen sparge was one hour. The nitrogen sparge was then replaced with a carbon dioxide sparge, and the pH reading usually equilibrated within minutes. The refractive index

FIGURE 4

The Refractive Index of
t-Butanol-Water Mixtures
at 25.°C

▲ experimental points

----computer-fitted curve

of the solution was tested after each pH reading was taken. Changes in refractive index were very small.

Vapour pressure measurements were made with a mercury manometer. The sample was prepared in the solubility cell (see Figure 5) by degassing under vacuum by several freeze-pump-thaw cycles. It was then stirred and thermostatted to $\pm 0.05^\circ\text{C}$ while the measurement of vapour pressure was made. The typical equilibration time was one hour.

The apparatus for the measurement of gas solubility is illustrated in Figure 5. It can be described as a modified BET-type apparatus used to measure the adsorption of gas on a solid. It consists of a constant-volume manometer, a seven-bulb gas burette and a solubility cell which contains the liquid sample. The volume of each bulb in the gas burette is known to $\pm 0.0002\text{ cm}^3$. The mercury level in the manometer is read with a cathetometer to $\pm 0.001\text{ cm}$. The solubility cell is constructed with a teflon "Rotoflo" stopcock so that there is no contamination of the sample by vacuum grease. the sample is stirred magnetically and thermostatted to $\pm 0.05^\circ\text{C}$. The vacuum in the system is monitored with a thermocouple gauge (Veeco type RG-31X).

The line volume, which is all capillary tubing, is defined as the volume bounded by the pointer of the manometer, the zero mark of the gas burette and stopcocks 1, 2 and 3. The dead volume is defined as the volume between stopcock 1 and the liquid level in the solubility cell. Both the line volume and the dead volume (with the sample cell empty) were determined with research grade helium to

FIGURE 5

GAS SOLUBILITY APPARATUS

0.1% precision, assuming He is an ideal gas. The line volume was $15.528 \pm 0.076 \text{ cm}^3$; the dead volume was $34.123 \pm 0.250 \text{ cm}^3$. The room (gas) temperature was monitored with a five junction chromel-alumel thermocouple referenced to an ice bath.

In order to check for any adsorption of CO_2 on the glass walls of the tubing, the line volume and dead volume were also calibrated with CO_2 , assuming ideal gas behaviour. The results were within 1.0% of those obtained with helium.

The procedure followed to measure the solubility of CO_2 in a water-t-butanol solution of a particular composition will be briefly outlined here. Solubility measurements were made at 25.00, 10.00 and 1.00°C.

The sample was first degassed under vacuum by at least five freeze-pump-thaw cycles, weighed, then stirred and thermostatted to the desired temperature. The line volume was evacuated, typically to 0.0001 torr, and then filled with CO to a pressure of about 700 torr with the mercury in the gas burette at mark 3. The number of moles of CO_2 in the line volume was determined from three volume/pressure measurements. The level of mercury in the manometer and the burette was then adjusted to give the desired pressure of CO_2 , and the gas was then admitted to the sample cell. The gas pressure was allowed to equilibrate for about one hour before the final reading was made. When equilibrium was reached, stopcock 1 was closed and the number of moles of CO_2 remaining in the line volume was measured. The number of moles of CO_2 remaining in the dead volume could be

calculated from the equilibrium pressure and the known dead volume. Thus the number moles of CO₂ absorbed by the sample could be calculated. This procedure was repeated until at least five points were obtained for the isotherm. At the end of the experiment the refractive index of the sample was measured. The typical time required for the complete experiment was 10 hours.

In calculating the number of moles of CO₂, the pressure was corrected for the non-ideality of the gas according to the equation:

$$n = PV/(RT+B(T)P) \quad 2.1$$

where B(T) is the temperature dependent second virial coefficient for CO₂ (73). For the calculation of the number of moles in the line volume, the gas temperature was assumed to be the room temperature. The temperature of the gas in the dead volume was assumed to be the temperature of the sample.

It was necessary to correct the dead volume for the volume of the sample, which required that the density of the solution be known. The densities of water-t-butanol solutions at 25°C and 10°C were found in reference 27. For measurements at 1°C the densities at 0°C found in reference 38 were used to correct the dead volume.

Clearly the gas contained in the dead volume consists of both carbon dioxide and sample vapour. It can be assumed that the line volume contains only carbon dioxide since it is capillary tubing and pressure changes from the line

volume to the dead volume were always positive. The gas pressure was corrected for the presence of sample vapour in the dead volume as follows. The mole fraction of dissolved carbon dioxide, X_{CO_2} , was first calculated from the total measured pressure. Then the partial pressure of carbon dioxide, P_{CO_2} , was calculated according to Raoult's law:

$$P_{\text{CO}_2} = P_{\text{TOT}} - (1 - X_{\text{CO}_2}) P_{\text{SOL}} \quad 2.2$$

where P_{SOL} is the vapour pressure of the solution. New X_{CO_2} values were then calculated using the partial pressure of CO_2 to determine the number of moles of CO_2 in the dead volume. This first-order correction was found to be sufficient.

3. RESULTS

3.1 Vapour Pressures of Water-t-Butanol Mixtures

The vapour pressures of water-t-butanol mixtures were measured over the concentration ranges where the samples were liquid at 10.00 C and 1.00°C. The vapour pressure of t-butanol was taken from reference 74. An attempt was made to measure the vapour pressure of water, but equilibrium could not be reached. This value was also taken from reference 74. The vapour pressures at 25.° C have been measured previously by Brown and Ives (75). The experimental vapour pressures are found in Table 1.

The shapes of the total vapour pressure curves at 10.° C and 1.°C are similar to that reported by Brown and Ives at 25.°C. The vapour pressure increases rapidly as small amounts of alcohol are added to water. In the region $X_{TBA}=0.05$ to 0.08 the curve abruptly flattens and changes slowly with further addition of alcohol. In the region $X_{TBA}=0.7$ the vapour pressure starts to decrease to the vapour pressure of pure t-butanol.

Brown and Ives used an equilibrium still to make their vapour pressure measurements and were able to determine the composition of the vapour phase with this apparatus. The partial pressure of each component was then calculated by an application of the Duhem-Margules relation. Using their data for the partial pressure of each component at 25.° C, the partial pressures at other temperatures could be calculated under the assumption that the activity

TABLE 1

Experimentally Measured Vapour Pressures of
t-Butanol-Water Mixtures

10.00°C		1.00°C	
x_{TBA}	P_{TOT}^a 10 ⁻³ atm	x_{TBA}	P_{TOT}^a 10 ⁻³ atm
±.0002		±.0002	
0.0101	12.96	0.0128	6.66
0.0197	14.72	0.0295	7.68
0.0443	18.37	0.0349	8.17
0.0478	18.99	0.0409	8.46
0.0603	20.84	0.0529	9.63
0.0705	21.57	0.0657	10.88
0.0916	22.32	0.0700	11.54
0.1037	22.50	0.0756	11.64
0.1630	23.08	0.0846	11.82
0.2860	23.55	0.1228	12.25
0.4510	23.82	0.1938	12.37
0.6230	23.95	0.2246	12.41
0.7020	23.89	0.3772	12.50
0.8140	23.34	0.4347	12.50
0.8600	22.58	0.4776	12.47
		0.5797	12.63
		0.6940	12.63
		0.7898	12.29

a. maximum experimental error: $\pm 0.07 \times 10^{-3}$ atm.

FIGURE 6

Vapour Pressures of t-Butanol-Water Mixtures
at 10.00°C

△ experimental points

— computer-fitted curves

r.m.s. deviation for computer fit of P_{TOT} :

1.00°C: $X_{TBA} = 0 - 0.1 : 0.088$

$X_{TBA} = 0.1 - 1 : 0.026$

10.00°C: $X_{TBA} = 0 - 0.1 : 0.094$

$X_{TBA} = 0.1 - 1 : 0.031$

25.00°C: $X_{TBA} = 0 - 0.1 : 0.174$

$X_{TBA} = 0.1 - 1 : 0.057$

(25.°C data from Brown and Ives (75))

$$Y = a_1 + a_2 X + a_3 X^2 + a_4 X^3 + a_5 X^4 + a_6 X^5 + a_7 X^6$$

$$X_{TBA} = 0-0.10$$

P_{TOT}	25°C	10°C	1°C
a_1	.23813E+2	.91749E+1	.49113E+1
a_2	.30166E+3	.68984E+2	.21937E+2
a_3	-.14553E+5	.11382E+4	-.13674E+4
a_4	.10259E+7	.17221E+5	.90542E+5
a_5	-.23372E+8	-.73180E+6	-.14961E+7
a_6	.21531E+9	.61463E+7	.96716E+7
a_7	-.70536E+9	-.15906E+8	-.21924E+8

$$X_{TBA} = 0.10-1.0$$

P_{TOT}	25°C	10°C	1°C
a_1	.40329E+2	.14698E+2	.84798E+1
a_2	.55989E+2	.33814E+2	.94359E+1
a_3	-.23548E+3	-.14407E+3	-.28872E+2
a_4	.55465E+3	.32049E+3	.17075E+2
a_5	-.67420E+3	-.38954E+3	.64041E+2
a_6	.38428E+3	.25777E+3	-.10262E+3
a_7	-.83599E+2	-.79217E+2	.39664E+2

coefficient of the water is relatively insensitive to temperature.

These calculations were done as follows. The activity coefficient of water, γ_{H_2O} , was calculated from the partial pressure of water at 25.°C according to Raoult's law:

$$\gamma_{H_2O} = \frac{P_{H_2O}}{X_{H_2O} P_{H_2O}^*} \quad 3.1$$

Then assuming that the activity coefficient of water changes negligibly (within experimental error) in the temperature range 1-25 °C, the partial pressure of water at a given temperature could be calculated from the relation:

$$P_{H_2O}(T) = \gamma_{H_2O}(25^\circ) X_{H_2O} P_{H_2O}^*(T) \quad 3.2$$

The partial pressure and activity coefficient of t-butanol could then be determined as:

$$P_{TBA}(T) = P_{TOT}(T) - P_{H_2O}(T) \quad 3.3$$

$$\gamma_{TBA}(T) = \frac{P_{TBA}(T)}{X_{TBA} P_{TBA}^*(T)} \quad 3.4$$

The activity coefficients calculated in this manner were found to be consistent with the Gibbs-Duhem relation to within about 0.8%. The smoothed data for the system at 25, 10 and 1°C are tabulated in Table 2.

The total vapour pressure and partial pressure curves at 10.°C are shown in Figure 6. The strong positive deviation

TABLE 2

Partial Pressures, Activity Coefficients and ΔG_M^E for
Water-t-Butanol Mixtures.

1.00°C						
X_{TBA}	P_{TOT}^a	P_{H_2O} 10 ⁻³ atm	P_{TBA}	γ_{H_2O}	γ_{TBA}	ΔG_M^E kJ mol ⁻¹
0	6.48	6.48	0	1.000	----	0
0.01	6.67	6.38	0.29	0.995	3.06	0.142
0.02	7.00	6.32	0.68	0.993	3.63	0.430
0.03	7.62	6.25	1.37	0.994	4.81	0.942
0.04	8.51	6.21	2.30	0.997	6.08	0.158
0.05	9.53	6.17	3.36	1.002	7.08	0.227
0.06	10.49	6.13	4.34	1.007	7.65	0.293
0.07	11.26	6.12	5.14	1.015	7.75	0.358
0.08	11.76	6.11	5.66	1.02	7.47	0.414
0.09	11.99	6.09	5.91	1.03	6.92	0.462
0.10	12.04	6.08	5.96	1.04	6.30	0.504
0.20	12.39	6.03	6.37	1.16	3.36	0.828
0.30	12.46	5.92	6.54	1.31	2.30	0.996
0.40	12.47	5.76	6.71	1.48	1.77	1.060
0.50	12.54	5.54	7.00	1.71	1.48	1.055
0.60	12.65	5.12	7.53	1.97	1.32	1.003
0.70	12.62	4.37	8.25	2.25	1.24	0.901
0.80	12.22	3.28	8.95	2.53	1.18	0.726
0.90	11.22	1.89	9.33	2.93	1.09	0.429
1	9.47	0	9.47	3.78	1.00	0

TABLE 2 (con't)

10.00°C

x_{TBA}	P_{TOT}	P_{H_2O} 10 ⁻³ atm	P_{TBA}	γ_{H_2O}	γ_{TBA}	ΔG_M^E kJ mol ⁻¹
0	12.12	12.12	0	1.000	----	0.032
0.01	13.14	11.93	1.21	0.995	6.60	0.032
0.02	14.54	11.80	2.74	0.993	7.47	0.076
0.03	16.16	11.68	4.47	0.994	8.12	0.130
0.04	17.83	11.61	6.22	0.997	8.48	0.188
0.05	19.37	11.53	7.84	1.002	8.54	0.249
0.06	20.63	11.47	9.16	1.007	8.32	0.305
0.07	21.55	11.43	10.12	1.015	7.87	0.361
0.08	22.09	11.41	10.68	1.02	7.28	0.410
0.09	22.34	11.38	10.96	1.03	6.64	0.454
0.10	22.43	11.37	11.08	1.04	6.03	0.494
0.20	23.32	11.28	12.04	1.16	3.28	0.817
0.30	23.61	1.08	12.53	1.31	2.28	0.988
0.40	23.72	10.78	12.93	1.48	1.76	1.055
0.50	23.84	10.36	13.49	1.71	1.47	1.049
0.60	23.90	9.55	14.41	1.97	1.31	0.986
0.70	23.92	8.16	15.75	2.25	1.23	0.878
0.80	23.39	6.13	17.26	2.53	1.18	0.718
0.90	21.86	3.55	18.30	2.93	1.11	0.455
1.00	18.36	0	18.36	3.78	1.00	0

TABLE 2 (con't)

25.00°C
(from reference 75)

X_{TBA}	P_{TOT}^a	P_{H_2O} 10 ⁻³ atm	P_{TBA}	γ_{H_2O}	γ_{TBA}	ΔG_M^E kJ mol ⁻¹
0	31.32	31.32	0	1.000	----	0.032
0.01	34.47	30.79	3.62	0.995	6.56	0.032
0.02	38.29	30.53	7.86	0.993	7.11	0.074
0.03	43.82	30.26	13.55	0.994	8.17	0.130
0.04	49.47	30.00	19.47	0.997	8.81	0.192
0.05	53.82	29.87	24.08	1.002	8.71	0.251
0.06	56.18	29.61	26.57	1.007	8.01	0.300
0.07	56.84	29.61	27.33	1.015	7.07	0.344
0.08	57.11	29.47	27.59	1.02	6.24	0.382
0.09	57.89	29.47	28.54	1.03	5.74	0.424
0.10	58.03	29.34	28.68	1.04	5.18	0.459
0.20	60.00	29.08	30.79	1.16	2.79	0.743
0.30	60.92	28.68	32.37	1.31	1.95	0.883
0.40	61.71	27.89	33.82	1.48	1.53	0.927
0.50	62.3	26.71	35.53	1.71	1.29	0.898
0.60	62.63	24.74	37.89	1.97	1.14	0.801
0.70	62.24	21.05	41.05	2.25	1.06	0.649
0.80	60.92	15.97	45.00	2.53	1.02	0.457
0.90	58.55	9.21	49.34	2.93	0.99	0.228
1	55.26	0	55.26	3.78	1.00	0

(*) data fitted to 6th-order polynomial, see Figure 6.

FIGURE 7

Activity Coefficients of t-Butanol and Water
determined from Vapour Pressures

$$\gamma_{H_2O} = a_1 + a_2 X + a_3 X^2 + a_4 X^3 + a_5 X^4 + a_6 X^5 + a_7 X^6$$

$$\begin{aligned} a_1 &= .99825 \\ a_2 &= -.50451 \\ a_3 &= .13966 \\ a_4 &= -.57113 \\ a_5 &= .12672 \\ a_6 &= -.13156 \\ a_7 &= .51280 \end{aligned}$$

from Raoult's law at low mole fractions of t-butanol can be seen to arise from the non-ideality of the alcohol. The water vapour pressure shows a small negative deviation in very dilute solutions but a larger positive deviation occurs as the alcohol content increases.

This behavior is more clearly seen by examining the activity coefficients of t-butanol and water as a function of solution composition. These curves are shown in Figure 7. The activity coefficient of t-butanol exhibits a temperature-dependent maximum at low mole fraction of alcohol. This maximum occurs at $X_{TBA} = 0.04$ at $25.^{\circ}\text{C}$, 0.05 at $10.^{\circ}\text{C}$ and 0.07 at $1.^{\circ}\text{C}$. The negative deviation from Raoult's law of the activity coefficient of water has also been detected by Knight (76) in freezing point depression measurements on this system.

It is possible to calculate the excess functions of mixing if the activity coefficients of the components are known. The excess Gibbs free energy of mixing, ΔG_M^E , is defined by the relation:

$$\Delta G_M^E = RT(X_1 \ln \gamma_1 + X_2 \ln \gamma_2) \quad 3.5$$

The excess enthalpy of mixing, ΔH_M^E , and excess entropy of mixing, ΔS_M^E , are then defined by the relations:

$$\Delta H_M^E = \frac{\partial (\Delta G_M^E / T)}{\partial (1 / T)} \quad 3.6$$

$$-T\Delta S_M^E = (\Delta G_M^E - \Delta H_M^E) \quad 3.7$$

FIGURE 8

Excess Free Energy of Mixing
of t-Butanol-Water Mixtures

The ΔG_M^E values calculated for water-t-butanol mixtures at different temperatures are illustrated in Figure 8. The maximum ΔG_M^E values occur at $X_{TBA} = 0.4$. It was not possible to calculate ΔH_M^E or ΔS_M^E from these values because the enthalpy of mixing is highly temperature dependent, and changes significantly in the temperature range 1-25°C. This can be seen in the closeness of the ΔG_M^E curves at 10. and 1.°C. This is also indicative of variations in the excess heat capacity of the system.

3.2 SOLUBILITY OF CARBON DIOXIDE IN t-BUTANOL-WATER MIXTURES

The accuracy of the experimental method used to calculate the solubility of CO_2 can be estimated by comparing the results for the solubility of CO_2 in pure water at 25.00°C with those available in the literature. This comparison is made in Table 3. (The Henry's Law constant was converted to Bunsen coefficient and Ostwald Solubility coefficient assuming ideal gas behaviour). The value obtained in this study agrees with the literature values to within 3%. Some of this deviation arises from the ideal gas assumption.

A possible source of error in the method employed is not applying a correction to the solution volume when the gas is dissolved. However, because of the very low concentrations of gas in solution, this correction is probably negligible. For the same reason, it is not necessary to take into account changes in vapour pressure of the solution caused by solution of the gas. A significant error source may be incomplete degassing of the sample solution. High purity gas was used in all experiments, so that no significant amounts of contaminated gas were present.

The main source of error in this experiment was the assumption that the gas in the dead volume is at the same temperature as the sample. This quite clearly is not true; a temperature gradient from the sample temperature to room temperature exists above the level of the sample bulb. It is difficult to apply a correction due to this thermal

TABLE 3

SOLUBILITY OF CO₂ IN WATER AT 25.00°C

Unit of measurement	This study	Other studies (ref)
Henry's law constant	(1.210±0.007) x 10 ⁶ torr	1.256 x 10 ⁶ torr (270-988 torr) (77)
Bunsen coefficient	0.7722±0.0046	0.7565 (78)
Ostwald coefficient	0.8428±0.0051	0.8267 0.8255 0.8256 (79)

gradient so it was necessary to make the assumption of uniform dead volume temperature. Such errors would however have negligible effects in considerations of relative solubilities.

A typical series of Henry's Law plots for the solubility of CO_2 at 10.00°C is illustrated in Figure 9. The intercepts of the lines do not pass through zero pressure as expected. The small positive intercept values are probably due to the error discussed above: assumption of uniform dead volume temperature. Incomplete degassing of the sample solution would also lead to a positive intercept in the Henry's Law plot.

The results for the determination of the Henry's Law constant, k_H , as a function of alcohol concentration and temperature are tabulated in Table 4. The average experimental error is less than 1.5%. The variation in k_H is more clearly seen in Figure 10.

These curves show that the addition of very small amounts of t-butanol initially has very little effect on CO_2 solubility; as further alcohol is added, the solubility suddenly begins to decrease (i.e. k_H begins to increase) up to an alcohol mole fraction of about 0.05. The solubility then increases rapidly up to $X_{\text{TBA}} = 0.10$, after which it increases less rapidly. The relative decrease in solubility becomes greater at lower temperatures.

This minimum in gas solubility at low alcohol concentration has been observed in other studies by Cargill (50-52) and Ben-Naim (47,49) in experiments using helium,

FIGURE 9

Henry's Law Plots for CO₂ Solubility
in t-Butanol-Water Mixtures
at 10.00°C

TABLE 4

Solubility of CO₂ in Water-t-Butanol Mixtures

T=25.00°C

x_{TBA}	k_{H} atm	$\Delta G_{\text{sol}}^{\circ}$ kJ mol ⁻¹
±.0002		
0	1592 ± 10	18.28 ± 0.11
0.0163	1599 ± 20	18.29 ± 0.23
0.300	1601 ± 17	18.29 ± 0.19
0.0346	1653 ± 30	18.37 ± 0.33
0.0567	1635 ± 27	18.34 ± 0.30
0.0623	1514 ± 26	18.15 ± 0.31
0.0840	1294 ± 3	17.76 ± 0.04
0.1300	941 ± 34	16.97 ± 0.61
0.2810	441 ± 2	15.09 ± 0.07
0.7752	156 ± 3	12.52 ± 0.24

TABLE 4 (con't)

x_{TBA} $\pm .0002$	$T = 10.00^\circ C$		ΔG_{sol}° $kJ\ mol^{-1}$
	k_H atm		
0	1066 $\pm$ 7		16.41 $\pm$ 0.11
0.0094	1084 $\pm$ 8		16.45 $\pm$ 0.12
0.0171	1077 $\pm$ 4		16.44 $\pm$ 0.06
0.0419	1086 $\pm$ 16		16.46 $\pm$ 0.24
0.0496	1234 $\pm$ 10		16.76 $\pm$ 0.14
0.0591	1360 $\pm$ 55		16.99 $\pm$ 0.69
0.0661	1205 $\pm$ 7		16.70 $\pm$ 0.10
0.1024	1045 $\pm$ 21		16.37 $\pm$ 0.33
0.1333	874 $\pm$ 13		15.95 $\pm$ 0.24
0.2371	521 $\pm$ 5		14.73 $\pm$ 0.14
0.2883	409 $\pm$ 4		14.16 $\pm$ 0.14

TABLE 4 (cont'd)

1.00°C		
x_{TBA} ± 0.0002	k_{H} atm	$\Delta G_{\text{sol}}^{\circ}$ kJ mol ⁻¹
0	760 $\pm$ 2	15.12 $\pm$ 0.04
0.0192	801 $\pm$ 9	15.24 $\pm$ 0.17
0.0318	966 $\pm$ 16	15.67 $\pm$ 0.26
0.0482	987 $\pm$ 8	15.72 $\pm$ 0.13
0.0536	1049 $\pm$ 80	15.85 $\pm$ 1.21
0.0587	1059 $\pm$ 10	15.88 $\pm$ 0.15
0.0591	1011 $\pm$ 24	15.77 $\pm$ 0.37
0.1576	699 $\pm$ 2	14.93 $\pm$ 0.04
0.2285	484 $\pm$ 5	14.09 $\pm$ 0.15

FIGURE 10

Henry's Law Constant, k_H , as a function
of t-Butanol Concentration at 25, 10, and 1°C

hydrogen, oxygen and argon as the solutes in water-methanol, water-ethanol and water-t-butanol mixtures. In these studies it was also observed that the gas solubility initially increased slightly with the addition of very small amounts of alcohol, up to about $X_{TBA}=0.03$ (see Figure 3). The relative amount of the increase depended on the specific gas and alcohol, and on temperature, the effect being more pronounced at lower temperatures. Generally, the initial increase was smaller in t-butanol solutions than ethanol solutions.

Cargill's results for the solubility of CO_2 in water-ethanol solutions (53) show a slight initial increase in solubility occurs with increasing X_{EtOH} at lower temperatures. At $4.7^\circ C$, the relative increase in solubility is 0.1%. This change is well within the experimental error of 1.5% for the present study. (The method used by Cargill gave results reproducible within 0.5%). Thus, if there was an initial slight increase in CO solubility in t-butanol-water mixtures the effect would probably be smaller than that observed in ethanol-water mixtures and would not be detected by the method used in the present experiment.

The free energy change of solution ΔG_{sol}° , was calculated from the Henry's Law constant and is included in Table 4. The enthalpy change, ΔH_{sol}° , and entropy change, ΔS_{sol}° of solution at $10^\circ C$ as a function of alcohol concentration are shown in Figure 11. The ΔG_{sol}° curve exhibits a slight maximum at $X_{TBA} = 0.06$. The ΔH_{sol}° and ΔS_{sol}° values are negative, and show a similar dependence on

alcohol concentration. Both the enthalpy and entropy of solution rapidly become less negative up to $X_{\text{BA}} = 0.15$. There is a slight inflection in the curves at this point, after which they change very slowly. The shapes of these curves are similar to those for the solution of CO_2 in water-ethanol (53).

FIGURE 11

Free Energy, Enthalpy and Entropy of Solution
of Carbon Dioxide in t-Butanol-Water Mixtures
at 10.00°C

3.3 pH Measurements

The pH measurements of t-butanol-water mixtures with a N_2 sparge and with a CO_2 sparge are tabulated in Table 5. These data are illustrated in Figure 12.

When the solution is sparged with N_2 gas, the pH changes very slowly with alcohol addition up to $X_{TBA} = 0.06$. At this point the pH increases rapidly from a value of 7 to a value above 9 pH units. The pH reaches a maximum around $X_{TBA} = 0.18$ and then decreases to pH 8 at $X_{TBA} = 0.3$, beyond which it changes more slowly. The pH curve indicates that the activity of hydrogen ion rapidly decreases at $X_{TBA} = 0.06$. This may be due to a change in proton concentration, and/or activity coefficient.

One difficulty in evaluating the results of pH measurements in mixed solvent media is in estimating the effect of addition of an organic cosolvent on the liquid junction potential of the electrode. The liquid junction potential, E_j , at the boundary where the reference electrolyte comes into contact with the test solution can be regarded as the effects due to the activity coefficients and ionic mobilities of species other than the proton in solution, so that (80):

$$pH = -\log a_H \pm \frac{E_j}{(RT \ln 10)/F} \quad 3.8$$

FIGURE 12

pH of t-Butanol-Water and t-Butanol-Water-CO₂
Mixtures at 25.°C

▲ predicted pH of t-butanol-water mixtures
from K values (see Table 6).

$$\log \gamma_{\pm} = - \frac{709.0 (d(s)/\epsilon(s))^{1/2} I^{1/2}}{1 + 8.876 (d(s)/\epsilon(s))^{1/2} I^{1/2}} \quad 3.9$$

where $d(s)$ and $\epsilon(s)$ are the density and dielectric constant of the solvent mixture and I is the ionic strength, which typically varied from 0.004 to 0.015m.

Using the pK_w values determined by Wooley et al., the change pH due to the change in K_w can be predicted. These calculations are summarized in Table 6. In Figure 12, the "predicted" pH curve is compared to the experimental pH curve obtained in this study. The two curves are initially similar in value and in slope. At the point where the experimental curve starts to rapidly increase at about $X_{TBA} = 0.06$, the slope of the "predicted" curve changes slowly in the opposite direction.

In aqueous media at pH less than 7, carbon dioxide species exist as $CO_2(aq)$, $H_2CO_3(aq)$ and $HCO_3^-(aq)$. The first dissociation constant of carbonic acid, K_1 , is given as:

$$K_1 = \frac{[H^+(aq)] [HCO_3^-(aq)]}{[H_2CO_3^*(aq)] - [HCO_3^-(aq)]} \quad 3.10$$

where $[H_2CO_3^*(aq)] = [CO_2(aq)] + [H_2CO_3(aq)]$. Since the ratio of $[H_2CO_3(aq)]$ to $[CO_2(aq)]$ is about 1:600 (64), K_1 can be written as:

$$K_1 = \frac{[H^+(aq)] [HCO_3^-(aq)]}{[CO_2(aq)] - [H^+(aq)]} \quad 3.11$$

In this experiment the measured pH values were not corrected for possible changes in E_j as the experimental method used did not allow estimation of this effect. However, some estimate of the change in E_j with addition of t-butanol may be made by titrating various buffer solutions with t-butanol.

Figure 13 illustrates the titration of pH 2, 7 and 10 buffers with t-butanol. The pH increases very slowly as the alcohol is added. If it is assumed that $\log(a_H)$ remains constant in the buffered solution, the change in pH can be related to the change in E_j and the change in the pK of the buffer due to the addition of t-butanol. This change is nearly linear and very small. It can be concluded that the change in pH in the unbuffered solutions, also shown in Figure 13, contains only a small, nearly linear contribution from E_j .

Also to be considered in this system is the effect on K_w of the changing dielectric constant of the system. Woolley et al. (81,82) have measured the ionization constant for water in various aqueous organic mixtures including t-butanol. The measurements were made in cells consisting of a glass electrode and a Ag/AgCl reference electrode. Small amounts of electrolytes were added to the solvent. By comparing the potentials of two such cells containing different amounts of electrolyte in the same solvent, the junction potential could be eliminated from the equations. The ionic activity coefficients were calculated from an equation based on the Debye-Huckel theory:

FIGURE 19

Titration of Aqueous (Phosphate) Buffers
with t-Butanol at 25.°C

○ titration of pure water with t-butanol

$$\log \gamma_{\pm} = - \frac{709.0 (d(s)/\epsilon(s))^{1/2} I^{1/2}}{1 + 8.876 (d(s)/\epsilon(s))^{1/2} I^{1/2}} \quad 3.9$$

where $d(s)$ and $\epsilon(s)$ are the density and dielectric constant of the solvent mixture and I is the ionic strength, which typically varied from 0.004 to 0.015m.

Using the pK_w values determined by Wooley et al., the change pH due to the change in K_w can be predicted. These calculations are summarized in Table 6. In Figure 12, the "predicted" pH curve is compared to the experimental pH curve obtained in this study. The two curves are initially similar in value and in slope. At the point where the experimental curve starts to rapidly increase at about $X_{TBA} = 0.06$, the slope of the "predicted" curve changes slowly in the opposite direction.

In aqueous media at pH less than 7, carbon dioxide species exist as $CO_2(aq)$, $H_2CO_3^*(aq)$ and $HCO_3^-(aq)$. The first dissociation constant of carbonic acid, K_1 , is given as:

$$K_1 = \frac{[H^+(aq)] [HCO_3^-(aq)]}{[H_2CO_3^*(aq)] - [HCO_3^-(aq)]} \quad 3.10$$

where $[H_2CO_3^*(aq)] = [CO_2(aq)] + [H_2CO_3(aq)]$. Since the ratio of $[H_2CO_3(aq)]$ to $[CO_2(aq)]$ is about 1:600 (64), K_1 can be written as:

$$K_1 = \frac{[H^+(aq)] [HCO_3^-(aq)]}{[CO_2(aq)] - [H^+(aq)]} \quad 3.11$$

TABLE 6

χ_{TBA}	pK_{w}	$\text{pH} = \text{pK}_{\text{w}}/2$
	± 0.06	
0	14.00	7.00
0.0094	14.12	7.06
0.0187	14.23	7.12
0.0278	14.32	7.16
0.0365	14.40	7.20
0.0455	14.46	7.23
0.0624	14.61	7.31
0.0787	14.70	7.35
0.0947	14.77	7.39
0.110	14.86	7.43
0.125	14.95	7.47
0.139	15.01	7.50
0.153	15.09	7.55
0.166	15.16	7.58
0.180	15.22	7.61
0.192	15.29	7.65

* from $\text{pK}_{\text{a}}/1$ values in reference (81)

Since $pK_1 = 6$ is much greater than pK_w , it is valid to write:

$$K_1 = \frac{[H^+(aq)]}{[CO_2(aq)] - [H^+(aq)]} \quad 3.12$$

where $[H^+(aq)] = [HCO_3^-(aq)]$. If the pH data is used; K'_1 can be defined as:

$$K'_1 = \frac{a_H^2}{[CO_2(aq)] - a_H} \quad 3.13$$

The concentration of $CO_2(aq)$ can be determined from the known solubility of CO_2 in t-butanol-water mixtures. Calculations of pK'_1 made from the results of this study are summarized in Table 7. The value of pK'_1 in pure water calculated from the experimental pH, 5.92, compares well with Harned and Davies' (64) value of 6.352 considering the simple method used in these pH measurements.

Figure 14 illustrates the dependence of pK'_1 on alcohol concentration. The increase in pK'_1 is to be expected since the dielectric medium of the solvent is decreasing. However there is a change in the slope of pK'_1 vs. X_{TBA} at about $X_{TBA} = 0.06$.

The distance of closest approach, d , of the two ions formed by the dissociation of carbonic acid; H^+ and HCO_3^- , can be estimated from the Fuoss equation:

$$K = \frac{3000}{4 \pi N d_{ab}^3} \exp \frac{z_a z_b e^2}{4 \pi \epsilon \epsilon_0 d_{ab} kT} \quad 3.14$$

where ϵ' is the relative permittivity of a vacuum, ϵ is the medium dielectric constant, z_a and z_b are ionic charges, e is the electronic charge and N is Avagadro's number. From the slope of $\ln K_1$ vs $1/\epsilon$ d_{ab} is determined from the slope to be 1.10 Å, which is a quite reasonable value. Since the initial pH slopes are the same in both the nitrogen and carbon dioxide sparged solutions, the distance of closest approach for the H^+ and OH^- ions will be approximately 1.1 Å also.

TABLE 7

pK'_1 of Carbonic Acid in t-Butanol-Water Mixtures
at 25.°C

X_{TBA} ± 0.0002	$X_{CO_2}^a$ $\times 10^4$	$[CO_2(aq)]$ $\times 10^2$	pH^b ± 0.01	a_H	pK'_1
0	6.281	3.478	3.69	2.04	5.92
0.0163	6.254	3.262	3.78	1.66	6.07
0.0300	6.246	3.110	3.85	1.41	6.19
0.0346	6.050	2.967	3.88	1.32	6.23
0.0567	6.116	2.887	4.00	1.00	6.46
0.0623	6.605	2.962	4.03	0.933	6.53
0.0840	7.728	3.237	4.14	0.724	6.79
0.1300	10.63	3.892	4.39	0.407	7.37
0.2810	22.68	5.866	5.20	0.063	9.17

a. calculated from k_H at 1 atm.

b. obtained by fitting experimental pH values to a straight line; std. dev. of fit = 0.09

FIGURE 14

pK_1 of Carbonic Acid in t-Butanol-Water Mixtures
at 25.°C

4. DISCUSSION

4.1 Component Activities in t-Butanol-Water Mixtures

Phenomena which are dependent on longer range, continuum properties of a fluid such as light-scattering and ultrasound absorption exhibit variations due to changes in the chemical potentials of the fluid components. Extrema in such properties are related to extrema in the chemical potential (or its derivatives) of a given component. From the occurrence of these extrema, some features of microscopic structure may be postulated; but no direct evidence of the microscopic structure can be obtained from macroscopic properties.

Many continuum properties of t-butanol-water mixtures have been experimentally measured and discussed in the literature. These properties are fundamentally related to the activities of the components; yet these have not yet been reported in the literature. On the basis of the occurrence of extrema in these properties, several models for the microscopic structure of these solutions have been advanced. A more rigorous proof of these models is to test them against fundamental properties such as activities rather than specific properties such as ultrasonic absorption or light scattering.

In Figure 15 the variation with composition of the activity coefficient of t-butanol in t-butanol-water

FIGURE 15

Properties of t-Butanol-Water Mixtures at 25.°C

- γ activity coefficient of TBA
- k_H solubility of carbon dioxide
- Φ_C apparent molal heat capacity of TBA
- Φ_V apparent molal volume of TBA
- R Rayleigh scattering intensity
- α^2/f ultrasonic absorption

mixtures at 25°C is compared to absorption of ultrasound (30), Rayleigh scattering intensity (32), apparent molal volume and heat capacity of t-butanol (27), and the Henry's law constant for the solubility of carbon dioxide determined in the previously described experiments. While all these properties exhibit extrema in the same region of low mole fraction alcohol, the concentrations of alcohol at which such maxima or minima occur are not coincident but range from $X_{TBA}=0.04$ to 0.10.

4.1.1 Light-Scattering in t-Butanol-Water Mixtures

A commonly cited study of the t-butanol-water system is the light-scattering experiment of Iwasaki and Fujiyama (32). From the Rayleigh scattering intensities they estimated the magnitude of the concentration fluctuations, CF, in the mixtures. The following features were observed : (i) in the range $X_{TBA}=0$ to 0.05, CF is slightly less than ideal; (ii) CF begins to increase abruptly near $X_{TBA}=0.05$; (iii) in the range $X_{TBA}=0.05$ to 0.16 CF increases linearly; and (iv) the slope of CF begins to decrease at about $X_{TBA}=0.15$.

On the basis of these trends in CF, the authors presented an associated complex model by which the local structure which exists in solution can be determined from concentration fluctuations. However, concentration fluctuations are due to non-ideal solution behaviour i.e. due to changes in the activity coefficients of the components. A more valid test of a model would be to use it

22

OF / DE

to predict the activities of the solution components. These can then be used as a basis for prediction of several properties, including light-scattering. Relating a molecular-level model directly to a property dependent on longer-range, continuum effects does not provide a thorough test of the model. As experimentally determined activity coefficients are now available for t-butanol-water solutions, the second stage of the calculation suggested above can be carried out, i.e. prediction of light-scattering from activities.

The contribution to the isotropic scattering due to concentration fluctuations, R_c , is given by (83):

$$R_c = (\pi^2 / 2 \lambda^4) kTVX_1 (\partial \epsilon / \partial X_2)^2 / (\partial \mu_2 / \partial X_2) \quad 4.1$$

where λ is the wavelength of incident light, V is the molar volume of the mixture, ϵ is the dielectric constant and μ_2 is the chemical potential of the solute. Since for an ideal solution:

$$\partial \mu_2 / \partial X_2 = RT / X_2 \quad 4.2$$

then in an ideal binary mixture, the R_c becomes:

$$R_{id} = (2 \pi^2 / \lambda^4) VX_1 X_2 (n \partial n / \partial X_2)^2 / N \quad 4.3$$

where n is the refractive index. Thus for a real solution, the ratio of ideal to real concentration scattering is:

$$\begin{aligned} R_{id} / R_c &= (X_2 / RT) (\partial \mu_2 / \partial X_2) \\ &= (X_2 / RT) (RT / X_2 + \partial \mu_2^E / \partial X_2) \quad 4.4 \end{aligned}$$

hydrophobic residue. The sharp maximum which occurs in γ_{TBA} at about $X_{TBA}=0.05$ at 25°C can be interpreted as corresponding to a maximum in the structuring effect on the surrounding water. At the concentrations of alcohol greater than this maximum, the addition of further alcohol causes a rapid destruction of the water structure up to about $X_{TBA}=0.10$. At higher concentrations the value of γ_{TBA} decreases less rapidly as it gradually approaches unity in the pure alcohol. The shift of the maximum to slightly higher alcohol concentrations as the temperature is lowered is indicative of the greater degree of water structure present at lower temperatures.

The model proposed by Mikhailov (15,16) for aqueous solutions of nonelectrolytes merits special attention. The activity coefficients of ethanol and water in mixtures predicted on the basis of this model agree remarkably well with the experimental curves obtained for t-butanol-water mixtures in the present study. The predicted activity coefficient curves for ethanol-water mixtures show the sharp maximum in γ_{ROH} in dilute solutions and the positive deviation of γ_{H_2O} in more concentrated solutions.

Information about intermolecular interactions can be obtained from the concentration dependence of thermodynamic functions. The composition dependence of the solvent activity coefficient can be expressed as a power series in terms of the solute mole fraction:

$$\ln \gamma_1 = BX_2^2 + CX_2^3 + DX_2^4 + \dots \quad 4.5$$

where $d\mu_2^E = RT d \ln \gamma_2$ with μ_2^E and γ_2 being the excess chemical potential and activity coefficient of the solute respectively.

From the values of γ_{TBA} obtained from vapour pressure measurements, the ratio R_{id}/R_c at 25°C can be calculated. R_{id} can be calculated from the refractive index data of Iwasaki and Fujiyama (32). Density data was taken from reference 27. In this way, R_c can be determined. The results of these calculations are illustrated in Figure 16. The concentration scattering R_c calculated from activity coefficients has the same features as the function $N\langle(LX)^2\rangle$ described by Iwasaki and Fujiyama. Any features of solution structure postulated on the basis of R_c would be better related to the behaviour of γ_{TBA} rather than directly to some pseudo-complex formation.

4.1.2 Hydrophobic Effects

Deviations from Raoult's law for mixtures indicate non-ideal interactions between the solute and solvent molecules. The strong positive deviation of γ_{TBA} in water-rich solutions indicates that the interaction between the water and alcohol molecules is on the average repulsive in nature despite the favourable formation of hydrogen bonds between the two components. This positive deviation arises from the strong interaction between the bulky, hydrophobic alkyl group of the alcohol and surrounding water molecules.

This phenomenon of hydrophic hydration is thought to increase the water structure in the immediate vicinity of the

hydrophobic residue. The sharp maximum which occurs in γ_{TBA} at about $X_{\text{TBA}}=0.05$ at 25°C can be interpreted as corresponding to a maximum in the structuring effect on the surrounding water. At the concentrations of alcohol greater than this maximum, the addition of further alcohol causes a rapid destruction of the water structure up to about $X_{\text{TBA}}=0.10$. At higher concentrations the value of γ_{TBA} decreases less rapidly as it gradually approaches unity in the pure alcohol. The shift of the maximum to slightly higher alcohol concentrations as the temperature is lowered is indicative of the greater degree of water structure present at lower temperatures.

The model proposed by Mikhailov (15,16) for aqueous solutions of nonelectrolytes merits special attention. The activity coefficients of ethanol and water in mixtures predicted on the basis of this model agree remarkably well with the experimental curves obtained for t-butanol-water mixtures in the present study. The predicted activity coefficient curves for ethanol-water mixtures show the sharp maximum in γ_{ROH} in dilute solutions and the positive deviation of $\gamma_{\text{H}_2\text{O}}$ in more concentrated solutions.

Information about intermolecular interactions can be obtained from the concentration dependence of thermodynamic functions. The composition dependence of the solvent activity coefficient can be expressed as a power series in terms of the solute mole fraction:

$$\ln \gamma_1 = BX_2^2 + CX_2^3 + DX_2^4 + \dots \quad 4.5$$

By the application of different solution theories the coefficients B, C, D etc. can be related to solute size, solute-solvent and solute-solute interactions. Franks (3) has discussed the relation between various theories and the signs and magnitudes of the virial coefficients.

Kozak et al. (84) have determined experimental values of B and C from the freezing point measurements of Knight (76) on various alcohol-water systems. The McMillan-Mayer theory (85) is suited to the analysis of pairwise and triplet solute-solute interactions since the virial coefficients are formally related to the radial distribution function in this theory.

The results of their analyses show that although attractive pairwise interactions do exist between solute molecules in very dilute solutions, these interactions are not strong enough to overcome the excluded volume effect of these molecules. Thus the activity coefficient of water does not deviate significantly from unity. The negative deviation from ideality was attributed to the solvation of the alcohol molecules by water, implying an increase in the structure of the solvating water molecules.

The temperature dependence of B indicates that solute-solute interactions become more positive with increasing temperature. The temperature dependence of $\ln \gamma_{H_2O}$ in i-propanol-water mixtures (23) shows a change from negative to positive deviation from ideality at low alcohol concentrations occurs at around 10°C. Thus there is a temperature at which the solution exhibits pseudo-ideality.

In more concentrated aqueous alcohol solutions, γ_{H_2O} exhibits a large positive deviations, and higher order coefficients in expression (5) become more important. This implies that the attractions between alcohol molecules in large clusters are strong. Kozak et al. suggested that the positive deviations from ideality observed are due to the formation of what were termed "alcohol micelles".

Mikhailov (15,16) has also advanced the idea that concentrated aqueous alcohol solutions, although macroscopically uniform, actually consist of a mosaic structure in which there are regions of ice-like structured water in equilibrium with regions in which this structure is broken down. In certain cases this may lead to complete phase separation, which is observed for the larger n-alcohols.

Desnoyers and colleagues (86,87) have suggested that the sharp extrema in various thermodynamic properties of t-butanol-water solutions are due to transitions which resemble micellization. These microphases cannot be called micelles however because alcohols do not form well-defined aggregates in water.

4.1.3 Proton Activity in t-Butanol-Water Mixtures

The pH measurements of t-butanol-water mixtures can also reveal some information about structural effects which occur as alcohol is added to water. In solution protons can be generated by the autodissociation of water and of the alcohol (88):

Since water is a much stronger acid than t-butanol in aqueous solution, the primary source of protons will be due to reaction 4.6. For the same reason the following reaction is unlikely:

The unfavourability of this reaction is due to solvation effects. In reaction 4.6, the OH^- ions formed will be well solvated by surrounding water molecules. In reaction 4.8, the Me_3CO^- ion is more poorly solvated by water molecules due to steric and hydrophobic effects.

Because t-butanol is also a weaker base than water (88):

it is also unlikely that the following reaction occurs in dilute solution:

It is thus reasonable to assume that in dilute t-butanol-water mixtures the primary source of protons is water self-dissociation. The protons formed will be solvated by the surrounding water molecules and will be sensitive to the structure of the solvation shell. Changes in the proton environment will be reflected by the proton activity coefficient. Changes in both the activity coefficient and

the concentration of the proton will effect the activity i.e. the pH of the mixture. In order to study the environment of the proton in various mixtures, the changes in pH due to concentration changes must be separated from changes due to the activity coefficient.

The problems of measuring pH in mixed solvent systems have been widely discussed (80,89,90), but a universal method for the measurement and reporting of pH has not yet been generally accepted. One approach is to define the "medium effect" ${}_m\gamma_i$ as:

$${}_m\gamma_i = \frac{{}_w\gamma_i}{{}_s\gamma_i} \quad 4.11$$

where ${}_w\gamma_i$, ${}_s\gamma_i$ refer to the activity coefficient of the solute in pure water and in the solvent mixture respectively. The free energy of transfer, ΔG_i° , of the solute between water and the solvent mixture is then:

$$\Delta G_i^\circ = RT \ln {}_m\gamma_i \quad (4.12)$$

The transfer free energy can be regarded as the sum of electrostatic charging effects related to the dielectric constant of the medium, $\Delta G_i^\circ(\text{el})$, and solvation effects, $\Delta G_i^\circ(\text{solv})$. The Born model predicts that the transfer of a charged species from one medium to another of lower dielectric constant results in a positive free energy change. Thus $\Delta G_i^\circ(\text{el})$ will be positive and increase as the

mixture dielectric constant is lowered so the $\gamma_{s i}$ is less than $\gamma_{w i}$.

The static dielectric constant in t-butanol-water mixtures has been measured by several researchers (75,91). If the pH of the mixtures is plotted against $1/\epsilon$, the curve is not linear but retains essentially the same features as the concentration dependence curve as shown in Figure 17. The rather dramatic increase in the pH which occurs at about $X_{TBA}=0.06$ is therefore related to solvation effects. The indication is that $\Delta G_i^*(\text{solv})$ for the proton in these mixtures is negative and becomes much greater in absolute magnitude than $\Delta G_i^*(\text{el})$ in the region beyond $X_{TBA}=0.06$. The failure of the data of Woolley et al. (81) to predict the observed pH is probably because the effects of the added electrolyte and the calculation method attenuated any solvation effects.

A possible interpretation of the observed pH is as follows. The activity of the proton decreases linearly as alcohol is added in the region $X_{TBA}=0$ to 0.06. This decrease is due mainly to dielectric effects. In fact if the experimental curve is compared to the curve derived from the data of Woolley et al., $\Delta G_i^*(\text{solv})$ may be positive in this region. This region corresponds closely to the region where water structure is increased by the presence of the alcohol as indicated by the increase in γ_{TBA} . Beyond $X_{TBA}=0.06$ the activity of the proton decreases rapidly as the water structure is destroyed by further additions of

FIGURE 17

pH versus $1/\epsilon$ in t-Butanol-Water Mixtures

alcohol up to about $X_{\text{TBA}}=0.18$. This increases the amount of unstructured water, which makes proton solvation more favourable. The activity then increases slowly as interactions of the proton with the alcohol become more important. Since the alcohol is less basic than water, these interactions are less favourable than those between the proton and water. Conductance measurements indicate that the standard free energy of solvation of the proton in alcohols is less negative than in water (92).

One approach for the calculation of $\Delta G_i^\circ(\text{solv})$ for the transfer of the proton from water to aqueous t-butanol solutions is that developed by Wells (93). He determined the equilibrium constant for the following reaction:

experimentally using spectrophotometric measurements on p-nitroaniline in t-butanol-water mixtures acidified with HCl. The values of $\Delta G_i^\circ(\text{solv})$ calculated from the experimental equilibrium constant were negative and reached a minimum value at $X_{\text{TBA}}=0.09$. Beyond this concentration $\Delta G_i^\circ(\text{solv})$ slowly increased. The maximum solvation effect observed by Wells probably occurs at lower alcohol concentration than in the present study due to the effect of the addition of HCl to the solutions.

4.2 Solubility of Carbon Dioxide in Water-t-Butanol Mixtures

4.2.1 Solubility of Carbon Dioxide in Water

Because it is a nonpolar molecule, carbon dioxide is only sparingly soluble in water. Hildebrand and Scott (6) point out that the "ideal" solubility of CO_2 is $X_{\text{CO}_2} = 1/f(\text{CO}_2) = 0.0257$; the solubility in water is about $X_{\text{CO}_2} = 7 \times 10^{-4}$ (at 1 atm partial pressure). However, carbon dioxide is more soluble in water than one would expect if its solubility is compared to that of other non-polar gases in water. This enhanced solubility is due to the formation of H_2CO_3 , HCO_3^- and CO_3^{2-} .

In Table 8 the thermodynamic functions for the solubility of various non-polar gases in water are compared. It is seen that as the solubility of gas increases, the enthalpy of solution becomes correspondingly more negative. This is also true of the entropy of solution except for the case of carbon dioxide. Thus the higher solubility of this gas is due to a more favourable enthalpy change and less unfavourable entropy change.

Cargill (53) has explained this as follows. The negative enthalpy and entropy of solution are due to the stabilization of water clusters by the neutral carbon dioxide molecules. The ions formed by the dissociation of carbonic acid will become hydrated, contributing to the negative enthalpy and entropy of solution. However, outside

the sphere of hydration, the ions cause the disruption of hydrogen bonding which is an endothermic and disordering process. The endothermicity due to hydrogen-bond breaking is probably much less than the exothermicity due to ion hydration so that the overall $\Delta H_{\text{sol}}^{\circ}$ is still quite large. However, the entropy gain at long range may be greater than the entropy loss on hydration, so that the overall $\Delta S_{\text{sol}}^{\circ}$ is reduced.

According to the Frank and Evans model for ionic hydration (7), as the size of the ion increases for a given charge, i.e. the more diffuse the charge, the long-range disordering is more important than the short-range ordering. Cargill has also pointed out (53) that if the thermodynamic functions for the solubility of various nonpolar gases in ethanol are compared, both $\Delta H_{\text{sol}}^{\circ}$ and $\Delta S_{\text{sol}}^{\circ}$ become uniformly more negative as solubility increases for all the gases.

TABLE 8

	D	α	S_0	$\Delta H_{\text{sol}}^{\circ}$	$\Delta S_{\text{sol}}^{\circ}$
	Å	10^{-24} cm^3		kJ mol^{-1}	$\text{J mol}^{-1} \text{K}^{-1}$
He	2.60	0.204	0.0323	-2.0	-106
H ₂	2.87	0.802	0.0679	-4.6	-110
O ₂	3.46	1.56	0.1056	-12.5	-127
Ar	3.40	1.63	0.1108	-12.4	-126
CO ₂	3.94	2.59	0.7745	-20.7	-131

(D is the molecular diameter, α is the polarizability and S_0 is the volume of gas dissolved per kg. of liquid at STP.)

FIGURE 19

ΔG_{sol}^E for CO_2 Solubility
in t-Butanol-Water Mixtures
at 25, 10, and 1°C

FIGURE 18

k_H for Carbon Dioxide Solubility in
t-Butanol-Water Mixtures at 25.00°C
as a function of mole fraction (Δ)
and volume fraction ($\square$) of t-Butanol

$x-x-k_H^{id}$ (see text for definition)

In Figure 18, the Henry's law constant for the solubility of carbon dioxide is shown as a function of volume fraction as well as mole fraction of t-butanol. The suggestion that the solubility may be a function of volume fraction of nonelectrolyte was made in a recent study of the solubility of electrolytes in aqueous nonelectrolyte solutions (94). The experimentally determined solubilities of several electrolytes were best fitted by an expression incorporating the volume fraction rather than the mole fraction of the organic cosolute.

The Henry's law constant for the solubility of carbon dioxide in an "ideal" t-butanol-water mixture, k_H^{id} , can be defined by:

$$\ln k_H^{id} = f_{TBA} \ln k_H^{TBA} + f_W \ln k_H^W \quad 4.14$$

where f refers to either mole or volume fraction; k_H^{TBA} and k_H^W are the Henry's law constants for CO_2 solubility in pure t-butanol and pure water respectively. The "ideal" solubility curve is also shown in Figure 18.

When k_H is plotted as a function of the volume fraction of t-butanol, the curve retains the same features as the mole fraction dependent curve. However, if each curve is compared to the "ideal" curve, there are important differences. The volume fraction curve is greater than the ideal curve over the whole concentration range, whereas the mole fraction curve is less than ideal beyond $X_{TBA} = 0.1$.

The volume fraction curve is greater than the ideal

curve because the excess molar volume of t-butanol-water mixtures is negative over the entire concentration range (27). These volume effects will be examined later.

In discussing the solubility of CO_2 in t-butanol-water solutions, it is useful to look at the excess free energy of solution, ΔG_{sol}^E . It is here defined as:

$$\Delta G_{\text{sol}}^E = \Delta G_{\text{sol}}^* - (X_W \Delta G_{\text{sol},W}^* + X_{\text{TBA}} \Delta G_{\text{sol},\text{TBA}}^*) \quad 4.15$$

where ΔG_{sol}^* is the experimental value for a given mixture composition, and $\Delta G_{\text{sol},W}^*$, $\Delta G_{\text{sol},\text{TBA}}^*$ are the free energies of solution of CO_2 in pure water and pure t-butanol respectively. Figure 19 illustrates the shape of this function at 25, 10 and 1°C.

The general shape of the ΔG_{sol}^E function is the same at each temperature in the range studied. At low mole fraction alcohol ΔG_{sol}^E reaches a maximum positive value. As the alcohol concentration increases, the sign of ΔG_{sol}^E changes and a minimum value is reached at intermediate concentrations of alcohol. The temperature effect is summarized below:

X_{TBA} for:	$\Delta G_{\text{sol}}^E = \text{max}$	$\Delta G_{\text{sol}}^E = 0$	$\Delta G_{\text{sol}}^E = \text{min}$
25°C	0.05	0.09	0.40
10°C	0.06	0.21	0.55
1°C	0.06	0.38	0.70

(At 10 and 1°C the curves have been extrapolated beyond $X_{\text{TBA}}=0.30$.) As the temperature is lowered, the

FIGURE 19

ΔG_{sol}^E for CO_2 Solubility
in t-Butanol-Water Mixtures
at 25, 10, and 1°C

concentrations of alcohol at which changes in the solubility are observed increases. Temperature also effects the absolute magnitude of ΔG_{sol}^E ; the height of the positive maximum increases as the temperature is lowered but the opposite effect is observed for the minimum value of ΔG_{sol}^E .

The shape and temperature dependence of the ΔG_{sol}^E functions of CO_2 in t-butanol-water mixtures are quite similiar to that observed for the solubility of Ar, H_2 , and He as calculated by Lucas and Cargill (62). The differences noted are that the data of Cargill shows that ΔG_{sol}^E exhibits a small negative minimum at very low mole fraction of alcohol; and the absolute magnitudes of ΔG_{sol}^E are less than in the carbon dioxide-water-t-butanol system.

4.2.3 Activity Coefficient of Carbon Dioxide in Water-t-Butanol Mixtures

It is clear from the ΔG_{sol}^E curves for the solubility of CO_2 in t-butanol-water mixtures that because the excess chemical potential of carbon dioxide is changing with the (solvent) mixture composition the activity coefficient of carbon dioxide also changes since:

$$\mu_i^E = RT \ln \gamma_i \quad 4.16$$

However only the ratio of the activity coefficients in two different mixtures can be calculated from the free energy change since

$$\Delta G_t^E = \mu_i^{E'} - \mu_i^E = RT \ln \frac{\gamma_i'}{\gamma_i} \quad 4.17$$

In order to obtain absolute values of a_{CO_2} , some reference standard state for the solute at which a_{CO_2} is unity must be chosen. The Henry's law reference state in the mole fraction scale is defined as the hypothetical state in which the solute exists if Henry's law is obeyed up to mole fraction equal to unity. In the reference state the solute vapour pressure, $P_{\text{CO}_2}^\circ$, is equal to the limiting value of the Henry's law constant for the system.

At X_2 equal to unity, P_2 may be less than or equal to P_2° , and a_2 is given by:

$$a_2 (X_2 = 1) = \frac{P_2}{P_2^\circ} \quad 4.18$$

Thus the activity of the pure solute may be less than one.

If the Henry's law reference state is chosen for the CO_2 - H_2O - t -butanol system, there are two problems. The first is that P_2° will be different for each solvent (mixture) composition. Secondly, it is clear from Figure 9 that the activity coefficients of carbon dioxide in the pressure range studied is unity for all mixtures. The activity may be less than unity at $X_2 = 1$, but it is impossible to predict what it may be from the available data. The conclusion is that Henry's law is not a good choice for a reference state for this particular system.

According to Raoult's law, the reference state for the solute is unit mole fraction so the the solute activity is defined as:

$$a_2 = \frac{P_2}{P_2^*} = \gamma_2 X_2 \quad 4.19$$

where P_2^* is the vapour pressure of the pure, supercooled, liquid solute. Thus the activity coefficient approaches unity as X_2 approaches the pure solute. In order to obtain information about the effect of changing solvent composition on γ_{CO_2} , the activity of carbon dioxide must be calculated at some arbitrary partial pressure. Here, γ_{CO_2} will be given by:

$$\gamma_{\text{CO}_2} = k_H / P_{\text{CO}_2} \quad 4.20$$

Using the value $P_{\text{CO}_2} = 63.49$ atm at $25.^\circ\text{C}$ (74) the activity coefficients of carbon dioxide in the alcohol-water mixtures were calculated from the experimental Henry's law constants at 25.00°C . These are tabulated in Table 9.

The results of these calculations show that the excess chemical potential of carbon dioxide in the mixture reaches a maximum at $X_{\text{TBA}} = 0.05$. The excess chemical potential in water is nearly eight times greater than in the alcohol. The activity coefficient of carbon dioxide in t-butanol is close to unity at 1 atm. partial pressure which indicates the affinity of the gas for the alcohol is similar to its affinity for liquid carbon dioxide. (Note that the excess chemical potential calculated is $\mu_{\text{CO}_2} - \mu_{\text{CO}_2}^*$ where $\mu_{\text{CO}_2}^*$ is the chemical potential of liquid carbon dioxide.)

TABLE 9

Activity Coefficient of CO₂ in CO₂-H₂O-TBA

Mixtures at 25.00°C

x_{TBA}	x_{CO_2} 10^4	γ_{CO_2}	$\mu_{\text{CO}_2}^E$	ΔG_t^E kJ mol ⁻¹
0	6.28	25.1	7.99	0
0.02	6.25	25.2	8.00	0.01
0.04	5.72	27.5	8.22	0.23
0.05	5.55	28.4	8.30	0.31
0.06	5.89	26.8	8.15	0.16
0.08	7.52	20.9	7.54	-0.45
0.10	8.59	18.3	7.21	-0.78
0.20	14.71	10.7	5.88	-2.11
0.40	28.82	5.5	4.23	-3.76
0.60	43.86	3.6	3.18	-4.81
0.80	65.79	2.4	2.17	-5.82
1.00	101.52	1.6	1.17	-6.82

4.2.4 Models of Solutions of Gases in Alcohol-Water Mixtures

Several experimenters have advanced models which involve changes in water structure to explain the experimental observations. Cargill (52) attributes the occurrence of maxima in the gas solubility to an increase in the hydrogen-bonded structure in the water due to the presence of both the gas and alcohol. The minima in gas solubility occurs because the increasing concentration of the alcohol progressively decreases the amount of water structure, therefore there are less interstitial sites available for the gas molecules.

Ben-Naim (49) also proposed that up to the maximum point of solubility both the gas and the alcohol stabilize water structure. In the region where the solubility goes from a maximum to minimum value, the gas molecules continue to stabilize the water structure but the alcohol has a destabilizing effect. Beyond the minimum solubility, both the gas and alcohol destroy water structure.

Glew (17) was able to explain argon solubility in water-alcohol solutions by invoking the clathrate-shell model he has developed for aqueous nonelectrolyte solutions. In very dilute solutions, each alcohol molecule is encaged by a complete water shell with the hydroxyl group hydrogen-bonded to the shell. As argon is introduced to solution, the water forms smaller shells around the gas molecule. The ternary argon-alcohol-water system is more stable than either of the binary water solutions. As the alcohol

FIGURE 20(A)

Excess Free Energy of Solution, ΔG_{exc} , of CO_2
in t-Butanol-Water Mixtures at 25.°C

- Curve 1. Experimental ΔG_{sol}^E , this study
Curve 2. $D_{CO_2} = 3.94 \text{ \AA}$; $D_{H_2O} = 2.733 \text{ \AA}$; $D_{TBA} = 5.29 \text{ \AA}$
Curve 3. $D_{CO_2} = 4.07 \text{ \AA}$; $D_{H_2O} = 2.733 \text{ \AA}$; $D_{TBA} = 5.29 \text{ \AA}$
Curve 4. $D_{CO_2} = 3.94 \text{ \AA}$; $D_{H_2O} = 2.76 \text{ \AA}$; $D_{TBA} = 5.29 \text{ \AA}$
Curve 5. $D_{CO_2} = 3.94 \text{ \AA}$; $D_{H_2O} = 2.733 \text{ \AA}$; $D_{TBA} = 5.96 \text{ \AA}$
(scale factor = 0.01)

FIGURE 20(b)

ΔG_{exc}^* (see text for definition) of Solubility of CO_2
in t-Butanol-Water Mixtures

See (a) for curve numbering. Scale factor for Curve 5 = 1.

4.2.5 A Model for Carbon Dioxide Solubility in Water-t-Butanol Mixtures

A possible explanation for the observed shapes of the ΔG_{sol}^E curves is as follows. Carbon dioxide has a much greater affinity for the alcohol than for water. Therefore if water and alcohol mixed ideally i.e. with no excess entropy effects, the solubility of carbon dioxide should increase simply because its chemical potential is lowered by the presence of the alcohol. However water and alcohol do not mix ideally; there are significant structure effects in dilute alcohol solutions. The increased water structure around the alcohol effectively forms a barrier against interaction of carbon dioxide with the alcohol. This would cause the chemical potential of the carbon dioxide to increase. This increase in chemical potential due to the unavailability of the alcohol will be compensated for by the increased lattice structure of the water which provides additional sites for carbon dioxide solution. Thus the chemical potential may remain relatively constant, as observed in this study, or decrease slightly as observed in other studies.

As the concentration of alcohol increases, the competition for interstitial cavity sites between the carbon dioxide and alcohol also increases. Because the alcohol is much more soluble in water than the non-polar gas molecule, as result of this competition the gas molecule is literally expelled from solution as it is displaced by the alcohol molecules. The solubility of the carbon dioxide decreases

because it is unable to compete any longer with the alcohol for interstitial sites and the alcohol is still effectively shielded from the gas molecule by its solvation shell. At the point of minimum solubility, the water is in a maximum structured state with the structured regions shared between the alcohol molecules and a very small number of gas molecules so that the water is literally "stretched to the limit".

As further alcohol is added, the water lattice structure is broken down and the separation of the alcohol molecules decreases i.e. a given alcohol molecule may have another alcohol as a nearest neighbour. Now the carbon dioxide can also interact directly with the alcohol, and the dramatic increase in solubility is an indication of the much greater affinity of the gas for the alcohol.

The observation that ΔG_{sol}^E becomes negative indicates that alcohol and water do not form "normal" solutions beyond the concentration range where the lattice structure begins to break down. Rather, these solutions possess a certain degree of microheterogeneity as suggested by Mikhailov (15,16) and Desnoyers (86,87).

The solutions consist of regions of structured water, and regions of alcohol aggregates mixed with monomer water species. The solubility of carbon dioxide is enhanced due to the existence of these "microphases" of water and alcohol, in which it has a greater solubility than in a regular solution of water and alcohol. The temperature

effects noted for ΔG_{sol}^E are due to the greater persistence of the structured water species as the temperature is lowered.

This model implies that the type of equilibria which were suggested by Mikhailov (16) exist in alcohol-water solutions, with the additional equilibrium:

It is important to point out that the regular solution formed by alcohol-water-carbon dioxide is not regular in the sense that the entropy of mixing is ideal. In alcohol-water mixtures $T\Delta S_M^E$ is negative over the entire concentration range.

4.2.6 Scaled Particle Theory Description of CO_2 Solubility

Each of the equilibria shown above involve not enthalpy and entropy changes but also significant volume changes. This suggests that volume effects may be another controlling factor in CO_2 solubility. The scaled particle theory provides a suitable method for investigating these volume effects because, although no assumptions are made regarding solvent structures, experimental densities and volumes are included in the calculations.

Lucas and Cargill (62) have compared the experimental free energies of solubility of He, H_2 and Ar in water-ethanol and water-t-butanol mixtures with values predicted by the scaled particle theory (SPT). The calculated excess

free energies of solution show a similar dependence on alcohol concentration as that observed experimentally. The expression for the free energy of solution, ΔG_{sol} , derived from SPT is (62):

$$\Delta G_{\text{sol}} = G_c + RT \ln(RT/V) + G_i \quad 4.22$$

The apparent molar volume of the mixture, V , is defined as:

$$V = \frac{X_w M_w + X_2 M_2}{d} \quad 4.23$$

where X_w and X_2 are the water and alcohol mole fractions; M_w and M_2 are the molecular weights of water and the alcohol respectively, and d is the experimental density of the water-alcohol mixture. G_c is the free energy of cavity formation, given by:

$$G_c/RT = \left\{ -\ln(1-y) + [3X/(1-y)]D + [3Y/(1-y)^2 + 4.5X^2/(1-y)^3]D^2 + \frac{N \Pi P}{6RT} D^3 \right\}$$

$$y, X, Y = \frac{N \Pi}{6} \frac{X_A D_A^p + X_W D_W^p}{X_W V_W^* + X_A \Phi_{V,A}} \quad 4.24$$

where p equals 3 for y , 2 for X , and 1 for Y ; D , D_w , D_A are the hard sphere diameters of the gas, water and the alcohol respectively; V_W^* is the molar volume of pure water and $\Phi_{V,A}$ is the apparent molal volume of the alcohol in the mixture.

In their calculations, Lucas and Cargill neglected the contribution from the free energy of interaction, G_i , on the

assumption that when a gas is transferred from water to another solvent, ΔG_i is very small. The calculated and experimental free energies of transfer from alcohol to water were found to be similar which lends some validity to this assumption.

The SPT and experimental values for the free energy of solution were compared by computing ΔG_{exc} , defined as:

$$\Delta G_{\text{exc}} = \Delta G_{\text{sol}}^* - (X_W \Delta G_{\text{sol},W}^* + X_A \Delta G_{\text{sol},A}^*) \quad 4.25$$

where ΔG_{exc} , $\Delta G_{\text{sol},A}^*$ and $\Delta G_{\text{sol},W}^*$ are the experimental (or calculated) values for the free energy of solution of the gas in the mixture, pure alcohol and pure water respectively. (Note that ΔG_{exc} is the same as ΔG_{sol}^E defined in 4.15). There was a qualitative agreement between the two sets of data.

This same treatment can be applied to the solution of carbon dioxide in water-t-butanol mixtures at 25°C. The computer program written to compute the SPT values of ΔG_{sol}^* and ΔG_{exc} is found in Appendix I. The densities and apparent molal volumes of t-butanol for the mixtures are taken from reference 27. In order to investigate the dependence on the hard sphere diameters, several values were taken from the literature. These are summarized in Table 10. The computer program was tested by calculating the free energy of transfer of argon from water to t-butanol at 25°C. The value of 9.21 kJ mol⁻¹ computed at 25°C compares well with the value of 9.1 kJ mol⁻¹ computed by Lucas and Cargill at 12°C.

TABLE 10

Molecular Diameters for SPT Calculations

MOLECULE	D / 10 ⁻⁸ cm	Reference, Method of Determination
H ₂ O	2.76	(95), Lennard-Jones parameter (63), density and compressibility data
CO ₂	3.94 4.06	(95), Lennard-Jones parameter (96), collision cross-section
TBA	5.29 5.70 5.96	(97), gas solubilities (98), computed vacuum potential (99), apparent molal volume

For the water-t-butanol system, the computed values of the free energy of solution are about three times larger than the experimental values. (Lucas and Cargill did not report the absolute values of their computed data.) The agreement with experimental free energy of transfer of carbon dioxide from t-butanol to water is better.

The excess free energy of solution, ΔG_{exc} , for carbon dioxide is illustrated in Figure 20(a) for the different molecular diameters. The best agreement with experimental data is found for the case $D_{CO_2}=3.94 \text{ \AA}$, $D_{H_2O}=2.733 \text{ \AA}$, and $D_{TBA}=5.29 \text{ \AA}$ (curve 2). The computed curve exhibits a small minimum at $X_{TBA}=0.015$ and a maximum at $X_{TBA}=0.05$. The minimum, which corresponds to a slight increase in carbon dioxide solubility, was not observed experimentally. This is possibly because the initial increase in solubility falls within the experimental error of the present study. The height of the computed maximum is much smaller than that observed experimentally. The computed values at other alcohol concentrations are more negative than the experimental values.

The dependence of ΔG_{exc} on the carbon dioxide diameter is not significant in dilute solutions but causes ΔG_{exc} to become slightly more negative as more alcohol is added if the diameter is increased (curve 3). The effect of increasing the water diameter is to make ΔG_{exc} slightly more negative (curve 4); however the shape of the curve is not changed. Increasing the alcohol diameter changes the magnitude and concentration dependence of ΔG_{exc} .

FIGURE 20(A)

Excess Free Energy of Solution, ΔG_{exc} , of CO_2
in t-Butanol-Water Mixtures at 25.°C

- Curve 1. Experimental ΔG_{sol}^E , this study
Curve 2. $D_{\text{CO}_2} = 3.94 \text{ \AA}$; $D_{\text{H}_2\text{O}} = 2.733 \text{ \AA}$; $D_{\text{TBA}} = 5.29 \text{ \AA}$
Curve 3. $D_{\text{CO}_2} = 4.07 \text{ \AA}$; $D_{\text{H}_2\text{O}} = 2.733 \text{ \AA}$; $D_{\text{TBA}} = 5.29 \text{ \AA}$
Curve 4. $D_{\text{CO}_2} = 3.94 \text{ \AA}$; $D_{\text{H}_2\text{O}} = 2.76 \text{ \AA}$; $D_{\text{TBA}} = 5.29 \text{ \AA}$
Curve 5. $D_{\text{CO}_2} = 3.94 \text{ \AA}$; $D_{\text{H}_2\text{O}} = 2.733 \text{ \AA}$; $D_{\text{TBA}} = 5.96 \text{ \AA}$
(scale factor = 0.01)

FIGURE 20(b)

ΔG_{exc}^* (see text for definition) of Solubility of CO_2
in t-Butanol-Water Mixtures

See (a) for curve numbering. Scale factor for Curve 5 = 1.

dramatically (curve 5).

It is of some interest to compute the solubility of carbon dioxide in "ideal" water-t-butanol mixtures; that is, assuming that the volume of mixing is zero. This "ideal" free energy of solution can be calculated from equation 4.22 by assuming that the molar volume of the mixture is given by:

$$V = X_W V_W + X_A V_A \quad 4.26$$

The difference between the computed "real" solubility and the computed "ideal" solubility, ΔG_{id} , is then:

$$\Delta G_{exc}^* = \Delta G_{sol}^* - \Delta G_{id} \quad 4.27$$

where ΔG_{sol}^* is the value of the free energy calculated previously using experimental densities and molal volumes.

The results of these calculation are shown in Figure 20(b). ΔG_{exc}^* is positive and reaches a maximum at $X_{TBA} = 0.1$ for the two more reasonable cases. The positive values of ΔG_{exc}^* arise from a negative excess molar volume of these mixtures, and indicate that the maximum volume effects for the solubility of carbon dioxide occur at $X_{TBA} = 0.1$. Since this is not the concentration at which the minimum solubility is observed, volume effects alone cannot explain the observed solubility.

4.2.7 Dissociation of Carbonic Acid in t-Butanol-Water Mixtures

The pH of solutions of t-butanol and water sparged with carbon dioxide increases linearly as alcohol is added in the region $X_{\text{TBA}} = 0$ to 0.4. However, the slope of pK'_1 increases at $X_{\text{TBA}} = 0.06$ (Figure 14). This concentration of alcohol corresponds to the dramatic decrease of the proton activity observed in t-butanol-water mixtures sparged with nitrogen. This increase in pK'_1 is probably related to the structural effects on the proton discussed previously, as the change in slope is also noted if pK'_1 is plotted as a function of $1/\epsilon$.

Atkinson et al. (100) have studied the effects of added alcohol on the carbonic acid dehydration reaction. The addition of t-butanol has a marked effect on the transition enthalpy and entropy. Both reach maximum values at about $X_{\text{TBA}} = 0.06$. This behaviour can be related to the effect of solvent structure on the transition state. At $X_{\text{TBA}} = 0.06$ the water structure is greatly enhanced. The change in the entropy of transition is indicative of the degree of water structure around the transition state.

CONCLUSIONS

The variation in the solubility of carbon dioxide in t-butanol-water mixtures as a function of alcohol concentration can be related to structural effects in these mixtures. The minimum in the gas solubility in very dilute alcohol solutions corresponds to the maximum in the activity coefficient curve of t-butanol. Extrema in the apparent molal volume, heat capacity, and other properties of t-butanol also occur in this concentration region, and indicate a maximum structuring effect on water. Thus the minimum gas solubility is observed at an alcohol concentration where water structure is maximized through the hydrophobic effect. As the alcohol concentration is increased, the gas solubility is enhanced due to the existence of microphases of alcohol molecules and regions of structured water.

The measurement of pH provides an effective method of probing solvation effects. It would be of interest to perform further pH measurements using different sparge gases, e. g. argon, and different combinations of electrodes, e. g. calomel, $\text{Pt}(\text{H}_2)$, to verify the results of this study.

There is much interest at the present time in the use of supercritical carbon dioxide for various industrial processes. Extending the present study to higher pressures and/or temperatures would provide a more complete description of a simple CO_2 -nonelectrolyte-water system.

6. REFERENCES

1. F. Franks, in "Water, A Comprehensive Treatise", F. Franks, ed. (Plenum Press, New York, 1973) vol. 2, ch. 1.
2. *ibid*, vol 2, ch. 5.
3. *ibid*, (1975) vol. 4, ch. 1.
4. C. Tanford, "The Hydrophobic Effect" (John Wiley and Sons, New York, 1973).
5. W. Kauzmann, *Advan. Protein Chem.* 14, 1 (1959).
6. J.H. Hildebrand and R.L. Scott, "The Solubility of Non-electrolytes", 3rd ed. (Dover, New York, 1964).
7. H.S. Frank and M.W. Evans, *J. Chem. Phys.* 13, 507 (1945).
8. D.W. Davidson, in "Water", F. Franks, ed. (Plenum Press, New York, 1973) vol. 2, ch. 3.
9. G. Nemethy and H.A. Scheraga, *J. Chem. Phys.* 36, 3382 (1962).
10. H.S. Frank and W.-Y. Wen, *Discuss. Faraday Soc.* 24, 133 (1957).
11. G. Nemethy and H.A. Scheraga, *J. Chem. Phys.* 36, 3401 (1962).
12. H.S. Frank and A.S. Quist, *J. Chem. Phys.* 34, 604 (1961).
13. L. Pauling, "The Nature of the Chemical Bond", 3rd ed. (Cornell Univ. Press, New York, 1960).
14. H.S. Frank and F. Franks, *J. Chem. Phys.* 48, 4746 (1968).
15. V.A. Mikhailov and L.I. Ponomarova, *J. Struct. Chem.* 9, 8 (1968).
16. V.A. Mikhailov, *J. Struct. Chem.* 9, 332 (1968).
17. D.N. Glew, H.D. Mak, and N.S. Rath, in "Hydrogen-Bonded Solvent Systems", A.K. Covington and P. Jones, eds. (Taylor and Francis, London, 1968) p195.
18. D.W. Wood, in "Water", F. Franks, ed. (Plenum Press, New York, 1979) vol. 6, ch. 6.

19. D.C. Rapaport and H.A. Scheraga, J. Phys. Chem. 86, 873 (1982).
20. W.Jorgensen, J. Chem. Phys. 77, 5757 (1982).
21. S. Okazaki, K. Nakanishi, and H. Touhara, J. Chem. Phys. 78, 454 (1983).
22. K. Nakanishi, K. Ikari, S. Okazaki, and H. Touhara, J. Chem. Phys. 80, 1656 (1984).
23. F. Franks and D.J.G. Ives, Quart. Rev. Chem. Soc. 20, 1 (1966).
24. F. Franks, in "Physico-Chemical Processes in Mixed Aqueous Solvents", F. Franks, ed. (Heinemann, London, 1967). p. 50.
25. F. Franks, in "Hydrogen-Bonded Solvent Systems", A.K. Covington and P. Jones, eds. (Taylor and Francis, London, 1968) p. 31.
26. Yu.I. Naberukhin and V.A. Rogov, Russ. Chem. Rev. 40, 207 (1971).
27. C. de Visser, G. Perron, and J.E. Desnoyers, Can. J. Chem. 55, 856 (1977).
28. K. Nakanishi, N. Kato, and M. Maruyama, J. Phys. Chem. 71, 814 (1967).
29. M.J. Blandamer, in "Water", F. Franks, ed. (Plenum Press, New York, 1973) vol. 2, ch. 9.
30. M.J. Blandamer, D.E. Clark, N.J. Hidden, and M.C.R. Symons, Chem. Comm., 342 (1966).
31. K. Tamura, M. Maekawa, and T. Yasunaga, J. Phys. Chem. 81, 2122 (1977).
32. K. Iwasaki and T. Fujiyama, J. Phys. Chem. 81, 1908 (1977).
33. K. Hallenga, J.R. Grigera and H.J.C. Berendsen, J. Phys. Chem. 84, 2381 (1980).
34. M.C.R. Symons and M.J. Blandamer, in "Hydrogen-Bonded Solvent Systems", A.K. Covington and P. Jones, eds. (Taylor and Francis, London, 1968) p. 211.
35. M.J. Blandamer and M.F. Fox, in "Water", F. Franks, ed. (Plenum Press, New York, 1973) vol. 2, ch. 8.
36. O.D. Bonner and Y.S. Choi, J. Solution Chem. 4, 457 (1975).

37. M. D. Zeidler, in "Water", F. Franks, ed. (Plenum Press, New York, 1973) vol. 2, ch. 10.
38. R.G. Anderson and M.C.R. Symons, Trans. Faraday Soc. 65, 2550 (1969).
39. J.R. Kuppers, J. Magn. Res. 4, 220 (1971).
40. W.-Y. Wen and H.G. Hertz, J. Solution Chem. 1, 17 (1972).
41. B. Kingston and M.C.R. Symons, J.C.S. Faraday II 69, 978 (1978).
42. M. Marciacq-Rousselot and M. Lucas, J. Phys. Chem. 77, 1056 (1973).
43. J. Oakes, J. C. S. Faraday II 69, 1311 (1973).
44. E. v. Goldammer and H.G. Hertz, J. Phys. Chem. 74, 3734 (1970).
45. R. Battino and H. L. Clever, Chem. Rev. 66, 395 (1966).
46. T. R. Rettich, R. Battino and E. Wilhelm, Ber. Bunsenges. Phys. Chem. 86, 1128 (1982).
47. A. Ben-Naim and S. Baer, Trans. Faraday Soc. 60, 1736 (1964).
48. A. Ben-Naim and G. Moran, Trans. Faraday Soc. 61, 821 (1965).
49. A. Ben-Naim, J. Phys. Chem. 71, 4002, (1967).
50. R.W. Cargill and T. J. Morrison, J.C.S. Faraday I 71, 618 (1975).
51. R.W. Cargill, J.C.S. Faraday I 72, 2296 (1976).
52. R.W. Cargill, J.C.S. Faraday I 74, 1444 (1978).
53. R.W. Cargill and D.E. MacPhee, J. Chem. Research (M), 2743 (1981).
54. H. Reiss, H.L. Frisch and J.L. Lebowitz, J. Chem. Phys. 31, 369 (1959).
55. R.A. Pierotti, J. Phys. Chem. 67, 1840 (1963).
56. R.A. Pierotti, J. Phys. Chem. 69, 281 (1965).
57. M. Lucas, J. Phys. Chem. 76, 4030 (1972).
58. M. Lucas, J. Phys. Chem. 77, 2479 (1973).

59. F.H. Stillinger, J. Solution Chem. 2, 141 (1973).
60. M. Lucas, J. Phys. Chem. 80, 359 (1976).
61. M. Lucas and A. Feillolay, Bull. Soc. Chim. Fr. 4, 1267 (1970).
62. M. Lucas and R.W. Cargill, J. Phys. Chem. 81, 703 (1977).
63. N. Desrosiers and J.E. Desnoyers, Can. J. Chem. 54, 3800 (1976).
64. W. Stumm and J.J. Morgan, "Aquatic Chemistry" (Wiley Interscience, New York, 1970).
65. D.M. Kern, J. Chem. Ed. 37, 14 (1960).
66. R.P. Bell, "The Proton in Chemistry", 2nd ed. (Chapman and Hall, New York, 1973) p. 38.
67. T.M. Abbott, G.W. Buchanan, P. Kruus, and K.C. Lee, Can. J. Chem. 60, 1000 (1982).
68. A. Patterson and R. Ettinger, Z. Elektrochem. 64, 98 (1960).
69. A.R. Davis and B. G. Oliver, J. Solution Chem. 1, 329 (1972).
70. B.G. Oliver and A.R. Davis, Can. J. Chem. 51, 698 (1973).
71. Y.K. Sze, W.A. Adams, and A.R. Davis, in "Chemistry and physics of aqueous gas solutions", W.A. Adams, ed. (Electrochemical Society, New Jersey, 1975).
72. J. A. Barbero, L. G. Hepler, K.G. McCurdy, and P.R. Tremaine, Can. J. Chem. 61, 2509 (1983).
73. American Institute of Physics Handbook, 3rd ed. (McGraw Hill, New York, 1972).
74. CRC Handbook of Chemistry and Physics, 58th ed. (CRC Press, Florida, 1979).
75. A.C. Brown and D.J.G. Ives, J. Chem. Soc., 1608 (1962).
76. W.S. Knight, Dissert. Abs. 24, 993 (1963).
77. International Critical Tables, volume III.
78. A.E. Markham and K.A. Kobe, Chem. Rev. 28, 519 (1941).

79. J. Livingstone, R. Morgan and H.R. Pyne, J. Phys. Chem. 34, 1578 (1930).
80. R.G. Bates, in "Solute-Solvent Interactions", Vol. 1, J.F. Coetzee and C.D. Ritchie, ed. (Marcel Dekker, New York, 1969) p. 45.
81. E.M. Woolley, D.G. Hurkot and L.G. Hepler, J. Phys. Chem. 74, 3908 (1970).
82. E.M. Woolley, J. Tomkins and L.G. Hepler, J. Solution Chem 1, 341 (1972).
83. H.H. Lewis, R.L. Schmidt and H.L. Clever, J. Phys. Chem. 74, 4377 (1970).
84. J.J. Kozak, W.S. Knight and W. Kauzmann, J. Chem. Phys. 48, 675 (1968).
85. W. McMillan and J. Mayer, J. Chem. Phys. 13, 176 (1945).
86. J.E. Desnoyers, Pure and Appl. Chem. 54, 1469 (1982).
87. G. Roux, D. Roberts, G. Perron and J.E. Desnoyers, J. Solution Chem. 9, 629 (1980).
88. J. March, "Advanced Organic Chemistry" (McGraw Hill, New York, 1977) ch. 8.
89. E.J. King, in "Physical Chemistry of Organic Solvent Systems", A.K. Covington and T.J. Dickinson, eds. (Plenum Press, New York, 1973) p.331.
90. R.G. Bates, in "Hydrogen-Bonded Solvent Systems", A.K. Covington and P. Jones, eds. (Taylor and Francis, London, 1968) p. 49.
91. T.L. Broadwater and R.L. Kay, J. Phys. Chem. 74, 3802 (1970).
92. R. de Lisi and M. Goffredi, Electrochimica Acta 16, 2181 (1971).
93. C.F. Wells, J.C.S. Faraday I 72, 601 (1976).
94. Z.B. Alfassi and J. Weiss, Ber. Bunsenges. Phys. Chem. 87, 890 (1983).
95. J.O. Hirschfelder, C.F. Curtiss and R.B. Bird, "Molecular Theory of Gases and Liquids" (Wiley, New York, 1966).
96. P.W. Atkins, "Physical Chemistry", 2nd ed. (W.H. Freeman, San Francisco, 1982) p. 873.

97. C.L. de Ligny and N.G. Van der Veen, Chem. Eng. Sci. 27, 391 (1972).
98. A.H. Clark, F. Franks, M.D. Pedley, and D.S. Reid, J.O.S. Faraday I 73, 290 (1977).
99. H.L. Friedman and C.V. Krishnan, J. Solution Chem. 2, 119 (1973).
100. G.A. Atkinson, R.C. Patel and R.J. Boe, in "Chemistry and physics of aqueous gas solutions", W.A. Adams, ed. (Electrochemical Society, New Jersey, 1975) p. 85.

APPENDIX I

Computer Program for Scaled Particle Theory Calculations

```

PROGRAM M1
  DIMENSION H(6)
  PI=3.141592653589793
  RHO=0.001
  D=0.001
  R=0.001
  RHO=0.001
  RHO=0.001

```

```

  DO 10 I=1,6

```

```

    H(I)=1

```

```

  END DO

```

```

  H(1)=1

```

```

  H(2)=1

```

```

  H(3)=1

```

```

  H(4)=1

```

```

  H(5)=1

```

```

  H(6)=1

```

```

  H(1)=1

```

```

  H(2)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

```

  H(1)=1

```

[illegible]

APPENDIX II

RESULTS OF SOME N.M.R. EXPERIMENTS

Several n.m.r. experiments were attempted on both water-t-butanol and water-t-butanol-carbon dioxide solutions in order to gain some information on a more fundamental level about solute-solvent and solute-solute interactions in these systems. Both proton and carbon-13 experiments were performed on the Varian XL-200 n.m.r. spectrometer in the Carleton Chemistry Department. Some measurements of the proton chemical shift in t-butanol-D₂O solutions were obtained, but the magnet developed a field inhomogeneity which made further proton experiments impossible.

Carbon-13 chemical shifts and T₁ times for t-butanol-D₂O solutions were also measured. The sample tubes were prepared by degassing the sample under vacuum, transferring it to the n.m.r. tube in a nitrogen atmosphere glovebag, and then sealing the glass tube with a torch. The results are shown on the following page.

Several difficulties were encountered in preparing aqueous solutions of carbon dioxide. (C-13 enriched (90%, Merk, Sharpe and Dome Isotopes) carbon dioxide was used.) Because the water-alcohol solutions expand when they melt, the carbon dioxide could not be frozen directly into a tube containing the solution without breaking the tube. After much trial and error, the following procedure was adopted.

The carbon dioxide (at 1 atm line pressure) was first frozen into an empty tube which was then clamped off and transferred, in a Dewar of liquid nitrogen, to a nitrogen atmosphere glovebag. The alcohol-water solution was then added dropwise to the tube while it was still in the liquid nitrogen. Using this method the solution froze in such a way that the tube would not break when the solution melted. The tube was then capped and allowed to thaw. It was possible to observe a peak due to carbon dioxide in the C-13 spectra of the samples prepared in this manner. Unfortunately, only a few samples could be prepared due to time limitations.

In the few spectra that were obtained of aqueous carbon dioxide solutions, the C-13 chemical shift of the t-butanol peaks were not affected by the presence of the carbon dioxide. There were changes in the frequency of the carbon dioxide peak, but too few samples were run to be able to identify any trend. One initial experiment was performed to determine the T_1 relaxation time of carbon dioxide in water. The T_1 time was on the order of 50 seconds which means overnight experiments will have to be performed to make these measurements due to the length of the experiment.

END

1	0	0	6	8	6
---	---	---	---	---	---

FIN