4

National Library
of Canada

Bibliothèque nationale
du Canada

UNIVERSITY THESES
ON MICROFICHE

THÈSES CANADIENNES
SUR MICROFICHE

0-315-01630-2

49487

NAME OF AUTHOR/NOM DE L'AUTEUR John Ernest Dawson

TITLE OF THESIS/TITRE DE LA THÈSE A Biosystematic Study of Section Rumex
in Canada and the United States

UNIVERSITY/UNIVERSITÉ Carleton

DEGREE FOR WHICH THESIS WAS PRESENTED/
GRADE POUR LEQUEL CETTE THÈSE FUT PRÉSENTÉE Ph.D.

YEAR THIS DEGREE CONFERRED/ANNÉE D'OBTENTION DE CE DEGRÉ 1979

NAME OF SUPERVISOR/NOM DU DIRECTEUR DE THÈSE Dr. J. McNeill

Permission is hereby granted to the NATIONAL LIBRARY OF
CANADA to microfilm this thesis and to lend or sell copies
of the film.

The author reserves other publication rights, and neither the
thesis nor extensive extracts from it may be printed or other-
wise reproduced without the author's written permission.

*L'autorisation est, par la présente, accordée à la BIBLIOTHÈ-
QUE NATIONALE DU CANADA de microfilmer cette thèse et
de prêter ou de vendre des exemplaires du film.*

*L'auteur se réserve les autres droits de publication; ni la
thèse ni de longs extraits de celle-ci ne doivent être imprimés
ou autrement reproduits sans l'autorisation écrite de l'auteur.*

DATED/DATE October 15, 1979 SIGNED/SIGNÉ *John E. Dawson*

PERMANENT ADDRESS/RÉSIDENCE FIXE 1276 Dorchester Street
D3
Ottawa, Ontario
K1Z 8E6

National Library of Canada
Collections Development Branch

Canadian Theses on
Microfiche Service

Bibliothèque nationale du Canada
Direction du développement des collections

Service des thèses canadiennes
sur microfiche

NOTICE

The quality of this microfiche is heavily dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to ensure the highest quality of reproduction possible.

If pages are missing, contact the university which granted the degree.

Some pages may have indistinct print especially if the original pages were typed with a poor typewriter ribbon or if the university sent us a poor photocopy.

Previously copyrighted materials (journal articles, published tests, etc.) are not filmed.

Reproduction in full or in part of this film is governed by the Canadian Copyright Act, R.S.C. 1970, c. C-30. Please read the authorization forms which accompany this thesis.

**THIS DISSERTATION
HAS BEEN MICROFILMED
EXACTLY AS RECEIVED**

AVIS

La qualité de cette microfiche dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

S'il manque des pages, veuillez communiquer avec l'université qui a conféré le grade.

La qualité d'impression de certaines pages peut laisser à désirer, surtout si les pages originales ont été dactylographiées à l'aide d'un ruban usé ou si l'université nous a fait parvenir une photocopie de mauvaise qualité.

Les documents qui font déjà l'objet d'un droit d'auteur (articles de revue, examens publiés, etc.) ne sont pas microfilmés.

La reproduction, même partielle, de ce microfilm est soumise à la Loi canadienne sur le droit d'auteur, SRC 1970, c. C-30. Veuillez prendre connaissance des formulés d'autorisation qui accompagnent cette thèse.

**LA THÈSE A ÉTÉ
MICROFILMÉE TELLE QUE
NOUS L'AVONS REÇUE**

A BIOSYSTEMATIC STUDY OF
RUMEX SECTION RUMEX IN CANADA
AND THE UNITED STATES

by

JOHN E. DAWSON
DEPARTMENT OF BIOLOGY

A thesis submitted in partial
fulfillment of the requirements
for the degree of
DOCTOR OF PHILOSOPHY
CARLETON UNIVERSITY
FALL 1979

The undersigned hereby recommend to the Faculty of
Graduate Studies acceptance of this thesis, submitted by John
Ernest Dawson, B.Sc., M.Sc., in partial fulfillment of the
requirements for the degree of Doctor of Philosophy.

.....
Chairman, Department of Biology

.....
Supervisor

.....
External Examiner

Abstract

A biosystematic study of Rumex section Rumex was carried out in order to resolve problems that have arisen since the last revision of the group in 1937. The introduction of new species and range extensions of others have resulted in the misidentification of many species. Two areas of infraspecific confusion were evident: (1) in R. maritimus and (2) in R. occidentalis. Also, little was known about the nature and extent of hybridization within the section in North America. The study utilized herbarium specimens from the major North American herbaria along with field observations and the study of cultivated material.

The usefulness and reliability of the various vegetative and fruit characteristics used to discriminate the species are discussed.

The investigation of the breeding biology of the species indicated high levels of pollen fertility and seed set which can be used as a baseline from which supposed hybrids can be recognized. Observations based on the flowering times of the species indicated that temporal barriers are of little importance in preventing gene flow between the species.

A review of the cytological literature on the species is presented. The chromosome numbers can vary from $2n = 20-200$, with most species being represented by the diploid, tetraploid and hexaploid levels. Pairing in meiosis of the species is normal.

Eleven interspecific hybrids were found to occur in North America; the pollen fertility and seed set in those hybrids examined are both extremely low. Observations made on meiotic material of one of the hybrids and information from the literature indicates abnormal pairing. A close assessment of hybrid morphology is useful in indicating the putative parents.

Numerical techniques are utilized to resolve the infraspecific problems and in examining interspecific associations.

R. maritimus is now recognized to consist of one introduced and three native varieties separable on tubercle size and shape. In R. occidentalis, two varieties are recognized separable on valve length. Canonical analysis did not suggest groups of species that could be easily delimited from each other or that corresponded directly to existing subsections.

A key to the species is given and species descriptions and distributions within North America are presented. Where possible type material of the species names was examined and it is shown that the name R. britannica a Linnaean name should be reinstated for the native species referred to as R. orbiculatus in recent works.

Acknowledgements

I would like to express my appreciation to Dr. C. Frankton who gave me the initial incentive to carry out this study of section Rumex.

The entire staff of the Biosystematics Research Institute, Saunders Building, deserves a great deal of thanks for their tolerance and assistance to a graduate student who was constantly underfoot. Technical aid was provided by Cliff Crompton in cytology and George White in photography.

I would like to thank the members of my advisory committee Drs. J. McNeill, J. Lambert, G. Carmody and K. Joy for their patience and guidance during the course of the study. My sincerest thanks go to Dr. G. Merriam, chairman of the Department of Biology, Carleton University for his co-operation in the completion of the study.

Dr. McNeill, my research supervisor deserves more than just my appreciation and thanks for his helpful suggestions and criticisms during the course of the research and during the preparation of the text. His guidance during the numerical aspects of the study was most invaluable.

Numerous friends have come to my aid at various times during the study and the preparation of the text and to all of them, I give my thanks.

Lastly, to my family and that of my girl friend Catherine
with whose assistance and constant support the project was
completed, I owe more thanks than can be expressed in words.

Table of Contents	Page
Abstract	ii
Acknowledgements	iv
List of Tables	viii
List of Figures	x
List of Maps	xii
Introduction	1
Taxonomic History	5
Morphological Characteristics of Section <u>Rumex</u>	12
Procedure	12
Leaf Characteristics	13
Fruit Characteristics	17
Valves	17
Tubercles	25
Pedicels	30
Achenes	31
Floral Structure	35
Inflorescence	36
Roots	46
Breeding Biology	48
Temporal Separation of Flowering Between Different Species	49
Procedure	49
Results	52
Pollen Fertility	52
Procedure	52
Results	60
Seed Germination and Seed Set in <u>Rumex</u> Species	66
Procedure	66
Results	69
Cytology	75
Hybrids	91
Distribution of Hybrids	93
Procedure	93
Results	93
Pollen Fertility	98
Procedure	98
Results	99
Seed Set and Seed Viability	105
Procedure	105
Results	106

	Page
Cytology of Hybrids	110
Procedure	112
Results	112
Morphological Characteristics of Hybrids	116
Procedure	116
Results	117
Numerical Taxonomy	147
Intraspecific Numerical Studies	152
Procedure	152
<u>R. maritimus</u>	160
<u>R. occidentalis</u>	169
Interspecific Relationships within Section <u>Rumex</u>	179
Procedure	182
Results	182
Key to the Species	190
Taxonomic Treatment and Description	
1. <u>R. hymenosepalus</u>	200
2. <u>R. densiflorus</u>	205
3. <u>R. pycnanthus</u>	209
4. <u>R. orthoneurus</u>	212
5. <u>R. praecox</u>	215
6. <u>R. occidentalis</u>	218
7. <u>R. arcticus</u>	229
8. <u>R. patientia</u>	235
9. <u>R. kernerii</u>	243
10. <u>R. crispus</u>	248
11. <u>R. confertus</u>	258
12. <u>R. alpinus</u>	261
13. <u>R. longifolius</u>	264
14. <u>R. pseudonatronatus</u>	264
15. <u>R. conglomeratus</u>	274
16. <u>R. sanguineus</u>	279
17. <u>R. britannica</u>	283
18. <u>R. obtusifolius</u>	294
19. <u>R. vulcher</u>	305
20. <u>R. dentatus</u>	313
21. <u>R. violascens</u>	319
22. <u>R. maritimus</u>	323
23. <u>R. palustris</u>	336
24. <u>R. stenophyllus</u>	340
25. <u>R. brownii</u>	346
Summary and Conclusions	349
Bibliography	353
Appendix I	363
Appendix II	365

List of Tables

Table	Page
1. Treatment of Genus <u>Rumex</u> sensu lato.	6
2. Characteristics Used to Separate Subgenera of <u>Rumex</u> .	8
3. General Assessment of the Development Changes observed in Valves of <u>Rumex</u> .	18
4. Comparison of Margin Teeth Characteristics.	24
5. Number of Tubercles Observed on <u>Rumex</u> Fruit.	26
6. Flowering Times.	53
7. Pollen Fertility.	62
8. Percentage Germination of Seed from Selfed and Open Pollinated Flowers.	70
9. Species Seed Set.	72
10. Chromosome Numbers.	81
11. Pollen Fertility of Hybrids.	100
12. Pollen Fertility of Nova Scotian Hybrids.	101
13. Hybrid Seed Set.	107
14. Comparison of <u>Rumex</u> Hybrids with Parent Species.	124
15. <u>R. maritimus</u> , List of Characters Used and the Methods of Measurement and Scoring.	154
16. <u>R. occidentalis</u> , List of Characters Used and the Methods of Measurement and Scoring.	157

Table	Page
17. Comparison of the Eight Main Characteristics Used to Discriminate Between Three Groups of <u>R. maritimus</u> .	168
18. Comparison of the Characteristics Used to Discriminate Between Two Groups in <u>R. occidentalis</u> .	177
19. Coding and Number of Specimens Used.	183
20. <u>Rumex</u> : Interspecific Relationships. Characteristics Used and the Methods of Measurement and Scoring.	184
21. Comparison of <u>R. arcticus</u> and <u>R. longifolius</u> .	233
22. Morphological Comparison of <u>R. crispus</u> , <u>R. obtusifolius</u> and <u>R. patientia</u> .	253
23. Morphological Comparison of <u>R. crispus</u> , <u>R. longifolius</u> and <u>R. pseudonatronatus</u> .	255
24. Comparison of <u>R. sanguineus</u> and <u>R. obtusifolius</u> .	298
25. Comparison of <u>R. obtusifolius</u> and <u>R. stenophyllus</u> .	299
26. Characteristics said to Distinguish <u>R. obtusifolius</u> subsp. <u>obtusifolius</u> from subsp. <u>transiens</u> compared with those observed on the North American Specimens of <u>R. obtusifolius</u> .	303
27. Comparison of <u>R. pulcher</u> and <u>R. obtusifolius</u> .	310
28. Comparison of <u>R. dentatus</u> with <u>R. pulcher</u> and <u>R. obtusifolius</u> .	316

List of Figures

Figure	Page
1. Indumentum Types.	15
2. Valves.	20
3. Tubercles.	28
4. Achenes.	33
5. <u>R. hymenosepalus</u> .	38
6. <u>R. occidentalis</u> .	39
7. <u>R. obtusifolius</u> .	40
8. <u>R. sanguineus</u> .	41
9. <u>R. conglomeratus</u> .	42
10, 11, 12. <u>R. maritimus</u> .	43
13. Flowering Times of <u>Rumex</u> .	54
14. Meiotic Irregularities in Hybrids.	114
15. Valves of Hybrids.	146
16. Group Average Dendrogram of <u>R. maritimus</u> .	161
17. Principal Component Analysis of <u>R. maritimus</u> .	163
18. Canonical Analyses of <u>R. maritimus</u> , Histogram.	164
19. Average-Linkage Dendrogram of <u>R. occidentalis</u> .	171
20. Principal Co-ordinate Analysis of <u>R. occidentalis</u> .	172
21. Canonical Analysis of <u>R. occidentalis</u> , Scatter Diagram.	174

Figure

Pa

22. Canonical Analysis of Rumex Species,
Scatter Diagram.

187

x1

List of Maps

Maps	Page
1. Distribution of <u>Rumex</u> hybrids.	96
2. Distribution of <u>R. crispus</u> x <u>obtusifolius</u> .	97
3. Distribution of <u>R. hymenosepalus</u> .	204
4. Distribution of <u>R. densiflorus</u> , <u>R. pycnanthus</u> , <u>R. orthoneurus</u> and <u>R. praecox</u> .	208
5. Distribution of <u>R. occidentalis</u> .	228
6. Distribution of <u>R. arcticus</u> .	234
7. Distribution of <u>R. patientia</u> .	242
8. Distribution of <u>R. alpinus</u> , <u>R. confertus</u> , <u>R. kernerii</u> and <u>R. sanguineus</u> .	247
9. Distribution of <u>R. crispus</u> .	257
10. Distribution of <u>R. longifolius</u> .	268
11. Distribution of <u>R. pseudonatronatus</u> .	273
12. Distribution of <u>R. conglomeratus</u> .	278
13. Distribution of <u>R. britannica</u> .	293
14. Distribution of <u>R. obtusifolius</u> .	304
15. Distribution of <u>R. pulcher</u> .	312
16. Distribution of <u>R. brownii</u> , <u>R. dentatus</u> , and <u>R. palustris</u> .	318
17. Distribution of <u>R. violas</u> <u>dens</u> .	322
18. Distribution of <u>R. maritimus</u> var. <u>fueginus</u> , var. <u>maritimus</u> , var. <u>pacificus</u> and var. <u>persicarioides</u> .	335
19. Distribution of <u>R. stenophyllus</u> .	345

Introduction

Rumex L., a genus of the family Polygonaceae, is estimated to include 150 (Lawrence 1959; Gleason 1966) to 250 species (Lawalree 1953; Sarkar 1958). It is cosmopolitan in distribution, but is most abundant in the north temperate regions. A sometimes indeterminable part of the range is due to introduction, as weeds or as adventives (Rechinger 1949; Good 1964). For example, R. crispus L., is considered to be one of the most successful colonizing species, as it is a weed of sixteen crops in thirty-seven countries (Holm et al. 1977).

The taxonomic treatment most commonly accepted at present (Rechinger 1964), divides the genus into four subgenera separable on a number of morphological and cytological characteristics: subgenus Rumex, subgenus Acetosella, subgenus Acetosa, subgenus Platypodium. Subgenus Rumex has been further divided into two sections; section Axillares and section Rumex (Rechinger 1937, 1949; Sarkar 1958). The present study is confined to section Rumex as represented in the United States and Canada.

Since the last formal treatment of section Rumex in North America (Rechinger 1937), areas of confusion have arisen in the status and identification of the species within the section. The purpose of this study is to resolve these problematic areas.

The introduction of new species and the occurrence of many range extensions have resulted in the misidentification of many of

2

the species, in particular, the weedy docks. An example is R. pseudohatronatus Borbas, an introduced species which was overlooked by Rechinger (1937). Although occupying an extensive range in the midwestern region of the United States, this species is not included in any major United States manual and hence was not recorded in the 1971 National List of Scientific Plant Names published by the Soil Conservation Service of the United States (McNeill 1976). In order to update both the number of species and the ranges of the taxa occurring in North America, a detailed examination was carried out of herbarium material. Specimens from the major North American herbaria and from other herbaria in regions of taxonomic concern were consulted.

The variation of many of the species across their range is poorly understood, resulting in a degree of confusion at and below the species level. Examples of this are problems in separating: R. stenophyllus Ledeb. from R. obtusifolius L.; R. crispus L. from R. obtusifolius L.; R. arcticus Trautv. from R. occidentalis S. Watson. Many of these problems are increased when dealing with vegetative, flowering or immature fruiting material. Poor understanding of the variation has led to a combined problem of an interspecific and intraspecific nature in the case of R. occidentalis s.l. This species has been separated into two species - R. occidentalis S. Watson and R. fenestratus Greene, or into a number of varieties under either species. A similar problem exists between R. maritimus L. var. fueginus (Phil.) Dusén and R. persicarioides L.; both taxa are native to North America, but the separation of R. persicarioides as a distinct species rather than as an intraspecific taxon of R. maritimus,

3

closely related to R. maritimus var. fueginus, is questioned.

Numerical studies, observations on herbarium, field and cultivated material will help to resolve these problems. In order to comment on the reliability of many of the characteristics used in determining the species of Rumex, their phenotypic plasticity was assessed by utilizing field and cultivated material from the same rootstock.

The following species were utilized in this section of the study:

- R. crispus L., R. britannica L., R. longifolius DC., R. pseudonatronatus Borbás and R. stenophyllus Ledeb.

Although far less extensive than the hybridization reported in Europe, hybrids are now known to be more common in North America than implied by Reehinger's (1937) report of only three interspecific hybrids from only fourteen locations. Hybridization adds to the confusion between the species involved. To provide an understanding of the hybrid situation as it occurs in North America, a study involving observations on flowering, pollen viability and self-compatibility utilizing herbarium specimens and material from field and cultivated collections of a hybrid (R. longifolius x R. obtusifolius) was carried out. A review of the known chromosome counts in conjunction with cytological investigations of meiotic material from cultivated hybrid specimens provides further information on the species-hybrid relationship.

This study will provide a better understanding of the species of section Rumex in North America, as well as of the extent and importance of hybridization. In addition, the up-to-date accounts

of species distribution and the better discrimination of the species
will help in the identification and eradication of the weedy species
which are often of considerable importance as contaminants of both
crop and pasture lands.

Taxonomic History

The generic name Rumex was employed by Linnaeus (1737) to replace earlier generic names; for example, for members of the present section Rumex, the generic name Lapathum (from the Greek lapazein to cleanse, from references to the plants medicinal properties (Grieve 1931)) had been applied. Species of Rumex can be distinguished from other North American members of the Polygonaceae by a combination of the following characteristics:

- (a) basal leaves forming a rosette which gradually dies back on flowering,
- (b) stem leafy and leaves alternate,
- (c) flowers not in involucrate clusters,
- (d) stamens six,
- (e) outer perianth parts unchanged in fruit and persistent at the base of the inner accrescent perianth parts which are enlarged in fruit,
- (f) achenes lenticular or usually three angled, wholly or mostly enclosed in the inner perianth parts,
- (g) embryo located along the middle of one side of the seed in cross-section (Martin and Barkley 1961; Musil 1963).

As noted earlier, the genus Rumex has been subdivided by various authors in subgenera and sections. Some of these subdivisions have been recognized as separate genera. These treatments are outlined in Table 1.

Table 1 Treatment of the Genus Rumex sensu lato
Outline of treatments of the genus Rumex by various authors.

TABLE 1

Treatment of the Genus Rumex L. sensu lato

Campestra (1819) Sections <u>Lapathum</u>	Heisner (1856) Sections <u>Lapathum</u>	Losin- Loninskaya (1936) Sections <u>Hololapathum</u> <u>Odontolapathum</u>	Reichinger (1937) Subgenera <u>Lapathum</u>	Löve (Löve and Löve 1948) Genera <u>Rumex</u>	
<u>Acetosa</u>	<u>Acetosa</u> <u>Acetosella</u>		<u>Acetosa</u> <u>Acetosella</u>	<u>Acetosa</u> <u>Acetosella</u>	<u>Lapathum</u> <u>Axillares</u> <u>Platyodi-</u> <u>um</u>
Rochinger (1954, 1968) Subgenera <u>Rumex</u>	Sections			Löve (Löve and Löve 1961, 1967) Genera <u>Rumex</u>	
<u>Acetosa</u> <u>Acetosella</u>	<u>Rumex</u> <u>Axillares</u> <u>Platypodium</u>		<u>Buccophalophora</u>	<u>Rumex</u> <u>Axillares</u>	

7

In Table 1, the early subgeneric name Lapathum and the sectional name Simplices, must be changed to Rumex to meet the requirement of Article 22 of the International Code of Botanical Nomenclature (Stafleu et al. 1978), that "the subgenus or section... including the type species of the correct name of the genus to which it is assigned bears that generic name unaltered as its epithet, but without citation of an author's name". R. patientia L. has been selected as the type species of the genus (Heller and Stearn 1958) making the subgenus and section in which it is placed Rumex rather than Lapathum or Simplices.

The characteristics used to delimit the subgenera of Rechinger (1954, 1964) or the genera of Löve (Löve and Löve 1961, 1967) are the same: the authors differ only in their choice of rank. Table 2 indicates the morphological and cytological features used to separate the genera/subgenera of Rumex s.l. Within Rumex sensu stricto, the treatments of Rechinger and Löve both recognize the existence of two sections - section Rumex and section Axillares. Rechinger (1937) dealt with both of these sections in North America, whereas Sarkar (1958) studied only the species of section Axillares. These sections are characterized as follows:

- Axillares - regular producing leafy axillary shoots that develop secondary inflorescences which often predominate over the main one
- the valve margins usually entire in fruit
- Rumex - usually producing a single unbranched stem with a branching terminal inflorescence, rarely (hybrids, damaged plants) with secondary shoots (these not distinctly leafy) predominating over the main one
- the valve margins entire or often strongly dentate in fruit

Table 2 Characteristics used to separate Subgenera
 of Rumex

The main features delimiting the subgenera are indicated.

TABLE 2

Characteristic used to separate Subgenera of
Rumex

<u>Subgenus</u>	<u>Growth Form</u>	<u>Flower Type</u>	<u>Fruit</u>	<u>Base of Leaf</u>	<u>Chromosome Number</u>
<u>Rumex</u>	Annual or perennial	All or most perfect	-Larger than the achene -with or without tubercles -with or without teeth	Cuneate, obtuse, truncate or cordate	$x = 10$ $2n = 20 - 200$ Small chromosomes (relative to the others)
<u>Acetosella</u>	Perennial	Diocious, rarely polygamous	-Seldom larger than the achene -lacking tubercles -lacking teeth	Hastate or sagittate	$x = 7$ $2n = 14, 28, 42, 56$ Medium chromosomes
<u>Acetosa</u>	Annual or perennial	Diocious or polygamous, rarely monocious	-Larger than the achene -with or without tubercles -lacking teeth	Usually hastate or sagittate	$x = 4, 5, 7, 8, 9, 10$ $2n = 8 - 40$ Medium to large chromosomes
<u>Platypodium</u>	Usually annual	Perfect	-Larger than the achene -with tubercles -with teeth	Cuneate or spatulate	$x = 6$ $2n = 16$ Large chromosomes

(Sarker 1958,
Ichikawa et al. 1971)

Sarkar (1958) in his treatment of section Axillares, reported that no hybrids between the two sections had been found. This situation has changed somewhat as Mulligan (1959) reported a hybrid between R. triangulivalvis (Danser) Rech.f. of section Axillares and R. pseudonatronatus Borbas of section Rumex from a single location near Kindersley, Saskatchewan. No further cases of hybridization between the sections have been reported. The relative lack of hybridization between the sections, as well as the morphological differences would tend to support the maintenance of their status.

Linnaeus (1753) reported only three species of section Rumex in North America: R. persicarioides L., R. britannica L., and R. sanguineus L.; as will be discussed later, R. sanguineus is not a native species. As settlement and botanical exploration continued in North America, additional introduced and native species both were reported. Prior to the treatment of Rumex in North America by Trelease (1892), accounts were predominantly regional and with the exception of Hooker's (1840) "Flora boreali - americana" dealt only with the north eastern portion of North America. Trelease included eleven species and one hybrid of section Rumex and of these, five were considered to be native. The native taxa recognized by Trelease were: R. britannica L., from northeastern North America; R. hymenosepalus Torrey from the southwestern portion of the United States, R. persicarioides (including R. persicarioides and R. maritimus var fueginus), R. occidentalis S. Watson var occidentalis from across North America and R. occidentalis var nanus (Hooker) Trelease (synonym of R. arcticus Trautv), from the northwestern portion of

Arctic America. The six introduced taxa recognized by Trelease were R. conglomeratus Murray from along the Atlantic coast and in California (more abundantly in the latter), R. crispus L., from across Canada and the United States, R. pulcher L., from along the southern portion of the Atlantic coast and on the Pacific coast, R. obtusifolius L., from across Canada and the United States, R. patientia L. (some of the specimens cited belong to R. longifolius DC) from sites across Canada and the United States, but predominantly the northeast, R. sanguineus from only a few sites on the Atlantic Coast (Trelease was the first to recognize that this species had a very restricted North American range and that the earlier widespread reports were based on misidentifications). The only hybrid reported was R. crispus x obtusifolius from six eastern states (DC, IL, MA, MI, MO, NY). When the hybrid was first described for North America by Meisner (1856), it was reported from St. Louis, Missouri. Floristic studies continued to improve the regional treatments of the species of Rumex and new species were named. Rechinger (1937) treated both section Rumex and section Axillares for Canada, the United States and Mexico, placing many of the new species in synonymy. He recognized twenty-two species and three hybrids for section Rumex; twelve of the species were recognized as being native. The native species were: R. arcticus Trautv. from Alaska and the Yukon, R. britannica L., from the northeast, R. densiflorus Osterh. from the southwest, R. fenestratus Greene (separated from R. occidentalis) from eastern and western areas, R. fueginus Phil. (synonym of R. maritimus var. fueginus (Phil.) Dusén) from across North America, R. hymenosepalus Torrey from the southwest, R. occidentalis S. Watson from across much of Canada and the United States, R. orthoneurus Rech. f., from the southwest, R. persicarioides L., from the northeast Atlantic

coast, R. praecox Rydb. from the southwest, R. pycnanthus Rech. f., from the southwest and R. violascens Rech. f., from the southwest.

In addition to providing a more up-to-date account of the ranges for the introduced species already reported as occurring in North America by Trelease, Rechinger recognized that the following also occurred on

the continent: R. alpinus L., from Nova Scotia, R. dentatus L.,

from Oregon and California, R. domesticus Hartm. (synonym of R.

longifolius DC.) from Alaska and the northeast and R. maritimus L.,

from New Jersey. In addition to adding further sites for R. crispus

x obtusifolius, Rechinger reported two further hybrids: R. britannica

x crispus from Massachusetts and R. crispus x pulcher from Virginia.

Since Rechinger's (1937) study, many changes have occurred within

Section Rumex. An additional six introduced species have been

reported, several of these now have extensive ranges in North

America; the ranges of many of the species treated by Rechinger have

expanded due to an active expansion by the species, and/or more

adequate collections of the species now being available. Furthermore,

eight more hybrids have been added to those already recognized as

occurring in North America.

This study will deal with the previously mentioned problem areas within section Rumex in North America.

Morphological Characteristics of Section Rumex

In Rumex, the morphological features that have been used to distinguish the species can for the most part, be placed in two categories, (a) basal or rosette and stem leaf characteristics, and (b) fruit characteristics. Here, the term fruit refers to "a matured ovary and its contents together with other flowering parts that may adhere to it" (Porter 1967). As noted by Lousley (1939), the fruit of Rumex consists of the achene ("nut" is the word used by Lousley), valves, sepals, receptacle and the pedicel down to the point where it breaks from the parent plant. Other features such as the floral structures, the nature of the inflorescence and the roots have also been used, but to a lesser degree.

Procedure

The observations made on the morphological characteristics used to distinguish the species have been derived from an examination of over 9000 herbarium specimens (the herbaria consulted are listed in Appendix I), supplemented by studies on field and cultivated material of R. crispus, R. obtusifolius, R. longifolius, R. britannica, R. pseudonatronatus and R. stenophyllus; the rootstocks of the first four species were obtained in 1974 and 1976 in Ontario, Quebec, New Brunswick and Nova Scotia and of the last two species in 1976 from Saskatchewan. Plants of R. conglomeratus, R. dentatus, R. pulcher and R. stenophyllus grown from seed obtained from Europe through the international seed exchange were also observed in cultivation. All plants were grown under natural lighting in

the greenhouse facilities of Agriculture Canada, Ottawa.

Observations were also made using the scanning electron microscope on both tubercle and pubescence features utilizing existing herbarium species and others collected by me. The tubercle and epidermal specimens were mounted on SEM stubs using gold paint and then gold-coated using a Hummer sputter coating device. Examination was carried out on an AMR 1000 scanning electron microscope at Agriculture Canada, Ottawa.

Leaf Characteristics

Generally most species can be recognized by characteristics of their basal or rosette leaves and their lower cauline leaves (size, shape, vein angle and indumentum); although the upper cauline leaves have not been utilized, the presence or absence of leaves subtending the branches and bracts can be diagnostic.

Observations have indicated that even in a single plant, the size of the basal leaves can be very variable. Jackowska (1964) found that under long-day photoperiod, R. obtusifolius produced smaller leaves than plants grown under short-day photoperiods; he also observed a decrease in leaf size with the age of the plant if it remained in the vegetative phase. However, the overall shape of the basal leaves of the species examined in this study was found to be relatively constant in cultivation from one season to the next. Jalas and Lindholm (1975) in their study of European plants of R. longifolius, found little change in the shape of the basal leaves over the period of their study. In R. pulcher, Lousley (1944) observed that in subsp. pulcher, the

panduriform leaves were less evident in moister habitats and most marked on dry, sandy or chalky soils. The shape of the apex and base have been found to be relatively constant for a given plant but within a species some variation can be evident. However, plants of R. britannica can produce basal leaves with cuneate or obtuse bases; in R. hydrolapathum, a European species very similar in habit and habitat to the North American R. britannica, Lousley observed a similar variation in plants producing leaves with alternate bases in the field produced leaves with broad bases when transplanted into the drier conditions of cultivation.

A measure of the angle of divergence of the veins from the midrib of the lamina (measured in the middle of the lamina) can aid in species recognition. In most species, the veins are raised but in R. arcticus, the veins are level with the lamina and scarcely evident, a feature which helps separate it from R. occidentalis, in which the main lateral veins are raised.

The petiole and abaxial veins of the basal leaves can possess some form of indumentum (Figure 1 a,b,e,f) or they can be glabrous (Figure 1 c,d). In section Rumex, the trichomes vary from being bumplike or papilliform (Figure 1 a,e) with the width equal to or greater than the length to cylindrical with the length several times greater than the width (Figure 1 b,f).

Observations have indicated that the type of trichome appears to be constant within a single plant and in general, although some variation can occur within some species, the type of trichome is characteristic of some species. Mature leaves must

Figure 1

Indumentum Types

	Magnification	Source
(a) <u>R. crispus</u>	115	All samples from cultivated material grown in Agriculture Canada facilities, Ottawa.
(b) <u>R. obtusifolius</u>	58	
(c) <u>R. conglomeratus</u>	115	
(d) <u>R. arcticus</u>	115	
(e) <u>R. crispus</u>	575	
(f) <u>R. obtusifolius</u>	575	

In R. conglomeratus (c), the veins are essentially glabrous.

In R. arcticus, the veins are glabrous and are barely evident in the abaxial surface.

The photomicrographs were taken using the SEM and illustrate the various indumentum types evident on the abaxial surface of the leaves.

a

b

c

d

e

f

be used as the pubescence characteristic of a species can be lacking in the early seedling or early basal rosette leaves.

In the species of Rumex, observed in the field or in cultivation, the basal leaves usually begin to die back once the plant has flowered. As noted by Lousley (1939) the stem bears cauline leaves which change in shape and form a continuous series from the radical leaves to those immediately below the panicle and even to the bracts within the panicle. The lowermost leaves of the stem, being very similar to the basal or radical leaves, can be used to identify the species. Numerous changes take place in the transition from basal to mid-cauline leaves: (a) a shortening of the petiole relative to blade length, with the upper leaves and bracts in some cases becoming sessile, (b) a narrowing of the blade, (c) the veins on the abaxial surface becoming less apparent the further up the stem the leaf or bract is found, (d) a decrease in the density and size of the trichomes in those species in which the basal leaves are papillose to pubescent.

In section Rumex, the branches can all be subtended by leaves while the whorls of flowers can all be subtended by bracts or some can lack these subtending structures. Two species which can be very similar in habit - R. sanguineus (Figure 9) and R. conglomeratus (Figure 10) can be distinguished even without mature fruit as in the former, only the lowermost whorls are subtended by leaf-like bracts, while in the latter, all but the

uppermost whorls are subtended by leaf-like bracts. Plants of R. conglomeratus retained this character in cultivation; in the case of R. sanguineus, this character is based on herbarium specimens and observations made on the species by Lousley (1939).

Fruit Characteristics

Of the fruit characteristics of Rumex, the following will be discussed: Valve Characteristics, Pedicel Characteristics and Achenes Characteristics.

Valves

Linnaeus (1753) referred to the inner perianth parts of Rumex as valves, but since then, numerous other terms have been applied. From a survey of the British treatments, Lousley (1939) reported that in addition to valve, the following terms have been used: fruiting sepals, sepals, enlarged petals, inner segment of the fruiting perianth, fruiting segments, enlarged petal in fruit, fruit petals, ripe inner perianth leaves, lockets. A survey of the American literature carried out in this study, indicated the following terms have been used in reference to the valves: tepals, fructiferous calyx, calyx, sepal wing, calyx-wings, fruiting calyx, inner sepals, sepals, inner perianth lobes, inner perianth parts. The term valve will be used in this study.

Rachinger (1949) described three main types of developmental changes observable during the maturation of the valves: "1. The

valves increase, they become membranous and a network of nerves develops. 2. The midrib of one or of all three valves thickens more or less and is transformed into a callosity. 3. The margin of the valve grows out into teeth of various sizes and form. These types may occur separately or variously combined." Within the species of section Rumex in North America, the combination of developmental changes observed are presented in the following table (Table 3).

Table 3 GENERAL ASSESSMENT OF THE DEVELOPMENTAL
CHANGES OBSERVED IN VALVES OF
RUMEX.

<u>SPECIES</u>	<u>Type 1</u>	<u>Type 2</u>	<u>Type 3</u>
<u>R. alpinus</u>	++	-	- to +
<u>R. arcticus</u>	++	-	- to +
<u>R. britannica</u>	++	+++	- to +
<u>R. brownii</u>	+		
<u>R. confertus</u>	++	+	- to +
<u>R. conglomeratus</u>	+	+++	-
<u>R. crispus</u>	++	+ to +++	- to +
<u>R. densiflorus</u>	++	-	- to +
<u>R. dentatus</u>	++	+++	+++
<u>R. hymenosepalus</u>	+++	-	-
<u>R. kernerii</u>	++	+++	+ to ++
<u>R. longifolius</u>	++	- to +	- to +
<u>R. maritimus</u>	+	+++	++ to +++

<u>SPECIES</u>	<u>Type 1</u>	<u>Type 2</u>	<u>Type 3</u>
<u>R. obtusifolius</u>	++	+ to +++	++ to ++
<u>R. occidentalis</u>	++ to +++	-	- to +
<u>R. orthoneurus</u>	++	-	- to +
<u>R. palustris</u>	+	+++	+++
<u>R. petientia</u>	++	+ to +++	- to +
<u>R. praecox</u>	++	-	- to +
<u>R. pseudonatronatus</u>	++	- to +	- to +
<u>R. pulcher</u>	++	+++	+ to +++
<u>R. sanguineus</u>	+	+ to +++	-
<u>R. stenophyllus</u>	++	+++	++ to +++
<u>R. violascens</u>	+	+++	+ to ++

a) + weak to +++ strongly developed or - not occurring

b) For Type 2, + = 1 tubercle to +++ three tubercles

Several features of the valve have been found in the present study to be useful in species recognition: the valve size, shape (apex, base, outline), margin, veining and tubercle characters. For most species, a combination of these is usually adequate for identifying a specimen.

The valves vary from 1.5 mm to 17.0 mm long and 0.5 to 15.7 mm wide (excluding the marginal teeth.) A degree of variation is apparent within a species and even within a single inflorescence as more terminal fruit on a branch tend to be somewhat smaller than the fruit lower down; for measurements, it is best to use a fruit from the middle of a branch, as it will be representative

Figure 2

Valves

Source (Herbarium, State, Collector)

- a) R. alpinus GH, NS, Fernald and Long 21052.
 b) R. arcticus DAO, BC, Harms 17130.
 c) R. britannica GH, ME, Corbett s.n.
 d) R. brownii NCU 227374, SC, Ahles and Haesloop 40466.
 e) R. confertus DAO, Germany, cultivated
 f) R. conglomeratus NCU 113728, NC Radford 36801.
 g) R. crispus DAO, ON, Dawson cultivated
 h) R. densiflorus NY, CO, Rydberg s.n.
 i) R. dentatus DAO, TX, Mosquin 5542.
 j) R. hymenosepalus COLO 3870, CO Weber 3848.
 k) R. kernerii CAS 415335, CA, Pollard s.n.
 l) R. longifolius DAO, NB, Dawson 74-50
 m) R. maritimus
 1) var. persicarioides GH, PE, Fernald and St. John 11034
 2) var. pacificus CAS 516004, CA, Howell 31867
 3) var. fueginus GH, MB, Scoggan 9933.
 n) R. obtusifolius ACAD, 50168, NB, Scoggan 12676.
 o) R. occidentalis
 1) var. occidentalis NY, AB, Breitung 17085.
 2) var. fenestratus CAS 132733, ID, Macbride and Payson 3812
 p) R. orthoneurus CAS 409795, AZ, Kearney and Peebles 15114
 q) R. palustris NY, NJ, Martindale s.n.
 r) R. patientia NY, NJ, Ruger s.n.
 s) R. pseudonatronatus DAO, SK, Dawson 76112.
 t) R. pulcher US 1971183, TX, Whitehouse 16008.
 u) R. pryanthus F, CO, Patterson s.n.
 v) R. sanguineus DAO, England, Bangerter and Groves 209
 w) R. stenophyllus DAO, SK, Frankton and Coupland 1547
 x) R. violascens US 243735, AZ, Dewey s.n.

One unit = 1 mm.

(R. praecox not included as only immature specimen available)
 The photographs indicate a typical valve.

a

b

c

d

e

f

g

h

i

Figure 2 Valves
(continued)

j

k

l

m1/m2

m3

n

o1

o2

Figure 2 Valves
(continued)

p

q

r

s

t

u

v

w

x

of the majority of the fruit of the inflorescence. Little variation was evident between the size of the valves in field grown plants and their cultivated counterparts, but only by varying individual environmental parameters can the extent of the phenotypic plasticity of the valve size be ascertained.

Valve shape is variable in the members of section Rumex in North America (Figure 2 a-y), narrowly oblong-ovate (e.g. R. sanguineus), ovate (e.g. R. crispus), orbicular (e.g. R. pseudonatronatus), reniform (e.g. R. longifolius), deltoid (e.g. R. dentatus). The valve apex can be hooked (e.g. R. brownii), narrowly acute (e.g. R. maritimus), acute (e.g. R. pulcher), subacute at the apex (e.g. R. crispus), narrowly obtuse (e.g. R. sanguineus) or obtuse (e.g. R. patientia). Four types of valve base are found amongst the species: cordate (e.g. R. hymenosepalus), truncate (e.g. R. pulcher), cuneate (e.g. R. maritimus) and obtuse (e.g. R. sanguineus). The degree to which the base is cordate can vary even in a single inflorescence. Within several species (e.g. R. occidentalis) cordate and truncate bases can be found, although one base type pre-dominates in an inflorescence. There is no evidence that other combinations of base types occur within the species studied. Observations on herbarium and cultivated specimens have indicated that the valve shape is a relatively constant feature for a given species.

The margin of the valves can vary from being entire (e.g. R. hymenosepalus) to distinctly spiny (e.g. R. maritimus); the teeth range from very small serrations of less than 0.1 mm in length to thin spines of up to 3.5 mm. In addition to the size

of the teeth, the number of teeth on each side of the valve can vary from 0 to 11. Within a species and even in an individual inflorescence, variation occurs in these features, but although some degree of overlap is shown between species, they can be helpful in species discrimination. Table 4 compares the characteristics of the valve margin in several species with teeth on the margin.

TABLE 4

COMPARISON OF MARGIN TEETH CHARACTERISTICS

SPECIES	NO. OF TEETH	LENGTH OF LARGEST (mm)	TYPE
<u>R. brownii</u>	3	0.6 - 1.3	spiny, recurved
<u>R. crispus</u>	0 - 6	0 - 0.3	entire to weakly serrate
<u>R. dentatus</u>	1 - 4	0.5 - 2.6	dentate - spiny
<u>R. maritimus</u>	1 - 3	0.5 - 2.0	thin, spiny
<u>R. obtusifolius</u>	1 - 5	0.6 - 2.0	dentate
<u>R. pulcher</u>	(0)5-11	0.5 - 2.0	dentate
<u>R. stenophyllus</u>	4 - 10	0.1 - 1.6	serrate to dentate

Study has shown that the type of margin is relatively constant within a species.

The veins on the valves tend to form a reticulate pattern on each side of the midrib. The veins leaving the midrib diverge at an angle ranging from less than 45° - 90° and a number of distinct anastomoses are usually evident. The number of anastomoses vary from 1 - 5 per side, but most species commonly have 2 - 3 anastomoses per side of the valves; the distance to the first

anastomosis also shows considerable variation, ranging from 0.2 - 4.0 mm from the midrib to the anastomosis. Although as a separate characteristic, the reticulate pattern is of little help in identifying most species, it can be useful when combined with other valve characteristics.

Tubercles

Four terms have been used to refer to the outgrowth of the midrib of the valves: bead, grain, callus (or callosity) and tubercle. In the North American literature, the choice is divided almost equally amongst the last three terms; in the present study, the term tubercle will be used. The tubercles are reported to aid in dispersal of the fruit by water, acting as floats (Lousley 1939); in this regard, Lousley (1944) observed that the tubercles are filled with large corky cells with big air-spaces. The typical internal structure can be clearly seen in Figure 3a. In carrying out floatability tests on Rumex species in England, Lousley (1944) reported that within R. crispus, the smaller tubercled fruit sank first; similar results were observed for other tubercled species.

Several features of the tubercle have been utilized in species identification; the number of tubercles on a fruit, and the size, shape, and surface patterning. Marek (1954) reported that in the Polish and Central European species, these tubercle features other than number are constant in any one species.

Within section Rumex, the number of tubercles on a fruit can vary from zero to three; usually the number and size (relative to the other tubercles) is relatively constant within a species. However, some variation in the number of tubercles can be observed within a species (e.g. R. crispus 1 - 3, R. longifolius 0 - 1) and often even on the fruit of a single inflorescence. Greenhouse observations on R. britannica, R. conglomeratus, R. crispus, R. dentatus, R. obtusifolius, R. pseudonatronatus, R. pulcher, R. stenophyllus, have indicated that the tubercle number and size are relatively constant character on a given plant. The following groups are based only on tubercle number and size (Table 5).

TABLE 5

NUMBER OF TUBERCLES OBSERVED ON
RUMEX FRUIT

0 Tubercles	0 - 1	1	1 - 3 (clearly unequal in size)
<u>R. alpinus</u>	<u>R. longifolius</u>	<u>R. confertus</u>	<u>R. crispus</u>
<u>R. arcticus</u>	<u>R. pseudonatronatus</u>		<u>R. obtusifolius</u>
<u>R. brownii</u>			<u>R. patientia</u>
<u>R. densiflorus</u>			<u>R. pulcher</u>
<u>R. hymenosepalus</u>			<u>R. sanguineus</u>
<u>R. occidentalis</u>			
<u>R. orthoneurus</u>			
<u>R. praecox</u>			
<u>R. pycnanthus</u>			

3 (clearly unequal in size)

R. violascens

R. kerneri

3 (approximately equal in size)

R. britannica

R. conglomeratus

R. dentatus (variable in Europe 1-3)

R. maritimus

R. palustris

R. stenophyllus

Tubercle size within the North American species of the section, varies from 0.5 - to 5.0 mm long, 0.2 - 2.5 mm wide and 0.1 - 2.5 mm deep. Because of the size variation within a species, size alone would not be sufficient to identify a species.

Tubercle shape in conjunction with these other features can be useful in determining the species, however, in some cases close similarities between some of the species require the use of additional valve characteristics. The tubercles tend to be orbicular, ovate, oblong or fusiform in shape. Observations made on the tubercled species of Rumex indicated that once the tubercle is mature, its shape is a constant feature; tubercles are not always fully developed even when the valve and the inner achene have completed their development. The variation between species in tubercle shape can be seen in Figure 3a-p.

The surface of the tubercles was examined under the SEM and most species showed some form of reticulate patterning, whereas R. dentatus and R. pulcher had a more verrucose surface. In general, the surface pattern was found to be of little diagnostic value due to the similarity between most of the species.

Figure 3

Tubercles

	Magnification	Source (Herbarium, State, Collector)
(a) <u>R. kernerii</u> (Cross-section)	12	US 2326168, CA, Pollard s.n.
(b) <u>R. britannica</u>	17	DAO, QU, Dore, Gillett and Jenkins 14992.
(c) <u>R. confertus</u>	22	DAO, ON, cultivated, Frankton s.n.
(d) <u>R. conglomeratus</u>	28	CAS 517502, CA, Hardman 1245.
(e) <u>R. crispus</u>	19	DAO, CA, Gillett and Crompton 12927.
(f) <u>R. dentatus</u>	17	DAO, ON, cultivated material
(g) <u>R. kernerii</u>	12	US 2326168, CA, Pol- lard s.n.
(h) <u>R. longifolius</u>	28	DAO, ON, Dore and Cody 47-366.
(i) <u>R. maritimus</u>	28	DAO, BC, Calder and MacKay 32648.
(j) <u>R. obtusifolius</u>	22	COLO 280809, AK, Thomas 20322.
(k) <u>R. patientia</u>	22	NDA, ND, Stevens 3310.
(l) <u>R. pseudonatronatus</u>	28	DAO, MB, Bassett and Kemp 3786.
(m) <u>R. pulcher</u>	22	DAO, LA, Smith s.n.
(n) <u>R. sanguineus</u>	28	US 780593, AL, Smith s.n.
(o) <u>R. stenophyllus</u>	28	DAO, ON, cultivated material
(p) <u>R. violascens</u>	28	US 243735, AZ, Dewey s.n.

(No illustration of R. Palustris - very similar to (i)
but larger)

The photographs illustrate a typical tubercle from a mature fruit.

a

b

c

d

e

f

g

h

i

Figure 3 Tubercles
(continued)

j

k

l

m

n

o

p

Pedicels

The pedicels of the Rumex species have been used in species discrimination (e.g. Gleason 1968, Rechinger 1964, and Scoggan 1978). Pedicel length has been observed to vary from 1.0 - 20.0 mm in length and 0.1 - 0.4 mm in width at the middle even within fruit of a single whorl, the size of a pedicel can vary and in general, it becomes smaller as the whorls are closer to the distal part of the stem or branches. With the exception of the terminal pedicels, the length of the pedicel relative to the valve length is relatively constant; although many species overlap in this character, it can be helpful in species recognition. Most species of section Rumex have filiform pedicels, broadening from the stem to the wide receptacle while in R. pulcher, the pedicel does not noticeably widen from the stem. In the case of R. pulcher, this provides an additional character separating it from R. dentatus and R. obtusifolius. Observations have indicated that the shape of the pedicel is a constant feature in a species. The main diagnostic feature of the pedicel is the type of articulation. The articulation can vary from being barely evident (e.g. in R. occidentalis) to being represented by a distinct swelling or ridge (e.g. in R. longifolius). Based on observations made both on cultivated and herbarium material, the type of articulation is a constant characteristic of a species while its position on the pedicel is more variable. The location of the articulation varies from being near the base to being at the middle of the pedicel. As the location of the

articulation is the same in most species, it provides little aid in species recognition. Greenhouse studies have indicated that on species with a swollen point of articulation, this feature usually becomes evident by the time of anthesis; however, in R. longifolius, the swollen articulation is often not clearly evident until the fruit is almost mature.

Achenes

In Rumex, the mature ovary with the adnate parts (pericarp) has been referred to as an achene, nut or nutlet. As defined by Lawrence (1959), an achene is a small dry indehiscent one-seeded fruit with a tight, thin pericarp, while a nut is an indehiscent 1-celled and 1-seeded hard and long fruit, even if resulting from a compound ovary. Since the pericarp of Rumex agrees more closely with the former definition, the term achene will be used in this study. Also, the majority of the North American floras and manuals (e.g. Musil 1963, Montgomery 1977) have preferred the term achene, the exception being Reehinger (1937), Radford et al. (1968), Munz and Keck (1959) and Hultén (1968).

The size of the trigonous achenes varies from 1.1 to 7.5 mm long, 0.6 to 5.0 mm wide and 0.4 to 3.8 mm deep. In general, the smaller the valve, the smaller the achene; this trend can be seen by comparison between R. hymenosepalus and R. sanguineus.

	Valve Length (mm)	Achene Length (mm)
<u>R. hymenosepalus</u>	9.0 - 17.0	4.0 - 7.5
<u>R. sanguineus</u>	2.0 - 3.0	1.5 - 2.0

Considerable overlap in achene size exists between the species in section Rumex in North America. Within a species, achene size can (as indicated in R. hymenosepalus in the above example), show considerable variation. As is the case with the valves, the achenes produced in more terminal fruit tend to be somewhat smaller than those produced lower down on the same branch; this has been observed in all of the species grown in this study and had been previously reported for R. crispus and R. obtusifolius (Holm et al. 1977). Maun and Cavers (1970) found a significant reduction in seed size (as measured by seed weight) for genetically identical plants of R. crispus when the soil temperature was increased from 26°C to 35°C, whereas between 10°C and 26°C, no significant difference was observed. Little difference was observed in achenes from cultivated plants as opposed to those collected in the field from the same rootstock.

The apex of an achene is always acute, whereas the base can be as acute as the apex or more broadly acute, or even distinctly obtuse to truncate. Within a species, little variation on the shape of the apex and base was apparent either from herbarium material or the field and cultivation studies. Figure 4 a-x illustrates the achene size and shape for the North American species.

Achene colour can vary from brown, red-brown or green-brown to almost black, provided that mature achenes are used, the colour appears to be a constant feature. Most species have red-

Figure 4

Achenes

Source (Herbarium, State, Collector)

a) <u>R. alpinus</u>	GH, ME, Parlin s.n.
b) <u>R. arcticus</u>	SASK 43337, BC, Harms 17218
c) <u>R. britannica</u>	DAO, NS, Gorham 451408
d) <u>R. brownii</u>	NCU 228858, SC, Ahles and Haesloop
53412
e) <u>R. confertus</u>	ALTA 30984, AB, Pegg 4132 and 4253
f) <u>R. conglomeratus</u>	UBC 765741, BC, Ashlee s.n.
g) <u>R. crispus</u>	NCU 404178, AK, Demaree 58175
h) <u>R. densiflorus</u>	US 433835, WY, Nelson 7711
i) <u>R. dentatus</u>	DAO, AB, Sexsmith 3
j) <u>R. hymenosepalus</u>	NY, TX, Cory 52195
k) <u>R. kernerii</u>	CAS, CA, Pollard s.n.
l) <u>R. longifolius</u>	GH, VT, Woodward s.n.
m) <u>R. maritimus</u>	US 2034156, MT, Cody and McCanso
3337
n) <u>R. obtusifolius</u>	NCU 168225, PA, Westerfield
12585
o) <u>R. occidentalis</u>	VLA 441, BC, Slocumb s.n.
p) <u>R. orthoneurus</u>	CAS 409795, AZ, Kearney and Peebles
15114
q) <u>R. palustris</u>	NY, NJ, Martindale s.n.
r) <u>R. patientia</u>	NY, NS, Brown s.n.
s) <u>R. pseudonatronatus</u>	DAO, SK, Alex and Gebhardt 1220
t) <u>R. pulcher</u>	CAS 313663, CA, Pollard s.n.
u) <u>R. pycnanthus</u>	F300619, UT, Jones 5957 ai
v) <u>R. sanguineus</u>	GH, OR, Suksdorf 1699
w) <u>R. stenophyllus</u>	SASK 10225, SK, Frankton and Coup-
land 1538
x) <u>R. violascens</u>	CAS 399205, CA, Twisselmann 2246

(R. praecox not included as only immature specimen available)

One unit = 1 mm.

The photographs illustrate two typical achenes.

a

b

c

d

e

f

g

h

i

j

k

l

Figure 4
(continued)

Achenes

m

n

o

p

q

r

s

t

u

v

w

x

brown coloured achenes, but in some cases (e.g. R. conglomeratus with dark brown to black achenes), it can be helpful in species recognition.

Externally, the pericarp of the achenes of Rumex are extremely smooth and provide no features useful in species recognition. However, the pericarps of R. occidentalis, R. arcticus, and R. britannica are more membranaceous and not hard as in the other species; the mature pericarp of these three species can be easily removed by rolling the achene between the fingers while in the other species, it remains firmly attached. Marek (1954) in studying the species of Rumex in Poland and Central Europe, found the size and shape of the achene to be sufficient to distinguish the species and in addition, he found the microscopic structure of the pericarp to be of value in species discrimination. Since the pericarp characteristics only succeeded in separating the species into a small number of groups, they were not employed here.

As is evident in Figure 4, most species are very similar in achene characteristics and based only on these features, identification of the North American species would be difficult. Some species, because of their distinct shape or size can be readily identified in this way (e.g. R. hymenosepalus Figure 4).

Floral Structure

In the bud or early flower stage, the flowers, because of their great uniformity (Lousley 1939, Rechinger 1949) between species, have not generally been used to provide characteristics

for species recognition. Pollen and anther characteristics have been used in a number of cases. In R. arcticus, Mulligan and Frankton (1972), found that the higher ploidy levels had pollen grains of 33.0 - 39.0 μ in diameter, while the lower ploidy levels had pollen grains of 28.5 - 31.5 μ . A preliminary study (Bassett, Crompton and Small, personal communication) of many of the Rumex species in North America, indicated that in most cases, species recognition using pollen characteristics would not be possible due to the overlap in size and morphology of their pollen. Within the North American species of Rumex, the length of the mature anthers have been found to vary from 0.3 - 2.9 mm. As the size ranges of most species overlap, anther lengths have not been used in species recognition except to distinguish R. maritimus from R. palustris.

Author	Anther Length (mm)	
	<u>R. maritimus</u>	<u>R. palustris</u>
Murbeck (1913)	0.45 - 0.62	0.9 - 1.3
Jorgenssen (1971)	0.4 - 0.6	0.9 - 1.2
Dawson (based on North American material)	0.3 - 0.5 (-0.7)	0.9 - 1.0

Inflorescence

Within the North American species of Rumex, the stem height can vary from several centimeters to approximately two metres; observations on field and cultivated material have indicated that stem height and subsequently the size of the panicle, can be variable, e.g. a plant of R. stenophyllus of less than 30 cm in

when collected in the field produced stems 1 - 2 metres high in the greenhouse. Field observations on R. maritimus var. fueginus in Alberta indicated that plants growing in the open edge of sloughs were much smaller than those growing more in the shade of the surrounding vegetation. Lousley (1939) reported the same situation for R. maritimus var. maritimus in Europe. Figures 10 - 12, although showing R. maritimus from three different sites indicate the type of variation that can be evident at a single site. Maun and Cavers (1969) found that in R. crispus, short periods (8 hr.) resulted in short, decumbent plants, whereas long photo-periods (15 hr.) resulted in tall, robust plants; this account for the shorter inflorescences observed in late summer and early autumn. Comparable results were obtained by Jackowska (1964) in R. obtusifolius. A significant reduction in panicle height was observed in R. crispus when soil temperatures were elevated from 26°C to 35°C (Maun and Cavers 1970).

The type of pubescence evident on the basal leaves is also commonly evident on the stem, either all along the stem or more usually in the inflorescence portion; the pubescence of the stem as the leaves can aid in species recognition.

The angle of divergence of the branches from the stem can also be important in species recognition. Although most species are very similar in this aspect, several species have widely divergent branches and here the angle aids in their identification, even if the basal leaves are gone and the fruit is immature. Figures 5-12 exhibit the variation present in the inflorescence and the type of

Figure 5

R. hymenosepalus - branches narrowly diverging,

- whorls contiguous,

- a compact inflorescence.

Figure 6

R. occidentalis - branches narrowly diverging,

- whorls contiguous,

- a compact inflorescence.

Figure 7

R. obtusifolius - branches moderately diverging,

- whorls becoming contiguous,

- a relatively open inflorescence.

61
Figure 8

R. sanguineus - branches widely diverging,

② - whorls remote,

- a very lax open inflorescence.

Note: whorls rarely subtended by a leaf-like bract

Figure 9

- R. conglomeratus - branches widely diverging,
- whorls remote,
- a lax open inflorescence.

Note: the majority of whorls are subtended by small leaf-like bracts. (contrast with Figure 8)

Figure 10,
(Figures 10, 11, 12)

R. maritimus - branches narrowly divergent,
- whorls generally contiguous,
- an open inflorescence.

Note: whorls subtended by leaf-like bracts.

The variation in habit observed at these three sites
can be seen occurring together in different portions
of a single site.

44

Figure 11 R. maritimus

(See page 43)

Figure 12 R. maritimus
(See page 43)

branching, ranging from narrowly to widely divergent. An example in which species can be distinguished in this way is R. sanguineus with a branching angle of 40° - 80° (Figure 9) and R. obtusifolius (Figure 7) in which it is (20°) 25° - 40° .

A further inflorescence characteristic that can aid in species recognition, is the position of the whorls on the branches and upper stem. Lower whorls are usually somewhat remote whereas the upper whorls can be remote or essentially contiguous. R. conglomeratus (Figure 8) and R. sanguineus (Figure 9) are examples of species with remote whorls, while R. occidentalis (Figure 5) and R. hymenosepalus (Figure 6), R. obtusifolius (Figure 7) and R. maritimus (Figures 10 - 12) exhibit the contiguous condition. Observations have indicated that the position of the whorls is constant for the species that have been observed under cultivation and this would be expected to be the case for the other species.

Roots

Within section Rumex, little taxonomic use has been made of the type or structure of the rootstock due to the fact that most species possess a deep taproot which can be very troublesome to dig up. However, several species have more distinctive rootstocks. In the case of R. orthoneurus, R. praecox, R. densiflorus, R. pycnanthus and R. alpinus, the rootstock are reported to be creeping (Rechinger 1937), being more horizontal in nature. This was confirmed for R. alpinus under greenhouse cultivation, but for

the other three species, further collections noting the root type are needed. Rumex maritimus has been observed to start off with the taproot typical of the species of the section, but as the plant matures, the root system becomes more fibrous in nature. The most distinctive root type in section Rumex in North America belongs to R. hymenosepalus, which has a tubercous root system; this has been confirmed on the basis of available herbarium collections.

The characteristics described have been utilized in constructing a key to the species of section Rumex in North America and are further detailed in the descriptions of the individual species. Since many of the characters show some degree of phenotypic plasticity with respect to absolute size, it is best to rely on characteristics such as leaf shape, the nature of the pubescence, the valve characteristics such as the shape, nature of the tubercles, etc., which are more constant. Detailed experiments using genetically identical plants grown under varying conditions are needed so that the range of phenotypic plasticity of particular characteristics can be observed.

Breeding Biology

In section Rumex, extensive information is available on the breeding biology of R. crispus and R. obtusifolius, largely due to the wide distribution of these species and their importance as agricultural weeds. However, little comparable information is available for other species within the section. A study of the flowering times, the pollen fertility, seed viability and seed production was carried out on those species of section Rumex in North America for which rootstocks or seed material was available. These aspects were studied to provide information on the species and to provide a guideline from which plants of non hybrid origin could be compared with plants of hybrid origin. The sections included in this study are: Temporal Separation of Flowering Between Different Species, Pollen Fertility and Seed Germination and Seed Set.

Temporal Separation of Flowering between Different Species.

The lack of coincidence of flowering periods is effective in restricting the crossings of related species of wind-pollinated or insect-pollinated species occupying the same area (Heslop-Harrison 1967). In section Rumex, the species are predominantly anemophilous and exhibit some degree of protandry (Knuth 1909). As the geographical ranges of many of the species of Rumex overlap, the differences in flowering periods of the Rumex species may represent a barrier to gene flow between the species. Information on the flowering periods of some of the species exists in various floras, e.g. Radford et al. 1963, Steyermark 1963, but these are too general to allow the existence or lack of temporal barriers between the species to be reduced with certainty.

Procedure

Herbarium specimens at the flowering stage representing most of the species of Rumex occurring in North America were examined in an attempt to determine the degree of temporal separation of the species flowering periods.

Five Nova Scotian sites at which two or more species occurred were examined during late June, 1976, to determine the flowering stage of such species.

Site	Species occurring at Site
1. South of Yarmouth	<u>R. crispus</u> <u>R. longifolius</u> <u>R. obtusifolius</u>
2. Overton, beside the United Baptist Church	same species as site 1
3. Town Point, cemetery	same species as site 1
4. Mateghan	<u>R. longifolius</u> <u>R. obtusifolius</u>
5. South of Mavillette	same species as site 1

Greenhouse studies were carried out on the available species of section Rumex to determine their flowering times when all were grown under identical conditions. The species used were:

Species	Source of Material
<u>R. britannica</u>	Ontario (No. 34 ^a , 40 ^a , 43 ^a ; 56 ^b ; Britannia Bay, Ottawa) Quebec (No. 81; Granby) ^b
<u>R. conglomeratus</u>	Botanical Gardens: Strassbourg (No. 22 ^a , 55 ^b), Versailles (No. 8 ^a , 12 ^a)
<u>R. crispus</u>	Ontario (No. 45 ^a , Prescott; 72 ^b , 83 ^b Morrisbourg, 87 ^b Kemptville) Quebec (No. 16; Granby ^a) New Brunswick (No. 39 ^a Van Buren; 42 ^a Hampton; 46 ^a Junc. Hy. 2 & 5; 38 ^a St. Leonard; 60 ^a , 61 ^a Fundy Nat. Pk; 73 ^b Garry) Nova Scotia (No. 10 ^a , 17 ^a Gaspereau)
<u>R. dentatus</u>	Botanical Garden: Versailles (No. 4 ^a , 18 ^a)
<u>R. kernerii</u>	Botanical Gardens: Versailles (No. 1 ^a , 15 ^a , 20 ^b , 52 ^a , 53 ^b , 103 ^a , 105 ^a), Udine (No. 2 ^a , 58 ^b , 101 ^a)
<u>R. longifolius</u>	Ontario (No. 48 ^c Kemptville, 55 ^a Iroquois, 89 ^a Kemptville) New Brunswick (No. 35 ^a Beechwood, 36 ^a Ste. Anne, 49 ^a Garry, 59 ^b St. Basil, 91 ^b Grand Falls) Nova Scotia (No. 25 ^b Sandy Cove, 50 ^b Rothsay)

Species	Source of Material	cont.
<u>R. maritimus</u>	British Columbia (No. 150 ^b)	
<u>R. obtusifolius</u>	Ontario (No. 51 ^a Morrisbourg) Quebec (No. 44 ^c , 68 ^a , 69 ^a , 70 ^a Church Hill) Botanical Gardens: Campo Alegre (No. 3 ^a) Paris (No. 110 ^b)	
<u>R. pseudonatronatus</u>	Alberta (No. 76112 ^b Edmonton) Botanical Gardens: Paris (No. 67 ^a), Vacratot (33 ^a , 62 ^a , 108 ^b , 113 ^b)	
<u>R. pulcher</u>	Botanical Garden: Montpellier (5 ^a , 6 ^a , 7 ^a , 99 ^b , 100 ^b)	
<u>R. stenophyllus</u>	Saskatchewan (No. 76151, 76152, 76153, 76156, 76168, 76169 Gull Lake ^b) Botanical Garden: Taschkent (No. 97 ^a , 118 ^b)	

a. used in the 1976 study

b. used in the 1977 study

c. used in both studies

Native North American material was grown from rootstocks collected during field trips carried out in the summers of 1974 and 1976. The botanical garden material was grown from seed obtained through the international seed exchange; the European material had been growing in the greenhouse facilities of Agriculture Canada since the spring of 1975.

In the fall of 1975 and 1976, the plants listed above were placed in a cold house at Agriculture Canada, Ottawa, under natural lighting with the temperature maintained at 4°C from November (1975/1976) to February (1976/1977); in March, the rootstocks were replanted and the plants were allowed to flower under natural lighting in the greenhouse facilities (daily watering and a monthly application of a nutrient solution was carried out). A previous test carried out in 1974-1975, had indicated that

control plants not provided with a cold treatment did not flower; later control tests in general agreed with this, but some plants of R. obtusifolius, R. stenophyllus, R. dentatus and R. maritimus were observed to flower without a cold treatment.

The time taken for the first buds to reach anthesis was determined as was the final date on which flowers were observed on this or subsequent inflorescences produced by the same rootstock. In both tests, R. pulcher flowered first and for comparative purposes, this time was considered as the point from which the flowering times of the remaining species would be measured. The results obtained are presented in Table 6 and Figure 13.

Results

An examination of the available herbarium material indicated that insufficient flowering material was available to be able to assess the flowering peak for a species throughout its range or to be certain that the earliest and latest flowering specimens are indicative of the actual flowering time of the species in any particular area.

Generally, the flowering periods of the species indicated little temporal separation of the species in a given region. However, one case of temporal separation did appear to be evident from the available specimens; Rumex hymenosepalus over its range in the southwestern United States showed little or no overlap with any of the other species occurring in the same regions, e.g. in Oklahoma R. hymenosepalus flowered in April to early May, while the earliest flowering time for the other species (R. crispus,

Table 6 Flowering Times of Rumex species Relative to
R. pulcher

The flowering times are based on observations made on
cultivated plants.

TABLE 6

FLOWERING TIMES OF Rumex SPECIES RELATIVE TO R. pulcher

SPECIES	NO. OF SPECIMENS	INITIAL AVERAGE	TIME OF FLOWERING (DAYS) RANGE	DURATION OF FLOWERING (DAYS)	
				AVERAGE	RANGE
<u>R. pulcher</u>	3	0.7	-2 to 4		(not recorded in 1976)
<u>R. kernerii</u>	3	1.5	0 - 6		
<u>R. longifolius</u>	8	4.0			
<u>R. conglomeratus</u>	3	4.0			
<u>R. dentatus</u>	2	7.5	4 - 14		
<u>R. obtusifolius</u>	6	11.6	7 - 14		
<u>R. crispus</u>	10	21.0			
<u>R. stenophyllus</u>	1	25.7	21 - 35		
<u>R. pseudonatronatus</u>	3	46.0			
<u>R. britannica</u>	3				
1977 ^a					
<u>R. pulcher</u>	2	1.0	-1 to 9	47	11 - 31
<u>R. maritimus</u>	1	3.0	5 - 15	102	24 - 62
<u>R. longifolius</u>	4	7.7	16 - 46	25	20 - 91
<u>R. kernerii</u>	7	33.8	35 - 46	43.1	59 - 68
<u>R. crispus</u>	4	39.3	13 - 86	54.5	41 - 100
<u>R. pseudonatronatus</u>	3	40.6		62.3	
<u>R. stenophyllus</u>	7	43.0		73.6	
<u>R. conglomeratus</u>	1	46.0		76	
<u>R. obtusifolius</u>	2	83.0	45 - 47	51	23 - 79
<u>R. britannica</u>	2		70 - 96	14	

¹R. pulcher flowered on June 1, 1976²R. pulcher flowered on June 4, 1977

Figure 13 Flowering Times of Rumex

A graphic illustration of the flowering times based on observations of cultivated plants.

Figure 13

Flowering Times of Rumex

Second point is 14 days from initially flowering, representing average flowering time for an inflorescence.

Third point is based on average duration of flowering of examined specimens; this includes flowering of further inflorescences produced by the same rootstock.

R. obtusifolius, R. patientia, R. pulcher, R. stenophyllus) was mid-May. More abundant collections of Rumex species in flower across their North American ranges, especially from sites at which more than one species occur together, are needed to clarify the existence or lack of temporal barriers between the species.

At the Nova Scotian sites examined in late June, 1976, the following general observations were made: a) at each site, R. longifolius was in full flower and on many plants, the lower whorls were in immature fruit; b) R. crispus was mainly in full bud with the more mature flowers at anthesis; c) R. obtusifolius was just coming into full bud and many whorls were still covered by an ochrea. Some degree of temporal separation of R. longifolius from the other species would exist as its peak of flowering would be completed before that of the other species. Although R. crispus was slightly more advanced than R. obtusifolius, the flowering peaks of these species would overlap greatly, providing little or no temporal separation. At these same sites, approximately two months later, a few plants of each species were in flower; these plants represented individuals that have flowered well after the majority of the plants of the respective species. Of the three species, R. longifolius had the fewest plants still flowering. These later flowering plants would tend to reduce the temporal separation of R. longifolius from the other species.

In the cultivated specimens used in this study, an inflorescence usually completed flowering in an average of two weeks; however, subsequent production of additional inflorescences can

extend the total time over which flowering takes place (Table 6). From Figure 13, it can be seen that of the plants examined in 1976, those of R. britannica were the only ones clearly separated temporally from the others. The flowering time of the plants of R. stenophyllus differs from that of the others except for those of R. crispus and R. pseudonatronatus; the plants of R. pseudonatronatus were separated temporally from all plants except those of R. stenophyllus. During the 1976 study, observations were based only on the initial inflorescences produced by a plant while in the 1977 study, the flowering on subsequent inflorescences were observed.

In the 1977 study, the average initial flowering times and the average flowering time of an inflorescence (two weeks) suggested that three groupings of plants temporally separated from each other existed (Figure 13); Group 1 consisted of the plants of R. pulcher, R. maritimus, R. longifolius and R. kernerii (within the group little separation exists between the species) Group 2 consisted of the plants of R. crispus, R. pseudonatronatus, R. stenophyllus, R. conglomeratus and R. obtusifolius (the plants of R. pseudonatronatus and R. stenophyllus do not show the separation observed in 1976; within Group 2, R. crispus plants show some temporal separation from the others and the most from the plants of R. obtusifolius); Group 3 consisted only of the plants of R. britannica, once again they were clearly separated from the others. However, in taking into consideration the total time during which flowers were observed on the specimens in the greenhouse (Table 6), it can be seen (Figure 13) that actually

considerable overlap in the total flowering times of the plants examined exists. Because of the subsequent flowerings of most of the plants, the temporal separation of the plants of R. britannica was observed to be far less complete. The plants of R. longifolius still exhibit temporal separation from the species of Groups 2 and 3.

In their study carried out in England, Cavers and Harper (1964) reported that frequently, plants of R. crispus and R. obtusifolius can produce many inflorescences from the same rootstock and ripe seed twice in one growing season, the first flowers appearing in late May or June and the second set in August and September. During the course of field studies in the Canadian prairies, Ontario, Quebec and the Maritimes, few plants of R. britannica, R. crispus, R. longifolius, R. maritimus, R. obtusifolius, R. occidentalis, R. pseudonatronatus and R. stenophyllus appeared to be undergoing a second flowering. With the exception of R. britannica and to a lesser extent R. longifolius, the species often produce many inflorescences from the same rootstock. From the greenhouse studies, it is apparent that for the species examined (with the exception of R. britannica and R. longifolius), subsequent flowerings from the same rootstock are possible. In the field, the absence of a second flowering period is probably due to a deterioration of the environmental conditions needed for flowering, e.g. most sites examined became much drier as the season progressed, limiting new growth.

The present study has indicated that only R. hymenosepalus and R. britannica exhibit a strong degree of temporal separation

from other species; in the case of R. britannica, the separation is somewhat weaker. From the field and greenhouse studies, it is evident that some temporal separation may exist between species. Only an examination of the species at the sites, where several exist will clarify this, but changes in environmental conditions from one season to the next could possibly affect the species differently, altering the times of flowering. In general, with the exception of the previously noted species, temporal separation between the North American species of section Rumex appears to be of little importance in preventing gene flow between the species.

Pollen Fertility of Rumex Species

In Section Rumex, few accounts of measurements of the pollen fertility within the species exist; Cavers and Harper (1964) reported that the individuals of R. obtusifolius that they studied, were highly fertile, with less than 5% of the grains conspicuously small and abortive. The pollen fertility of the species of section Axillares in North America is reported to be very high, ranging from 87 to 99 percent (Sarkar 1958). Within Rumex, a high level of pollen fertility has been generally considered to be indicative of plants that were not of hybrid origin, in contrast to low levels of pollen fertility in interspecific hybrids (Lousley 1939, Cavers and Harper 1964, Sarkar 1958).

Procedure

Pollen fertility was assessed using the lactophenol cotton blue method (Watkins 1925, modified by using 0.5 percent cotton blue). In this technique, empty sterile grains do not take up the stain while fertile grains containing an abundance of chromatin material are readily stained (Lawrence 1959).

In assessing pollen fertility within a species, specimens whose identification had previously been confirmed (excluding any of possible hybrid origin) were selected from across the North American geographic range of the species. To calculate the pollen fertility of a specimen, the mean of two counts of 100 grains was taken, each count being made on a preparation from a single specimen.

Anthers were taken from buds that had fully opened, but in which the anthers had not yet dehisced. Pollen fertility was assessed for 24 of the species of Rumex; in several cases, European and/or cultivated material was also employed. The values obtained are arranged in Table 7.

Result

In general, plants of the species of section Rumex based on the present sampling, have a high pollen fertility (Table 7). In contrast to section Axillares as noted above, a wide fertility range was observed (58 to 100 percent). Based on a total of 165 samples, the distribution of the percentage fertility measurements was as follows:

Percentage Fertility	51-60	61-70	71-80	81-90	91-100
Percentage of Specimens	1.8	2.4	3.6	16.4	75.8

In re-examining the 7 counts (4.2 percent) from specimens having a percentage fertility below 71 percent, it was apparent that they were from specimens in which the inflorescence was predominantly past flowering with only a few whorls still in flower or are very old collections. However, high counts were obtained from other specimens in similar conditions. Possibly, these low counts are due to environmentally induced failure in pollen development, but the actual cause is uncertain. The morphological characteristics were also re-examined, but none of the specimens revealed any indication of morphological intermediary that would suggest hybridization.

Provided that interspecific hybrids in section Rumex produce pollen of a much lower percentage fertility, a high pollen fertility can be considered as indicative of a plant of non hybrid origin. The results obtained here will be used as a basis with which to compare plants of suspected hybrid origin and will be referred to in the study of Rumex hybrids.

TABLE 7

POLLEN FERTILITY
SOURCE OF MATERIAL AND PERCENTAGE STAINABLE POLLEN

SPECIES

<u>R. alpinus</u>	NS 98	Scotland 96	Austria 99	$\bar{x} = 96$
<u>R. arcticus</u>	AK 79, 81 92, 100	T 98, 99	MT 80, 100	$\bar{x} = 91$
<u>R. britannica</u>	AB 76 JU 59 ME 33	MN 58 NK 82, 85 NJ 61	OH 94, 98 NS 86	$\bar{x} = 78$
<u>R. brownii</u>	SC 83, 95			$\bar{x} = 89$
<u>R. confertus</u>	Cultivated 82			
<u>R. conglomeratus</u>	VA 97	NC 95	CA 98, 100	$\bar{x} = 98$
<u>R. crispus</u>	BC 80, 82, 100 ME 95 Cultivated 100	B 100 NS 92, 93	OH 77, 92, 100 NF 90	$\bar{x} = 92$

2

SPECIES, cont.

<u>R. densiflorus</u>	WY 90	CO 95		$\bar{x} = 93$
<u>R. dentatus</u>	CA 98	AZ 98	AB 99	
	Cultivated 98			$\bar{x} = 98$
<u>R. hymenosepalus</u>	CA 86	AZ 87		$\bar{x} = 87$
<u>R. kernerii</u>	CA 99			
<u>R. longifolius</u>	NS 94 ³ , 96 ² , 97, 98, 99 ⁵ , 100 ³	ON 60, 53	ME 97	$\bar{x} = 95$
<u>R. maritimus</u>	YT 82, 99	BC 86, 100 ²	AB 100	
	SK 97	MB 99	ON 98	
	QU 100 ³	NS 93	PE 100	
	CA 100	WY 99	ND 100	
	MB 92	MN 96	MA 100	$\bar{x} = 92$
<u>R. obtusifolius</u>	WI 100	IL 100	ON 30, 95, 98	
	NS 96, 97, 99 ³			$\bar{x} = 96$

SPECIES, Cont.

<u>R. occidentalis</u>	BC 74, 86	AB 97, 98 ² , 99 ³ , 100 ³	DE 92 ²	
	WU 97	AK 73, 85, 94, 95, 96, 99, 100	WA 90	
	OR 62, 80, 99, 100 ²	CA 95, 100	ID 72, 99	
	WY 94	AL 86		$\bar{x} = 93$
<u>R. orthoneurus</u>	AZ 87, 92, 100	NEI 99		$\bar{x} = 95$
<u>R. palustris</u>	CA 94	NJ 100		$\bar{x} = 97$
<u>R. patientia</u>	WU 98	NC 100		$\bar{x} = 99$
<u>R. praecox</u>	CO 59	WY 100		$\bar{x} = 80$
<u>R. pseudonatronatus</u>	AB 74	SK 86, 96	ND 90, 100	$\bar{x} = 89$
<u>R. pulcher</u>	CA 97	OK 99	SC 97	$\bar{x} = 98$
<u>R. sanguineus</u>	HIJ 89	England 97		$\bar{x} = 93$

SPECIES, cont.

R. stenophyllusSK
61, 90, 95, 97MB
36, 90²MB
100SD
100Cultivated
96, 99, 100² $\bar{x} = 94$ R. violascensCA
100AZ
100TX
100 $\bar{x} = 100$

Index equals number of specimens with that count
 Specimens utilized were obtained from herbaria cited in Appendix I
 $\bar{x}$ = mean value

Seed Germination and Seed Set in Rumex Species

In general, the docks have a high rate of seed production (Chancellor 1956). R. crispus was reported by Salisbury (1942) to produce 25,000 to 40,000 seeds per plant, with an average of 30,000. Salisbury's germination tests indicated a 97 - 98 percent fertility, giving a viable seed set of 29,000 seeds. Cavers and Harper (1964) found that a large plant of R. obtusifolius can produce 60,000 seeds of which more than 80 percent are viable; smaller plants of both R. crispus and R. obtusifolius can produce as few as 100 seeds up to the higher levels previously noted. Under conditions suitable for germination, Cavers and Harper (1966) found that all their samples of R. crispus gave a 95 to 100 percent germination, while in R. obtusifolius, a range of 88 to 100 percent was observed. Although information of this nature is readily available for R. crispus and R. obtusifolius, this has not been the case for the other species of section Rumex.

Procedure

Plants representing ten species of section Rumex (R. britannica, R. conglomeratus, R. crispus, R. dentatus, R. kernerii, R. longifolius, R. obtusifolius, R. pseudonatronatus, R. pulcher, R. stenophyllus) were grown under controlled environmental conditions at Agriculture Canada, Ottawa. The sources of the plant material are indicated in the table of results.

As described earlier, (Temporal Separation of Flowering between Different Species, p. 51) flowering was induced by previously providing the plants with a cold treatment. In addition to allowing

the plants to set seed by open pollination, portions of the inflorescence were bagged to prevent this. The bags were placed over the branches prior to the most mature buds showing any signs of opening; sufficiently large bags were used to ensure that normal inflorescence development was not prevented by overcrowding. The bags were sealed by folding up the opening and stapling it shut and then allowed to remain on the plant until the inflorescence was seen to be fully mature. Samples from the unbagged portions of the inflorescence on the same plant were also collected. Fruits from these open-pollinated inflorescences were collected for comparative purposes. Initially, the samples were examined by comparing the outer appearance of the selfed-fruit with that of the open-pollinated fruit and by observing the fullness of the achenes (i.e. well-developed viable achenes are firm and fall; in contrast, achenes that are not well developed appear empty and are sterile).

The viability of the self-pollinated achenes was tested and compared to that of the open-pollinated achenes. Prior to testing the viability of the seeds, the samples were stored dry in paper bags at room temperature for approximately 1 year. The effect of storage conditions on viability has been discussed by Cavers (1974) and has been found to be very variable. Holm et al. (1977) in reviewing the literature on seed germination of R. crispus and R. obtusifolius stated that "there is general agreement that seeds of both species require light, alternating temperatures or both for germination". In examining seeds of R. obtusifolius and R. sanguineus, Thompson et al. (1977) found that a fluctuation of

2.0 - 2.5°C was sufficient to obtain 50 percent of the maximum germination recorded. The present study tested the viability by attempting to germinate the achenes on dampened filter paper at room temperature (20°C ± 5°C) and under available lighting (approximately 14 hr. daylight and 10 hr. darkness). A germination sample consisted of twenty seeds from which the perianth parts had been removed by hand. Cavers and Harper (1966) reported that when testing seed viability of R. crispus, seeds which had the perianth segments attached were not suitable for precise comparison as germination was intermittent (60 percent germination after 3 months) while naked seeds germinated quickly (95 percent germination after 1 week). In the present study, if the sample (or any portion of it) did not germinate after three weeks, it was given a two week long cold treatment at a constant temperature of 9°C in an attempt to induce germination as Cavers and Harper (1966) stated that pretreatment of R. crispus at low temperatures had been reported to increase the percentage germination. The percentage germination of the self and open-pollinated samples are listed in Table 8; the samples requiring a cold treatment are marked with CT as an index.

Four species (R. crispus, R. dentatus, R. pseudonatronatus, R. stenophyllus) were examined more closely to determine the actual seeds set. At least one whorl from the self and open-pollinated samples were selected and the total number of potential flower units were counted (i.e. the number of flowers and the number of pedicels from which the flowers had aborted and fallen off). The number of fruits that appeared to be fully developed externally

was counted and the number of fully developed fruits was expressed as a percentage of the potential flowering units (% Full Fruit).

Each of the fully developed valves was examined to observe whether the achene was fully developed and from this, the percentage of full fruit with fully developed seed (% Full Fruit with Full Seed) and the total percent seed set (No. of Fruit with Full Seed/ No. of Flowers) were calculated. The values obtained are recorded in Table 9; the corresponding seed viability measurements are in Table 8.

Results

No differences in the outward appearance of any of the samples of selfed fruit as against the open pollinated ones were observed; a visual assessment of both the selfed and pollinated fruit indicated that the majority of fruit had fully developed (few aborted or not fully developed fruit) with the exception of R. pseudonatronatus, which had sufficient fruit that were not fully developed to be noticeable.

The achenes removed from the twenty fully developed fruit of each sample to be tested for viability were all fully developed with firm achenes that did not appear to be empty. Five of the species tested (R. britannica, R. conglomeratus, R. dentatus, R. pulcher, R. stenophyllus) had samples requiring a cold treatment to obtain maximum germination (Table 5). In all but one of the tested samples of R. britannica and R. pulcher, a cold treatment was required, while in the other three species, the majority of

Table 8 Percentage Germination of Seed from Selfed
and Open Pollinated Flowers

The source of the seeds is indicated for each sample: in the case of the foreign material, the plants used were grown from seed obtained from the international seed exchange.

TABLE C

PERCENTAGE GERMINATION OF SEED FROM SELFED AND OPEN POLLINATED FLOWERS

SPECIES AND SOURCE	POLLINATED		SPECIES AND SOURCE	POLLINATED	
	SELF	OPEN		SELF	OPEN
<i>R. britannica</i>			<i>R. conglomeratus</i>		
No. 37 PE	70ct	20ct, 100ct	No. 22 France	100ct	95
No. 40 OH	100ct	25ct, 100ct	No. 26 France	100ct, 100	100
No. 120 OH	100ct	100ct	No. 132 France	100	100 ²
No. 43 OH	not selfed	20ct, 85ct, 100			
<i>R. crispus</i>			<i>R. dentatus</i>		
No. 16 H	100	100	No. 23 France	100 ²	100
No. 45 HK	100 ²	100	No. 18 France	100ct, 100	100
No. 46 HK	95, 100 ²	50			
No. 47 OH	100	100			
No. 61 HK	100	95, 100			
No. 81 OH	100ct	100			
No. 7624 HS	100	100			
<i>R. kernerii</i>			<i>R. longifolius</i>		
No. 50 France	100	100	No. 25 HS	100	100
			No. 36 HK	100	100
			No. 48 OH	100	90
			No. 50 HS	100	100
			No. 55 OH	100	100 ⁴
			No. 59 HK	100	100
<i>R. obtusifolius</i>			<i>R. pseudonatronatus</i>		
No. 3 Portugal	100	100 ³	No. 33 Hungary	100	100 ²
No. 13 Portugal	100	100	No. 62 Hungary	100	100 ²
No. 51 OH	100 ²	95, 100 ²	No. 67 Hungary	100	100
No. 68 QU	100	100	No. 113 Hungary	100	100
No. 69 QU	100	100 ²			
No. 7622 HS	100	100			

Table 8 Percentage Germination of Seed from
Selfed and Open Pollinated Flowers
(continued)

TABLE 2

PERCENTAGE GERMINATION OF SEED FROM SELFED AND OPEN POLLINATED FLOWERS, cont.

<u>SPECIES AND SOURCE</u>	<u>POLLINATED</u>		<u>SPECIES AND SOURCE</u>	<u>POLLINATED</u>	
	<u>SELF</u>	<u>OPEN</u>		<u>SELF</u>	<u>OPEN</u>
<u>R. pulcher</u>			<u>R. stonophyllus</u>		
No. 5 France	90 ^{ct}	100 ^{ct}	No. 21 USSR	100 ^{ct}	100 ^{ct}
No. 7 France	100	100 ^{ct}	No. 200 USSR	100	100
			No. 76151 SK	100 ^{ct}	100 ^{ct}
			No. 76152 SK	100 ^{ct}	100 ^{ct}
			No. 76168 SK	100 ^{ct}	100 ^{ct}

ct = Cold treatment required to complete germination.
Numerical Index equals number of samples tested.

Table 9 Species Seed Set

The table presents an assessment of the total percentage seed set of four species utilizing cultivated plants.

TABLE 9

SPECIES SEED SET

SPECIMEN NO. AND LOCATION	POLLINATION	NO. OF FLOWERS	1/3 FULL FRUIT WITH FULL SEED	1/3 FULL FRUIT WITH FULL SEED	TOTAL % SEED-SET
<i>Rumex crispus</i> 80, OH	Open	44	100	100	100
<i>Rumex crispus</i> 80, OH	Self	20	100	90	90
<i>Rumex crispus</i> 7624, NS	Open	20	100	100	100
<i>Rumex crispus</i> 7624, NS	Self	32	100	100	100
<i>Rumex dentatus</i> 23, France	Open	23	100	100	100
<i>Rumex dentatus</i> 23, France	Self	21	100	90.5	90.5
<i>Rumex dentatus</i> 23, France	Self	28	100	100	100
<i>Rumex pseudonatronatus</i> 113, Hungary	Open	129	47.3	98.4	46.5
<i>Rumex pseudonatronatus</i> 113 Hungary	Open	204	59.3	95.0	56.4
<i>Rumex stenophyllus</i> 76151 SK	Open	29	68.9	100	68.9
<i>Rumex stenophyllus</i> 76151 SK	Self	72	88.9	100	88.9
<i>Rumex stenophyllus</i> 76152 SK	Open	64	98.4	100	98.4
<i>Rumex stenophyllus</i> 76152 SK	Self	42	95.0	100	95.0
<i>Rumex stenophyllus</i> 76168 SK	Open	27	96.3	100	96.3
<i>Rumex stenophyllus</i> 76168 SK	Self	30	93.3	100	93.3

the samples did not require a cold treatment. The requirement of a cold treatment to obtain germination is probably indicative of a dormancy mechanism in these species; this factor was not pursued further as the present study was only concerned with the total germinability.

Only in the case of R. britannica was there a noticeable difference in the percentage germination between open and selfed achenes; in R. britannica there was much lower germination in achenes from the open-pollinated samples, although wide variation exists in samples from a single plant. It is possible that in R. britannica, conditions for consistently high germination were not provided. In general, viable seeds were produced under both the self and open-pollinated conditions and these experiments indicate that the ten species of Rumex examined are self-compatible.

In the four species examined more closely for seed set (Table 9), the number of flowers producing "full fruit" (based on external appearance) varied from 47.3 to 100 percent, with only R. pseudonatronatus being consistently at the lower end of the range. Achenes produced by the fruit that appeared fully developed externally (Full Fruit) were for the majority fully developed as 90 to 100 percent of the number of full fruit produced full achenes (% Full Fruit with Full Seed). The earlier germination tests on the full seeds indicated that these achenes were viable. For the most part, the total seed set was equal to the percentage of full fruit. Within a species, little difference was apparent between the self and open-pollinated samples tested. As R. pseudonatronatus

was the only species of the ten species involved in the viability study in which a large number of fruit did not develop fully, its seed set of 46.5 percent is assumed to represent the lower end of the potential seed set for the other species with most ranging from this value to 100 percent.

Based on the species involved in this study, members of section Rumex produce viable achenes under experimental conditions that prevent cross-pollination. The percentage germination reported for Rumex species in the literature may not be clearly indicative of the reproductive capacity of the plants unless the study indicates how the achenes were selected. This is evident from the large divergence of the seed set as compared to the seed germination values in R. pseudonatronatus. The values in this study were obtained to provide a baseline from which plants of hybrid origin could be compared with those in which no hybridization was suspected; a high seed set being indicative of plants of non hybrid origin while a low seed set would be indicative of plants of hybrid origin. Later references to these results will be made in the study of hybrid seed set.

Cytology

Within the genus Rumex, chromosome numbers and sizes are characteristics of the major groups: the subgenera of Rechinger (1954, 1964) and the genera of Löve (1967) (Ichikawa et al. 1971; Degraeve 1974, 1975b). In section Rumex, the basic chromosome number is $x = 10$, with only one contradictory report, that of Segiura (1936; cited in Ichikawa et al. 1971) who recorded $2n = 18$ in R. conglomeratus. This would seem to be based either on a misidentified specimen or is an error in counting as the many later counts on this species are all of $2n = 20$. The North American species of section Rumex have been found to represent the following ploidy levels: 2, 4, 6, 8, 10, 14, 16, 17, 18 and 20, with the majority of the species being represented by the first three levels. Ichikawa et al. (1971) and Degraeve (1975) found the chromosomes of section Rumex and section Axillares to be very similar in size. Within section Axillares, Sarkar (1958) found that karyotypic analysis could not be undertaken with any degree of certainty due to the small size of the chromosome and Ichikawa et al. (1971) made no attempt at karyotyping species of the genus Rumex, including section Rumex for the same reason. A more recent study by Degraeve (1975) has attempted this by creating an average karyotype of the basic haploid set utilizing the averages of the chromosome length for the species studied in his investigations. He found that the size varied primarily with the degree of polyploidy and he obtained the following averages: 1.50 for diploids, 1.11 for tetraploids, 1.13 for hexaploids, 1.30 for octaploids, 1.06 for decaploids

and 1.43 for higher ploidy levels.

Numerous counts have been made on the majority of the North American species of section Rumex. Only four, R. praecox, R. pycnanthus, R. orthoneurus and R. violascens, all rather restricted in range, have no reported counts. To provide a basis for comparison of the species and assessment of hybrid counts, a review of the known counts is presented in Table 10. When available, the source of the seeds or plants upon which the counts were made has been given. For the chromosome counts, by Mulligan, the voucher specimens have been examined and the identifications confirmed. (see Table 10). In addition, the fruits and often the plants used by Ichikawa et al. (1971) were identified by Dr. C. Frankton, Biosystematics Research Institute, Ottawa, and these counts can be considered to be based on material whose identification is reliable.

In addition to the counts summarized from the literature, Table 10 includes four counts made by me during this investigation. Root tips used for mitotic chromosome counts were collected in 8-hydroxyquinoline and left for two hours before fixing in an absolute alcohol-glacial acetic acid mixture (3:1) for at least 24 hours. The fixative was washed out with three rinses of 70% ethanol and at this point, the material was refrigerated and stored. Portions of the inflorescences (at varying stages prior to opening of the flowers), collected for meiotic chromosome counts, were fixed in an absolute alcohol, chloroform, glacial acetic acid mixture (6:3:1) prepared just before use. After at least 24 hours, the

fixative was washed out as for the root tips using 70% ethanol. For both the root tips and the buds, squashes were prepared using Snow's alcoholic hydrochloric acid-carmin staining procedure (Snow 1963). Mitotic counts were made on R. crispus, R. longifolius, and R. obtusifolius; a meiotic count was obtained from R. longifolius.

From the table, it can be seen that the chromosome counts for the majority of the species are in agreement with one another, with the exception of some counts of the following species: R. arcticus, R. britannica, R. confertus, R. conglomeratus, R. longifolius, R. occidentalis, R. palustris and R. pulcher.

In R. arcticus, the chromosome number has been found to vary from $2n = 40$ to $2n = ca\ 200$. The count of $2n = ca\ 200$ from Finland is questionable as the species is not known to occur there (Rechinger 1964; Mulligan 1969). A detailed discussion of the chromosome numbers of this species is given by Mulligan and Frankton (1972). They found the North American material to be tetraploid with the exception of plants from the northwest coast of Alaska which, like those from the north-western U.S.S.R., had higher chromosome numbers. A recent count of $2n = 40$ by Löve (1975) from Kamchatka indicates that the situation may not be clear cut. Mulligan and Frankton (1972) did not recognize the different chromosome races taxonomically on account of their great morphological similarity.

In R. confertus, R. palustris and R. pulcher, only one

divergent count has been reported. For these species there is no evidence to indicate whether the divergent counts are due to misidentified specimens, inaccurate counts or the occurrence of more than one chromosome number in the species.

In R. longifolius, the count of $2n = 80$ by Kihara and Ono (1926; cited in Ichikawa et al. 1971) has not been repeated. The existence of two chromosome races in Europe, one with $2n = 40$, and the other with $2n = 60$, has been discussed by Jalas and Lindholm (1975). In North America, the numerous counts suggest that only plants with $2n = 60$ occur.

R. britannica and R. occidentalis both North American native species, are reportedly represented by several counts at very high ploidy levels. An examination of available herbarium material of R. britannica indicated a relatively uniform species morphologically making the possibility of correlation of morphological differences with chromosome counts unlikely. In the case of R. occidentalis, an examination of the four voucher specimens of Mulligan (1957) ($2n = 140$) and Taylor and Mulligan (1968) ($2n = ca 140$) collected from across much of the Canadian range of the species indicated that although all apparently at the same ploidy level, these specimens represented much of the morphological variation reported for the species. The divergent counts of Degraeve (1975) ($2n = 160$), Love and Ritchie (1966) ($2n = ca 200$) and Love (1967) ($2n = ca 200$) are from plants collected in the same areas as others from which counts of $2n = ca 140$ have been obtained. Although most counts made

are of $2n = 140$ and $2n = ca\ 140$ and these come from throughout the species' range, many additional counts would be needed to confirm the existence and extent of possible chromosome races.

The few meiotic counts available are all from R. crispus ($2n = 60$), R. longifolius ($2n = 60$) and R. obtusifolius ($2n = 40$). These indicate regular pairing during meiosis in these species. It is not wise to extrapolate from these results to species of other ploidy levels, especially the higher ones. These three species are however, all involved in the more common hybrids in North America; these hybrids show irregular pairing which will be discussed in the section on hybridization.

In summary, chromosome numbers within each species of section Rumex in North America are generally in agreement with each other, with the exception of R. arcticus, R. britannica and R. occidentalis, in which major cytological studies would be needed to determine the full range of cytological variability within each species. At present, the lack of karyotypic analysis has prevented the use of cytological data in providing further insight into interspecific or intersubgeneric relationships in Rumex. Further studies such as that by Degraeve (1975) may help to change this and might provide additional evidence that the present groupings are natural ones. The different ploidy levels within section Rumex would be expected to act as at least partial barriers to hybridization between species that are at different levels as pairing at meiosis would be irregular, preventing the development

or survival of a hybrid. Even in hybrids between species of the same ploidy level, eg. R. crispus X patientia, or R. crispus X longifolius, sterile pollen and few viable seeds are produced (Loualely and Williams 1974) suggesting that other barriers to hybridization exist between species of section Rumex. Knowledge of the chromosome counts of the parental species allows chromosome numbers of a hybrid to be predicted, which can help to resolve questions of hybrid parentage.

TABLE 10

Chromosome Numbers - section Rumex

<u>Species</u> <u>Chromosome Number</u>	<u>Author</u>	<u>Source</u>
<u>R. alpinus</u> <u>2n = 20</u>	Love (1967)	Romania: near Sibiu, A. Love A. Love RO2461
	Ichikawa et. al (1971)	Botanic Garden origin (Copen- hagen, Rouen)
	Degraeve (1975)	Botanic Garden origin, (Oslo, Liège)
	Additional reports in A, B,C,D	
<u>R. arcticus</u> <u>2n = 40</u>	Knaben (1968 in B,E)	Alaska: Central River Dist., White Mtns., Gjaervoll 623 (TRH)
	Mulligan and Porsild (1969)	Yukon Territory: Orilgive Mts., R.T.P. 155 (CAN)*
	Mulligan and Frankton (1972)	Mackenzie Dist., N.W.T. Cody 16969 (DAO)*; Cody 17505 (DAO)*; Johanssen s.n. (DAO)*
	Love ^I (1975)	U.S.S.R.: Kamchatka, Love and Löve G.67
<u>2n = 120</u>	Ichikawa et al (1971)	U.S.S.R. Novosibirsk, D.A.J. Kerkis; Botanic Garden origin (Ottawa)
	Mulligan and Frankton (1972)	U.S.S.R.: Chukhotsk Pen, Frankton 2332 (DAO)*
	Pond (cited in Mulligan and Frankton 1972)	U.S.S.R.: Chukhotsk Pen, Frankton 2332 (DAO)*
	Löve (1975)	U.S.S.R.: Kamchatka, Love and Löve, G 2232

<u>Species</u> <u>Chromosome Number</u>	<u>Author</u>	<u>Source</u>
<u>2n</u> = 150	Zhukova (1965 in B,E)	U.S.S.R.: Chukhotsk Pen.
<u>2n</u> = ca 170	Johnson and Packer (1965)	Alaska: Ogotoruk Creek Valley
<u>2n</u> = ca 200	Löve (Löve and Löve 1948 in A ² , B)	Finland
	Sokolovskaya (1968 in A ² , b, E ³)	U.S.S.R.: Kamchatka Pen
	Reported under ¹ R. <u>Kamtschadalus</u> Kam.; ² R. <u>aquaticus</u> ssp <u>arcticus</u> (Trautv) Hiit; ³ R. <u>arcticus</u> Trautv. var. <u>kamtschadalus</u> (Kam.) Rech. f.	
<u>R. britannica</u> <u>2n</u> = 160, ca 170, 180	Ichikawa et al. (1971) Additional report in A,B	Botanic Garden origin (Ottawa)
<u>2n</u> = ca 200	Löve (1967) All reported under R. orbiculatus	Canada: Manitoba, Winnipeg, A. Löve R05367
<u>R. brownii</u> <u>2n</u> = 40	Degraeve (1975) Additional report in A,B	Botanic Garden origin (Brisbane)
<u>R. confertus</u> <u>2n</u> = 40	Löve (Löve and Löve, 1961, in A,B)	
<u>2n</u> = 100	Mulligan, 1969 Ichikawa et al. 1971	Rumania: Cluj Bot. Gard. Frankton 1962 (DAO)*, Austria; Wein Rangierbahnhof Floridsdorf; Rechinger 249, 1949 (DAO)* Botanic Garden origin (Copen- hagen, East Berlin; Riga Latvian S.S.R.)

<u>Species</u> <u>Chromosome Number</u>	<u>Author</u>	<u>Source</u>
<u>R. conglomeratus</u> <u>2n = 18</u>	Seguira (1936 in B)	
<u>2n = 20</u>	Löve (1967 ^a , ^b)	^a Sweden: Scania, Lund. A. Löve . R059 ^b Spain: Sierra Nevada, near Cazorla, A. Löve R0211
	Ichikawa et al. (1971)	Botanic Garden origin (Copen- hagen, Rouen)
	Degraeve (1975)	Botanic Garden origin (Bruxelles Coimbra, Portugal; Vacratot, Hungary)
	Löve (1976)	France: Littoral languedocien Mauguio, Labadie
	^a reported under ssp. <u>conglomeratus</u> ; ^b reported under ssp <u>nevadensis</u> H. Lindb.	
	Additional reports in A,B,C	
<u>R. crispus</u> <u>2n = 60</u>	Kihara (1927)	
	Löve (1967)	Sweden: Scania, Lund, A. Löve R02111
	Mulligan (1957)	Canada: Ottawa, C. Frankton, 2132 (DAO)*; Sask., Crichton, A.C. Budd 2222 (DAO)*; (B.C., Langley Prairie, Mulligan 2394 (DAO)*, counted in 1958, not reported)
	Taylor and Mulligan (1966)	Canada: B.C. Queen Charlotte Islands, (2n = 30) Calder and Taylor 36938 (DAO 98530)*; (2n = 30II) Calder and Taylor 35449 (DAO 98533)*; Calder and Taylor 35333 (DAO 98536)*.
	Ichikawa et al. (1971)	Botanic Garden origin (Ottawa; Upton, New York)
	Degraeve (1975)	Botanic Garden origin (Coimbra, Portugal, Istanbul; Brisbane)

<u>Species</u> <u>Chromosome Number</u>	<u>Author</u>	<u>Source</u>
	Dawson	Canada: N.S. 13 mi. N. of Oxford, Hy 2, Dawson 74-74.
	Additional reports in A,B, C,D	
<u>R. densiflorus</u> <u>2n = 100</u>	Degraeve (1975)	Botanic Garden origin (Boulder)
<u>R. dentatus</u> <u>2n = 40</u>	Löve ^{b-h} (1967)	^b Sweden: Gottenburg, Bot. Gard., A. Löve R0101 ^c Georgian S.S.R.: Tbilisi Bot. Gard., A. Löve R0245 ^d India: near Calcutta, F.S. Merrisor, A. Löve R0201 ^e Iraq: Near Baghdad, M.S. Smith, A. Löve R0153 ^f India: near Madras, A.G. John, A. Löve R01149 ^g Taiwan: Taipei, C.C. Chang, A. Löve R02593 ^h Iraq: near Baghdad M.S. Smith, A. Löve R0155
	Ichikawa et al. (1971)	Botanic ^g Garden origin (Copen- hagen)
	Degraeve (1975)	Botanic Garden origin (London)
	Löve (1975)	Botanic Garden origin (West Bengal, India)
	Additional reports A,B,C,D	
	reported under ^a ssp. <u>callosissimus</u> (Meisn.) Rech.f.;	
	^b ssp. <u>dentatus</u> : ^c ssp. <u>halacsyi</u> ; (Rech. f.) Rech.f.;	
	^d ssp. <u>kotzschianus</u> (Meisn.) Rech. f.; ^e ssp. <u>mesopotamicus</u>	
	Rech. f.; ^f ssp. <u>nigricans</u> (Hook.) Rech. f.; ^g ssp.	
	<u>nipponicus</u> (Franch. and Saur.) Rech. f.; ^h ssp. <u>reticulatus</u>	
	(Bees.) Rech. f.	
<u>E. hymenosepalus</u> <u>2n = 40</u>	Love and Patil (1967)	U.S.A.: New Mexico, near Gab- rillo Dam Hy 85, Löve and Löve

<u>Species</u> <u>Chromosome Number</u>	<u>Author</u>	<u>Source</u>
	Mulligan (1969)	U.S.A.: Texas, 29 mi. E. of El Paso, T. and L. Mosquin 5672 (DAO)*
<u>R. kernerii</u> <u>2n = 60</u>	Degraeve (1975)	Botanic Garden origin (Dallas, Liege)
<u>R. longifolius</u> <u>2n = 40</u>	Hamet-Ahti and Virrankowski (1970 in C, D)	northern Finland
	Jalas and Lindholm (1975)	Norway: Akerhus, Sundig; Finland South Hame Saarisalo, Kuusamo, Ahti
<u>2n = 60</u>	Mulligan (1959)	Canada: Que., Bedford, Frankton, Mulligan 2369 (DAO)*; Ont., Temagami, Bassett and Bragg 3075 Mulligan 2241 (DAO)*; Ottawa, Frankton, Mulligan 2392 (DAO)*
	Love (1967)	Sweden: Scania Lund, A. Löve R073
	Ichikawa et al. (1971)	Botanic Garden origin (Copenhagen, Ottawa)
	Degraeve (1975)	Botanic Garden origin (Lund, Oslo, Ottawa)
	Jalas and Lindholm (1975)	Spain: Cataluna, Jalas (three different collections); Norway; Hordaland, Wielgolaski; Akershus Sunding; Sweden: Skane, Vailin; Halland, Bothmer; Vastergotland; Bothmer; Stockholme, Bothmer; Dalarne, Bothmer; Finland: Varsinais-Suomi, Kytovuori; Urisinau, Uotila; Vusimaa, Saarisalo; South Hame, Saarisalo South Hame, Yotila, North Hame, Saarisalo; Kuusamo, Ahti
	Dawson	Canada: (2n = 30II) N.B. St. Anne, Dawson 74-1; Ontario, Kemptville, Dawson 74-210

<u>Species</u> <u>Chromosome Number</u>	<u>Author</u>	<u>Source</u>
<u>2n</u> = 80	Kihara and Ona (1926 in B)	Asia
	Additional reports in A,B	
<u>R. maritimus</u> <u>2n</u> = 40	Mulligan (1959) ^c	Canada: P.E.I. Brackley Beach, Erskine, Mulligan 2184 (DAO)*
	Löve and Löve (Löve 1967 ^{b,d})	Canada: Man. Winnipeg, Löve and Löve 5083; Que. Ile d'Orleans, Löve and Löve 7341
	Ichikawa et al (1971) ^a	Botanic Garden origin (Ottawa, Santa Ana, California)
	Degraeve (1975) ^b	Botanic Garden origin (Dallas, London, Ottawa)
	reported under ^a <u>R. maritimus</u> L., ^b <u>R. fueginus</u> Phil., ^c <u>R. maritimus</u> L. var. <u>fueginus</u> (Phil.) Dusen, <u>R. persicarioides</u> L.	
	Additional reports in A,B, C.	
<u>R. obtusifolius</u> <u>2n</u> = 40	Mulligan (1959) ^a	Canada: B.C. Langley Prairie, Jenkins, Mulligan 2242 (DAO)*
	Löve (1967) ^c	Sweden: Scania near Svalöv, A. Löve RQ407
	Taylor and Mulligan (1968)	Canada: B.C. (2n = 20II) Graham Is., between Skidegate and Skidegate Mission, Calder and Taylor 35840 (DAO)*; Floresby Is., Sandspit, Calder and Taylor 35355 (DAO)*
	Ichikawa et al. (1971)	^{a-c} Botanic Garden origin (Copenhagen, Ottawa) ^a Costa Rica, Dr. C.C. Hah
	Degraeve (1975)	Botanic Garden origin (Copenhagen; Lund; Vacratot, Hungary)
	Dawson	Canada: N.S. 5 mi from Gaspereau on road to Black River, Dawson 74-93

Species
Chromosome Numbers

Author

Source

reported under R. obtusifolius ssp. obtusifolius (subsp. agrestis (Fries) Danser); ssp. sylvestris (Wallr.) Rech.; ssp. transiens (Simonakal) Rech. f.; ssp. subalpinus (Schur) Celak.

Additional reports in A,B,C,D

R. occidentalis
2n = ca 140

Mulligan (1957)

Canada: Man. Macdonald, Frankton, Mulligan 2223 (DAO)*; Sask. 18 mi N. of Swift Current, Budd, Mulligan 2183 (DAO)*; Sask. Crichton, Budd, Mulligan 2182 (DAO)*

Taylor and Mulligan (1966)

Canada: B.C. Moresby Is., Ana Inlet, Calder, Taylor, Sherk 34926 (DAO)*

2n = 140

Ichikawa et al (1971)

Botanic Garden origin (Ottawa)

2n = 160

Degraeve (1975)

Botanic Garden origin (Ottawa, Dallas)

2n = ca 200

Löve and Ritchie (1966)

Canada: Manitoba

Löve (1967)^e

Canada: B.C., near Nanaimo, A. Löve R05936

^acited as R. fenestratus Greene.

R. palustris
2n = 40

Degraeve (1975)

Botanic Garden origin; (London; Uppsala; Vacratot, Hungary)

2n = 60

Mulligan (1969)

Botanic Garden origin (Cambridge Frankton 2030 (DAO)*

Ichikawa et al. (1971)

Botanic Garden origin (Copenhagen) Frankton 2029 (DAO)* (Cambridge)

<u>Species</u>	<u>Chromosome numbers</u>	<u>Author</u>	<u>Source</u>
<u>R. patientia</u> <u>2n = 60</u>		Löve (1967) ^{a-g}	<p>^aU.S.S.R.: Kamtschatka, R. Malaise, A. Löve RO497; ^bChina: Chia-Cheng, Pashui-ko, H. Smith 7534, A. Löve RO536 ^cSyria: Mt. Amanus, M.B. Sears, Löve RO499 ^dKashmir, near the Cherab, F.S. Grant, A. Löve RO507 ^eGeorgian SSR, Tblisi Bot. Gard., A. Löve RO521 ^fHungary: Budapest Bot. Gard., A. Löve RO498 ^gKashmir: Setpar Valley, F.S. Grant, A. Löve RO511</p>
		Ichikawa et al (1971) ^{c,d,e}	Botanic Garden origin (^a , ^d Stockholm, ^c Bremen, ^e Copenhagen)
		Degraeve (1975)	Botanic Garden origin (Coimbra, Portugal; Copenhagen: Liege)
		cited as ^a <u>Rumex patientia</u> L. ssp. <u>callosus</u> (Schmidt) Rec. f., ^b ssp. <u>interruptus</u> (Rech. f.) Rech. f., ^c ssp. <u>orientalis</u> (Bernh.) Danser, ^d ssp. <u>pamiricus</u> Rech. f., ^e ssp. <u>patientia</u> fssp. <u>recurvatus</u> (Rech. f.) Rech. f., ^g ssp. <u>tibeticus</u> (Rech. f.) Rech. f.	
		Additional reports in A,B,C,D	
<u>R. pseudonatronatus</u> <u>2n = 40</u>		Mulligan (1957) (1958 not reported) ¹	<p>Canada: Man. Macdonald; Frankton, Mulligan 2224 (DAO)*; Man. Meadow, Frankton, Mulligan 2225 (DAO)*; Sask. Gravebourg Frankton and Bassett 1579, Mulligan 2226 (DAO)*; Sask. Crichton Budd, Mulligan 2181 (DAO)*; U.S.A.: South Dakota Conde Brenckle, Mulligan 2228 (DAO)* ¹Ont., Ottawa, Frankton, Mulligan, 2391 (DAO)*</p>
		Löve (1967)	Finland: Helsinki Bot. Gard. A. Löve RO479
		Ichikawa et al. (1971)	Botanic Garden origin (Ottawa)
		Degraeve (1975)	Botanic Garden origin (London; Vacratot, Hungary)
		Additional reports in A,C	

<u>Species</u> <u>Chromosome Numbers</u>	<u>Author</u>	<u>Source</u>
<u>R. pulcher</u> <u>2n = 20</u>	Love (1967) ^{a-e}	^a Israel: Mt. Scopus, F. Saunders A. Löve R0555 ^b Israel: Mt. Corbel, F. Saunders A. Löve R0560 ^c Italy: near Lucca, A. Löve R05634 ^d Sweden: Lund, A. Löve R023 ^e Greece: Samos, M.N. Dyson: A. Löve R0691
	Ichikawa et al. ^{a,d} (1971)	Botanic Garden origin (Rouen, Copenhagen)
	Degraeve (1975)	Botanic Garden origin (Coimbra, Portugal; Istanbul: Liege)
<u>2n = 40</u>	Jaretsky (1928 in c)	
	reported under ^a ssp. <u>anodontus</u> (Hausskn.) Rech. f., ^b ssp. <u>cassius</u> (Boiss.) Rech. f., ^c ssp. <u>divarcatulus</u> (L.) Harb, ^d ssp. <u>pulcher</u> , ^e ssp. <u>rauland</u> (Boiss.) Rech. f.	
	Additional reports in A,B,C,D	
<u>R. sanguineus</u> <u>2n = 20</u>	Löve (1967)	Sweden: Scania, Svalou. A. Löve R057
	Ichikawa et al. (1971)	Botanic Garden origin (Copen- hagen, Rouen)
	Degraeve, 1975	Botanic Garden origin (Copen- hagen, Lund; Vacratot, Hungary)
	Additional reports in A,B,C	
<u>R. stenophyllus</u> <u>2n = 60</u>	Palya (1950 in A,B,C)	Hungary
	Mulligan (1957)	Canada: Sask: Swift Current, Budd, Mulligan 2136 (DAC)*
	Sarkar (Löve and Bernard 1958)	Canada: Man.
	Löve (1967)	Canada: Man. near Steinbech, A. Löve R05346

Species
Chromosome Numbers

Author

Source

Ichikawa et al. (1971)

Botanic Garden origin (Copenhagen, Ottawa)

Degraeve (1975)

Botanic Garden origin (London; Vacratot, Hungary)

A Löve and Kapoor, 1967

B Ichikawa et al., 1971

C Degraeve, 1975

D Moore, 1973

E Mulligan and Frankton, 1972

* voucher specimen examined and found to be correctly determined.

Hybrids

In 1856, Meisner reported the first Rumex hybrid in North America: R. crispus X obtusifolius from the state of Missouri. Later, studies by Trelease (1892) extended the range of this hybrid (DC, IL, MA, MI, MO, NY). No additional hybrids were reported until 1937, when Reehinger identified two further hybrids - R. britannica X crispus from Massachusetts and R. crispus X pulcher from Virginia; at the same time, the known distribution of R. crispus X obtusifolius was further extended to include seven additional states (CA, LA, OH, OR, PA, TX, WI) and one Canadian province (NF) and by then was essentially distributed from coast to coast. More recently, Boivin (1968) has listed three hybrids as occurring in Canada - R. crispus X obtusifolius from Newfoundland, R. pseudonatronatus X triangulivalvis from Saskatchewan and R. maritimus X stenophyllus also from Saskatchewan. R. pseudonatronatus X triangulivalvis first reported by Mulligan (1959) represents the only known hybrid between section Rumex and a member of another section (Sect. Axillares, to which R. triangulivalvis belongs).

Within Rumex, the following characteristics have been used to detect hybrid plants: pollen produced by the hybrids has a low percentage of fertile grains (Lousley 1939, Cavers and Harper 1964, Lousley and Williams 1975); the majority of flowers do not set fruit, but fall from the inflorescence with little enlargement of the perianth parts (Trelease 1892, Lousley 1939, Cavers and Harper 1964, Lousley and Williams 1975); the inflorescence of hybrids takes

on a somewhat ragged appearance due to the irregular fruit development (Trelease 1892, Lousley 1979); in any whorl, the following features can be observed:

- (a) bases of peduncles of flowers that have fallen off,
- (b) dried up remains of perianths which have not developed,
- (c) partially enlarged perianth segments,
- (d) a few mature valves containing apparently good fruit;

finally, most reported hybrids have morphological characteristics intermediate between those of the parental species involved (Lousley and Williams 1975).

In the present study, hybridization has been examined in section Rumex in North America from the standpoint of the Distribution of Hybrids, Pollen Fertility, Seed Set and Seed Viability, Hybrid Cytology and the Morphological Characteristics of Hybrids in an attempt to provide an up-to-date assessment of the present situation.

Distribution of Hybrids in section Rumex in North America

Procedure

With a view to detecting hybrids, the morphological characteristics reported earlier (Hybrids, p.91) were examined on approximately 9,000 herbarium specimens from throughout North America. (The herbaria utilized are cited in the Appendix I). The localities from which hybrid specimens were found were mapped; field work in the Maritimes of Provinces of Canada during the summers of 1974 and 1976 provided further distributional information.

Results

The following interspecific hybrids are now recognized as occurring in North America with the accompanying distributions.

R. britannica X crispus (NB; CT, MA, PA, RI) (Map 1)

R. crispus X kernerii (CA) (Map 1)

R. crispus X longifolius (ON, MB, NS*) (Map 1)

* R. crispus X obtusifolius (BC, NF, NS*, ON, PE; CA, DC, IL, LA, MA, MI, MO, NC, NH, NY, OH, OK, OR, PA, TN, VA, WV, WI, TX)
This is a much wider geographic range in Canada and the United States than was previously reported. (Map 2)

R. crispus X patientia (ON, UT) (Map 1)

R. crispus X pulcher (CA, VA) (Map 1)

R. conglomeratus X pulcher (VA) (Map 1)

R. longifolius X obtusifolius (NS*) (Map 1)

R. maritimus X stenophyllus (SK) (Map 1)

R. occidentalis X pseudonatronatus (SK) (Map 1)

R. pseudonatronatus X triangulivalvis (SK) (Map 1)

The asterisk indicates hybrids discovered during my field studies and not previously reported from these areas.

R. longifolius X obtusifolius, although known in Europe (Lousley and Williams 1975) has not previously been reported from North America. From field studies, it appears that this hybrid is relatively common in southern Nova Scotia, occurring wherever large patches of the parent species were located. In a number of sites, there were as many plants of the hybrid as of the parental species.

It is interesting to note that seven of the eleven hybrids involve R. crispus or R. obtusifolius and that the most common hybrid is between these two species. R. crispus X obtusifolius is also the most common Rumex hybrid in Europe. (Lousley and Williams 1975). Since Rechinger's (1937) study, eight additional hybrids have been reported for North America, but none of these have wide distributions. Of the eleven North American hybrids, only R. crispus X obtusifolius is widespread but from the distribution of the parents, one would expect it to be even more prevalent than the present study suggests. The ranges of the majority of Rumex species in Canada and the United States overlap but only half of the recognized species have been involved in hybridization. This lack of

Hybrids may be due to one or more of the following factors:

- (a) the species exhibit habitat preferences, even though their overall ranges overlap;
- (b) the lack of extensive collections of Rumex by collectors who would recognize the hybrids as such;
- (c) internal barriers preventing or limiting interspecific hybridization.

Map 1 Distribution of the hybrids of section
Rumex in Canada and the United States.

Map 1

Map 2 Distribution of R. crispus x obtusifolius
(the most common hybrid of the section) in
Canada and the United States.

Map 2

Pollen Fertility of Hybrids

As noted in the introduction to the study of Rumex hybrids, the hybrids have been reported to produce only a small proportion of fertile pollen grains. Murbeck (1913) and Lousley (1939) noted that non-hybrid plants of Rumex produced grains of a uniform size, whereas those of interspecific hybrids had pollen grains of various sizes, with the majority being much smaller than those of the parents and empty or shrunken.

Procedure

Rumex specimens identified as hybrids (by using the morphological characteristics discussed earlier) and in a flowering stage were tested for their pollen fertility, using the lactophenol cotton blue method as outlined earlier (Breeding System - Pollen Fertility of Rumex Species p.59). To calculate the percentage fertility, two counts of 100 grains were made, each count based on a preparation using two anthers from a single flower. The anthers were taken from buds that had fully opened but in which the anthers had not yet dehisced. Herbarium material was available of the following hybrids:

- R. crispus ($2n = 60$) X obtusifolius ($2n = 40$)
- R. crispus ($2n = 60$) X patientia ($2n = 60$)
- R. crispus ($2n = 60$) X pulcher ($2n = 20$)
- R. longifolius ($2n = 60$) X obtusifolius ($2n = 40$)
- R. pseudonatronatus ($2n = 40$) X triangulivalvis ($2n = 20$)

European material was utilized in a number of cases in order to provide additional data, especially when suitable North American specimens were lacking. The percentage fertility of these hybrids is listed in Table 11.

Pollen fertility was also examined in flowering material of R. crispus X obtusifolius and R. longifolius X obtusifolius from field collections made in Nova Scotia in 1976; several counts were obtained using material collected in the field, but cultivated in the greenhouse. The percentage fertility of the Nova Scotian hybrids is listed in Table 12.

Results

The size disparity and the high percentage of shrunken, non-staining grains were evident in the majority of the samples³ examined; Tables 11 and 12 indicate the low levels of pollen viability found in the hybrid.

R. crispus X obtusifolius (utilizing both tables) exhibited a range of 0 - 12 (69)% fertility and a mean value of 7%, whereas from Table 7, R. crispus is shown to have a range of 77 - 100% pollen fertility and a mean value of 92%; correspondingly, R. obtusifolius exhibits a range of 80 - 100% fertility and a mean value of 96%. With the exception of the high count of 69% from Koriuge in Sweden, the pollen fertility of this hybrid is substantially below that of either parent. The pollen fertility of R. crispus X obtusifolius as measured by Cavers and Harper (1964) was less than 1% whereas the values in Tables 11 and 12 are for the most part slightly higher. Cavers and Harper also noted that the larger grains found in the pollen of the hybrid were 1.5 times the diameter of those of the

TABLE 11.

POLLEN FERTILITY OF HYBRIDS

<u>HYBRID</u>	<u>LOCATION AND SOURCE</u>	<u>% FERTILITY</u>
<u>R. crispus</u> X <u>obtusifolius</u>	Hespeler, Ont. (DAO)	2
	Chatsworth, Ont. (DAO)	3
	Sault Ste. Marie, Ont. (DAO)	8
	Binghamton, N.Y. (F 79479)	4
	Oak Bluffs, Mass. (GH)	12
	Marblehead, Mass. (GH)	2
	Gloucester, Mass. (GH)	7
	vicinity of Washington, D.C. (GH)	2
	N.W. of Sleepy Cr., W.V. (NCU 372756)	2
	Rocky Mount, H.C. (NCU 204245)	0
	N.N.W. of Winton, N.C. (NCU 204241)	1
	N.N.W. of Winton, N.C. (NCU 204242)	3
	S.E. of Big Laurel, N.C. (NCU 204236)	0
	S.E. of Big Laurel, N.C. (NCU 204240)	0
	W. of Hunt Dale, N.C. (NCU 204243)	6
	Svenstrup, Denmark (DAO)	2
<u>R. crispus</u> X <u>patientia</u>	Korluge, Sweden (DAO)	69
	Ronneby, Sweden (DAO)	1
	Skane, Sweden (DAO)	6
<u>R. crispus</u> X <u>pulcher</u>	Murray, Utah (GH)	7
	Warwick, Bermuda (GH)	0
	Warwick Pond, Bermuda (GH)	0
	Smilyhs Parish, Bermuda (GH)	0
<u>R. longifolius</u> X <u>obtusifolius</u>	Sakkola, Finland (DAO)	1
<u>R. pseudonatronatus</u> X <u>triangulivalvis</u>	Kindersley, Sask. (DAO)	59
	cult. from seed of above, (Mulligan, 1635, DAO)	92

TABLE 12

POLLEN FERTILITY OF NOVA SCOTIAN HYBRIDS

<u>HYBRID</u>	<u>LOCATION AND COLLECTION NO.</u>	<u>% FERTILITY</u>
<u>R. crispus</u> X <u>obtusifolius</u>	E. of Clementsport 7658	1
	just S. of Yarmouth 7666	7
	just S. of Yarmouth 7668	1
	just S. of Yarmouth 76206	14*
	just S. of Yarmouth 76209	12*
	Baptist Church, Overton 76214	9*
<u>R. longifolius</u> X <u>obtusifolius</u>	Yarmouth 7616	7*
	on rd to Cape Forchu 7622	3
	on rd to Cape Forchu 76215	2
	Baptist Church, Overton 7646	2
	7647	1
	7648	1
	7649	1
	7650	2
	7651	2
	7653	13*
	76193	2
	76196	2
	76197	4
	76203	3
	76205	24
	just S. of Yarmouth 7669	0
	just S. of Yarmouth 76220	2
	just S. of Yarmouth 76222	2
	S. of Mavillette 7671	2
	S. end of Meteghan 7674	1
	Middle of Meteghan 7679	0
	Middle of Meteghan 7680	2
	Middle of Meteghan 7681	3
	Middle of Meteghan 76249	6
	Middle of Meteghan 76251	3
	Comeauville 76254	5
	Comeauville 76258	6

* = from cultivated material,
root collected in field.

parent species. Lousley and Williams (1975) reported that an artificially produced F1 hybrid between R. crispus and R. obtusifolius had pollen of 80% fertility; however, no further information about this plant was given.

R. crispus X patientia (Table 11) produced only 7% fertile pollen, a much lower level of fertility than that of R. crispus (Table 7); correspondingly, R. patientia shows a range of 98 - 100% pollen fertility and a mean value of 99%. This hybrid represents a cross between species of the same ploidy level; such a low pollen fertility relative to the parents suggests irregularities during meiosis due to genetic differences between the haploid complement of chromosomes from each parent.

R. crispus X pulcher (Table 11) did not produce any fertile pollen; R. pulcher shows a range of 97 - 99% pollen fertility and a mean value of 98%; the values in R. crispus are also much higher as indicated above.

R. longifolius X obtusifolius (Tables 11 and 12) shows a range of 0 - 24% pollen fertility and a mean value of 4%, whereas R. longifolius exhibits a range of 60 - 100% pollen fertility and a mean value of 95% (Table 7); correspondingly, R. obtusifolius shows high pollen fertility as noted above. Even the upper value of 24% for this hybrid is well below that of either parent.

The hybrid R. pseudonatronatus X triangulivalvis in addition to being a cross between members of section Rumex (R. pseudonatronatus) and section Axillares is of interest as the plant examined had a very high pollen fertility (59%). R. pseudonatronatus exhibits a

range in pollen fertility between 74 and 100% with a mean value of 89% (Table 7), whereas for R. triangulivalvis, Sarkar (1958) reports a range of 91 - 97% pollen fertility and a mean value of 97%. Although still less fertile than either parent, the hybrid is within the lower part of the ranges of fertility of some species (Table 7). Until further collections of this hybrid are made, it is impossible to determine whether such a relatively high fertility level is typical of this hybrid combination, or an uncommon occurrence as in the case of R. crispus X obtusifolius from Sweden. The cultivated specimen of R. pseudonatronatus X triangulivalvis with a pollen fertility of 92% represents an F2 plant grown by G. A. Mulligan (B.R.I., Ottawa) from one of the few viable seeds collected from the F1 specimen; whether such a limited seed production in nature would result in the production of a mature plant is doubtful. In addition to the F2's pollen fertility (92%) being substantially higher than that of the F1, the pollen fertility of the F2 was clearly within the ranges of that of both parents and indeed of all of the other species of Rumex that were examined. The cytology of this hybrid as well as R. crispus X obtusifolius and R. longifolius X obtusifolius will be discussed in the section on the Cytology of the Hybrids.

In summary, the study has confirmed that those plants of Rumex identified as hybrids based on external morphological characteristics commonly produce pollen of a very low fertility. However, as noted in the cases of R. crispus X obtusifolius and R. pseudonatronatus X triangulivalvis, exceptionally high pollen fertility can occur; in

such instances, chromosome counts from meiotic material would indicate the hybrid status. In the case of the majority of hybrids, the typically low pollen fertility would be expected to drastically reduce the number of successful fertilizations and subsequent seed set on a hybrid plant. The low pollen fertility of the hybrids must be considered as a definite factor contributing to the rarity of Rumex hybrids and to the distinctness of the species.

Hybrid Seed Set

As noted earlier, irregular fruit enlargement and low seed set have been considered indicative of Rumex hybrids. Although agreeing that there is generally low seed set in Rumex hybrids, Lousley (1939) noted that, occasionally, plants of R. crispus X obtusifolius produce a large number of well formed achenes. Williams (1971), in investigating germination polymorphism in R. crispus and R. obtusifolius, studied seed germination in a population consisting of both the species and their hybrids and found that the percentage germination of seed from hybrid plants ranged from zero to a level similar to that of the parental species; the latter he interpreted as being due to backcrossing with one of the parents. In contrast with most Rumex hybrids, Lousley and Williams (1975) reported that in Europe, R. longifolius X obtusifolius often appeared fertile as most valves matured and noted that in 1868, Syme claimed it was fertile. Prior to the present study, little detailed information on fruit development, seed set or seed viability of Rumex hybrids existed in the literature.

Procedure

Rootstocks identified as R. crispus X obtusifolius and R. longifolius X obtusifolius were collected in Nova Scotia in 1976 and cultivated in the greenhouse facilities of Agriculture Canada, Ottawa. R. crispus X obtusifolius was represented by two plants (76206a and 76206b) from the same original rootstock and R. longifolius X obtusifolius by three plants (7647, 7649, 76214); the source of

the original material is given in Table 12. The plants were placed in a Cold House (temperature maintained at 4°C) from November 1976 to February 1977 to induce flowering; in March, the rootstocks were repotted and the plants were allowed to flower. Employing the same methods used earlier (Seed Germination and Seed Set, p60), seed set and seed viability was examined. However, for the hybrids, the number of potential fruiting units was determined for the entire bagged portion of the branch and a comparable sized open-pollinated branch. The results are presented in Table 13.

Results

From Table 13, it can be seen that in both hybrids, few fully developed fruits are produced, resulting in the inflorescences having the typically ragged appearance of a Rumex hybrid. Although R. longifolius X obtusifolius produced approximately four to five times the number of fully developed fruit in the open and selfed states respectively than did R. crispus X obtusifolius, the plants still had a ragged appearance as most valves do not mature, contrary to the findings of Lousley and Williams (1975). An even smaller proportion of these fully developed fruit (Table 13, % Full Fruit with Full Seed) actually contained well-developed seed. When selecting seed from plants of various Rumex species (open and bagged samples) (see Table 9), it was found that fruit which appeared externally to be fully developed always contained a well-developed achene. In R. crispus, R. longifolius and R. obtusifolius, the % Full Fruit with Full Seed was around 100% compared with the mean values of 17.8 to 35.1% exhibited by the hybrids between the species.

Table 13 Hybrid Seed Set

The total percentage seed set for R. crispus x obtusifolius and R. longifolius x obtusifolius for cultivated plants of these hybrids is presented.

TABLE 13

HYBRID SEED SET

Rumex crispus x obtusifolius

SPECIMEN NO.	POLLINATION	NO. OF FLOWERS	% FULL FRUIT	% FULL FRUIT WITH FULL SEED	TOTAL % SEED SET	NO. OF SEEDS TESTED	SEED GERMINATION - % VIABILITY
76206a	Open	160	5.0	50.0	2.5	2	100
76206a	Open	300	3.0	33.3	1.0	3	67
76206b	Open	466	4.9	13.0	0.6	2	50
			$\bar{x}=4.6$	$\bar{x}=25.0$	$\bar{x}=1.2$		$\bar{x}=71$
76206a	Self	440	5.7	28.0	1.6	4	75
76206a	Self	245	8.2	30.0	2.4	4	100
76206b	Self	480	2.5	58.3	1.5	4	50
			$\bar{x}=4.7$	$\bar{x}=35.1$	$\bar{x}=1.7$		$\bar{x}=75$

Rumex longifolius x obtusifolius

7647	Open	140	35.7	10.0	1.8	2	50
7647	Open	100	30.0	13.3	4.0	2	100
7647	Open	513	18.3	28.7	4.8	10	90
7647	Open	225	13.3	16.7	2.2	2	100
7649	Open	125	16.0	10.0	1.6	2	100
76214	Open	230	17.4	25.0	1.7		
			$\bar{x}=19.8$	$\bar{x}=17.8$	$\bar{x}=2.5$		$\bar{x}=89$
7647	Self	219	23.3	19.6	4.1	10	90
7649	Self	261	14.1	43.0	10.7	10	90
7649	Self	217	18.9	34.1	6.5	10	100
76214	Self	160	40.0	15.6	6.3	10	40
			$\bar{x}=25.8$	$\bar{x}=28.1$	$\bar{x}=7.2$		$\bar{x}=80$

 $\bar{x}$ = average value

As reported earlier(Seed Germination and Seed Set, p:72), the total percent seed set (No. of Fruit with Full Seed/ No. of Flowers) exhibited by species of Rumex was found to vary from 46.5 to 100%; even the lower limit is still 4.3 times higher than the upper level of 10.7% shown by the hybrids (Table 13). From Table 13, it can be seen that the bagged inflorescences had a somewhat higher seed set than the open ones, but the percentage viability of seed from the bagged samples was lower than that from the open pollinated samples. A comparison of the percentage viabilities with those of the parent species in Table 8 indicates that the seed of the majority of the hybrid samples was equally viable.

Salisbury (1942) records an average production of 30,000 seeds for individual plants of R. crispus; in R. crispus, the percent seed set and percentage viability closely approximate each other, but in the hybrids examined, this is not the case. Using the value of 30,000 as the base number for seed production, the highest total percentage seed set shown by R. crispus X obtusifolius - 2:53 would result in only 750 well developed seeds; similarly, in R. longifolius X obtusifolius, the highest seed set of 10.7% would result in 3,210 well developed seeds, while even the lowest seed set of a Rumex species (46.5%) would result in an average of 13,950 seeds. Since in nature, only a small portion of the viable seeds would be expected to survive, the relatively small quantity of hybrid seed will be further diminished and will make up a very small proportion of the total viable seed in areas where the parents also occur. The production of seed by F1 hybrids selfing will be very unlikely; backcrosses with the parental species are much more likely but none

of the hybrid sites examined in Nova Scotia showed evidence of successful backcrossing with either parent. Introgression, were it to occur, would tend to make species delimitation more difficult. In Britain, Lousley (1939) believed that such secondary crosses were comparatively rare since he had been unable to find any locality with a series of intermediates connecting two parent species and also if the secondary crosses were more common, the difficulties in recognizing the species would be increased. However, Williams (1971) has studied several British populations of R. crispus, R. obtusifolius and their hybrids in which by means of a hybrid index, introgression has been indicated to occur; introgression in the direction of R. crispus was more common than towards R. obtusifolius.

The fact that with such low pollen fertility (Table 12), any viable seed at all is produced by the hybrids is of interest. As observed in this study, the percentage seed set of a hybrid is typically very low while the viability of the seed produced is high (based on testing only full seed); simply noting the seed viability for a Rumex hybrid would be clearly misleading unless a clear indication is made as to the selection of the seed to be tested. Although a low seed set or a low pollen fertility clearly indicates plants of hybrid origin, a close examination of morphological and cytological information may be required to confirm a plants hybrid status if introgression is suspected.

Cytology of Hybrids

Cytological studies can be very valuable in indicating the hybrid status of a plant by means of the chromosome number or by observation of meiotic irregularities. Jensen (1936) in examining metaphase in microsporogenesis of R. crispus X obtusifolius, found 15 bivalents and 10 univalents with meiosis proceeding very irregularly in the few dyads formed, rarely was there the exact diploid number of 50. Lousley and Williams (1975) reported counts in R. crispus X obtusifolius of $2n = 44 - 56$. A report by G.A. Mulligan of $2n = 50$ appears on the label of a specimen from London, Ontario, collected by Dr. P. Cavers; this was confirmed to be R. crispus X obtusifolius.

Boivin (1968) reported a specimen collected in Saskatchewan (C. Frankton 1543) to be R. crispus X occidentalis, in fact it is R. occidentalis ($2n = 140$) X pseudonatronatus ($2n = 40$). Mitotic counts of $2n =$ ca. 112, ca. 114, ca. 122, from root tips of seedlings representing the F2 generation of the hybrid were reported by G. A. Mulligan on the label of the specimen of the hybrid.

Mulligan (1959) reported a mitotic count of $2n = 30$ in a hybrid between R. pseudonatronatus ($2n = 40$, section Rumex) and R. triangulivalvis ($2n = 20$, section Axillares). Voucher specimens (Frankton 1635 and Mulligan 2239, 2398 at DAO) were examined and their identification confirmed. On Mulligan 2398, the collector

indicated that meiosis in the F1 was very irregular and that the probable genome was T P1 P2 (parental genomes would be R. triangulivalvis - TT and R. pseudonatronatus - P1 P1 P2 R2). Further study on the hybrid was described by Mulligan (1965). The only two seeds found on the F1 plant were germinated and the resultant plants examined cytologically with the following results:

<u>Voucher Specimen</u>	<u>Mitotic Count</u>	<u>Metaphase I Meiotic Count</u>	<u>Probable Genome</u>
Mulligan 2399	2n = 5Q	10 univalents 20 bivalents	T P1 P1 P2 P2
Mulligan 2460	2n = 6Q	30 bivalents	T T P1 P1 P2 P2

The seeds must therefore have been the product of unreduced gametes with Mulligan 2460 representing an amphiploid. Although the plant represented a potential new species, Mulligan noted that with only two viable seeds formed, the chances of such a plant being produced and surviving under field conditions would be very remote. However, he felt that "if two barely crossable species have the capacity to colonize newly disturbed areas, this inherent capacity can apparently increase the opportunities for these species to be involved in abrupt speciation".

An investigation of the cytology of the hybrids was carried out to determine, (a) the regularity of meiosis and (b) the correlation between the details of chromosome behavior observed at meiosis and other features of the hybrid such as pollen fertility and seed set.

Procedure

Buds of cultivated plants of R. crispus X obtusifolius and R. longifolius X obtusifolius originating in Nova Scotia were collected and prepared following the method reported earlier (Cytology, p. 8). Two plants of R. crispus X obtusifolius (76206a and 76206b) from the same original rootstock were used as the source of buds and four plants of R. longifolius X obtusifolius (7647, 7648, 7653a, 7653b - the latter two from the same original rootstock). Even though numerous collections of buds were made, it was very difficult to obtain any at the metaphase stage of microsporogenesis because in any one whorl, only a few flowers were found to contain anthers and hence, it was difficult to determine the bud size that indicated meiotic activity.

Results

Preliminary observations of the cells undergoing meiosis indicated that in the plants of both hybrids, meiosis appeared to be very irregular. In R. longifolius X obtusifolius, a large number of univalents and multivalents were evident and only a few normal bivalents, whereas in R. crispus X obtusifolius, the meiotic cells were past the stages in which the chromosomes could be accurately interpreted and counted. However, counts of $2n = 50$ from mitosis in somatic tapetal cells were obtained; one count was from 76206a and two from 76206b. $2n = 50$ is the expected number for the F1 hybrid between R. crispus ($2n = 60$) and R. obtusifolius ($2n = 40$).

In R. longifolius X obtusifolius, meiosis was clearly irregular in the meiotic material that was examined. To illustrate

the irregularities observed, several cells were interpreted at metaphase I.

Specimen <u>7847</u>	<u>Univalents</u>	<u>Bivalents</u>	<u>Trivalents</u>	<u>Quadrivalents</u>	<u>Pentavalents</u>
Cell 1	8	2	3	1	
Cell 2	4	7	1	1	
Cell 3	3	5	3		1 (a chain

Specimen <u>7849</u>	<u>Univalents</u>	<u>Bivalents</u>	<u>Trivalents</u>	<u>Quadrivalents</u>	<u>Pentavalents</u>
Cell 1	4	6	3		
Cell 2	5	4	4		
Cell 3	13	3	2		

Specimen <u>7853</u>	<u>Univalents</u>	<u>Bivalents</u>	<u>Trivalents</u>	<u>Quadrivalents</u>	<u>Pentavalents</u>
Cell 1	8	3	1	2	
Cell 2	6	6	1	1	
Cell 3	2	5	3	1	

The diploid number in all cases would be 50. Examples of these irregularities are illustrated in Figure 14. As in the previous hybrid, $2n = 50$ is the expected number based on the chromosome counts of the parents - *R. longifolius* ($2n = 60$) and *R. obtusifolius* ($2n = 40$).

The meiotic irregularities observed in these plants support their identification as hybrids. Since meiosis does not proceed normally as it has been shown to do in the species (Cytology, p. 79), pollen fertility and seed set will obviously be adversely affected; from the previous tables (Tables 11 and 12), it can be observed that

Figure 14 Meiotic Irregularities
in R. longifolius x obtusifolius
Top No. 7647 Cell 3 (See page 113)
Bottom No. 7653 Cell 2
Magnification 3400

pollen viability and seed set are very low for the plants in which meiosis was examined. It would appear that at least in the Rumex hybrids examined, a major factor preventing or at least severely limiting their widespread occurrence exists at the chromosomal level. If meiotic material had been available, it would have been of interest to observe meiosis in hybrids between species of the same ploidy level, e.g. R. crispus X patientia, since at least in the cases examined, pollen viability was also very low, indicating at least the probability of meiotic irregularities in these hybrids also.

From the available data on hybrid cytology, it is clear that in most cases, the observation of meiotic irregularities and abnormal chromosome numbers is probably not only indicative of a plant of hybrid origin, but responsible for the low pollen viability and low seed set reported earlier. This is supported by the fact that the amphiphloid of R. pseudonatronatus X triangulivalvis with $2n = 60$, also had a high pollen fertility (92%, see Table 11). The meiotic irregularities would also make most backcrosses with the parents or other species relatively unsuccessful unless the chance production of an unreduced gamete occurred.

Morphological Characteristics of Rumex Hybrids

In addition to the high degree of pollen sterility and the irregular development of the fruit noted earlier (Hybrids, p.91) being indicative of Rumex hybrids, an examination of the morphological characteristics of the suspected hybrids will also help identify a hybrid. Often, the parent species involved in the formation of the hybrid can be deduced as the hybrids tend to exhibit features common to the parents, e.g. "most hybrids of R. obtusifolius inherit something of its cordate basal leaves and toothed and acuminate valves" while hybrids of R. pulcher "usually show something of the divaricate branching, remote whorls, short pedicels and thick venation of the valves" (Rechinger 1964). To allow for easier recognition of the hybrids, the present study has compared the main morphological characteristics of the hybrids and the parent species.

Procedure

A comparison of the following characteristics of the hybrids and parent species was made:

Basal Leaves - Shape, L:W, Apex, Base, Angle of Veins, Vestiture (on abaxial veins); Inflorescence - Angle of Branches, Subtention of Branches and Whorls; Pedicel - Size, Type of Articulation and its Location; Valve - Shape, Size, Apex, Base, No. of Teeth/side, No. of Anastomoses/side; No. of Tubercles; Large Tubercles - Shape, Size, Apex; Small Tubercle - Shape, Size; Achene - Size,

Apex, Base, Colour. In the case of the parent species, the observations are based on the specimens employed in producing the species descriptions whereas for the hybrids the observations are based on the hybrid specimens found in the herbarium loans and in the case of R. crispus x obtusifolius, material collected in the field studies made during the present study were also used. For the hybrid R. longifolius x obtusifolius which was found during this study, the observations are based entirely on the material collected during the Nova Scotian field studies.

Binomials will not be used for the Rumex hybrids as the application of such a name implies that the plants are consistent and persistent in nature which need not apply to hybrid plants. Synonymy will be given only for hybrids named in North America and references will be given for the synonymy of the others.

Results

The hybrid R. britannica x crispus, involving the native North American species, R. britannica was given the binomial R. X dissimilis by Rechinger (1937). In Table 14-No. 1, it can be seen that the hybrid more closely resembles R. britannica than R. crispus. The influence of R. crispus is most apparent in the distinctly swollen articulation of the pedicel, the wider and more ovate tubercles and the harder seed coat. The irregular fruit development typical of the hybrids is apparent in that only the large-valved fruit indicated in Table 14.1 were found to contain fully developed achenes while the smaller valves representing the

majority of the fruit did not; the smaller valved fruit did not have distinctly swollen articulations on their pedicels. The somewhat longer spines on the valves of the hybrids might suggest the involvement of R. obtusifolius rather than R. crispus; however, none of the other characteristics typical of R. obtusifolius hybrids are present.

R. conglomeratus x pulcher (Table 14-No. 2) more closely resembles the latter parent. The influence of R. conglomeratus is evident in the narrower valve of the hybrid, the somewhat entire valve margin, the size of the small tubercles and in the longer pedicels. Two additional characteristics indicative of R. conglomeratus not listed in the table are the smoother surface of the tubercles as R. pulcher's are bumpy and the achenes protruding beyond the edge of the valve while in R. pulcher the achene is totally covered; further characteristic indicating R. pulcher is the distinctly raised veins of the valves while in R. conglomeratus, they are barely evident. The valve of this hybrid is illustrated in Figure 15-a. Although producing a high proportion of small fruit with poorly developed achenes, many whorls appeared to contain at least one fruit with fully developed achenes. The characteristics observed for this hybrid in North America agree with those given by Lousley and Williams (1975) in Britain where they report the hybrid to occur frequently when the parents occur together. The synonymy for R. conglomeratus x pulcher is dealt with by Ascherson and Graebner (1908), Rechinger (1932) and Lousley and Williams (1975).

R. crispus x kernerii more closely resembles R. kernerii in overall appearance (Table - no. 3). R. crispus appears to have brought about a reduction in the size of the valve, tubercles and achenes as well as an ovate valve, more typical of R. crispus than R. kernerii. A valve of this hybrid is illustrated in Figure 15b. Most fruit produced by this hybrid is smaller than the values listed in the table and did not contain fully developed achenes; this characteristic tends to make the larger fruit more conspicuous. In the small-valved fruit, the three valves making up the fruit appear to be somewhat mismatched as they are of different sizes. The synonymy for this hybrid can be found in Rechinger's (1933) study of the section in central Europe where both species are native.

As noted by Lousley and Williams (1975), R. crispus x longifolius resembles the latter parent more than R. crispus. The influence of R. crispus is indicated by the narrower basal leaves, by the valves usually possessing three tubercles which are larger than the tubercle of R. longifolius and by the smaller red-brown achenes. Superficially, most of the fruit appears to be fully developed but the smaller valved fruit have mismatched valves and sterile achenes. In this hybrid, few of the larger valved fruit indicated in the table contained fully developed achenes. A typical valve is illustrated in Figure 15c. The works of Ascherson and Graebner (1908) and Lousley and Williams (1975) give the synonymy of R. crispus x longifolius.

The hybrid, R. crispus x obtusifolius is the most common hybrid in the section and morphologically is intermediate in most characteristics (Table 14-No. 5). Only the large fully developed fruit contained fully developed achenes (Figure 15d), but even many of these contained empty achenes; the majority of the fruit produced by this hybrid are very small, poorly developed and with sterile achenes. In Europe, Williams (1971) carried out a detailed study of several populations consisting of plants of the hybrid and the parents; he observed the occurrence of introgression with either parent making identification difficult. No cases of introgression have been recognized in this study. The synonymy of R. crispus is dealt with by Ascherson and Graebner (1908), Rechinger (1937) and Lousley and Williams (1975).

As observed by Rechinger (1964) and Lousley and Williams (1975), the hybrid between R. crispus and R. patientia more closely resembles the latter parent - (Table 14-No. 6). The influence of R. crispus is evident in the type of vestiture on the basal leaves, the shape of the base of the basal leaves, the reduced valve size and the shape of the tubercles. The hybrid stands out because of the large number of fruit that are smaller than those indicated on the table; the valves of the small fruit are mismatched in size and the small fruit contains poorly developed achenes as does many of the larger fruit. The synonymy of this hybrid can be found in Ascherson and Graebner (1908), Rechinger (1933) and Lousley and Williams (1975).

R. crispus x pulcher, as noted by Lousley and Williams (1975) reflects the characteristics of the latter parent more so than R. crispus (Table 14-No. 7). The involvement of R. crispus is indicated by the lack of subtention of most whorls, the longer pedicels, the tendency to have cordate valves, the shorter teeth; the tubercles are also smoother than those found in R. pulcher, reflecting the smooth tubercles of R. crispus. None of the fruit of the specimens of this hybrid were fully developed or produced fully developed achenes. Synonymy for this hybrid can be found in Ascherson and Graebner (1908), Rechinger (1932) and Lousley and Williams (1975).

R. longifolius x obtusifolius (Table 14-No.8), prior to this study had not been reported for North America. The North American specimens agree closely with the observations made by Lousley and Williams (1975). The basal leaves more closely resemble those of R. longifolius but they have the vestiture of R. obtusifolius. The inflorescence of the hybrid was intermediate in nature as it was denser than R. obtusifolius but not as dense as R. longifolius. As indicated in the table, the valve of the hybrid reflects characteristics of both parent species whereas the achenes were closer in appearance to R. obtusifolius. (Figure 15e) As in most Rumex hybrids, the majority of fruit were small, abnormally developed and containing poorly developed achenes. The small number of achenes produced were found in the larger valves indicated in the table. Synonymy for this hybrid can be found in Ascherson and Graebner (1908) and Lousley and Williams (1975).

In general, the hybrid R. maritimus var. fueginus x stenophyllus

more closely resembles the former parent. (Table 14-no.9). The influence of R. stenophyllus is apparent in the wider valves and the greater number of teeth on the margin of the valves. No full developed achenes were evident on the only available specimen of this hybrid. This hybrid did not exhibit the same discrepancy in fruit sizes that is often observed in Rumex hybrids. In describing this hybrid, Boivin (1966) gave it the binomial R. X Alexidis.

R. occidentalis x pseudonatronatus, more closely resembles the former parent than R. pseudonatronatus (Table 14-No.10). The fruit tends to show the influence of R. pseudonatronatus in the valve shape, the presence of a tubercle and the distinctly swollen articulation on the pedicel. The majority of the fruit are smaller than those indicated in the table and have mismatched valves. In the only available specimen, few fully developed achenes were produced in the larger fruit and none in the smaller ones. This specimen had been previously identified as R. crispus x occidentalis but R. pseudonatronatus represents a better choice as the putative parent than R. crispus because of its orbicular valves and its single oblong tubercle.

The hybrid R. pseudonatronatus x triangulivalvis represents the only known hybrid between members of section Rumex and section Axillares. Its basal leaves more closely resemble R. pseudonatronatus than R. triangulivalvis while the habit of the plant reflects the influence of R. triangulivalvis. The fruit characteristics of the hybrid are somewhat intermediate in nature with respect to the parent species. The hybrid fruit can be distinguished from R. pseudo-

natronatus by its ovate valves, less cordate base, by the greater number of tubercles, by the shape and size of the tubercles; it can be distinguished from R. triangulivalvis by its ovate valves, the smaller ovate tubercles and the larger achenes. In describing this hybrid, Boivin (1966) applied the binomial R. X Franktonis. (Table 14.10) From the single field specimen of this hybrid, only two fully developed achenes were collected. These were grown on by Mulligan (Cytology, of Hybrids P. 111) and achenes from the resulting plants were also grown on. These plants could be easily distinguished from the parent species on the basis of their chromosome numbers but the fruit characteristics now more closely resemble R. pseudonatronatus; provided that the fruit still possessed three tubercles, they could still be distinguished morphologically from R. pseudonatronatus but two specimens (Mulligan 2558, 2559), clearly resembled R. pseudonatronatus. All of the plants produced from the original two achenes had inflorescences with normally developing fruit and fully developed achenes. Fortunately, these plants represent cultivated specimens and did not occur in the field.

A comparison of a suspected hybrid with the characteristics of the putative parents helps to indicate whether it is a hybrid and whether or not the suspected parents are involved. If examples of introgression similar to that described by Williams (1971) are found in North America, then the more detailed approach of setting up a hybrid index to assess the suspected hybrids would be required. Further cultivated studies are needed on Rumex hybrids in order to determine the nature of introgression and its possible effects on species delimitation.

Table 14. Comparison of Hybrids with Parent Species
No. 1 R. britannica x criscus
(continued on p. 125)

No. 1

CHARACTERISTICS

Basal Leaves
Shape

L:W
Apex
Base
Angle of Veins
Venititure on
Axerial Veins

R. britannica

Lanceolate

3-4:1
Acute
Cuneate to obtuse
650-800
Glabrous to indi-
stinctly papillose,
trichomes papilli-
form, less than
0.1 mm. long

Hybrid

Lanceolate

2.6-2.8:1
Acute
Obtuse to truncate
600-800
Indistinctly papil-
lose, trichomes papil-
liform, less than
0.1 mm. long

R. crispus

Elliptic, oblong
or linear - lanceolate
4-6:1
Acute
Cuneate to obtuse
500-700(-800)
Indistinctly papil-
lose, trichomes papil-
liform, less than
0.1 mm. long

Inflorescence

Angle of Branches
Subtention of
Branches and whorls

(200-)300-500(550)
Only lower branches
subtended, no whorls
subtended

300-400

Only lower branches
subtended, no whorls
subtended

200-400

All branches sub-
tended by leaves or
bracts, occasionally
lowermost whorls sub-
tended by bracts

Pedicel
Size

Usually longer than
valve
Not swollen, in the
lower third

As in R. britannicaAs in R. britannica

Distinctly swollen,
in the lower two-
fifths

Distinctly swollen,
in the lower third,
only evident on
pedicels of large
fruit

Table 14

Comparison of Hybrids with Parent Species

No. 1 continued

CHARACTERISTICS

R. britannica

Hybrid

R. crispus

Valve

Shape
Size (mm.)

Orbicular
{3.3-4.5-6.0(-7.5)}x
{2.8-4.0-6.9(-7.2)}

Orbicular
7.0-7.8x6.5-8.5

Ovate
{2.8-3.5-5.0(-6.0)}x
{2.5-3.5-5.5(-7.5)}

Apex

Base

Obtuse to subacute
Truncate to cordate,
indented 0.1-0.7 mm.

Obtuse to subacute
Cordate, indented
0.9-1.2 mm.

Cordate, indented
0.2-0.5(-1.0)mm.

No. of Teeth/side

Length of Teeth (mm.)

1-9
Usually less than
0.3(-0.4)

2-7
Usually less than
0.5(-0.7)

1-8 to entire
Less than 0.3

No. of Anastomoses/
side

1-2

(1-2)(-3)

(1-12)(-3)

No. of Tubercles

3, equal in size

3, equal in size

1-3, usually 3,
unequal in size

Large Tubercle

Shape
Size (mm.)

Fusiform
{1.6-2.5-3.5(-4.8)}x
{0.6-0.8-1.2(-1.4)}

Fusiform to ovate
3.6-4.3x1.3-1.4

Ovate
{1.4-1.8-3.0(-3.3)}x
{0.8-1.0-1.5(-2.0)}

Apex

Small Tubercle

Shape

Size (mm.)

Acute

Acute

Obtuse to acute

Obtuse to acute

Ovate

Size (mm.)

1.0-2.0(-2.5)x
0.5-1.2

Achene

Size (mm.)

{2.8-3.2-4.0(-4.5)}x
{1.5-1.7-2.3(-2.7)}

3.1-3.4x1.6-1.8

{2.0-2.2-2.8(-3.0)}x
{1.2-1.4-1.6(-2.0)}

Apex

Base

Acute

Acute

Acute

Acute

Acute

More broadly acute than apex
to somewhat obtuse

Seed Coat

Easily removed

Not easily removed

Not easily removed

Table 14. Comparison of Hybrids with Parent Species
No. 2 R. conglomeratus x pulcher
(continued on p. 127)

No. 2

CHARACTERISTICS

	<u>R. conglomeratus</u>	Hybrid	<u>R. pulcher</u>
Basal Leaves			
Shape	Oblong to oblong-lanceolate	None on specimens	As in <u>R. conglomeratus</u>
L:W	4-5:1		2-3:1
Apex	Obtuse to acute		Subacute
Base	Cuneate		Broadly cuneate, truncate r weakly cordate
Angle of Veins			500-750
Vestiture	500-700 Glabrous to indistinctly papillose, trichomes papiliform, less than 0.1 mm. long	(stem leaves showed vestiture of <u>R. conglomeratus</u>)	Glabrous to distinctly pubescent, trichomes papiliform to cylindrical, less than 0.2 mm.
Inflorescence			
Angle of Branches		500-700	(400-)500-700(-800)
Substitution of Branches and Whorls	300-600 All branches subtended by leaves or bracts, all but very upper whorls subtended by bracts	As in <u>R. conglomeratus</u>	As in <u>R. conglomeratus</u>
Pedical			
Size	Usually longer than valve	Shorter to longer than valve	Shorter to as long as valve
Type of Articulation and its Location	Distinctly swollen, in the lower two-fifths	As in <u>R. conglomeratus</u>	As in <u>R. conglomeratus</u>
Shape	Piliform	Uniformly thickened	Uniformly thickened

No. 2 continued

CHARACTERISTICS

	<u>R. conglomeratus</u>	Hybrid	<u>R. pulcher</u>
Valve Shape	Ovate to oblong	Deltoid	Ovate-deltoid to oblong-deltoid
Size (mm.)	{1.8-}2.3-2.9(-3.0)x {1.0-}1.2-1.5(-1.8)	3.0-4.3x2-2.5	{2.7}3.5-4.8(-5.5)x 1.8-3.0(-3.5)
Apex	Obtuse	Acute	Acute
Base	Obtuse	Cuneate	Truncate
No. of Teeth/side	Entire to rarely 1-2	Entire to 1-3	{0-}5-9(-11)
Length of Teeth (mm.)	Less than 0.1	Less than 0.3	{0.2-}1.0-1.6(-2.0)
No. of Anastomoses/side	1(-2)	2	1-2(-3)
No. of Tubercles	3, equal in size	3, unequal in size	1-3, unequal in size
Large Tubercle	Ovate to oblong	As in <u>R. conglomeratus</u>	Ovate
Shape	{0.9-}1.3-1.8(-2.0)x {0.6-}0.8-1.0(-1.1)	2-2.3x1-1.2	{1.5-}1.8-2.5(-3.0)x {0.5-}0.8-1.3(-1.7)
Size (mm.)	Obtuse	Obtuse	Acute
Apex		Ovate to oblong	Ovate
Small Tubercle			1.0-1.5x0.5-0.6
Shape			
Size (mm.)			
Achene			
Size (mm.)	1.3-}1.5-1.9(-2.1)x {1.0-}1.2-1.4	2.0-2.3x1.4-1.5	{1.8-}2.2-2.8x {1.4-}1.5-2.0
Apex	Acute	Acute	Acute
Base	Obtuse	Obtuse	Obtuse to almost truncate

Table 14 Comparison of Hybrids with Parent Species
No.3 R. crispus x kernerii
(continued on page 129)

CHARACTERISTICS		<u>R. crispus</u>	Hybrid	<u>R. kernerii</u>
Basal Leaves				
Shape		Elliptic, oblong or linear lanceolate	Oblong	Oblong-lanceolate
L:W		4-6:1	2.8:1	2-3:1
Apex		Acute	Broadly acute	Broadly acute
Base		Cuneate to obtuse	Cuneate	Obtuse to weakly cordate
Angle of Veins		50°-70°(-80°)	60°-70°	500-700
Vestiture		Indistinctly papillose, trichomes papilliform, less than 0.1 mm. long	Indistinctly papillose, trichomes papilliform, less than 0.2 mm. long	Glabrous to indistinctly papillose to pubescent trichomes papilliform or cylindrical, less than 0.3 mm. long
Inflorescence				
Angle of Branches		200-400	300-450	300-500
Subtention of Branches and Whorls		All branches subtended by leaves or bracts, occasionally lowermost whorls subtended by bracts	All branches subtended by leaves or bracts, no whorls subtended	Same as <u>R. crispus</u>
Pedical				
Size		Usually longer than valve	Same as <u>R. crispus</u>	Same as <u>R. crispus</u>
Type of Articulation and its Location		Distinctly swollen, in the lower two-fifths	Same as <u>R. crispus</u>	Distinctly swollen, at or near the middle

No. 3 continued			
CHARACTERISTICS		<u>R. crispus</u>	<u>R. kernerl</u>
Valve Shape		Hybrid	
Size (mm.)		Ovate	Approximately orbicular
Apex		{2.8-}3.5-5.0(-6.0)x	(6.0-}6.5-9.0x
Base		{2.5-}3.5-5.5(-7.5)	(6.0-}6.5-9.0
		Subacute indented	Abruptly acute
		Cordate indented	Cordate, indented
		0.2-0.5(-1.0) mm.	0.8-1.3 mm.
No. of Teeth/side		1-8	3-8
Length of Teeth (mm.)		Less than 0.3	0.3-1.0
No. of Anastomoses/side		(1-}2(-3)	3-4(-5)
No. of Tubercles		1-3, usually 3, unequal in size	3, unequal in size
Large Tubercle Shape		Ovate	Oblong-ovate
Size (mm.)		{1.4-}1.8-3.0(-3.3)x	2.5-5.0x
		{0.8-}1.0-1.5(-2.0)	(0.5-}1.0-2.5
Apex		Obtuse to acute	Obtuse to acute
Small Tubercle Shape		Ovate	Oblong-ovate
Size (mm.)		1.0-2.0(-2.5)x	1.5-4.0x
		0.5-1.2	(0.5-}1.0-2.0
Achene Size (mm.)		{2.0-}2.2-2.8(-3.0)x	3.0-3.5(-3.8)x
		{1.2-}1.4-1.6(-2.0)	(1.8-}2.1-2.7(-2.8)
Apex		Acute	Acute
Base		More broadly acute than apex to somewhat obtuse	Obtuse

Table 14. Comparison of Hybrids with Parent Species
No. 4 R. crispus x longifolius
(continued on page 131)

No. 4	CHARACTERISTICS	<u>R. crispus</u>	Hybrid	<u>R. longifolius</u>
Basal Leaves				
Shape		Elliptic, oblong or linear-lanceolate	Oblong-lanceolate	Lanceolate to oblong-lanceolate
L:W		4-6:1	4.2-5.5:1	2-4:1
Apex		Acute	Acute	Acute to subacute
Base		Cuneate to obtuse	Cuneate	Broadly cuneate, obtuse to weakly cordate
Angle of Veins		50°-70° (-80°)	60°-70°	(40°-)50°-70°
Vestiture		Indistinctly papillose, trichomes papilliform, less than 0.1 mm. long	As in <u>R. crispus</u>	Indistinctly to moderately papillose, trichomes papilliform, less than 0.1 mm. long
Inflorescence				
Angle of Branches		20°-40°	20°-50°	25°-40°
Subtention of Branches and Whorls		All branches subtended by leaves or bracts, occasionally lowermost whorls subtended by bracts	As in <u>R. crispus</u>	As in <u>R. crispus</u>
Pedical				
Size		Usually longer than valve	As in <u>R. crispus</u>	As in <u>R. crispus</u>
Type of Articulation and its Location		Distinctly swollen, in the lower two-fifths	As in <u>R. crispus</u>	As in <u>R. crispus</u>

No. 4 continued			
CHARACTERISTICS		<u>R. crispus</u>	Hybrid
Valve			<u>R. longifolius</u>
Shape			
Size (mm.)		Ovate {2.8-3.5-5.0(-6.0)}x {2.5-3.5-5.5(-7.5)}	Orbicular to reniform {3.5-5.2-6.5(-7.5)}x {4.0-6.0-7.5(-9.5)}
Apex		Subacute	Obtuse
Base		Cordate, indented	Cordate, indented
		0.2-0.5(-1.0) mm.	(0.3-1.0-1.5 mm.
No. of Teeth/side		Entire to 1-8	Entire to 2-6
Length of Teeth (mm.)		Less than 0.3	Less than 0.3
No. of Anastomoses/ side		{1-12(-3)}	2-3
No. of Tubercles		1-3, usually 3, unequal in size	0-1
Large Tubercle			
Shape			
Size (mm.)		Ovate {1.4-1.8-3.0(-3.3)}x {0.8-1.0-1.5(-2.0)}	Oblong {0.3-0.5-1.3(-1.4)}x {0.3-0.4-1.0}
Apex		Obtuse to acute	Obtuse
Small Tubercle			
Shape			
Size (mm.)		Ovate 1.0-2.0(-2.5)x 0.5-1.2	
Achene			
Size (mm.)		{2.0-2.2-2.8(-3.0)}x {1.2-1.4-1.6(-2.0)}	{2.5-2.8-3.3(-3.5)}x {1.3-1.5-1.8(-2.0)}
Apex		Acute	Acute
Base		More broadly acute than apex to some- what obtuse	As in hybrid and <u>R. crispus</u>
Colour		Red-brown	Brown

Table 14. Comparison of Hybrids with Parent Species
No. 5 R. crispus x obtusifolius
(continued on p. 133)

No. 5	CHARACTERISTICS	<u>R. crispus</u>	Hybrid	<u>R. obtusifolius</u>
Basal Leaves Shape	Elliptic, oblong or linear - lanceo- late 4-6:1 Acute Cuneate to obtuse 50°-70° (-80°)	Oblong-lanceolate	Oblong-ovate	
L:W	4-6:1	2.2-4.5:1	1.7-2.5	
Apex	Acute	Acute	Acute	
Base	Cuneate to obtuse	Obtuse to truncate	Distinctly cordate	
Angle of Veins	Indistinctly papil- lose, trichomes	Distinctly pubes- cent to papillose,	45°-80°	
Vestiture	lose, trichomes papilliform, less than 0.1 mm. long	Distinctly pubes- cent to papillose, trichomes cylin- drical, less than 0.1 mm. long	Distinctly pubescent, trichomes cylin- drical, less than 0.3 mm. long	
Inflorescence Angle of Branches Subtension of Branches and Whorls	20°-40° All branches sub- tended by leaves or bracts, occasion- ally lowermost whorls subtended by bracts	30°-55° As in <u>R. crispus</u>	(20°-)25°-40° As in <u>R. crispus</u>	
Pedicel Size	Usually longer than valve	As in <u>R. crispus</u>	As in <u>R. crispus</u>	
Type of Articula- tion and its Location	Distinctly swollen, in the lower two- fifths	As in <u>R. crispus</u>	As in <u>R. crispus</u>	

No. 5 continued		<u>R. crispus</u>		<u>R. obtusifolius</u>	
CHARACTERISTICS		Hybrid			
Valve Shape Size (mm.)		Ovate {2.8-}3.5-5.0(-6.0)x {2.5-}3.5-5.5(-7.5)	Ovate to reniform 4.0-5.5x4.0-6.0	Deltoid {3.0-}3.5-5.0(-6.0)x {2.0-}2.5-3.5(-4.0)	
Apex Base		Subacute Cordate, indented 0.2-0.5(-1.0) mm.	Acute Cordate, indented 0.3-0.8 mm.	Acute Truncate to weakly cordate, indented to 0.5 mm. {1-}3-4(-5) {0.8-}1.0-1.8(-2.0)	
No. of Teeth/side Length of Teeth (mm.) No. of Anastomoses/ side		1-8 to entire Less than 0.3 (1-}2(-3)	5-8 0.3-1.0 2	1-2	
No. of Tubercles		1-3, usually 3, unequal in size	3	1-3, usually 1	
Large Tubercle Shape Size (mm.)		Ovate {1.4-}1.8-3.0(-3.3)x {0.8-}1.0-1.5(-2.0)	Ovate 1.5-2.8x1.0-1.8 Obtuse	Ovate {1.1-}1.5-2.5(2.9)x {0.4-}0.7-1.5(-1.8) Obtuse to acute	
Apex Small Tubercle Shape Size (mm.)		Ovate 1.0-2.0(-2.5)x 0.5-1.2	Ovate 0.8-2.0x0.5-1.0	Ovate- 0.5-1.5x0.3-0.5	
Achene Size (mm.)		{2.0-}2.2-2.8(-3.0)x 1.2-}1.4-1.6(-2.0)	1.5-2.8x1.0-1.8	{1.8-}2.3-2.9(-3.0)x {1.2-}1.3-1.7(-1.8)	
Apex Base		Acute More broadly acute than apex to some- what obtuse	Acute More broadly acute than apex to some- what obtuse	Acute More broadly acute than apex	

Table 14. Comparison of Hybrids with Parent Species
No. 6 R. crispus x patientia
(continued on p. 135)

No. 6

CHARACTERISTICS

R. crispus

Hybrid

R. patientia

Basal Leaves

Shape

Elliptic, oblong
or linear - lanceo-
late
4-6:1

Oblong-lanceolate

L.W.

Apex

Base

Angle of Veins

Vestiture

3.2-5.7:1

Acute

Cuneate

700

As in R. crispus

3-4:1

Acute

Obtuse

500-700

Usually glabrous to
indistinctly papil-
lose, trichomes
papilliform, less
than 0.1 mm. long

Inflorescence

Angle of Branches

Subtention of

Branches and Whorls

200-400

All branches sub-
tended by leaves or
bracts, occasionally
flowermost whorls sub-
tended by bracts

200-300

As in R. crispus

(200-300-400

As in R. crispus

Pedicel

Size

Usually longer than
valve

As in R. crispus

Type of Articulation

and its Location

Distinctly swollen,
in the lower two-
fifths

As in R. crispus

No. 6 continued

CHARACTERISTICS

	<u>R. crispus</u>	Hybrid	<u>R. patientia</u>
Valve Shape	Ovate	Approximately orbicular	Orbicular to reniform
Size (mm.)	(2.8-3.5-5.0(-6.0))x (2.5-3.5-5.5(-7.5))	5-7x6.0-8.0	(4.0-6.0-8.5x (4.5-6.0-9.0)
Apex	Subacute		Obtuse
Base	Cordate, indented		Cordate, indented
No. of Teeth/side	0.2-0.5(-1.0) mm.		(0.5-0.8-1.5(-1.8)
Length of Teeth (mm.)	Entire or 1-8	Entire or 1-4	Entire or 3-11
No. of Anastomoses/side	Less than 0.3	Less than 0.4	Less than 0.3
	(1-2(-3))		2-3
No. of Tubercles	1-3, usually 3, unequal in size	3, unequal in size	1-3, unequal in size
Large Tubercle			
Shape	Ovate	Same as <u>R. crispus</u>	Oblong-ovate
Size (mm.)	(1.4-1.8-3.0(-3.3))x (0.8-1.0-1.5(-2.0))	2.0-2.8x1.2-1.8	(1.1-1.6-2.8(-3.5))x (0.5-1.0-1.5(-1.8))
Apex	Obtuse to acute	Obtuse	Obtuse to acute
Small Tubercle			
Shape	Ovate	Ovate	Ovate
Size (mm.)	1.0-2.0(-2.5)x	0.9-1.8x0.4-1.0	0.8-2.0x0.3-0.9
Apex	0.5-1.2		
Base	(2.0-2.2-2.8(-3.0))x (1.2-1.4-1.6(-2.0))	3.3-3.5x1.8-2.0	2.9-3.8x(1.4)1.8-2.3(-2.5)
No. of Teeth/side	More broadly acute than apex to somewhat obtuse	Acute	Acute
Length of Teeth (mm.)		More broadly acute than apex	More broadly acute than apex to somewhat obtuse

Table 14. Comparison of Hybrids with Parent Species
No. 7 R. crispus x rulcher
(continued on p. 137)

No. 7

CHARACTERISTICS

Basal Leaves
Shape

L:W
Apex
Base

Angle of Veins
Vestiture

Inflorescence
Angle of Branches
Substitution of
Branches and Whorls

Pedicel
Size

Type of Articula-
tion and its
Location
Shape

R. crispus

Elliptic, oblong
or linear-lanceo-
late
4-6:1
Acute
Cuneate to obtuse

50°-70°(-80°)
Indistinctly papil-
lose, trichomes
papilliform, less
than 0.1 mm. long

20°-40°
All branches sub-
tended by leaves
or bracts, occasion-
ally lowermost whorls
subtended by bracts

Usually longer than
Valve
Distinctly swollen,
in the lower two-
fifths
Filiform

Hybrid

Oblong

3.4-4.9:1
Acute
Cuneate

40°-70°
Moderately to dis-
tinctly pubescent
to papillose, tri-
chomes papilliform
to cylindrical,
less than 0.1 mm.
long

(30°-)40°-50°
As in R. crispus

As in R. crispus

Somewhat uniformly
thickened

R. pulcher

Oblong to oblong-
lanceolate

2-3:1
Subacute
Broadly cuneate,
truncate or weakly
cordate
50°-75°

Glabrous to distinctly
pubescent, trichomes
papilliform to cylindri-
cal, less than 0.2 mm.
long

(40°-)50°-70°(-80°)
All branches subtended
by leaves or bracts,
all but very upper
whorls subtended by
bracts

Shorter to as long
as valve
As in R. crispus

Uniformly thickened

No. 7 continued

CHARACTERISTICS	<u>R. crispus</u>	Hybrid	<u>R. pulcher</u>
Valve Shape	Ovate	Oblong-ovate	Ovate-deltoid to oblong-deltoid
Size (mm.)	{2.8-}3.5-5.0(-6.0)x {2.5-}3.5-5.5(-7.5)	4-4.5x3.5-4.0	{2.7-}3.5-4.8(-5.5)x 1.8-3.0(-3.5)
Apex	Subacute	Acute to subacute	Acute
Base	Cordate, indented	Truncate to cordate, indented to 0.5 mm.	Truncate
No. of Teeth/side	0.2-0.5(-1.0) mm.	5-7	{0-}5-9(-11)
Length of Teeth (mm.)	Entire to 1-8	Less than 0.3	{0.2-}1.0-1.6(-2.0)
No. of Anastomoses/side	Less than 0.3		
	(1-)2(-3)	2-3	1-2(-3)
No. of Tubercles	1-3, usually 3, unequal in size	1-3, unequal in size	1-3, unequal in size
Large Tubercle			
Shape	Ovate	Ovate	Ovate
Size (mm.)	{1.4-}1.8-3.0(-3.3)x {0.8-}1.0-1.5(-2.0)	1.3-3.0x0.8-1.5	{1.5-}1.8-2.5(-3.0)x {0.5-}0.8-1.3(-1.7)
Apex	Obtuse to acute	Acute	Acute
Small Tubercle			
Shape	Ovate	Ovate	Ovate
Size (mm.)	1.0-2.0(-2.5)x	1.0x0.5	1.0-1.5x0.5-0.6
Achene	0.5-1.2		
Size (mm.)	{2.0}2.2-2.8(-3.0)x {1.2-}1.4-1.6(-2.0)	No fully-developed achenes	{1.8-}2.2-2.8x {1.4-}1.5-2.0
Apex	Acute		Acute
Base	More broadly acute than apex to somewhat obtuse		Obtuse to almost truncate

Table 14. Comparison of Hybrids with Parent Species
No. 8 R. longifolius x obtusifolius
(continued on p. 139)

No. 8

CHARACTERISTICS

Basal Leaves
Shape

L.W.
Apex
Base

Angle of Veins
Vestiture

Inflorescence
Angle of Branches
Substantion of
Branches and Whorls

Pedicel
Size

Type of Articula-
tion and its
Location

R. longifolius

Lanceolate to
oblong - lanceo-
late

2-4:1

Acute to subacute
Broadly cuneate,
obtuse to weakly
cordate

(40°-)50°-70°

Indistinctly to
moderately papil-
lose, trichomes
papilliform, less
than 0.1 mm. long

25°-40°

All branches sub-
tended by leaves or
bracts, occasionally
lowermost whorls
subtended by bracts

Usually longer than
valve

Distinctly swollen,
in the lower two-
fifths

Hybrid

Oblong -
lanceolate

2-3:1

Acute
Truncate, obtuse
to weakly cordate

42°-72°

Distinctly pubes-
cent, trichomes
cylindrical, less
than 0.3 mm. long

(20°-)30°-50°

As in R. longifolius

As in R. longifolius

As in R. longifolius

R. obtusifolius

Oblong - ovate

1.7-2.5

Acute

Distinctly cordate

45°-80°

As in Hybrid

(20°-)25°-40°

As in R. longifolius

As in R. longifolius

As in R. longifolius

No. 8 continued

CHARACTERISTICS

	<u>R. longifolius</u>	Hybrid	<u>R. obtusifolius</u>
Valve Shape	Orbicular to reniform	Reniform	Deltoid
Size (mm.)	(3.5-5.2-6.5(-7.5))x (4.0-6.0-7.5(-9.5))	4.8-7.4x4.7-8.3	(3.0-3.5-5.0(-6.0))x (2.0-2.5-3.5(-4.0))
Apex	Obtuse	Abruptly acute	Acute
Base	Cordate, indented (0.3-0.6-1.5 mm.)	Cordate, indented 0.4-1.8 mm.	Truncate to weakly cordate, indented to 0.5 mm.
No. of Teeth/side	Entire to 2-6	Entire to 3-7	(1-3-4(-5))
Length of Teeth (mm.)	Less than 0.3 mm.	0.1-0.7	(0.8-1.0-1.8(-2.0))
No. of Anastomoses/side	2-3	2	1-2
No. of Tubercles	0-1	1	1-3, usually 1
Large Tubercle Shape	Oblong	Ovate	Ovate
Size (mm.)	(0.3-0.5-1.3(-1.4))x (0.3-0.4-1.0)	0.8-1.9x0.5-1.0	(1.1-1.5(-2.9))x (0.4-0.7-1.5(-1.8))
Apex	Obtuse	Obtuse to acute	Obtuse to acute
Small Tubercle Shape			
Size (mm.)			
Achene			
Size (mm.)	(2.5-2.8-3.3(-3.5))x (1.3-1.5-1.8(-2.0))	2.3-3.1x1.3-1.8	Ovate 0.5-1.5x0.3-0.5
Apex	Acute	Acute	(1.8-2.3-2.9(-3.0))x (1.2-1.3-1.7(-1.8))
Base	More broadly acute than apex to somewhat obtuse	More broadly acute than apex	Acute As in hybrid
Colour	Red-brown	Red-brown as in <u>R. obtusifolius</u>	Red-brown

Table 14. Comparison of Hybrids with Parent Species
No. 9 R. maritimus var. fueginus x stenophyllus
(continued on p. 141)

No. 9

CHARACTERISTICS

R. maritimus
var. fuscus

Hybrid

R. stenophyllus

Basal Leaves

Shape

Narrowly lanceolate
to narrowly oblong -
lanceolate

No Basal Leaves
on Specimen

Narrowly lanceolate
to oblong

L:W

Apex

Base

3-7:1
Acute to acuminate
Cuneate, truncate
or weakly cordate

Angle of Veins

Vestiture

45°-70°
Distinctly pubescent,
trichomes cylindrical,
less than 0.3 mm. long

(Stem leaves with
vestiture of R.
maritimus)

Glabrous to indistinctly
papillose, trichomes
papilliform, less than
0.1 mm. long

Inflorescence

Angle of Branches

Subtention of

Branches and Whorls

(200°-)300°-60(-80°)
All branches sub-
tended by leaves or
bracts, all whorls
subtended by bracts

45°

As in R. maritimus

200°-400°

All branches subtended
by leaves or bracts,
all but upper whorls
subtended by bracts

Pedicel

Size

Usually longer than
valve

As in R. maritimus

As in R. maritimus

Type of Articula-

tion and its

Location

Not swollen, near
base

Distinctly swollen,
in the lower fifth

Distinctly swollen,
in the lower two-
fifths

No. 9 continued			
CHARACTERISTICS		<u>R. maritimus</u>	<u>R. stenophyllus</u>
Valve			
Shape		Deltoid	Deltoid
Size		(1.5-2.3-3.0(-3.5)x (0.8-1.1-1.5(-1.6)	(2.5-3.5-4.8(-5.3)x (2.5-3.3-4.5(-5.0)
Apex		Acute	Acute
Base		Cuneate	Truncate to weakly cordate
No. of Teeth/side		1-3	5-9(-10)
Length of Teeth (mm.)		(1.0-1.5-2.8(-3.5)	(0.2-1.0-1.0(-1.3)
No. of Anastomoses/side		1-2, usually 1	(1-2-3
No. of Tubercles		3, equal in size	3, equal in size
Large Tubercle		Fusiform	Ovate
Shape		(1.0-1.2-1.6(-2.0)x	(1.2-1.6-2.1(-2.5)x
Size (mm.)		0.3-0.5(-1.0)	0.9-1.2
Apex		Acute	Obtuse
Achene			
Size (mm.)		(1.1-1.3-1.7(-2.0)x (0.6-0.7-0.9(-1.1)	(1.9-2.0-2.6(-3.0)x (1.0-1.2-1.4(-1.5)
Apex		Acute	Acute
Base		More broadly acute than apex	Acute
		As in <u>R. maritimus</u>	

Table 14 Comparison of Hybrids with Parent Species
No. 10 R. occidentalis x pseudonatronatus
(continued on p. 143)

No. 10

CHARACTERISTICS

Basal Leaves
Shape

L:W

Apex

Base

Angle of Veins
Vestiture on
Axial Veins

R. occidentalis

Ovate-lanceolate
to oblong-lanceo-
late

2-4:1

Acute

Truncate to cordate

50°-90°

Glabrous

Hybrid

Oblong

4:1

Acute

Truncate

60°

Glabrous

R. pseudonastrenatus

Lanceolate

greater than 4:1

Acute

Attenuate (approx.
equal to apex)

50°-70°

Glabrous to indistinctly
papillose, trichomes
papilliform, less
than 0.1 mm. long

Inflorescence

Angle of Branches

Substitution of

Branches and Whorls

20°-40°

Lower to middle

branches subtended

by leaves, upper

branches and all

whorls not sub-

tended by leaves

or bracts

30°-40°

As in R. occiden-

talis

20°-40° (-50°)

All branches subtended

by leaves or bracts,

occasionally some

lower whorls subtended

by bracts

Pedical

Size

Usually longer

than valve

Not swollen, in

lower third

As in R. occiden-

talis

Distinctly swollen,

in lower quarter

Somewhat less to

longer than valve

Distinctly swollen,

in lower half

Type of Articulation and its

Location

No. 10 continued

CHARACTERISTICS

Valve

Shape
Size (mm.)

Apex
Base

No. of Teeth/side

Length of Teeth (mm.)
No. of Anastomoses/
side

No. of Tubercles
Large Tubercle
Shape
Size (mm.)
Apex

Achene
Size (mm.)

Apex
Base

R. occidentalis

Ovate to orbicular
(4.0-14.8-9.0(-12.5)x
(3.9-5.0-8.0(-1.3)
Obtuse to subacute
Truncate to cordate,
indented 0.3-1.0(-1.5)
Entire to weakly
serrate, 1-5
Less than 0.1 mm.
1-2(-3)

0

Oblong
0.9-1.3x0.5-0.7
Obtuse

(2.8-13.5-4.7(-4.8)x
1.6-2.5(-2.6)
Acute
Acute

Hybrid

Orbicular to ovate
3.3-6.2x3.9-5.5
Obtuse to subacute
Cordate, indented
0.3-0.7 mm.
Entire to weakly
serrate, 1-6
As in R. occidentalis
1-2

1

Oblong
0.9-1.3x0.5-0.7
Obtuse

2.1-3.3x1.0-1.3
Acute
Acute

R. pseudonatronatus

Orbicular
(2.0)3.8-4.9(-6.0)x
(2.3-14.0-5.3(-6.0)
Obtuse to subacute
Cordate, indented
(0.2-)0.4-0.8(-1.0) mm.
Entire to weakly
serrate, 1-3
As in R. occidentalis
1-3

0-1

Oblong
0.4-1.1x0.2-0.8
Obtuse

(1.3-)2.1-2.6(-2.9)x
(0.8-)1.2-1.4(-1.6)
Acute
More broadly acute
than base

Table 14 Comparison of Hybrids with Parent Species
No. 11 R. pseudonatronatus x triangulivalvis
(continued on p. 145)

No. 11

CHARACTERISTICS

R. pseudonatronatus

Hybrid

R. triangulivalvis
(from Rechinger 1937)

Basal Leaves

Shape

L:W

Apex

Base

Lanceolate
Greater than 4:1
Acute
Attenuate (approx.
equal to apex)

Linear-lanceolate
ca. 5:1
Acute
Acute

Angle of Veins

Vestiture on

Abaxial Veins

Glabrous to
indistinctly papil-
lose trichomes
papilliform, less
than 0.1 mm. long

ca. 60°
Glabrous

ca. 45°
Glabrous

Inflorescence

Angle of Branches

Subtension of

Branches and Whorls

20°-40° (-50°)
All branches sub-
tended by leaves
or bracts, occasion-
ally some lower
whorls subtended
by bracts

35°-50°
Lower to middle
branches subtended
by leaves, upper
branches not sub-
tended, no whorls
subtended by bracts

30°-40°
As in hybrid

Pedicel

Size

Longer than valve

Type of Articulation and its

Location

As in R. pseudonatronatus
Distinctly swollen,
in the lower quarter

Distinctly swollen,
near base

No. 11 continued

CHARACTERISTICS

	<u>R. pseudonatronatus</u>	Hybrid	<u>R. triangulivalvis</u> (from Reehinger 1937)
Valve Shape Size (mm.)	Orbicular {2.0-3.8-4.9(-6.0)x 2.3-4.0-5.3(-6.0)}	Ovate 2.9-5.2x3.3-4.8	Deltoid (1-1.3(-4)x2.5-3
Apex Base	Obtuse to subacute Cordate, indented (0.2-10.4-0.8(-1.0 mm.	Subacute	Acute
No. of Teeth/side	Entire to weakly serrate, 1-3	Subcordate, indented to 0.5 mm.	Subcordate to truncate
Length of Teeth (mm.)	Less than 0.1 mm.	Entire to weakly serrate Less than 0.1	Entire to crenulate
No. of Anastomoses/ side	1-3	1-2	1(-2)
No. of Tubercles	0-1	1-3, usually equal in size	3, equal in size
Large Tubercles Shape Size (mm.)	Oblong 0.4-1.1x0.2-0.8	Ovate 1.3-1.7x0.5-0.9	Fusiform 1.8-2.5x0.6-0.9
Apex Small Tubercle Shape Size (mm.)	Obtuse	Obtuse	Acute
Achene Size (mm.)	{1.3-2.1-2.6(-2.9)x 0.8-1.2-1.4(-1.6)}	Ovate 1.2-1.6x0.4-0.7	2.5-3
Apex Base	Acute More broadly acute than base	Acute Acute	Acute Acute
Colour	Red-brown	Red-brown as in <u>R. pseudonatronatus</u>	Dark red-brown to black

Figure 15

Valves of Hybrids

(source (Herbarium, State,
Collector)

- (a) R. conglomeratus x pulcher GH. VA. Randolph 377.
- (b) R. crispus x kernerii CAS 415202, CA,
Pollard s.n.
- (c) R. crispus x longifolius DAO, NS, Dawson
76225.
- (d) R. crispus x obtusifolius DAO, ON, cultivated
material from Dawson
76209.
- (e) R. longifolius x obtusifolius DAO, ON, cultivated
material from Dawson
7649.

The photographs illustrate the intermediate nature of the hybrid
valves.

1 unit = 1 mm.

a

b

c

d

e

Numerical Taxonomy

Numerical taxonomy is defined as "the numerical evaluation of the affinity or similarity between taxonomic units and the ordering of these units into taxa on the basis of their affinities" (Sokal and Sneath 1963). The chief aims of numerical taxonomy are repeatability and objectivity obtained through reducing the individual subjective biases of taxonomists. The use of electronic computers has allowed the widespread employment of multivariate statistics which are concerned with the joint distribution of several variates: previously the calculations involved were impossibly long unless the number of variables was very small. Marriott (1974) notes that the purpose of any multivariate analysis is to gain some insight into the structure of the measurements and the main features of the data. An important way of gaining a better understanding of the structure is by graphical presentation of the data in a reduced number of dimensions.

Techniques have been developed for grouping objects on the basis of observations on many attributes considered together and these are referred to as cluster analysis. Discussions of cluster analysis can be found in Marriott (1974), Sokal and Sneath (1963) and Sneath and Sokal (1973). Cluster analysis is employed to create groups (or clusters) whose members share as many common characteristics as possible (natural groups), either in the hope that these will represent groups that have a reality measured in terms of external criteria or of discontinuities in the variation

or merely for the convenience of dividing a large group into smaller units. The production of dendrograms based on these techniques provides a helpful pictorial representation of the data.

Many clustering procedures have been developed (Sneath and Sokal 1973). Three used in the studies on Rumex are single linkage (nearest neighbour), group average (UPGMA) and flexible sorting. Each method starts by uniting a pair of objects that are most similar or in a distance model, closest together. The methods differ in the way in which they calculate the distance from a cluster to other objects. In single linkage, the distance of an object to a cluster is represented by the distance to the nearest point in that cluster. In group average, the arithmetic mean of the distance to each point in a cluster is taken. The form of average linkage used in this study was that which is termed unweighted (UPGMA) in that when clusters are fused with other clusters, the average taken is the average of the distance between the individual points of each cluster. In the case of flexible sorting, the distance between an object L and a cluster formed by the fusion of objects J and K is defined as $D_{(J,K)L} = \alpha D_{JL} + \beta D_{KL} + \lambda D_{JK}$ where D_{JL} etc. represents the distance between J, L etc. and α and β are constants chosen such that $2\alpha + \beta = 1$. (Sneath and Sokal 1973). When $\beta = 0$, flexible sorting reduces to the weighted group average method (WPGMA) in which the distances between clusters are averages without regard to the number of objects in a cluster.

Single linkage is space contracting in that once a cluster is formed, it is the peripheral points that define the distance of the cluster from other objects and other clusters. The space within the periphery is in effect contracted out of existence. As a result, objects tend to link onto clusters one at a time forming long straggly clusters in which the distances between the extremities may be very great. Group average is space conserving since by taking an average of the distances to the objects, the spatial configuration of the objects is conserved. Flexible sorting is at the other extreme in that if $\beta < 0$, the space around a cluster will be dilated so that depending on the value of β , it becomes relatively easier for new clusters to be formed than for an object to join an existing cluster. It has been found that with $\beta = -0.25$, a balance is achieved between no clear grouping and an excessive constraint to form equal sized clusters.

Ordination procedures can be used to display some of the relationships of the data in a reduced number of dimensions. Two such methods are principal co-ordinate analysis (Gower 1966) and principal component analysis. In general, these methods accommodate the most important features of a set of data in a smaller number of dimensions. The similarities are discussed by Marriott (1974); briefly, the first operates on an object by object matrix and the second on a character by character matrix. When Euclidean distance is used to compare the objects and there are no missing data, the projections of the objects onto the principal components will be identical to the projection of the objects onto the principal co-ordinates. The space defined by these principal axes

(components or co-ordinates) preserves in total all the original distances between the objects but as each successive orthogonal axis represents less of the total distance, the major portion of the variation is represented in the first two axes. Hence two dimensional graphic representations allows a visualization of relationships of objects to each other. Whereas clustering methods as represented in the dendrograms, faithfully reproduce the relationships of very similar objects with increasing distortion at the higher levels of clustering, projections onto the principal axes preserve the major pattern of variation at the expense of an accurate portrayal of inter object distances.

These numerical techniques provide useful tools to the taxonomist but when the analysis is completed, it is still necessary to interpret it in terms of the original taxonomic problem as there are no internal tests of significance for these techniques.

Canonical analysis can be carried out if groups are definable on a priori grounds. One can then examine how far it is possible to recognize these groups on the basis of other observations made on the members of them. (Marriott 1974). The analysis allows for an estimate to be made of the probability of correct allocation to the designated groups. This method acts as an extension to multivariate observations of the ordinary analysis of variance within and between groups. By minimizing the within-group to between-group variance, canonical axes are obtained upon which the maximal separation of the groups is

expressed. The number of canonical (or discriminant) axes is one less than the number of groups recognized. Two examples of canonical analysis were used in this study. In the first case, taxa that had been defined on a single character were identified as groups and the discriminant axes were obtained. The significance of the difference between the group means on these axes was assessed. If no significant difference exists between the group means then it can be assumed that the single character separation is not supported by the other characters used in the study and the taxon represents the recognition of an artificial grouping. In the second type of canonical analysis, the analysis was carried out not to discriminate between groups but to obtain new axes (canonical axes) to provide a better representation of the inter group similarities.

In the studies of the intraspecific variation, the clustering procedures, the ordination procedures and the first type of canonical analysis were employed while in the study of the interspecific relationships, the ordination procedures and the second type of canonical analysis were carried out.

Intraspecific Numerical Studies

Within section Rumex in North America, two areas of infraspecific taxonomic problems were recognized as being ones in which numerical techniques might help to provide a better understanding of the variation and its subsequent taxonomic recognition. The two areas are in the R. maritimus and R. occidentalis species complex.

Procedure

For the study of R. maritimus and R. occidentalis approximately one thousand specimens of each species from the major Canadian and American herbaria were examined. After confirming their identification as a member of one of the two species complexes, 81 specimens of R. maritimus and 100 specimens of R. occidentalis representing the geographic and morphological ranges of their respective species complex in North America, were selected for use in the numerical portion of the study. Vegetative and floral or fruit characters were scored, resulting in an initial set of 49 characters for R. maritimus (Table 15) and 98 characters for R. occidentalis (Table 16). For each study, the pairwise similarity assessment was made using Gower's general coefficient of similarity (Gower 1971).

In R. maritimus, clustering was performed using the group average method (UPGMA) (Sneath and Sokal 1973) which is considered as a space conserving method and the groupings formed are less likely to be imposed by the clustering method such as flexible

sorting (with $B < 0$) or incremental sum of squares (see McNeill 1975). In R. occidentalis, clustering was performed using: single-linkage, group-average and flexible sorting (Sneath and Sokal 1973; Lance and Williams 1971). To portray the overall pattern of variation within the entire group of OTUs, a principal component analysis was carried out on R. maritimus, whereas on R. occidentalis, a similar method, a principal coordinate analysis (Gower 1966) was used; in each case, the representation on the principal axes is done at the expense of a faithful representation of the individual inter-OTU distance.

A discriminant analysis of the groups recognized by the previous analyses was carried out in both R. maritimus and R. occidentalis. In the discriminant analysis of both species, a reduced character set was used (Tables 15, 16).

TABLE 15 Rumex maritimusList of Characters used and the methods of Measurement
and Scoring

	Methods	
1. Valve length	apex to base (mm)	*
2. Valve width	at widest place (mm)	*
3. Length of longest spine	Edge of valve to apex of spine (mm)	*
4. Distance to first anastomosis	From midrib to first anastomosis (mm)	*
5. Sepal length	Apex to attachment on receptacle (mm)	*
6. Pedicel length	Receptacle to attachment on whorl	*
7. Pedicel width	At middle (mm)	*
8. Achene length	Apex to base (mm)	*
9. Achene width	At widest place (mm)	*
10. Achene depth	(mm)	*
11. Length of lowest branch	From stem to end of branch (cm)	*
12. Length of highest branch	From stem to end of branch (cm)	*
13. Greatest distance between adjacent branches of inflorescence	(cm)	*
14. Branch angle	From stem to branch (degrees)	*
15. Stem trichome length	Apex to base (average of three)	*
16. Stem trichome width	At base (average of three)	*
17. Middle Leaf Apex	(degrees)	
18. Middle Leaf Base	(degrees)	

TABLE 15, cont.

19. Middle Leaf Pubescence Length	Apex to base (cm)
20. Middle Leaf Pubescence Width	At widest place (cm)
21. Middle Leaf Petiole Length	(cm)
22. Middle Leaf Vein Angle	Middle of lamina (degrees)
23. Upper Leaf Pubescence Length	as in 15
24. Upper Leaf Pubescence Width	as in 16
25. Diameter of Stem	At widest point (mm)
26. Diameter of Stem at Base	(mm)
27. Diameter of Stem at 20cm	20cm down from top of inflorescence (mm)
28. Length of Stem	Top of inflorescence to base of stem (cm)
29. No. of teeth per side of valve	(number, average of two sides) *
30. No. of anastomoses per side of valve	(number) *
31. No. of branches on inflorescence	(number)
32. Cordateness of Middle Leaf	Indentation measured from base up *
33. Location of valve width	Base upwards 0 - 0.2 (1), 0.2 - 0.4 (2), 0.4 - 0.6 (3), 0.6 - 0.8 (4), 0.8 - 1.0 (5)
34. Stem Colour	Brown (1), Brown-red brown (2), Red brown (3), Red brown-red (4), Red (5) *
35. Stem trichomes density	Densely (3), moderately (2), glabrous (1)
36. Middle stem leaf width location	(same as 33)
37. Middle Stem Leaf Base	Cordate (1), Truncate (2), Obtuse-acute (3)

TABLE 15, cont.

38. Margin of Middle Leaf	Entire (1), entire-weakly crisp(2), crisp(3)	
39. Ochrea Description	Not persistent (0), weakly (1), moderately (2), persistent (3)	
40. Middle Stem Leaf Pubescence	(as in 35)	
41. Variation in spines/side of valve on a plant	1 (1), 1 - 2 (2), 2 (3), 2 - 3 (4), 3 (5)	*
42. Tubercle Apex	Acute (1), Subacute (2), obtuse (3)	*
43. Tubercle Width Location	(same as 33)	
44. Length of Valve Apex	From point of narrowing to apex (mm)	*
45. Tubercle Length	Apex to base (mm)	*
46. Tubercle Width	At widest place (mm)	*
47. Tubercle Depth	(mm)	*
48. Tubercle Base Location	Location of tubercle base from valve base (mm)	*
49. Anther Length	(average of three) (mm)	*

* Complete for all OTUs and used for discriminant analysis

Fruit characteristics based on one fruit per specimen; one that appeared representative of the majority of the fruit in the inflorescence. Unless the inflorescence was fully mature, upper whorls were not used as these are the last to mature; usually, the fruit was taken from the mid-whorls of a mature branch.

TABLE 16 Rumex occidentalisList of characters used and the methods of measurement
and scoring

Character	Methods of Measurement and Scoring
1. Valve length	Apex to base (mm) *
2. Valve width	At widest place (mm) *
3. Valve base indentation	From base to tope of indentation (mm) *
4. Length of longest serration	Edge of valve to apex of serration (mm) *
5. Distance to first anastomosis	From midrib to first anastomosis (mm) *
6. Sepal length	Apex to attachment on receptacle (mm) *
7. Pedicel length	Receptacle to attachment in whorl (mm) *
8. Pedicel width	at middle (mm) *
9. Achene length	Apex at base (mm) *
10. Achene width	at widest place (mm) *
11. Achene depth	(mm) *
12. Longest inflorescence branch length	From stem to end of branch (cm)
13. Shortest inflorescence branch length	From stem to end of branch (cm)
14. Greatest distance between adjacent branches of inflorescence	(cm)
15. Branch angle	From stem to branch (degrees)
16. Stem trichome length	Apex at base (mm) (average of three) *
17. Stem trichome width	At base (mm) (average of three) *

TABLE 16, cont.

18. Basal leaf length	Apex to base (cm)
19. Basal leaf width	At widest place (cm)
20. Basal leaf apex	(degrees)
21. Basal leaf base	(degrees)
22. Basal leaf basal indentation	From base to top of indentation (cm)
23. Basal leaf trichome length	Apex to base (mm) (average of three)
24. Basal leaf trichome width	At base (mm)
25. Basal leaf petiole length	(cm)
26. Basal leaf vein angle	Middle of lamina (degrees)
27.- 35 Lower leaf characteristics	as in 18 - 26
36 - 44 Middle Leaf characteristics	as in 18 - 26
45 - 51 Upper Leaf characteristics	as in 18 - 26
52. Diameter of stem	At widest point (mm)
53. Diameter at base of complete stem	(mm)
54. Diameter at 20cm	20cm down from top of inflorescence (mm)
55. Stem length	Top of inflorescence to base of stem (cm)
56. Teeth per side of valve	(number, average of two sides) *
57. No. of anastomoses on each side of valve	(number) *
58. No. of branches in inflorescence	(number)
59. Location of valve width	see Table 15
60. Valve apex	Subacute (1), Obtuse (2)
61. Valve margin	Entire (1), Weakly serrate (2)

TABLE 16, cont.

62. Description of valve reticulation	veins at 90° to the midrib (1), veins at 90° - 45° (3), veins at 45° (5)
63. Sepal declination	Reflexed upwards (1), horizontal (2), reflexed downwards (3)
64. Point of pedicel articulation	(as in number 59)
65. Type of articulation	none evident (0), faint line (1), distinct swelling (3)
66. No. of fruit in whorl	0 - 9 (1); 10 - 20 (2), 20 + (3)
67. Stem colour	see Table 15
68. Stem trichome density	See table 15
69. Basal leaf width location	(as in number 59)
70. Basal leaf apex	Acute (1), subacute (2), obtuse (3)
71. Basal leaf base	(see table 15)
72. Basal leaf trichomes on petiole	Absent (0), Present (1)
73. Basal leaf trichomes on veins	(as in 72)
74. Basal Leaf trichome density	(as in 68)
75. Basal Leaf margin	(see table 15)
76 - 82 Lower leaf characteristics	as in 69 - 75
83 - 89 Middle leaf characteristics	as in 69 - 75
90 - 96 Upper leaf characteristics	as in 69 - 75
97. Type of subtention of branches and whorls by leaves or bracts	No branches or whorls subtended (0), lower to middle branches subtended (2), all branches except very upper ones (3), all branches (4), all branches and rarely some lower whorls (5)
98. Ochreae description	(see table 15)

* Complete for all specimens and used in discriminant analyses.

Fruit selected as described in Table 15.

Rumex maritimus

In North America, past treatments have recognized (a) a North American variety, var. fueginus of the otherwise Eurasian R. maritimus, along with a closely related native species, R. persicarioides (St. John, 1915), (b) two North American native species (R. persicarioides and R. fueginus) loosely related to R. maritimus (Rechinger 1937), or (c) two native varieties, (var. persicarioides and var. fueginus) within R. maritimus (Mitchell 1977). In each case, the only consistent characters separating the two North American taxa are those of tubercle size and shape. In one, the tubercle is large with a rounded apex, while the other tends to have a fusiform tubercle with an acute apex. In Rechinger's treatment, a number of varieties were also recognized within R. fueginus with acute fusiform tubercles, based on spine length.

Results

The group average clustering technique (UPGMA) indicated two groups (Figure 17); the first and smaller group consisted of six specimens, all of which possessed larger tubercles and with one exception had an obtuse apex. This group consisted of specimens from coastal British Columbia, Oregon and California. The remaining specimens forming the second group were from across Canada and the United States; they had acute to obtuse tubercles that were variable in size. Within this group, the five specimens with obtuse tubercles did not cluster together, but they were all from

Figure 16 Group Average Dendrogram of R. maritimus
The dendrogram represents the relationships of 81
OTUs of R. maritimus.

Figure 16

Group Average
Rumex maritimus
 Tubercle Apex
 O Oblique
 A Acute
 S Subacute

coastal areas in Eastern North America (Quebec, Nova Scotia, Prince Edward Island and Massachusetts). The specimens in which the tubercles were subacute did not occur in any specific geographic area. The principal component analysis (Figure 16, showing a plot of the first two components) shows that the large tubercled west coast group (predominantly with obtuse apices) appears to represent a loose association of OTUs somewhat distinct from the remainder of the OTUs. Within these remaining OTUs, those with subacute tubercles, also tended to cluster loosely, however, no clear separation of these OTUs was evident as was the case for the west coast OTUs already discussed. An examination of the OTUs with acute tubercles failed to indicate any evident groupings with the varieties that Reehinger recognized on the basis of spine length.

Canonical analyses were carried out utilizing the 25 characters that were complete for all 81 OTUs. Separation into groups was carried out on the basis of the shape of the tubercle apex; three groups were formed: Group A with an obtuse apex (10 OTUs), Group B with a subacute apex (15 OTUs) and Group C with an acute apex (56 OTUs). Initially, Group B was not included in the analysis; both Hotelling's multivariate T^2 test and Wilks test indicated a significant difference between the mean vectors of groups A and C. Hotelling's multivariate T^2 test indicated significance at the 95% and the 99% level.

The canonical axis is calculated to maximize the separation between the groups and in this case, the members of A and C, are all correctly classified on this axis. An analysis of variance

Figure 17 Principal Component Analysis of Rumex maritimus

Plot of the 81 OTUs of R. maritimus on the first two axes recognized by a principal component analysis (axis I: horizontal; axis II: vertical).

Figure 17.

Figure 18 Canonical Analyses of R. maritimus

Histograms plotted on horizontal axes calculated from two canonical analyses carried out on the 81 OTUs of R. maritimus.

Figure 16

Canonical Analyses

Rumex maritimus

Canonical Axis Calculated Using Groups A(1+2) and C

Canonical Axis Calculated Using Groups A₂ and C

indicated that the following characters accounted for the majority of the variation within the groups (in descending order of importance): tubercle width, trichome length, valve width, achene width, tubercle length, length of longest spine, pedicel length, number of teeth/ side of valve. When Group A was plotted on the canonical axis, two groups emerged, A, consisting of 5 west coast OTUs and A2 consisting of 5 east coast OTUs. A further canonical analysis was carried out using only the members of Group A2 and Group C to determine the relationship of A2 to C without the effect of the large tubercled A1 specimens interfering. Hotelling's T^2 test and Wilks test indicated only a slight degree of significance between the mean vectors. Hotelling's multivariate T^2 test was just barely significant at the 95% level.

One specimen of Group C was reported as being misclassified and placed in Group A2. This specimen was however not geographically associated with the OTUs in A2, as it was from inland Alaska. An analysis of variance indicated that the same eight characters reported from the previous analysis accounted for the majority of the variation; the only difference was that the length of the longest spine was now more important than tubercle length.

By applying canonical loadings for each character, the OTUs of Group B were assigned to the canonical axes calculated from Groups A and C, and similarly, those for groups B and similarly those for groups B and A, on the canonical axes calculated from Groups A2 and C. A histogram of the distribution of all the OTUs along the two canonical axes is shown in Figure 19. In each case,

two of the OTUs scored as having subacute tubercles, have been placed in the region of specimens of Group A2. Of these, one is a maritime specimen from a coastal habitat in Prince Edward Island and geographically associated with Group A2, while the other is a specimen from an inland habitat from northern Ontario. Only one OTU with subacute tubercles was placed with the west coast Group A, and this was a west coast OTU from British Columbia, which had also been associated with the obtuse-tubercled west coast specimens in the earlier cluster analyses. The majority of the OTUs with subacute tubercles (13) were plotted amongst the OTUs of group C. From Figure 18, it can be seen that Group A2 (eastern obtuse tubercled OTUs) is more closely associated with Group C, than is Group A (western obtuse tubercled OTUs).

The range and mean of each of the eight important characteristics (not including the tubercle apex) separating the large tubercled western specimens, the large tubercled eastern specimens and the smaller tubercled wide ranging specimens are in Table 17.

In addition to being geographically isolated, the Western group was found to be separated from the Eastern group and the Acute-subacute group, by the following pair-wise combination of the above characters: Valve Width/ Valve Length; Seed Width/ Tubercle Length; Tubercle Length/ Tubercle Width. However, no single pair-wise combination of the eight characters was found to clearly separate the Eastern group from the Acute-subacute group. Further analyses using reduced sets of characters are needed.

The present study recognizes three infraspecific taxa within

R. maritimus, in addition to the introduced var. maritimus: a western taxon and an eastern taxon with large tubercles, commonly obtuse at the apex and a wide ranging taxon with small tubercles, acute to subacute at the apex. The taxonomic recognition of these taxa will be discussed under the taxonomic treatment of R. maritimus in North America (Canada and the United States).

TABLE 17

Comparison of the Eight Main Characteristics
used to Discriminate between the
three Groups of *R. maritimus*

	Minimum	Maximum	Mean
1. Tubercle Width (mm)			
Eastern (A2)	0.4	0.7	0.5
Western (A1)	0.7	1.0	0.8
Acute & Subacute (C & B)	0.3	0.5	0.4
2. Stem Pubescence Length (mm)			
Eastern	0.05	0.17	0.116
Western	0.1	0.25	0.175
Acute & Subacute	0.02	0.42	0.195
3. Valve Width (mm)			
Eastern	1.0	1.4	1.1
Western	1.2	1.7	1.3
Acute & Subacute	0.9	1.7	1.3
4. Seed Width (mm)			
Eastern	0.7	0.9	0.8
Western	0.8	1.0	0.9
Acute & Subacute	0.6	0.9	0.7
5. Tubercle Length (mm)			
Eastern	1.3	1.8	1.4
Western	1.9	2.2	2.0
Acute & Subacute	1.0	1.8	1.4
6. Length of Longest Spine (mm)			
Eastern	1.2	2.8	1.9
Western	2.0	2.5	2.2
Acute & Subacute	1.0	3.5	2.8
7. Pedicel Length (mm)			
Eastern Group	2.0	4.0	3.1
Western Group	2.5	6.5	4.5
Acute & Subacute Group	1.8	6.5	3.7
8. No. of Teeth/side			
Eastern Group	1.0	2.0	1.9
Western Group	1.0	2.0	1.3
Acute & Subacute Group	1.0	3.0	2.2

Rumex occidentalis

Within the R. occidentalis species complex, the members of which are native to North America, two species were recognized by Rechinger (1937); these were based almost entirely on valve length, the one having valves up to 5 mm long and the other comprising specimens with valves greater than 7 mm long. Lepage (1954) recognized Rechinger's two species as varieties, but named an intermediate variety comprising those specimens with valves greater than 5 mm and up to 6 mm long; this intermediate variety was appropriately named var. perplexus. Although Lepage had allowed for the identification of some specimens not considered by Rechinger, there are many plants in which the valves are between 6 and 7 mm in length so that even using this single character criterion, some specimens of R. occidentalis were still not assignable to a variety. Calder and Taylor (1968) on examining a large series of specimens from western Canada found it very difficult to segregate two species from this polymorphic complex and recognized only one taxon, R. occidentalis. Boivin (1968) recognized two varieties within R. occidentalis, var. occidentalis, with valves 4 to 6 mm long and var. labradoricus, with valves 6 to 8 (to 11) mm long. In the present study, a combination of traditional taxonomic methods with numerical ones was used to provide an assessment of the variation within R. occidentalis.

Results

Initially, all of the scored characteristics were utilized in the analyses. The dendrograms obtained from the three clustering procedures produced very similar results; from the dendrogram

(Figure 19) obtained by the average-linking algorithm, it can be seen that no very distinct groups emerge; however, by marking the valve length characters presently used to distinguish infraspecific taxa, it can be seen that many of those specimens with valves greater than 7 mm, cluster loosely together. All of the specimens with valves greater than 7 mm, were found to be western in geographic origin, occurring in British Columbia, Washington, Oregon, California, and Idaho; the specimens with valves less than or equal to 7 mm, were from across the range of R. occidentalis (including the regions in which the larger valved specimens occurred. A similar loose association of specimens with valves less than or equal to 5 mm, can be observed, but as in the previous case, they are not in a distinct cluster. These OTUs indicated no characteristic geographical range. A plot of the OTUs based on their first two principal axes from the principal co-ordinate analyses (Figure 20) indicated two loosely clustered groups connected by a large number of intermediates; the two groups corresponding to those OTUs with valves less than or equal to 5 mm and those greater than 7 mm. As many of the vegetative characters were lacking on the majority of the specimens, these characters were not used in subsequent analyses; a greatly reduced character set (15 characters, Table 16) using only characters complete for all OTUs was employed. Since it appeared that valve length was producing some groupings within R. occidentalis, the valve length character used to separate taxa in R. occidentalis was investigated more directly.

The hypothesis to be tested was that the taxa recognized within R. occidentalis were not significantly different and represented

Figure 19 Average-Linkage Dendrogram of R. occidentalis
The dendrogram represents the relationships of 100
OTUs of R. occidentalis.

Figure 19.

AVERAGE LINKAGE

Rumex occidentalis

Figure 20 Principal Co-ordinate Analysis of
R. occidentalis

Plot of the 100 OTUs of R. occidentalis on the first
two axes recognized by a principal co-ordinates
analysis (axis I: horizontal; axis II: vertical).

Figure 20

Principal Co-ordinate Analysis *R. occidentalis*

portions of a continuum that had been arbitrarily chosen. Separation into groups was made on the basis of valve length, which as noted, was the character upon which three varieties had been based. 20 OTUs of valve length up to 5.0 mm formed Group A, 38 OTUs of valve length greater than 5.0 to 6.0 mm formed Group B, 19 OTUs of valve length greater than 7 mm, formed Group C. OTUs of valve length > 6 to 7 mm were initially omitted from the analysis as they had not been clearly assigned to a variety excepting in Boivin's (1966) treatment. A canonical analyses was carried out on these three groups. A one-way analysis of variance indicated that the most significant characters distinguishing these groups were: Valve width, sepal length, distance to the first anastomosis, pedicel length, achene length and achene width. Based on Hotelling's multivariate T^2 test for equality of mean vectors, the difference between the means of groups A and B were barely significant at the 95% level while the differences between the means of B and C, A and C and A + B and C were significant at the 95% and 99% levels.

A scatter diagram plotted on the canonical axes (Figure 21) indicates the relationships between these groups. Two specimens from Group A were considered misclassified and placed in Group B and 10 OTUs originally in Group B were assigned by the discriminant function to Group A. Correspondingly, 1 OTU from Group C was assigned to Group B, but no OTUs of Group A or B were considered as being misclassified and placed in Group C. As valve width was obviously closely associated with valve length, a further canonical analysis was performed on the three groups with both the length and

Figure 21 Canonical Analysis of R. occidentalis

A scatter diagram based on the canonical analysis of the 77 OTUs of R. occidentalis.

A: OTUs of valve length up to 5.0 mm

B: OTUs of valve length greater than 5.0 to 6.0 mm

C: OTUs of valve length greater than 7.0 mm

OTUs of valve length greater than 6.0 to 7.0 mm were omitted from this analysis.

Figure 21.

Canonical Analysis Rumex occidentalis

width characters removed; the one-way analysis of variance indicated that the groups were now separated mainly on sepal length, distance to the first anastomosis, pedicel length, achene length and achene width. An examination of Hotelling's T^2 test indicated the same relationships between the groups as was evident in the previous analysis with the exception that the difference in means between groups A and B was now not significant at the 95% level. The scatter diagram plotted on the canonical axis showed little difference from that of Figure 21. Two OTUs of Group A were misclassified and placed in Group B (these were found to be the same ones as in the earlier case). Eleven OTUs of Group B were considered misclassified and placed in Group A (seven of these were also previously misclassified). One specimen of Group C was misclassified and placed in Group B (it was not misclassified earlier). Once again, no specimens of Groups A and B were misclassified and placed in Group C. Because of the lack of significance between their means, A and B cannot be considered as distinct groups, while Group C is a distinct group and separable even when the valve length and width characters are removed. The group of OTUs with valves above 6 mm and up to and including 7 mm in length, were now added to the other OTUs in order to examine Group C, on its own versus the other OTUs. In this analysis, Group 1 represents all OTUs with valves less than or equal to 7 mm in length (81 OTUs) while Group 2 represents the remaining 19 OTUs with valves greater than 7 mm in length. Initially, valve length was removed from this analysis and then both the valve length and width were removed; a significant difference was observed at the 95% and 99% levels using the

Hotelling's T^2 test but in practise, groups 1 and 2 would be difficult to discriminate. The one-way analysis of variance indicated that most of the separation was based on: distance to the first anastomosis; sepal length, pedicel length, valve base, achene length and width. Of the 100 OTUs, only three were considered misclassified; two from Group 1 and one from Group 2; the OTU misclassified in Group 2 had not been previously misclassified, while the two misclassified OTUs from Group 1 were from the OTUs in the group of OTUs with valves above 6 mm and up to and including 7 mm in length and were from the same geographic area as members of Group 2. It is interesting to note that the majority of the OTUs with valves above 6 mm and up to and including 7 mm in length were considered in the analysis to be correctly assigned to Group 1, rather than Group 2 as would be expected if the initial separation between the groups existed at the 6 mm valve length as indicated by Boivin (1968).³

The present study indicates that within R. occidentalis, two groups are recognizable, but the point of separation between the predominantly western large valved OTUs (valves greater than 7 mm long (group C)) and the east to west ranging OTUs (valves less than or equal to 7 mm long (groups A and B)) has been decided arbitrarily and must be considered as tentative. The ranges for the 15 characteristics used in discriminating the two groups and their means are presented in Table 10.

TABLE 18

Comparison of the Characteristics used to Discriminate
Between the Two Groups in A. occidentalis

Character	Minimum	Mean	Maximum
1. Valve Length (mm)			
Group 1	4.3	5.7	7.0
Group 2	7.3	8.7	12.0
2. Valve Width (mm)			
Group 1	3.9	5.7	7.4
Group 2	5.7	7.9	11.3
3. Valve Base Indentation (mm)			
Group 1	0.05	0.4	0.9
Group 2	0	0.7	3.0
4. Length of Longest Tooth (mm)			
Group 1	0	0.1	0.3
Group 2	0	0.11	0.25
5. Distance to First Anastomosis (mm)			
Group 1	0.8	1.2	1.7
Group 2	1.0	1.8	3.0
6. Sepal Length (mm)			
Group 1	1.2	1.9	3.0
Group 2	1.9	2.4	3.0
7. Pedicel Length (mm)			
Group 1	3.5	8.5	14.0
Group 2	7.3	11.4	15.0
8. Pedicel Width (mm)			
Group 1	0.1	0.15	0.25
Group 2	0.15	0.17	0.25
9. Achene Length (mm)			
Group 1	2.8	3.7	4.75
Group 2	3.7	4.2	4.8
10. Achene Width (mm)			
Group 1	1.6	2.0	2.5
Group 2	1.8	2.2	2.7
11. Achene Depth (mm)			
Group 1	1.2	1.7	2.3
Group 2	1.5	1.9	2.3

TABLE 18, cont.

Character	Minimum	Mean	Maximum
12. Trichome Length (mm)			
Group 1	0	0.03	0.1
Group 2	0	0.05	0.3
13. Trichome Width (mm)			
Group 1	0	0.037	0.13
Group 2	0	0.044	0.09
14. No. of Teeth per side of Valve			
Group 1	0	2.2	6
Group 2	0	2.6	9
15. No. of Anastomoses per side of Valve			
Group 1	1	1.46	3
Group 2	1	1.52	3
Group 1 = Specimens with valves less than or equal to 7 mm in length corresponding to groups A and B in the text.			
Group 2 = Specimens with valves greater than 7 mm in length, corresponding to group C in the text.			

With the exception of the valve length, the character upon which the groups were defined, no other character appeared to be able to clearly separate the two groups.

The appropriate taxonomic recognition of these two groups will be discussed under the taxonomic treatment of R. occidentalis.

Interspecific Relationships within Section Rumex

In his treatment of the North American species of section Rumex, Rechinger (1937) arranged the species of the section in subsections; in 1949 when discussing relationships within section Rumex, Rechinger stated that Meisner's (1856) groups based entirely on whether the valves were entire or dentate were artificial whereas "the section is easily divided into natural groups in respect of the characters of the valves combined with the annual or perennial growth".

Utilizing Rechinger's treatment, the North American species recognized in this study are grouped into 15 subsections as follows:

- 1) Subsection Hymenosepali Rechinger f., Publ. Field Mus. Bot. 17:7. 1937.
R. hymenosepalus
- 2) Subsection Densiflori Rechinger f., Publ. Field Mus. Bot. 17:7. 1937.
R. densiflorus, R. pyrenanthus, R. orthoneurus,
R. praecox
- 3) Subsection Aquatici Rechinger f., Repert. Sp. Nov. 31:232. 1933.
R. occidentalis, R. arcticus
- 4) Subsection Rumex (= Subsection Patientiae Rechinger f., Repert. Sp. Nov. 31:230. 1933.
R. patientia, R. kernerii

- 5) Subsection Crispi Rechinger f., Publ. Field
Mus. Bot. 17:8. 1937.
R. crispus
- 6) Subsection Conferti Rechinger f., Candollea 12:61. 1949.
R. confertus
- 7) Subsection Alpini Rechinger f., Candollea 12:60. 1949.
R. alpinus
- 8) Subsection Longifolii Rechinger f., Candollea 12:49. 1949.
R. longifolius, R. pseudonatronatus
- 9) Subsection Conglomerati Rechinger f., Publ. Field Mus.
Bot. 17:8. 1937
R. conglomeratus, R. sanguineus
- 10) Subsection Hydrolapatha Rechinger f., Pub. Field
Mus. Bot. 17:8. 1937.
R. britannica
- 11) Subsection Obtusifolii Rechinger f., Publ. Mus.
Bot. 17:8. 1937.
R. obtusifolius, R. pulcher
- 12) Subsection Dentati Rechinger f., Publ. Field
Mus. Bot. 17:8. 1937.
R. dentatus, R. violascens
- 13) Subsection Maritimi Rechinger f., Publ. Field
Mus. Bot. 17:8. 1937.
R. maritimus, R. palustris
- 14) Subsection Stenophylli Rechinger f., Candollea 12:89. 1949.
R. stenophyllus

- 15) Subsection Hamati Rechinger f., Candollea 12:112. 1949.

R. brownii

Most studies in Europe have accepted Rechinger's division into sections, however, Borodina (1979) divided section Rumex into two sections; section Rumex and section Orientales A. Baran. et B. Skvortz. Within her section Rumex, she recognizes four subsections and within section Orientales, she recognizes two subsections:

Section Rumex

- (1) Subsection Hydrolapatha Rechinger f.
- (2) Subsection Patientiae Rechinger f., including Rechinger's subsections Crispi, Longifolii, Alpini, Conferti, Stenophylli, Aquatici.
- (3) Subsection Obtusifolii Rechinger f., she excludes R. pulcher but includes Rechinger's Conglomerati.
- (4) Subsection Pulchri Borodina, she creates a new subsection for R. pulcher.

Section Orientales

- (1) Subsection Dentati Rechinger f.
- (2) Subsection Maritimi Rechinger f.

Although not mentioned in Borodina's account, the two endemic North American subsections, Hymenosepali and Densiflori would seem to fall within her subsection Patientiae.

In order to determine the utility of these subsections so far as the North American representatives are concerned and to assess the relative merits of Rechinger's or Borodina's treatments,

a numerical study designed to indicate the interspecific relationships within section Rumex was carried out.

Procedure

From the specimens measured for the species descriptions, several specimens representing the morphological variation of each species were selected (R. praecox had to be omitted as no mature specimens exist) (Table 19). Initially 46 characteristics (Table 20) available on a mature inflorescence were used. (No leaf characteristics were used as they are not always available in the field or on herbarium specimens.) In order to observe the associations of the species, both a principal components analysis and a canonical analysis were utilized. Since a canonical analysis can not without an estimation of their values employ characters that are not present on all specimens, and as the programs used did not provide for such estimates, all tubercle characteristics were removed from this analysis. All programs were from Agriculture Canada, Ottawa, and were utilized through the courtesy of the Biosystematics Research Institute.

Results

The associations of the species could not be ascertained using the principal components analysis since the species were not clearly separating in the projection of the OTU's on the first few principal components due to the high degree of within species variability shown by the species. Various transformations

Table 19

Symbol	Species	No. of Samples
A	<u>R. orthoneurus</u>	8
B	<u>R. hymenosepalus</u>	9
C	<u>R. pycnanthus</u>	8
D	<u>R. brownii</u>	6
E	<u>R. arcticus</u>	7
F	<u>R. britannica</u>	9
G	<u>R. longifolius</u>	9
H	<u>R. densiflorus</u>	9
I	<u>R. pseudonatronatus</u>	9
J	<u>R. violascens</u>	9
K	<u>R. dentatus</u>	9
L	<u>R. confertus</u>	9
M	<u>R. kernerii</u>	9
N	<u>R. alpinus</u>	9
O	<u>R. pulcher</u>	9
P	<u>R. conglomeratus</u>	9
Q	<u>R. stenophyllus</u>	9
R	<u>R. obtusifolius</u>	9
S	<u>R. occidentalis</u>	9
T	<u>R. sanguineus</u>	8
U	<u>R. crispus</u>	9
V	<u>R. maritimus</u> v. <u>fueginus</u> & var. <u>pacificus</u>	9
W	<u>R. palustris</u>	3
X	<u>R. maritimus</u> var. <u>maritimus</u>	2
Y	<u>R. patientia</u>	9
	Total	204

The coding and number of samples used for each species in the canonical analysis of the species of section Rumex in Canada and the United States are given.

Table 20 Rumex: Interspecific RelationshipsCharacteristics Used and the Methods
of Measurement and Scoring

Character	Methods of Measurement and Scoring
1. Valve length	see table 15
2. " width	" " "
3. " base measure	Truncate to cordate (0-4), cuneate (8), obtuse (12)
4. Length of longest serration, tooth or spine	see table 15
5. Distance to first anastomosis	" " "
6. Sepal length	" " "
7. Pedicel length	" " "
8. " width	" " "
9. Achene length	" " "
10. " width	" " "
11. Largest tubercle length	" " "
12. " " width	" " "
13. " " depth	" " "
14. " " base location	" " "
15-18. Small tubercle characteristics	as in 11-14
19. Branch angle	see table 15
20. Stem trichome length	" " "
21. " " width	" " "
22. Size of articulation	width of articulation (mm.)
23. Number of teeth per side of valve	see table 15
24. Number of anastomoses per side of valve	" " "
25. Largest tubercle width location	" " "
26. Shape of apex of large tubercle	acute (1), subacute (2), obtuse (3)
27. Small tubercle width location	as in 25
28. Shape of apex of small tubercle	as in 26
29. Number of tubercles usually on fruit of inflorescence	0(1); 0-1(2), 1(3), 1-2(4), 2(5), 2-3(6), 3(7)
30. Location of valve width	see table 15

Character	Methods of Measurement and Scoring
31. Shape of valve apex	narrowly acute (1), acute (2), subacute (3), obtuse (4), narrowly obtuse (5)
32. Valve margin	entire (1), entire-weakly crenate (2), weakly crenate (3), weakly crenate-weakly serrate (4), weakly serrate (5), weakly serrate-dentate (6), dentate (7), dentate-spiny (8), spiny (9)
33. Description of valve reticulation	see table 15
34. Sepal declination	" " 16
35. Type of articulation	" " "
36. Location of seed width	" " "
37. Achene apex	widely acute (1), moderately acute (2), narrowly acute (3)
38. " base	acute (1), obtuse (2), truncate (3)
39. " edge	keeled (1), acute (2), rounded (3)
40. " colour	black - dark-red-brown (1), light red-brown (2), light brown (3), brown (4), green-brown (5)
41. No. of fruit in whorl	see table 16
42. Stem colour	" " 15
43. Stem ridging	distinctly ridged (1), weakly ridged (2), not ridged (3)
44. Stem trichome density	see table 15
45. Point of pedicel articulation	" " "
46. Type of subtention of branches and whorls by bracts or leaves	" " 16

were also used on the data to attempt to minimize this variation and to reduce the spread of the OTU's within a species when projected on the principal components; the transformations did not help for all the species as the direction of the variation in all species was not the same.

In using the canonical analysis, the specimens are pre-assigned to their species and the method minimizes the variation within and maximizes the variation between species. Based on Bartlett's test for significant axes, it was found that of the 24 axes all but the last three axes were significant at the 95% level, indicating that most of the species are independently discriminated using the characteristics employed. Scatter diagrams based on the plots of axis 1 vs 2, 1 vs 3.....1 vs 24, were produced in order to see what the main trends in the associations of the species are. Figure 22 shows a scatter diagram of axes 1 and 2. In this plot, only one group corresponding directly to a subsection of Rechinger was indicated - the association of R. conglomeratus (P) and R. sanguineus (T) which is clearly distinct from the other species and would correspond to subsection Conglomerati. This does not correspond to Borodina's subsection as the Conglomerati were placed in the Obtusifolii and no such association is apparent. The group of R. dentatus (K), R. palustris (W), R. maritimus var. maritimus (I) and R. maritimus (native varieties - V) corresponds in part to the section Orientalis of Borodina except that R. violascens (J) belonging to the subsection Dentati does not indicate an association with this group. No clear separation of the Dentati and Maritimi was apparent and

Figure 22 Canonical Analysis of Rumex Species

A scatter diagram based on the canonical analysis of the species of Rumex in Canada and the United States. The coding for the species is listed in Table 19.

Figure 22

Canonical Analysis of Rumex Species
Scatter Diagram

49 OTUs were coincident
155 OTUs plotted

Borodina's section Orientales indicated a closer association with the other species than did subsection Conglomerati. The other groups do not correspond to any of the existing subsections of Borodina or Rechinger, nor did they appear to represent useful taxonomic units.

In the other plots of axes 1 vs 3 ... 24, the group of R. conglomeratus and R. sanguineus remained separated from the other species while the species making up Borodina's section Orientales tend to associate together but close to the remaining species. By axes 1 and 16, and subsequent plots, only two groups of species are apparent, the Conglomerati and a loose grouping of the remaining species. The first few plots also indicated that R. obtusifolius and R. pulcher tend to be associated together and the value of Borodina's placing of R. pulcher in its own section is questionable. The groups of species formed in the plots of axes 1 vs 3 24, as in axes 1 and 2, did not correspond to existing subsectional groups of Borodina or Rechinger, nor could they be easily delimited from each other.

The present study does not support the existence of the two sections of Borodina as the species of Rechinger's Conglomerati which she includes in her Obtusifolii are even more recognizable as a group than are the species of her section Orientales. The species of her section Orientales do tend to group together but close to the other species. Borodina's subsection Patientiae encompasses too much variation and would also

be of little taxonomic value.

In re-examining the sections of Rechinger in light of tubercle and valve characteristics and the groupings observed in this study, Rechinger's subsections do appear to associate species with similar vegetative and fruit characteristics. However, several of his subsections (e.g. (a) Rumex, Crispi and Longifolii or (b) Dentati and Maritimi) based on their close association in the canonical analysis may be better represented by one group. Although grouping species with similar characteristics together, the canonical analysis did not suggest groups that could be easily delimited from each other or that correspond directly to existing subsections. Consequently, this study has not employed the subsectional taxa. The acquisition of additional data which provides practical means of discriminating subsections might change this situation. Since in North America, many subsections of Rechinger are represented by only a few species, their recognition as distinct groups might have been obscured. Only a study of all of the species recognised in section Rumex would actually indicate the naturalness and taxonomic utility of the subsections.

Key to the Species

A key to the species of section Rumex was set up with the main points of separation based on fruit and basal leaf characteristics. Specimens from across the North American range of the species were employed in the construction of the key so that it would be capable of dealing with the variation evicent in the species.

For the taxonomic descriptions of the species following the key, most vegetative characteristics were of a somewhat general nature due to their inherent variability with respect to size and the fact that most herbarium specimens of Rumex do not include good leaf material. The methods of measuring the characteristics used in the descriptions can be found in Tables 15, 16, 20. For the more common or well represented species, 19 selected specimens in mature fruit were used in order that the ranges given for most of the quantitative characteristics would provide more information by expressing the 10th centile, median and 90th centile as suggested by Jardine and Sibson (1970). Measurements given in brackets represent observed values outside of the given range. For several species, R. dentatus, R. densiflorus, R. kernerii and R. violascens, the same approach was taken for the quantitative characteristics but only 9 specimens were used, largely due to the lack of mature specimens. When fewer than 9 mature specimens were available as in R. alpinus (5), R. arcticus (3), R. brownii (3), R. confertus, R. palustris (2), R. praecox (3-all immature), R. pycnanthus (3), R. sanguineus (3) and R. orthoneurus (4), the range simply represents the smallest and largest values observed.

1. Fruiting valves all lacking tubercles; pedicels with or without distinctly swollen points of articulation.
2. Valves distinctly spiny, the spines 0.8 - 1.3 mm long with recurved tips; inflorescences very lax, spreading; basal leaves narrowly lanceolate, often contracted above the base of the leaf. 25. R. brownii
2. Valves entire or at most with a few small straight teeth, 0.4 mm long; inflorescences compact; basal leaves orbicular to narrowly lanceolate, not contracted above the base of the leaf.
3. Pedicels with the point of articulation clearly swollen in fruit.
4. Valves ovate-triangular, acute, basal leaves orbicular, about as long as wide, the base deeply cordate; abaxial leaf veins distinctly pubescent with trichomes cylindrical several times longer than wide. 12. R. alpinus
4. Valves orbicular to reniform, obtuse; basal leaves lanceolate, oblong-lanceolate or narrowly lanceolate, several times longer than wide, the base cuneate; abaxial leaf veins glabrous to papillose with trichomes papilliform, shorter than to as long as wide.
5. Achenes red-brown, less than 1.5 mm wide; basal leaves 4 - 8 times as long as wide, widest at or near the middle, the base narrowly cuneate; valves usually less than 5.0 mm wide. 14. R. pseudonatronatus

5. Achenes deep brown; greater than 1.5 mm wide; basal leaves less than 4 times as long as wide, widest at the middle to near the base, the base broadly cuneate; valves usually greater than 5.0 mm wide. 13. R. longifolius
3. Pedicels with the point of articulation no wider than the rest of the pedicel, even in fruit.
6. Valves deltoid, widest at base, the apices narrowly acute. 3. R. pycnanthus
- * 6. Valves ovate to orbicular, widest at or near middle the apices acute to obtuse.
7. Valves ovate, acute at apex.
8. Valves abruptly contracted toward the apex, widest at or near middle; pedicels less than twice as long as the valves; basal leaves with large lateral nerves alternating with shorter ones; achene deep brown to red-brown. 2. R. densiflorus
8. Valves gradually narrowing to the apex, widest in the lower third; pedicels usually twice as long as the valves; basal leaves with lateral nerves all $\pm$ equal in size; achenes usually green-brown to red-brown. 4. R. orthoneurus
7. Valves orbicular to oblong-orbicular, obtuse to subacute at apex.
9. Valves deeply cordate, the indentation greater than 1.5 mm; achenes rounded at base, greater than 2.5

mm wide; basal leaves cordate, truncate, rounded or broadly cuneate, the basal angle greater than 90° , roots not tuberous.

10. Basal leaves oblong to elliptic, less than 4 times as long as wide, widest at middle, rounded or narrowly cuneate at base. 5. R. praecox

10. Basal leaves lanceolate to oblong-lanceolate, greater than 4 times as long as wide, widest near base, broadly cuneate to truncate or cordate at base.

11. Basal and lower leaves truncate to cordate at base; lateral leaf veins raised on abaxial surface in living material; inflorescences usually with long lower branches, more than 8cm long; veins of valve diverging at approximately 90° to the midrib.

6. R. occidentalis

11. Basal and lower leaves broadly cuneate at base; lateral leaf veins not raised and barely evident on abaxial surface in living material; inflorescences usually with small lower branches, less than 8cm long; veins of valve diverging at approximately 45° to the midrib.

7. R. arcticus

1. At least one fruiting valve with a tubercle; pedicels with distinctly swollen points of articulation.

12. Only one fruiting valve with a tubercle, the others with at most weakly enlarged midribs.

13. Valve margins entire or, if possessing teeth, the longest tooth less than 0.4 mm long.

14. Valves oblong to oblong-ovate, approximately twice as long as wide; tubercle occupying ca 80% of valve width.

16. R. sanguineus

14. Valves orbicular to reniform or ovate, as long as or shorter than wide; tubercle occupying ca 20% of valve width.

15. Valves reniform, abruptly acute; basal leaves deeply cordate, abaxial veins of leaves and inflorescence branches with cylindrical trichomes, several times as long as wide.

11. R. confertus

15. Valves orbicular, reniform or ovate, obtuse to subacute; basal leaves cuneate, truncate, obtuse or very weakly cordate; abaxial veins of leaves and inflorescence branches glabrous or with small papilliform trichomes, shorter than to only slightly longer than wide.

16. Valves ovate, subacute, truncate to weakly cordate.

10. R. crispus.

16. Valves orbicular to reniform, obtuse, weakly to distinctly cordate.
17. Basal leaves 4 - 6 times as long as wide, widest at or near the middle, the base narrowly cuneate. 14. R. pseudonatronatus
17. Basal leaves less than 4 times as long as wide, widest at middle to above base, the base abruptly and broadly cuneate.
18. Tubercle usually 0.5 - 1.3 mm long, 0.3 - 0.8 mm wide, 0.3 - 0.6 mm deep; abaxial veins of cauline leaves distinctly papillose. 13. R. longifolius
18. Tubercle usually 0.6 - 2.8 mm long; 1.0 - 1.5 mm wide, 0.6 - 1.5 mm deep; abaxial veins of cauline leaves usually glabrous, if papillose, not distinctly so. R. patientia
13. Valve margins dentate to spiny, the longest spine or tooth greater than 0.4 mm long.
19. Pedicels shorter than the valves, thick throughout, not noticeably widening from the point of attachment on the stem to the receptacle; tubercle usually warty. 19. R. pulcher

19. Pedicels longer than the valves, filiform and widening from the point of attachment on the stem to the receptacle; tubercle smooth. 18. R. obtusifolius

12. All three fruiting valves with a tubercle, the tubercles approximately equal in size or one larger than the other two.

20. Tubercles on each valve approximately equal in size and appearance.

21. Valves entire or possessing a few small teeth, 0.3 mm long.

22. Valves $\pm$ orbicular, about as long as wide; tubercles fusiform not reaching the valve base; pedicel with the point of articulation no wider than the rest of the pedicel, even in fruit. 17. R. britannica

22. Valves oblong to oblong-ovate, approximately twice as long as wide; tubercles oblong to ovate, usually reaching the base of the valve; pedicel with a distinctly swollen point of articulation.

15. R. conglomeratus

21. Valves distinctly serrate or spiny, teeth 0.3 mm long.

23. Valves about as long as wide, usually with 4 or more teeth on each side. 14. R. stenophyllus

23. Valves about twice as long as wide, usually with less than 4 teeth on each side.

24. Pedicel shorter than the valve, moderately thick and not noticeably widening from stem to fruit,

ca 0.3 mm wide at middle; tubercles greater than 2mm long; achene with a rounded base. 20. *R. dentatus*

24. Pedicel as long or longer than the valve, thin at base and widening from stem to fruit, ca 0.2 mm wide at middle; tubercles less than 2mm long; achene with a cuneate base.

25. Leaves and inflorescence branches glabrous; anthers 0.6 - 1.0 mm long. 22. *R. palustris*

25. Leaves and inflorescence branches papillose-pubescent with the trichomes papilliform or cylindrical and less than 0.4 mm long; anthers 0.3 - 0.5 (-0.7) mm long.

22. *R. maritimus*

20. Tubercles unequal, one noticeably larger than the other two.

26. Valves entire or possessing a few small teeth, less than 0.4 mm long.

27. Valves oblong-ovate, more than twice as long as wide; sepals curved upwards between the valves; achenes often showing beyond the valve edge; inflorescence branches lax usually diverging at 40° - 80° to the stem.

16. *R. sanguineus*

27. Valves orbicular, reniform or rotund-triangular, as long or shorter than wide; sepals spreading or slightly reflexed; achenes not showing beyond valve edge; inflorescence branches compact, diverging at 20° - 40° to the stem.

23. Valves ovate, subacute, truncate to weakly cordate; basal leaves linear-lanceolate, 4 - 6 times as long as wide, the margin usually distinctly crisped; achenes (2.0-) 2.2 - 2.6 (-3.0) mm long. 10. R. crispus
23. Valves usually orbicular to reniform, obtuse, weakly to distinctly cordate; basal leaves oblong-lanceolate, 3 - 4 times as long as wide, the margin weakly crisped, achenes 2.9 - 3.8 mm long. 8. R. patientia
26. Valves serrate to spiny, the largest tooth or spine usually greater than 0.4 mm long.
28. Valves usually more than 6.0 mm wide, as wide as long, suborbicular, abruptly contracted to an acute apex; anastomoses on each side of the valve 3 - 4(-5), the first at a distance of 1.0 - 1.5 (-1.8) mm from the midrib. 9. R. kerneri
29. Valves usually less than 4mm wide, narrower than long, deltoid, gradually narrowing to an acute apex; anastomoses on each side of the valve, 1 - 2, the first at 0.4 - 1.0 (-1.3) mm from the midrib.
30. Branches of the inflorescence diverging at an angle greater than 50° , at maturity often tangled with each other; pedicel shorter than valve; tubercles verrucose. 19. R. pulcher
30. Branches of the inflorescence diverging at an angle of less than 50° , not tangled at maturity; pedicel longer than valve; tubercles smooth.

31. Longest tooth of valve rarely greater than 0.5 mm; smallest tubercle greater than 0.5 mm wide and clearly recognizable as a tubercle; base of valve broadly cuneate; basal leaves oblanceolate to obovate, broadly cuneate at base; abaxial leaf veins glabrous. 21. R. violascens

31. Longest tooth of valve greater than 0.8 mm; smallest tubercle less than 0.5 mm wide and often merging with the midrib and scarcely recognizable as distinct; base of valve truncate to weakly emarginate, basal leaves ovate to oblong-ovate, distinctly cordate at base, the abaxial leaf veins usually pubescent with cylindrical trichomes.

12. R. obtusifolius

1.

Rumex hymenosepalus Torrey, Bot. Mex. Bound. Surv, 177. 1853.

Type: Texas: E. Paso - Hudspeth Co. line, March 1851, G. Thurber.
(Holotype, GH!)

R. saxei Kellogg, Pacif. Rural Press 7:371. June 1879.

R. tuberosus A. Nelson, Bull. Torr. Bot. Club 25:283. 1898.

R. salinus A. Nelson, Bull. Torr. Bot. Club 25:549. 1898 #

R. tuberosus A. Nelson.

Type: Wyoming: Bitter Creek, 3 June 1897, A. Nelson 3114
(NY!)

R. arizonicus Britton, Trans. N.Y. Acad. Sci. 7:73. 1899.

Type: Arizona: Fort Verde, 29 Feb. 1883, E.A. Mearns 300
(NY!)

Perennial. Root distinctly tuberous. Stem 27 - 90 cm tall, typically ca 50 cm, ca 1 cm wide at base, leafy, alternately branched above middle, sand brown, erect, sulcate, glabrous to indistinctly papillose; trichomes papilliform, most evident on branches, less than 0.1 mm long, slightly longer than to as long as wide; ochraceous membranous, white, weakly to strongly persistent on mature plants. Branches diverging at 20° - 30° - 40° to the main stem. Basal leaves oblong to oblong-elliptic, acute to obtuse, narrowly cuneate to cuneate, 2 - 4 times as long as wide, widest at or near middle, largest observed leaf 32 cm long, 8.5 cm wide, crisped; middle veins of lamina diverging at 40° - 60° to the midrib; lamina glabrous above, glabrous to indistinctly papillose on veins below; trichomes papilliform, less than 0.1 mm long, shorter than to as

long as wide; petiole shorter than lamina, glabrous to indistinctly papillose. Lower cauline leaves very similar to basal but becoming progressively smaller up the stem. Upper cauline leaves narrowly lanceolate to narrowly oblong-elliptic, acuminate to acute, becoming progressively smaller up the stem; vestiture as basal leaves; petiole much shorter than lamina, the leaf often sessile. Inflorescence a narrow dense panicle, occupying the upper half of the stem, the lower to middle branches subtended by leaves, the upper branches without subtending leaves. Whorls commonly consisting of 5 - 20 flowers, arranged on branches and portion of main stem above branches, rarely subtended by leaf-like bracts, the lower whorls remote, the upper almost contiguous. Pedicels filiform, (4.5-) 5.5 - 9.0 - 12.8 (-20.0) mm long, less than 0.3 mm wide at middle, articulated at the middle to the lower third; articulation indistinct not swollen. Sepals oblong, obtuse, angled downwards, 1.6 - 2.3 - 2.5 (-2.8) mm long and about half as wide. Valves oblong-orbicular to ovate, obtuse to acutely obtuse, cordate, indented (1.0-) 1.5 - 3.0 - 3.5 mm at base, (9.0-) 11.2 - 13.5 - 13.8 (-17.0) mm long, (7.8-) 9.5 - 11.0 - 14.8 (-15.7) mm wide, widest at middle to lower third, the margin entire; veins weakly raised, diverging from the midrib at ca 90°, rarely to 45°, with 2 - 3 - 4 (-5) anastomoses on each side of the valve; distance to first anastomosis from midrib 1.5 - 2.5 - 3.0 (-4.0) mm. Tubercles lacking. Anthers 1.5 - 2.9 mm long. Mature achenes, ovate, brown to red-brown, (4.0-) 4.3 - 5.0 - 7.5 mm long, (2.5-) 2.6 - 3.1 - 4.5 (-5.0) mm wide, (2.0-) 2.3 - 2.5 - 3.0 (-3.8) mm deep, widest in lower two-fifths; acute at

apex, obtuse at base.

Distribution: Native in the west and southwestern United States and Mexico.

U.S.A.: AZ, CA, CO, MT, NV, NM, OK, TX, UT, WY. (Map 3)

Habitat: Occurring on dry, alkaline or sandy soils of plains, basins and hillsides.

Discussion: Rechinger (1937) recognized two varieties distinguishable only in the fruiting state: var. hymenosepalus with achenes ca 5mm long, valves elliptic-cordate, apex slightly acute, longer than broad and var. salinus (A. Nelson) Rechinger f., with achenes to 7mm long, valves cordate-orbicular, often longer than wide. In an examination of the available herbarium material (including a number of specimens cited by Rechinger (1937)), the distinction in valve shape claimed to separate the varieties was not apparent. Wyoming specimens had large achenes (6.5 - 7.5 mm long) but as noted no correlating valve shape was apparent; achenes from specimens in other parts of the range were 4 - 5.8 mm long, even in areas where var. salinus was expected, based on Rechinger's (1937) study. As most specimens from across the range of R. hymenosepalus are not fully mature and as the distinguishing characteristics of the varieties are not reliable, the present study will not recognize the infraspecific taxa.

Due to the distinctness of its tuberous roots, basal leaves, large anthers and large valved fruit, it would be very difficult to confuse this species with any other native or introduced

species of section Rumex. Some confusion might exist with the native R. venosus Pursh of section Axillares, occupying a mid-west to western range in the United States and extending into the southern portions of the Canadian prairies (Rechinger 1937; Sarkar 1958). R. venosus is clearly separable due to its (a) smaller, narrowly cuneate and somewhat leathery leaves, (b) axillary branches, (c) shorter stem (15 - 30 cm tall), (d) large reniform valves, 24 - 30 mm wide, and (e) lack of tuberous roots.

204

Map 3 Distribution of R. hymenoseratus.

2.

Rumex densiflorus Osterhout, *Erythraea* 6:13, 1898.

Type: Wyoming: North Park from near the Colorado-Wyoming border in the Laramie River drainage, G.E. Osterhout, 1678 (Holotype, UC; Isotypes, US 3489201, GH1)

Perennial. Rootstock creeping. Stem ca 1m tall, greater than 1cm wide at base, erect, sulcate, brown to red-brown, leafy, branched alternately above middle, glabrous to indistinctly papillose; trichomes papilliform, most evident on upper branches, less than 0.1 mm long, shorter than to as long as wide, rarely to twice as long as wide; ochreae commonly deciduous to partially persistent on mature plants.

Branches diverging at 20° - 30° to the main stem and often weakly arching. Basal leaves large, largest observed 53.5 cm long, 12.0 cm wide, oblong to oblong-lanceolate, acute to obtuse, commonly obtuse to very weakly cordate, indented to 0.5 cm at base, 2 - 4.5 times longer than wide, widest at middle to lower third, entire; middle veins of lamina diverging at 60° - 80° to the midrib; lateral veins of alternating sizes, long and short; lamina glabrous above, glabrous to indistinctly pubescent on veins below; trichomes cylindrical, less than 0.3 mm long, 1.5 - 4 times as long as wide; petiole somewhat shorter than lamina, glabrous to indistinctly pubescent.

Lower cauline leaves very similar to basal but becoming progressively smaller up the stem; petiole shorter than lamina. Upper cauline leaves narrowly lanceolate to lanceolate, acute to acuminate, obtuse to narrowly cuneate, entire; glabrous above, glabrous to indistinctly papillose on veins below; trichomes papilliform, less

than 0.1 mm long, shorter than to as long as wide; petiole much shorter than lamina. Inflorescence a narrow panicle occupying the upper half of the stem, only the branches in the lower half of the panicle subtended by leaves. Whorls commonly consisting of 10 - 20 flowers, arranged on branches and portion of main stem above branches, not subtended by bracts, the lower whorls remote, the upper contiguous. Pedicels filiform, (4.5-) 8.5 - 11.0 - 16.0 mm long, less than 0.3 mm wide at middle, articulated in the lower two-fifths to the lower third; articulation a faint line or else scarcely visible. Sepals oblong, obtuse, spreading at right angles to pedicel to reflexed slightly downwards, 1.5 - 2.3 - 2.5 mm long and about half as wide. Valves ovate-triangular, abruptly acute, weakly cordate, indented 0.1 - 0.6 - 1.0 mm at base, (4.5-) 5 - 6.5 - 8.3 mm long, (4.8-) 5.3 - 7.0 - 9.0 mm wide, widest at middle to lower third, the margin below the apex weakly serrate to erose; distinct teeth 3 - 7 per side, the largest less than 0.3 mm long; veins raised, diverging from the midrib at ca 90° with 1 (2) anastomoses on each side of the valve; distance to the first anastomosis from midrib 1.0 - 1.5 - 2.5 mm. Tubercles lacking. Anthers ca 1.5 mm long. Mature achene ovate, green-brown (2.5-) 3.0 - 3.5 - 4.7 mm long, (1.6-) 1.8 - 2.0 - 2.3 (-2.5) mm wide, 1.5 - 1.6 - 2.0 mm deep, widest at or near middle, acute at the apex and base, the base tending to be somewhat broadly so.

Distribution: Native in the S.W. United States, not widespread.

U.S.A.: CO, WY (Map 4)

Habitat: Montane to alpine, along streams and valleys; according to Harrington (1954) found from 1500 - 3000 m, mostly above 2700 m.

Discussion: Although more specimens of this species are now available than were cited by Rechinger (1937), few of them represent fully mature plants. Additional collections, especially of mature fruiting plants are needed in order to fully understand the variability within the species; Osterhout (1898) reported that the rootstock seemed to be horizontal (creeping) but this cannot be ascertained from herbarium material. The present treatment recognizes R. densiflorus as a separate species but the lack of mature specimens seriously restricts our understanding of its variability and interrelationships.

Mar 4 Distribution of R. densiflorus, R. orthoneurus,
R. praecox and R. rycnanthus.

Map 4

3.

Rumex pycnanthus Reehinger f., Repert. Spec. Nov. Regni Veg.

38:372. 1935; replacing R. subalpinus M.E. Jones, Proc. Cal.

Acad. ser. 2,5:1895, non R. subalpinus (Schur) Simonkai.

Type: Utah: Brigham Peak, near Marysville, 10,500 feet (3450 m)

in gravel, in the bed of a subalpine stream. M.E. Jones 5957

(Lectotype, UC; Isotype, F 300619!)

Perennial. Roots fusiform or rootstock creeping. Stems ca 1m tall, greater than 1cm wide at base; erect, sulcate, brown to red-brown, moderately leafy, branched alternately above the middle, glabrous to indistinctly pubescent; trichomes cylindrical most evident on branches, less than 0.1 mm long, longer than to as long as wide; ochreous deciduous to partially persistent on mature plants. Branches diverging at ca 30° to the main stem and often weakly arching. Basal leaves large, oblong-lanceolate, acute, obtuse, ca 2 - 3 times as long as wide, entire to weakly crisped; middle veins of lamina diverging at ca 60° to the midrib, glabrous above and below; petiole somewhat shorter than the lamina, glabrous. Lower cauline leaves similar to basal but becoming progressively smaller up the stem, crisped, glabrous above, indistinctly pubescent on veins below; trichomes cylindrical, less than 0.2 mm long, 1 - 4 times as long as wide; petiole shorter than lamina, indistinctly pubescent. Upper cauline leaves linear-lanceolate, acuminate, narrowly cuneate, crisped, glabrous above, weakly papillose on veins below; petiole much shorter than lamina, weakly papillose. Inflorescence a narrow, somewhat dense panicle

occupying the upper half of the stem, only the lowermost branches subtended by leaves. Whorls commonly consisting of 10 - 20 flowers, arranged on branches and portion of main stem above branches, not subtended by leaf-like bracts, the lower whorls remote, the upper contiguous.. Pedicels filiform, 2.5 - 11.5 mm long, less than 0.3 mm wide at middle, articulated in lower half; articulation at most weakly evident. Sepals oblong, obtuse, spreading at right angles to pedicel, 1.3 - 2.5 mm long and about half as wide. Valves weakly deltoid, acute, truncate to weakly emarginate at base, 4 - 7 mm long, 3.5 - 5.3 mm wide, widest at or near the base, the margin below the apex weakly serrate; distinct teeth 2 - 6 per side, the largest typically less than 0.5 mm long; veins raised, diverging at ca. 90° to the midrib; with 1 (2) anastomoses on each side of the valve; distance to the first anastomosis from the midrib 0.8 - 1.5 mm. Tubercles lacking. Mature achenes, ovate, brown to red-brown, 3.0 - 3.8 mm long, 1.8 - 2.2 mm wide, widest at or just below the middle, apex and base acute, the base somewhat broadly so.

Distribution: Restricted to the S.W. United States, apparently rare.

U.S.A. CO, UT. (Map 4)

Habitat: Montane to alpine, along streams.

Discussion: Rechinger (1937) examined specimens of R. pycnanthus from two sites in Utah; since then, only one specimen from a site

in Colorado has become available. Additional specimens and field studies are needed in order to understand better both the within species variability and the relationships with R. densiflorus, R. orthoneurus and R. praecox.

4.

Rumex orthoneurus Rechinger f., Repert. Spec. Nov. Regni Veg. 40: 294. 1936.

Type: Arizona: Chiricahua Mts., Barefoot Park, rolling, andesitic pineland, recently lumbered, 8000 - 8250 feet (2430 - 2510 m), Blumer 1449. (Holotype, NY! Isotypes, F. 242311! Neb 659791, US 5616991, US 5633071, W, U, Z)

Perennial. Stem ca 1m tall, greater than 1cm wide at base, erect, sulcate, red-brown, moderately leafy, branched alternately above middle, glabrous or indistinctly papillose; trichomes weakly cylindrical to papilliform, evident on upper stem and branches, less than 0.1 mm long, shorter than to twice as long as wide; ochreous deciduous to partially persistent on mature plants.

Branches diverging at ca 30° to the main stem and often weakly arching. Basal leaves large, largest observed 49.0 cm long, 18.0 cm wide, oblong to oblong-lanceolate, acute to subacute, widely cuneate to obtuse or weakly cordate, 3 - 4 times as long as wide, widest at or near middle, entire; lateral veins approximately equal in length; middle veins of lamina diverging at (50°-) 60° - 90° to the midrib, commonly ca 80°; lamina glabrous above, glabrous to moderately papillose to pubescent on veins below; trichomes papilliform to cylindrical, less than 0.1 (-0.3) mm long, shorter than to 3 times as long as wide; petiole somewhat shorter than to longer than the lamina, glabrous to indistinctly papillose, to pubescent. Lower cauline leaves similar to basal but soon lanceolate, acute to acuminate, cuneate, becoming progressively shorter up the stem; petiole shorter than the lamina. Upper cauline leaves lanceolate to linear-lanceolate, acuminate, narrowly cuneate,

entire, the lamina glabrous above and below; petiole much shorter than lamina, glabrous, the leaf sometimes subsessile. Inflorescence a narrow panicle, occupying the upper half of the stem, all but the upper branches subtended by leaves. Whorls commonly consisting of 10 - 20 flowers, arranged on branches and portion of main stem above branches, not subtended by bracts, the lower whorls remote, the upper becoming somewhat contiguous. Pedicels filiform, (4.5-) 13.3 - 17.0 (-18.5) mm long, less than 0.3 mm wide at middle, articulated in the lower two-fifths; the articulation at most scarcely visible. Sepals oblong, obtuse, 1.3 - 2.5 mm long and about half as wide, commonly reflexed slightly downwards to spreading at right angles to the pedicel. Valves ovate-deltoid, acute to occasionally abruptly so at apex, truncate at base or more commonly emarginate and then indented to 0.5 mm, 4.3 - 7.0 mm long, 3.8 - 7.0 mm wide, widest in the lower third, the margin below the apex entire to weakly serrate; distinct teeth 2 - 7 per side, the largest to 0.5 mm long, commonly less than 0.3 mm; veins raised, diverging midrib at an angle of 90° , with 1 (2) anastomoses on each side of the valve; first anastomosis 1.0 - 1.8 mm from midrib. Tubercles lacking. Anthers 1.3 - 2.0 mm long. Mature achene ovate, brown, 2.6 - 4.0 mm long, 1.5 - 2.5 mm wide, 1.4 - 2.0 mm deep, widest at or near middle, acute at apex and base, somewhat more broadly so at base.

Distribution: Native only in the S.W. United States; apparently very rare.

United States: AZ (Map 4)

Habitat: Along streams, originally reported from rolling andesitic pineland, recently lumbered.

Discussion: R. orthoneurus was described on the basis of material from one Arizona collection. Rachinger (1937) noted that it was very closely related to R. densiflorus, a species also occurring in the southwest. His treatment was accepted in the floristic work of Kearney and Peebles (1951). At present, fifteen specimens of the species from only four localities in Arizona are known; only four of these have mature fruit. Because the material is so limited, the present treatment accepts R. orthoneurus with the reservation that additional material is needed to be certain of the delimitation of the species from R. densiflorus, R. pycnanthus, and R. praecox and its relationships to them.

5.

Rumex praecox Rydberg, Bull. Torrey Bot. Club 33: 137. 1906.

Type: Colorado: Grayback mining camps, 1900, Rydberg and Vreeland 6328 (NY!)

Perennial. Rootstock fleshy, tuber-like, creeping. Stems up to 45 cm tall, less than 1 cm wide at base, leafy, branched alternately, erect, sulcate, red-brown, glabrous; ochreae deciduous to partially persistent on mature plants. Branches short, less than 6.0 cm long, diverging at ca 15° to the main stem. Basal leaves small, oblong or elliptic, obtuse to subacute, rounded to broadly cuneate, shorter than 6 (-14.5) cm long, 2 - 3.2 times as long as wide, widest at or near the middle, entire to weakly crisped; middle veins of lamina diverging at (45° - 60° - 70°) to the midrib; lamina glabrous above, commonly glabrous to weakly papillose on veins below; trichomes cylindrical, less than 0.2 mm long, twice as long as wide; petioles longer than to as long as lamina, glabrous. Lower cauline leaves similar to basal but progressively smaller up the stem; petiole much shorter than the lamina. Upper cauline leaves ovate, obtuse to subacute, rounded to broadly cuneate at base, entire; glabrous above, commonly glabrous to indistinctly papillose on veins below; trichomes papilliform, less than 0.1 mm long, shorter than wide; petiole much shorter than lamina, glabrous. Inflorescence a narrow panicle occupying much of the stem, the lower branches subtended by leaves, the upper branches without subtending leaves. Whorls consisting of 10 - 20 flowers, arranged on branches and portion of main stem above branches; none subtended by bracts, the lower and upper whorls remote. Pedicels filiform, 3.5 - 7 mm

long, less than 0.2 mm wide at middle, articulated below middle; articulation not marked by any swelling, scarcely evident.

Sepals oblong, obtuse, 1.6 - 2.5 mm long and about half as wide.

Immature valves ovate, to oblong, obtuse, ca 4.5 mm long, ca 2.5 mm wide, widest at or near middle, the margin entire to very weakly serrate; distinct teeth 2 - 5 per side, the largest ca 0.1 mm long; veins weakly raised, diverging at ca 45° to the midrib, with 1 anastomosis on each side of the valve; distance to the first anastomosis from the midrib ca 0.8 mm.

Tubercles lacking. Anthers 1.2 - 1.9 mm long. Immature achenes ovate, green-brown, 2.8 - 3.8 mm long, 1.5 - 2.0 mm wide, widest at or near middle, acute at apex and base, more broadly so at base.

Distribution: Native only to the S.W. United States; apparently very rare.

U.S.A. CO, WY. (Map 4)

Habitat: open ground, along streams and valleys, at about 10,000 feet (ca 3000m) (Harrington 1954).

Discussion: Rydberg in naming R. praecox cited two Colorado specimens whereas Rechinger (1937) because of some vegetative differences between them, only accepted the Rydberg and Treland 6328 specimen as R. praecox. Rechinger assigned the other specimen - Baker, Earle and Tracy 270 to R. densiflorus. An examination of both specimens and of a further specimen collected in nearby Wyoming suggests that they are all referable to this species. However, none of the collections is mature and a

completely satisfactory assessment of the status of this taxon cannot be made until further material is available. Reehinger (1937) suggested that R. densiflorus, R. pycnanthus, R. orthoneurus and R. praecox could possibly be referred to one collective species. He maintained them however, as separate species because the material available to him was inadequate to justify making such an extreme taxonomic change. More than forty years later, the same situation exists.

6. Rumex occidentalis S. Watson, Proc. Amer. Acad. 12:253. 1877.

Locality cited: "From Alaska to northern California, eastward to the Saskatchewan and Labrador and northward in the mountain to Colorado and New Mexico."

Lectotype: Saskatchewan, 1858, E. Bourgeau. (GH!) (see description)

R. aquaticus sensu Pursh, Fl. Amer. Sept. 248. 1816, non Linnaeus (1753).

R. domesticus sensu Hooker, Fl. Bor. Amer. 2. 129, 1840, non Hartman (1820)

R. longifolius sensu Meisner ex de Candolle Prodr. 14:44 1856, non de Candolle (1815)

Perennial. Root fusiform. Stem up and to greater than 1m tall, ca. 1cm wide at base, smooth to weakly sulcate, brown to red-brown, often with red-purple areas, moderately leafy, branched alternately in upper two-thirds, or often only in the upper half, glabrous to indistinctly papillose, rarely distinctly pubescent; trichomes papilliform to cylindrical, most widest on branches, less than 0.1 mm long, rarely to 0.2 mm, shorter than to 2½ times as long as wide; ochreae deciduous to partially persistent on mature plants. Branches diverging at 20° - 30° - 40° to the main stem and often weakly arching. Basal leaves large, largest observed 36.5 cm long, 12.5 cm wide, ovate-lanceolate to oblong-lanceolate, acute to subacute, truncate or weakly to distinctly cordate, the indentation less than 2.2 cm, 2 - 4 times as long as wide, widest at or near base, entire to weakly crisped; middle veins of lamina

leaving midrib at 50° - 90° , commonly at 60° ; lamina glabrous above, glabrous to indistinctly papillose or rarely pubescent on vein below; trichomes papillate to rarely cylindrical, less than 0.3 mm long, commonly less than 0.1 mm long, shorter than to as long as wide, rarely to 5 times longer than wide; petiole much longer to somewhat shorter than lamina, glabrous to indistinctly papillose, rarely pubescent. Lower cauline leaves similar to basal but progressively smaller up the stem, more weakly cordate to rounded at base; petiole shorter than the lamina. Upper cauline leaves narrowly lanceolate to linear-lanceolate, acute to acuminate, truncate to rounded or narrowly cuneate, entire to weakly crisped, glabrous above, glabrous to indistinctly papillose or rarely pubescent on the veins below. Inflorescence a narrow panicle, occupying the upper two-thirds or often only the upper half of the stem, the lower to middle branches subtended by leaves, the upper branches not subtended by leaves. Whorls consisting of (8-) 12 - 16 - 26 (-28) flowers, on branches and portion of main stem above branches, whorls not subtended to rarely the lowermost whorls subtended by leaf-like bracts, the lower whorls remote, the upper whorls contiguous. Pedicels filiform, (3.5-) 6.5 - 9.6 - 13.0 (-17.5) mm long, less than 0.3 mm wide at middle, articulated in the lower third; articulation weakly evident, not swollen. Sepals oblong, obtuse, at right angles to pedicel, to weakly reflexed, (1.2-) 1.6 - 2.0 - 2.6 (-3.0) mm long and about half as wide. Valves orbicular to ovate, obtuse to subacute, truncate to weakly cordate or emarginate, indented (0-) 0.3 - 0.5 - 1.0 (-1.5) mm at base, (4.0-) 4.8 - 6.1 - 9.0 (-12.5) mm long,

(3.9-) 5.0 - 5.8 - 8.0 (-11.3) mm wide, widest at middle to lower third, entire to very weakly serrate; distinct teeth 1 - 4(5) per side, the largest less than 0.3 mm long; veins raised at ca. 90° to the midrib with 1 - 2 - 2 (-3) anastomoses on each side of the valve; distance to first anastomosis 0.9 - 1.4 - 1.8 mm. Tubercles lacking. Anthers ca. 1mm long. Mature achene ovate, red-brown, (2.8-) 3.0 - 4.0 - 4.4 (-4.8) mm long, (1.4-) 1.7 - 2.0 - 2.5 (-2.6) mm wide, (1.0-) 1.5 - 1.7 - 2.1 (-2.3) mm deep, widest at or near the middle, acute at the apex and base, somewhat narrower at the apex than the base.

Within R. occidentalis, two varieties are recognizable:

Valves up to 7mm long, base tending to be truncate to weakly emarginate. var. occidentalis.

Valves greater than 7mm long, base tending to be weakly emarginate to subcordate. var. labradoricus.

Rumex occidentalis var. occidentalis.

R. bakeri Greene, Pl. Baker. 3:15. 1901

Type Specimen: Colorado: Gunnison. 22 Aug. 1901. Baker 903 (US 412516; US 959010; RM 163066; NY!)

R. gracilipes Greene, Pittonia 4:304. 1901.

Type Specimen: Nevada, most meadows at the Pine Creek Hay Ranch, above Palisade. 25 July, 1893. E. L. Greene. Not High Ranch as reported by Rechinger (1937).

R. polyrhizus Greene, Pittonia 4:305. 1901.

Type Specimen: Wyoming: near Sherman. July 1893. Greene.

R. occidentalis var perplexus Lepage, Nat. Can. 81:257.
1954.

Type Specimen: Ontario: embouchure de la riviere Octamapow,
54° 25' Lat. N. 21 Oct 1953. Dutilly and Lepage 31491.

(Holotype, CAN 232336!)

Distribution:

Canada: AB, BC, NT, LB, MB, NF, ON, QU, SK, YT;

United States: AK, AZ, CA, CO, ID, MN, NV, NH, ND, OR, SD, UT,
WA, WY. (Map 5).

This variety is found across the entire range of the species. The reports of this variety from New Brunswick (Macoun 1883), Nova Scotia (Roland and Smith 1969) and the New England States (Seymour 1969) are doubtful as no specimens of this species have been seen from these regions. Also, several specimens previously identified as R. occidentalis from the maritimes, including a Macoun specimen from Campbellton, New Brunswick, have been identified as immature specimens of R. longifolius.

Rumex occidentalis var. labradoricus (Rechinger f.) Lepage, Nat.
Can. 81:66. 1954.

R. confinis Greene, Pittonia 4:306. 1901.

Type Specimen: Idaho: wet meadows around Lake Pend
d'Oreille. Leiberg 562.

R. fenestratus Greene, Pittonia 4:306. 1901.

Type Specimen: British Columbia: near Comox, in salt
marshes, Vancouver Island. 23 June 1893. Macoun. Herbarium
of Geological Survey Dept. No. 1570 (Holotype, CAN 434001)

R. procerus Greene, Pittonia 4:305. 1901.

No specimens cited.

Type locality: wet, boggy depressions among the coast
hills about San Francisco and Monterey.

Lectotype: California: Solano Co. common in marshes,
Goodyear. Baker 3219 (indicated as type by Rechinger 1937).

R. fenestratus var. labradoricus Rechinger f. Field Mus.
Nat. Hist. 17:115. 1937.

Lectotype: Labrador, Blanc Sallon, marshy belt back of
the strand, Sept. 3, 1910. Fernald and Wiegand 3283
(CAN 434131)

R. fenestratus subsp. puberulus Hulten, Lunds Univ. Arssk.
N.F. Avd, 2:40, nr 1:602. 1944.

Type Specimen: Alaska: Burroughs Bay, grassy river
flat. Walker 1019 (US 12004401)

R. occidentalis S. Watson var. procerus (Greene) Howell,
Leaflets of Western Botany 5:105. 1948.

R. occidentalis var. fenestratus (Greene) Lepage,
Naturalist Canadien 81(12):66. 1954.

Distribution:

Canada: BC, LB, QU, (NF). (The variety has been reported from Newfoundland (Rechinger 1937) but no specimens belonging to the variety as now delimited have been seen.)

United States: AK, CA, ID, OR, WA. (Map 5).

Habitat: Wet areas such as bogs, marshes, wet meadows, sloughs, along streams, etc. No differences in habitat appear to exist between the two varieties.

Discussion:

S. Watson (1877) proposed the name R. occidentalis for a native North American species that had previously been referred to as R. aquaticus not Linnaeus, R. longifolius not De Candolle or R. domesticus not Hartman. No specimens were cited in the original description and the lectotype was suggested by Rechinger in a 1953 annotation on the Bourgeau specimen; an examination of this specimen indicates its close agreement with Watson's description and it is here designated as the lectotype of R. occidentalis. Trelease (1892) accepted R. occidentalis in his treatment of North American Rumex; he also recognized its similarities to R. arcticus since he placed R. domesticus var. nanus Hooker (a synonym of R. arcticus) as a variety of R. occidentalis. In 1901, Greene described six western species with relationships to R. occidentalis: R. fenestratus, R. procerus, R. gracilipes, R. confinis, R. polyrhizus and R. bakeri; these names found little acceptance in taxonomic works of the period. However,

Rechinger (1937) recognized two closely related species in this complex: R. occidentalis occupying the widest range ("New England in the east to Montana (and occasionally still further) in the west and in the Rocky Mountains south to New Mexico") represented by "forms with an average length and breadth of valves of 5mm" and R. fenestratus occupying "exclusively or prevailing" a western zone "from western middle California in the south to Alaska in the north" represented by "forms with large fruits, the valves averaging 9mm in size". In Labrador and Quebec, he recognized variants with large fruits as a variety of R. fenestratus. R. fenestratus var. labradoricus, reportedly differing only slightly from the western forms as the valves were said to be often somewhat subtriangular in outline and frequently acute. Greene's R. procerus was reduced by Rechinger to a variety of R. fenestratus being limited in range to the western mid-section of California (i.e. southwestern part of R. fenestratus' range). Rechinger noted that both of these varieties were not clearly distinguished from the typical one. Of the other taxa described by Greene, Rechinger placed R. confinis as a synonym of R. fenestratus, whereas R. gracilipes, R. bakeri, R. polyrhizus were reduced to synonymy under R. occidentalis. Vegetative characters were found by Rechinger to be of no use in delimiting the two species or the various varieties of the species. In his key (Rechinger 1937) separated the species as follows:

Valves more than 7mm long

R. fenestratus

Valves to 5mm long

R. occidentalis

Additional characteristics on which the species were reported to differ were:

R. occidentalis: Valves truncate to weakly emarginate; achenes 3mm long by 1.5 mm wide.

R. fenestratus: Valves cordate to deeply emarginate; achenes 3.5 - 4mm long by 2mm wide.

A further infraspecific taxon was named within R. fenestratus by Hulten (1944) - R. fenestratus subsp. puberules differing from the typical subspecies the abaxial veins of the leaves being pubescent; it was said to be confined to Alaska. As this characteristic is observable on the type of R. fenestratus although to a lesser degree, Hulten's subspecies is reduced to a synonym of R. fenestratus. Howell (1948) recognizing the close relationship of R. fenestratus var. procerus with R. occidentalis, transferred it to that species as R. occidentalis var. procerus.

Lepage (1954) observed specimens that did not appear to be clearly referable to either species and noted that achene size did not correspond with base or valve size as had been reported. He believed that R. occidentalis and R. fenestratus represented the two extremes of one species consisting of a series of intermediate races. As R. occidentalis was the older name, the two species were included under that name and the intermediates were recognized as varieties. Lepage recognized: R. occidentalis var. occidentalis, var. fenestratus, var. labradoricus, var. procerus and var. perplexus. The last named

variety described by Lepage represented specimens with valves 5 - 6mm long and emarginate to truncate bases and achenes 3.5 - 4.0 mm long by 1.5 - 2.0 mm wide; in noting that it had the large achenes of var. fenestratus and the valves of var. occidentalis, Lepage suggested that this intermediate race might represent a hybrid between these varieties.

Boivin (1968) recognized two varieties under R. occidentalis: var. occidentalis with valves 4 - 6mm long and achenes 3 - 4mm long and var. labradoricus with valves 6 - 8 (-11) mm long and achenes 4 - 5mm long. He placed var. procerus in synonymy under var. labradoricus making it a disjunct variety represented in the eastern and western parts of R. occidentalis range.

As previously reported in the infraspecific numerical study, two closely related groups are recognizable within R. occidentalis. These groups have been recognized at the varietal level - the groups with valves up to 7mm long represented by R. occidentalis var. occidentalis and the group with valves greater than 7mm long represented by R. occidentalis var. labradoricus with the specimen of Fernald and Wiegand 3288 (CAN 43413) designated as the lectotype. The latter name was chosen over var. procerus (both var. procerus and var. labradoricus having been designated as varieties of R. fenestratus by Rechinger 1937)) because of the combination made by Boivin (1968) in which var. procerus was included as a synonym of var. labradoricus.

Although as noted in the numerical study, additional characters were found to be involved in the recognition of these taxa, but considerable overlap in the ranges of these characters (Table 10) was observed and individually, they can not be used to recognize the varieties.

Mar 5 Distribution of R. occidentalis.

4

Map 5

7.

Rumex arcticus Trautvetter ex Middendorff, Reise Nord. Osten Sibir.
1:29, 1847.

Type Specimen: (U.S.S.R.) ad fl. Boganida 71°0', Exped. Sibir.

Acad. 1843, Herb. Acad. Petrop. (LE, DAO photo!)

R. domesticus var. nanus Hooker, Fl. Bor. Amer. 2:129, 1840.

Type Specimen: Alaska: Arctic Sea coast, Dr. Richardson.

R. longifolius var. nanus (Hooker) Meisner ex de Candolle,

Prodr. 14:44, 1856.

R. occidentalis var. nanus (Hooker) Trelease, Miss. Bot.

Gard. Rep. 3:82, 1892.

Perennial. Root fusiform. Stem usually less than 1m tall, less than 1cm wide at base, moderately leafy, branched alternately above middle, erect, smooth to weakly sulcate, brown to red-brown or red-purple, glabrous to indistinctly papillose; trichomes papilliform, less than 0.1 mm long, shorter than wide; ochreae deciduous to partially persistent on mature plants. Branches short, the largest less than 5cm long, diverging at 30° - 40° to the main stem and often weakly arching. Basal leaves narrowly lanceolate, to narrowly oblong or narrowly ovate, acute to subacute, usually broadly cuneate to obtuse or rarely very weakly cordate, 3 - 7 times as long as wide; widest at or near base, weakly crisped; middle veins of lamina diverging at 40° - 65° to the midrib; veins commonly not raised to barely raised above the abaxial surface of the lamina; lamina glabrous above, commonly glabrous to indistinctly papillose on veins below; trichomes papilliform, less than 0.1 mm long, shorter

than to 2.5 times as long as wide; petiole shorter than lamina, commonly glabrous to indistinctly papillose. Lower cauline leaves similar to basal, but becoming progressively smaller up the stem. Upper cauline leaves lanceolate to linear-lanceolate, acuminate, narrowly cuneate, weakly crisped; lamina glabrous above and below; petiole much shorter than the lamina, the leaf sometimes subsessile. Inflorescence a narrow panicle, occupying the upper half of the stem, the lower to middle branches subtended by leaves, the upper branches without subtending leaves. Whorls consisting of ca 10 flowers, arranged on branches and portion of stem above branches, not subtended by bracts, the lower whorls remote, the upper whorls becoming somewhat contiguous. Pedicels filiform, 4.5 - 15.0 mm long, less than 0.2 mm wide at middle, articulated in the lower third; articulation indistinct, not swollen. Sepals oblong, obtuse, commonly at right angles to the pedicel. Valves ovate, obtuse to subacute, truncate to weakly emarginate, indented less than 0.8 mm at base, 4.3 - 7.6 mm long, 4.1 - 7.0 mm wide, widest at middle to just above the base, the margin entire to very weakly serrate; distinct teeth less than 4 per side, the largest less than 0.3 mm long; veins weakly raised, diverging from the midrib at ca 45° (-90°) with 1 - 2 anastomoses on each side of the valve; distance to the first anastomosis from the midrib 1.0 - 1.8 mm. Tubercles lacking. Anthers 1.2 - 1.4 mm long. Mature achenes ovate, red-brown, 3.5 - 4.1 mm long, 1.7 - 2.0 mm wide, 1.5 - 1.7 mm deep, acute at apex and base, somewhat more broadly so at base; the seed coat easily removed.

Distribution: Native to northern U.S.S.R. and North America, where it is confined to the northwest.

Canada: Northern BC, NT, YT. United States: AK. (Map 6)

Habitat: Occurring in marshes, snow beds, arctic and alpine tundra and woods.

Discussion: As discussed in the cytology section, R. arcticus is predominantly a tetraploid in North America, while in the Russian portion of the geographic range, higher ploidy levels appear to be more common. Mulligan and Frankton (1972) suggested that further cytological information on the Asiatic populations would be needed before the chromosome races could be recognized taxonomically. The present study is in agreement with Welsh (1974) that additional North American specimens are needed in order to better understand the within species variability of R. arcticus, since present herbaria material consists of very few mature basal rosettes or especially mature fruiting specimens.

R. arcticus has been often misidentified as R. occidentalis or even R. longifolius. Cytologically, R. arcticus ($2n = 40$) can be distinguished from R. occidentalis ($2n = 140$) and R. longifolius ($2n = 60$). The morphological characteristics separating R. arcticus from R. occidentalis, especially the basal leaves are set out in the key. A very limited survey of the pollen of these species (Crompton and Small, personal communication) indicated that pollen size may also be helpful in differentiating the two species.

	Pollen Diameter (μ)
<u>R. arcticus</u>	25.5 - 32.6
<u>R. occidentalis</u>	32.4 - 37.2

However, as reported by Mulligan and Frankton (1972), the pollen of the Asiatic populations of R. arcticus ($2n = 120 - ca\ 200$) was 33.0 to 39.5 in diameter. Morphologically, R. arcticus can be readily distinguished from R. longifolius. A comparison of the characteristics that can be used is presented in the following table (Table 21).

TABLE 21

Comparison of R. arcticus and R. longifolius

<u>Characteristic</u>	<u>R. arcticus</u>	<u>R. longifolius</u>
Basal Leaves	Narrowly lanceolate to narrowly oblong or ovate, 3-7 times as long as wide; abaxial veins commonly not raised and glabrous.	Lanceolate to oblong-lanceolate, 2-4 times as long as wide; abaxial veins raised and weakly to distinctly papillose.
Pedicel	Point of articulation not swollen.	Point of articulation distinctly swollen.
Valve Base	Truncate to weakly emarginate, indented less than 0.6 mm.	Cordate, indented (0.3-0.6 - 1.0 - 1.5 mm.
Tubercles	No tubercles present on a fruit.	Usually one small tubercle present on a fruit.
Seed Coat	Membranaceous and easily removed	Hard and not easily removed.

234

Map 6. Distribution of R. arcticus.

Map 6

8.

Rumex patientia Linnaeus, Sp. :l. 333. 1753.

Locality cited: "in Italia".

Type: not known (See discussion)

Perennial. Root fusiform. Stems commonly more than 1m tall, tallest, observed 165 cm, commonly more than 1cm wide at base, moderately leafy, branched alternately above middle, erect, sulcate, brown to red-brown or red-purple, indistinctly papillose to commonly glabrous; trichomes papilliform, evident on branches, less than 0.1 mm long, shorter than wide; ochreae deciduous to partially persistent on mature plants. Branches diverging at (20°-) 30° - 35° - 40° to the main stem and often weakly arching. Basal leaves large, largest observed 46.0 cm long, 12.4 cm wide, ovate-lanceolate to oblong-lanceolate, acute to subacute, broadly cuneate to truncate or rarely weakly cordate, 3 - 4 times as long as wide, weakly undulate, entire; middle veins of lamina diverging at 50° - 70° to the midrib, lamina glabrous above, glabrous to very indistinctly papillose on veins below; trichomes papilliform, less than 0.1 mm long, shorter than to as long as wide; petiole somewhat shorter than lamina. Upper cauline leaves elliptic to narrowly lanceolate, acute to acuminate, cuneate to narrowly cuneate, entire; glabrous above, very indistinctly papillose to commonly glabrous on veins below; petiole much shorter than lamina, the leaf sometimes sessile. Inflorescence, a narrow, dense panicle, occupying the upper half of the stem, all or all but the very upper branches subtended by leaves.

Whorls commonly of 10 - ca 20 flowers, arranged on branches and portion of main stem above branches, none or rarely the lowermost whorls subtended by bracts, the lower whorls remote, the upper contiguous. Bracts leaf-like, similar to the uppermost cauline leaves. Pedicels, filiform, (3.0-) 6.0 - 8.0 - 10.8 (-13.5) mm long, less than 0.3 mm wide at middle, articulated in the lower two-fifths; the articulation distinctly swollen. Sepals oblong, obtuse, (1.3-) 1.8 - 2.0 - 2.8 mm long and about half as wide, reflexed slightly downwards. Valves orbicular to weakly reniform, obtuse, subacute or rarely acute at apex, distinctly cordate at base, indented (0.5-) 0.8 - 1.3 - 1.5 (-2.0) mm at base, (4.0-) 6.0 - 7.3 - 8.5 (-9.5) mm long, (4.5-) 6.0 - 7.5 - 9.0 mm wide, widest at or near the middle, the margin entire to very weakly serrate; teeth 3 - 11 per side, the longest less than 0.1 mm long; veins raised, diverging from the midrib at 45° - 90° with 2 - 3 anastomoses on each side of the valve; distance to the first anastomosis from the midrib 1.0 - 1.5 - 1.8 mm. Tubercles 1 - 3, oblong to ovate, obtuse to acute at apex, obtuse at base, unequal in size, surface smooth but faintly reticulate; largest tubercle (1.1-) 1.6 - 2.1 - 2.8 (-3.5) mm long, (0.5-) 1.0 - 1.4 - 1.5 (-1.8) mm wide, (0.5-) 0.8 - 1.0 - 1.5 mm deep, widest in lower third to just below middle, base of tubercle at the valve base or up to 0.5 mm above, smallest tubercle 0.8 - 2.0 mm long, 0.3 - 0.9 mm wide, 0.5 - 1.3 mm deep, widest near base to below middle, base of tubercle at the valve base or up to 0.5 mm above. Anthers 1.0 - 1.3 mm long. Mature achene ovate, brown, 2.9 - 3.5 - 3.8 mm long, (1.4-) 1.8 - 2.0 - 2.3 (-2.5) mm wide, (1.3-) 1.5 -

1.8 - 1.9 (-2.2) mm deep, widest in the lower two-fifths, acute at apex, more broadly acute to obtuse at base.

Distribution: Native in eastern Europe, the Mediterranean region and southwestern Asia; introduced in the rest of continental Europe, the British Isles and North America (Lawalree 1953). In North America, infrequent in Canada and the United States.

Canada: ON, NU; United States: CT, ID, IL, IA, KS, ME, MA, MI, MO, NB, NH, NJ, NY, NC, ND, OK, PA, RI, UT, VA, WI. (Map 7)

Habitat: Occurring in waste places, fields, roadsides and along railroad tracks.

Discussion:

The type of this species is not known, nor can one readily be designated. An examination of the photograph of LINN No. 464.1 and the observations made on it by Dr. J. McNeill in 1975, confirm that it is referable to the species to which the name R. patientia has been applied. Although a specimen of R. patientia, it does not agree with all aspects of the protologue in the Species Plantarum in that it lacks tubercles and has cuneate leaves, whereas Linnaeus (1753) describes one valve as being tubercled and the leaves as cordate at the base. Consequently, Linnaeus cannot have based his account of R. patientia on LINN No. 464.1. An examination of the microfiche of the Linnaean specimens in the herbarium in the Riskmuseet of Stockholm failed to indicate a specimen of R. patientia. Other collections of Linnaean material must be examined to determine if a specimen of R. patientia suitable for lectotypification exists.

Rechinger (1937) reported that most American specimens of R. patientia "belong to subsp. patientia with only one valve grain-bearing; but some of them approaching by the still larger fruiting perianths, with the grains of unequal size, the subsp. orientalis (Bernh.) Danser." Subsp. patientia was reported to have valves 5 - 7 mm wide by Rechinger (1933, 1949) and 5 - 6 mm wide by Lousley (1936), whereas subsp. orientalis is said by these authors to have valves 8 - 10 mm wide. One valve of the fruit of subsp. patientia was said to have a tubercle, ca. 1.5 mm long and the other valves might possess rudimentary tubercles (a weakly thickened midrib), whereas subsp. orientalis was described as having fruit with one to three tubercles, the largest being 2 - 3 mm long (Rechinger 1933, 1949). In the most recently published key to the subspecies in Europe (Rechinger 1964), subsp. patientia is distinguished from subsp. orientalis only by its valves being less than 8mm wide and about the same long, with no mention being made of the previously noted differences in tubercle size and number.

Of the specimens identified by Rechinger as subsp. patientia, several were annotated as "approaching subsp. orientalis" - these have tubercles characteristic of subsp. orientalis as well as valves typically 8mm wide or slightly greater; other specimens exhibiting similar variation in valve width and tubercle size were annotated by Rechinger simply as subsp. patientia. In both sets of specimens, the valves are predominantly orbicular, a feature said to be typical of subsp. patientia; however, utilizing

the key in Rechinger (1964), these specimens would be referable to subsp. orientalis. As Rechinger (1933, 1949, 1964) did not vary the reported size range of the valve (length and width) of subsp. orientalis, the orbicular shape of the aforementioned specimens must have been sufficient to place them in subsp. patientia. Specimens with valves less than 8mm wide but with their tubercles more characteristic of subsp. orientalis have been examined, but based on Rechinger's (1964) key, these would key to subsp. patientia. Finally, specimens in agreement with the descriptions of subsp. patientia (Rechinger 1933, 1949, 1964) in both tubercle size and number and in valve size exist; these specimens give no indication of the two additional tubercles found in the other specimens. An examination of available European material referable to subsp. patientia did not indicate the tubercle variation apparent in the North American specimens annotated by Rechinger. Until the fruit characteristic delimiting each species within their native Eurasian ranges are more adequately described, it would be inadvisable to use subsp. orientalis and subsp. patientia for the North American specimens.

In 1953, Rechinger identified a specimen from New York as R. patientia subsp. recurvatus (Rechinger) Rechinger f., but unfortunately, the inflorescence is not fully mature. Subsp. recurvatus is reportedly separable from the two subspecies discussed above in that the valves are less than 8mm wide (ranging from 4.5 - 6.0 mm), longer than wide, acute at the apex and truncate or subcordate at the base, (Rechinger 1933, 1949, 1964). Although

the New York specimen had some mature valves that were acute at the apex, the valves varied from being longer than wide to shorter than wide, the width varied from 4.5 - 5.0 mm and the bases were typically cordate. This specimen exhibits more variation than has been reported for this subspecies in Europe; whether this is due to the specimen being atypical or reflects the normal variation of the subspecies has not been determined. No other specimens even approaching subsp. recurvatus have been observed in the North American material. As was the case with the subspecies discussed previously, a clearer delimitation of the variation of the subspecies in Europe is needed before a subspecific classification can be applied to the North American material.

At the present time, it is uncertain whether the existing subspecific classification is unsatisfactory or whether the subspecific differentiation that may be clear in Europe has become blurred in the North American population. In the differentiation of subsp. patientia and subsp. orientalis, the examination of available European material tends to indicate that at least in part, the latter may be the case. A detailed examination of the range of variation occurring within the native geographic range of R. patientia would be required to resolve this problem.

R. patientia has often been confused with R. crispus, R. longifolius and R. pseudonatronatus. Cytologically, R. patientia ($2n = 60$) differs from R. pseudonatronatus ($2n = 40$) but not from the other species which are also hexaploid. The morphological

characteristics distinguishing R. patientia from these species can be seen in the key to the species and in Table 22, under the discussion of R. crispus.

242

Map 7 Distribution of R. ratientia.

Map 7

9.
Rumex kerneri Borbas, Temeş Megye Vegetavja (Flora comitatus
 temesiensis) 34. 1884.

Type: Romania: Timişara: "Lunkasagi", Borbas? (See discussion)

Perennial. Root fusiform. Stem usually ca 1m tall, less than 1cm wide at base, leafy alternately branched in upper two-thirds, erect, sulcate, olive-green to red-brown, glabrous to indistinctly papillose trichomes most evident near whorls or branches, papillate to cylindrical, less than 0.3 mm long, usually less than 0.1mm long, 1/2 to 5 times as long as wide; ochreous deciduous to partially persistent on mature plants. Branches diverging at 30° - 40° - 50° to the main stem but often weakly arching. Basal leaves oblong-lanceolate, acute, usually obtuse to weakly cordate, 2 to 3 times as long as wide, entire to crisped; middle veins of lamina at 50° - 70° to the midrib; lamina glabrous above, indistinctly to moderately papillose on veins below; trichomes papilliform to cylindrical, less than 0.3mm long, shorter than to four times as long as wide; petiole as long as or longer the lamina, glabrous to indistinctly papillose near the lamina. Lower cauline leaves very similar to basal but becoming progressively smaller up the stem, petiole shorter than the lamina. Upper cauline leaves, oblong-elliptic to narrowly lanceolate; narrowly acute, narrowly cuneate, becoming progressively smaller up the stem; lamina glabrous above, indistinctly papillose to glabrous on veins below; petiole much shorter than lamina, indistinctly papillose to glabrous. Inflorescence a wide, open, lax panicle occupying up to two-

thirds of stem, the branches subtended by leaves. Whorls commonly consisting of ca 20 flowers arranged on branches and portion of main stem above branches, the lower whorls remote and often subtended by leaf-like bracts, the upper contiguous and not subtended by bracts. Bracts leaf-like, very similar to uppermost cauline leaves by becoming smaller up the stem. Pedicels filiform, (4.0-) 6.0 - 7.5 - 12.5 mm long, less than 0.3mm wide at middle, articulated at or near the middle articulation distinctly swollen. Sepals oblong, obtuse, (1.5-) 2.0 - 2.0 - 2.5 (-2.8)mm long and about half as wide, reflexed slightly downwards. Valves approximately orbicular, abruptly acute, cordate, indented 0.3 - 1.0 - 1.3 mm at base, (6.0-) 6.5 - 7.2 - 9.0 mm long, (6.0-) 6.5 - 7.0 9.0 mm wide, the margin below the apex serrate to weakly dentate; teeth 3 - 7 - 8 per side, the longest 0.3 - 0.5 - 1.0 mm long; veins raised, diverging from the midrib at ca 90°, with 3 - 4 (-5) anastomoses on each side of the valve; distance to the first anastomosis from the midrib 1.0 - 1.5 - 1.5 (-1.3) mm. Tubercles 3, oblong usually obtuse to acute at pex, obtuse at base, unequal in size, surface verrucose and faintly reticulate; largest tubercle 2.5 - 3.0 - 5.0 mm long, (0.5-) 1.0 - 1.4 - 2.5 mm wide, 1.0 - 1.3 - 2.5 mm deep, widest at or near the middle, base of tubercle at the valve base or up to 0.5 mm above; smallest tubercle 1.5 - 1.6 - 4.0 mm long, (0.5-) 1.9 - 1.2 - 2.0 mm wide, 0.6 - 1.3 - 2.0 mm deep, widest at or near the middle, base of tubercle 0.5 - 0.8 mm above the base of the valve. Anthers 1.3 - 2.0 mm long.

Mature achene ovate, red-brown, 3.0 - 3.3 0 3.5 (-3.8) mm long, (1.3-) 2.1 - 2.3 - 2.7 (-2.6) mm wide, (1.5-) 1.6 - 2.0 - 2.5 mm deep, widest just below the middle, acute at the apex, rounded at the base.

Distribution: Native in Europe, from Hungary and Romania to Central Greece. (Rechinger 1964). A rare introduction in North America only in southwestern California.

United States: CA (Map 8)

Habitat: Occurring in mainly ruderal areas, roadsides.

Discussion: In the type description of R. kernieri, Borbas included the following abbreviations "Vs., Vg., Bz., Hk., Lk!" for Versecz, Vinga, Buzias, Hidegkut = Guttebrun es Zabran, Lunkasag, Vinga and Buzias are towns in the region of Timisoara, the area of Borbas' flora while the other names probably represent towns/localities in the same area. The ! indicates a Borbas specimen from Lunkasag and this has been indicated as the probable Type location. Based on the type description, it is evident that the type must be a specimen from the Timisoara region in mid-western Romania.

R. Kernerii, is a relatively recent introduction to North America, being first collected in California by Pollard

in 1952; these specimens were identified as R. kernerii by Rechinger. Munz and Keck (1959) reported only one locality: Hope Ranch Santa Barbara Co., but examination of herbarium material now available has indicated that although still restricted to the same general region, R. kernerii now occupies several sites in Santa Barbara Co., as well as several sites in San Luis Obispo Co.

247

Map 8 Distribution of R. alpinus, R. confertus,
R. kernerii and R. sanguineus.

Map 6

10.

Rumex crispus Linnaeus, Sp. Pl., 335. 1753.

Lectotype: Hb. LINN 464.7 "9 crispus, Fl. Suec." (Linn, DAO photo!) (See discussion)

Perennial. Root fusiform. Stem to 150 cm tall, commonly ca 1m, less than 1cm wide at base, erect, smooth to weakly sulcate, brown to red-brown, often with red-purple areas, leafy, glabrous to indistinctly papillose; trichomes papilliform, most evident on upper stem and branches, less than 0.1 mm long, shorter than to as long as wide; ochreae totally deciduous to partially persistent on mature plants. Branches diverging 20° - 30° - 40° to the main stem and often weakly arching. Basal leaves, large, largest observed 29.5 cm long, 5.3 cm wide, oblong to linear-lanceolate, commonly acute to obtuse, commonly abruptly cuneate to rarely obtuse or very weakly cordate, 4 - 6 times as long as wide, widest at or near middle, undulate, crisped; middle veins of lamina leaving midrib at 50° - 70° (-90°); lamina glabrous above, indistinctly to moderately papillose on veins below; trichomes papilliform, less than 0.1 mm long, shorter than to as long as wide; petiole usually somewhat shorter to rarely longer than lamina, glabrous to indistinctly papillose. Lower cauline leaves similar to basal but progressively smaller up the stem. Upper cauline leaves linear-oblong, acuminate, narrowly cuneate, crisped, glabrous above, glabrous to indistinctly papillose on veins below; petiole much shorter than the lamina, the leaf sometimes subsessile. Inflorescence a narrow to wide, often dense panicle, occupying the upper half of the stem, all branches subtended by leaves. Whorls commonly consisting of ca 20 flowers arranged on branches and portion of main stem above branches, usually non-subtended

by bracts or occasionally only the lowermost subtended by leaf-like bracts; the lower whorls remote, the upper contiguous. Bracts leaf-like, very similar to uppermost cauline leaves, but becoming progressively smaller up the stem. Pedicels filiform, (2.0-) 4.5 - 6.3 - 8.0 (-9.5) mm long, ca 0.2 mm wide at middle, articulated in the lower two-fifths to the lower fifth, commonly in the lower third; articulation distinctly swollen. Sepals oblong, obtuse, commonly spreading at right angles to pedicel to reflexed weakly downwards. Valves orbicular-ovate to ovate-deltoid, commonly subacute to obtuse, truncate to commonly weakly cordate at base, indented at base 0.2 - 0.3 - 0.5 (-1.0) mm, (2.8-) 3.5 - 4.5 - 5.0 (-6.0) mm long, (2.5-) 3.5 - 4.5 - 5.5 (-7.5) mm wide, widest at middle to the lower third, the margin entire to very weakly serrate; distinct teeth 1 - 8 per side, the largest less than 0.3 mm long; veins raised, diverging from the midrib at ca 45° with (1-) 2 (-3) anastomoses on each side of the valve; distance to the first anastomosis from the midrib (0.8-) 0.9 - 1.0 - 1.3 mm. Tubercles 1 - 3, commonly 3, but often variable on same plant, oblong-ovate, obtuse to acute at apex, obtuse at base, unequal in size, surface smooth but faintly reticulate; largest tubercle (1.4-) 1.6 - 2.4 - 3.0 (-3.3) mm long, (0.6-) 1.0 - 1.3 - 1.5 (-2.0) mm wide, (0.5-) 0.6 - 1.0 - 1.3 (-1.5) mm deep, widest in lower two-fifths to lower fifth, the base of the tubercle usually at the valve base or protruding past base; smallest tubercle 1.0 - 2.0 (2.5) mm long, 0.5 - 1.2 mm wide, 0.5 - 1.0 mm deep, widest in lower third or rarely at middle, the base of the tubercle at the base of the valve to 0.5 mm above. Anthers ca 1 mm long. Mature achene ovate, red-brown (2.0-) 2.2 - 2.5 - 2.8 (3.0) mm long, (1.2-) 1.4 - 1.5 - 1.6 (-2.0) mm wide, 1.2 - 1.3 - 1.5 (-2.0) mm deep, widest

in the lower two-fifths, rarely to middle, acute at apex more broadly acute to obtuse at base.

Distribution: Native in the temperate regions of Eurasia, but the geographic limits are not known with certainty (Lawalree 1953); "introduced to all parts of the world" (Rechinger 1937); even occurring above the Arctic Circle in Norway and the Soviet Union (Holm et al. 1977) and just below the Arctic Circle in Alaska; in equatorial regions found "in the valleys and high places" (Holm et al. 1977). Rechinger (1937) noted that although R. crispus was considered as an introduced species in North America, he questioned whether or not it could be native in the Rocky Mountain areas of North America. Based on Hulten's (1968, 1971) maps, it appears that R. crispus is very rare in western Russia, Siberia and northern Asia; moreover, Hulten (1971), considers it an introduction in south-eastern Asia. The present Eurasian range of R. crispus would not appear to lend support to its being native in the Rocky Mountain region. R. crispus is widespread in Canada and the United States, but is less common in the Canadian prairies than in the other provinces and very rare in the far north.

Canada: AB, BC, MB, NK, NF, NS, ON, PE, QU, SK:

United States: AL, AK, AZ, AR, CA, CO, CT, DE, DC, FL, GA, ID, IL, IN, IA, KS, KY, LA, ME, MD, MA, MI, MN, MS, MO, MT, NB, NV, NH, NJ, NM, NY, NC, ND, OH, OK, OR, PA, RI, SC, SD, TN, TX, UT, VT, VA, WA, WV, WI, WY. (Map 9).

Habitat: Present on almost all types of soils, but less often on peat and rarely on acid soils (Holm et al. 1977); commonly occupying seasonally, most usually disturbed ground, e.g. around dwellings, on roadsides, in ditches, meadows, cultivated fields, etc., but also occasionally, on wet clearings, shores about ponds, lakes, streams, sloughs and in marshes.

Discussion: The lectotype of R. crispus (LINN 464.7) was designated by Lousley (1941). An examination of the Linnaean photograph of the specimen and observations made by Dr. J. McNeill in 1975, confirmed that this specimen agreed with the protologue (Linnaeus 1753).

R. crispus, an introduced species, is the most common and widespread member of section Rumex in North America. The earliest reference to the species occurring in North America appears to be that of Pursh (1814). Ascherson and Graebner (1900) attempted to deal with the infraspecific variation in Central Europe by recognizing ten varieties; Lousley (1941) stated that as few of the varying features observed in the species are consistently associated with other varying features, the classification of Ascherson and Graebner was purely artificial and of little value. Danser's (1921; cited in Lousley 1941) comment was "for the moment a natural classification of R. crispus seems to me to be an impossible task and an artificial superfluity". Lousley (1941) recognized four varieties as occurring in Britain. He did so largely because these four were ecologically distinct but even then, he considered this to be only a tentative treatment. In attempting to classify the variation, Rechinger (1949) recognized six varieties and later (Rechinger 1961) describes R. crispus as a "polymorphic species evidently comprising several races characterized by slight morphological differences, some of them also

ecologically different". He goes on to say: "All these races are so evidently interfertile and some are extremely invasive and by their extensive spread and interbreeding, they seem to have obscured the original distributional pattern of the other races". In *Flora Europaea*, Rechinger (1964) does not recognize any of his earlier infraspecific taxa in R. crispus, probably for the fore-mentioned reasons. Since R. crispus is an introduced species and as the infraspecific taxa have not been accepted in the area of origin, no attempt has been made to recognize such taxa in North America.

Confusion exists between R. crispus and R. longifolius, R. patientia, R. pseudonatronatus and R. obtusifolius. R. crispus ($2n = 60$) differs cytologically from R. pseudonatronatus and R. obtusifolius, each with $2n = 40$; the other two species, like R. crispus, have $2n = 60$.

The morphologically characteristics used to distinguish R. crispus from the other species have been tabled (Table 22, 23). Of the four species, R. obtusifolius, is the most readily distinguishable from R. crispus, followed by R. longifolius and R. pseudonatronatus. R. patientia is the most difficult to separate from R. crispus.

TABLE 22

Morphological Comparison of R. crispus
and R. obtusifolius and
R. patientia

Characteristic	<u>R. crispus</u>	<u>R. obtusifolius</u>	<u>R. patientia</u>
Basal Leaf Shape	Oblong to linear-lanceolate, 4-6 times as long as wide	*Broadly ovate to oblong-ovate, 1.7-2.5 times as long as wide	*Ovate-lanceolate to 3-4 times as wide
Basal Leaf Base	Commonly abruptly cuneate	*Commonly distinctly cordate	Commonly broadly cuneate to truncate
Basal Leaf Pubescence (on abaxial veins)	Indistinctly to moderately papillose, trichomes papilliform, less than 0.1 mm long	*Commonly distinctly pubescent to papillose, trichomes papilliform to cylindrical, less than 0.3 mm long	Glabrous to very indistinctly papillose, trichomes papilliform, less than 0.1 mm long
Valve Outline	Ovate	*Deltoid	*Orbicular to weakly reniform
Valve Base	Commonly weakly cordate, indented at base 0.2-0.5 (-1.0) mm	Truncate to weakly emarginate indented at base to 0.5 mm	*Distinctly cordate indented at base (0.5-) 0.8-1.5 (-2.0) mm
Valve Width	(2.5-) 3.5-5.5 (-7.5) mm	*(2.0-) 2.5-3.5 (-4.0) mm	*(4.5-) 6.0-9.0 mm
Valve Margin	Entire to weakly serrate with 1-5 teeth per side and the largest less than 0.3 mm long	*Dentate-spiny, with 3-5 teeth per side and the largest usually 1.0-1.5 mm long	Entire to weakly serrate with 3-11 teeth per side and the largest less than 0.1 mm long
Tubercle No.	1 - 3	1 - 3	1 - 3

TABLE 22, cont.

<u>Characteristic</u>	<u>R. crispus</u>	<u>R. obtusifolius</u>	<u>R. patientia</u>
Large Tubercle Size	(1.4-)1.6-3.0 (-3.3) mm long by (0.8-)1.0-1.5 (-2.0) mm wide	(1.1-)1.5-2.5 (-2.9) mm long by (0.5-)0.8-1.3 (-1.5) mm wide	(1.1-)1.6-2.8 (-3.5) mm long by (0.5-)1.0-1.5 (-1.8) mm wide
Small Tubercle Size	1.0-2.0(-2.5) mm long by 0.5-1.2 mm wide	*0.5-1.5 mm long by 0.3-0.5 mm wide	0.8-2.0 mm long by 0.3-0.9 mm wide
Achene Colour	Red-brown	Red-brown	*Deep brown
Achene Size	(2.0-)2.2-2.6 (-3.0) mm long by (1.2-)1.4-1.6 (-2.0) mm wide	(1.8-)2.3-2.9 (-3.0) mm long by (1.2-)1.3-1.7 (-1.8) mm wide	*2.9- 3.8 mm long by (1.4-)1.8-2.3 (-2.5) mm wide

* = good separation from R. crispus.

TABLE 23

Morphological Comparison of R. crispus and
R. longifolius and
R. pseudonatronatus

Characteristic	<u>R. crispus</u>	<u>R. longifolius</u>	<u>R. pseudonatronatus</u>
Basal Leaf Shape	Oblong to linear lanceolate, 4-6 times as long as wide	*Lanceolate to oblong-lanceolate, 2-4 times as long as wide	*Narrowly lanceolate, greater than 4 times as long as wide
Basal Leaf Base	Commonly abruptly cuneate	Commonly broadly cuneate	*Narrowly cuneate
Basal Leaf Pubescence (on abaxial veins)	Indistinctly to moderately papillose, trichomes papilliform, less than 0.1 mm long	Distinctly papillose, trichomes papilliform, less than 0.1 mm long	Indistinctly papillos, trichomes papilliform, less than 0.1 mm long
Valves Outline	Ovate	*Orbicular to weakly reniform	*Orbicular to weakly reniform
Valve Base	Commonly weakly cordate, indented at base 0.2-0.5 (-1.0) mm	Distinctly cordate, indented at base (0.3-)0.6-1.5 mm	*Cordate, indented at base, (0.2-) 0.4-0.6 mm
Valve Width	(2.5-)3.5-5.5 (-7.5) mm	*(4.0-)6.0-7.5 (-9.5) mm	(2.3-)3.0-4.9 (-6.0) mm
Valve Margin	Entire to weakly serrate with 1-3 teeth per side and the largest less than 0.3 mm long	Entire to weakly serrate with 2-6 teeth per side and the largest less than 0.3 mm long	Entire to weakly serrate with 1-3 teeth per side and the largest less than 0.1 mm long
Suberle No.	1 - 3	*0 - 1	*0 - 1

TABLE 23, cont.

Characteristic	<u>P. crispus</u>	<u>P. longifolius</u>	<u>P. pseudonatronatus</u>
Large Tubercle Size	(1.4-)1.6-3.0 (-3.3) mm long by (0.6-)1.0-1.5 (-2.0) mm wide	*(0.3-)0.5-1.3 (-1.4) mm long by (0.3-)0.4-1.0 mm wide	*0.4-1.1 mm long by 0.2-0.8 mm wide
Small Tubercle Size	1.0-2.0(-2.5) mm long by 0.5- 1.2 mm wide		
Achene Colour	Red-brown	*Deep-brown	Red-brown
Achene Size	(2.0-)2.2-2.8 (-3.0) mm long by (1.2-)1.4-1.6 (-2.0) mm wide	*(2.5-)2.6-3.3 (-3.5) mm long by (1.3-)1.5-1.8 (-2.0) mm wide	(1.3-)2.1-2.6 (-3.6) mm long by (0.8-)1.2-1.4 (-1.5) mm wide

* = good separation from P. crispus

257

Map 9 Distribution of R. crispus.

Map 9

11.

Rumex confertus Willdenow, Enum. Plant. Hort. Reg. Bot. Berol., 397, 1809.

Type: a cultivated specimen taken from the Berlin Garden.

Perennial. Rootstock fusiform. Stems ca 1m tall, ca 1cm wide at base, erect, sulcate, red-brown, leafy, alternately branched above the middle, weakly to distinctly pubescent; trichomes cylindrical, evident on upper stem and branches, less than 3mm long, 4 - 10 times as long as wide; ochreous usually deciduous on mature plants. Branches diverging at 25° - 40° to the main stem and often weakly arching. Basal leaves triangular-ovate, obtuse to subacute, deeply and widely cordate, 1.3 - 1.6 times as long as wide, widest at or near base, weakly crisped; middle veins of lamina diverging at (50° -) 60° (-80°) to midrib; lamina glabrous above, distinctly pubescent on veins below; trichomes cylindrical, less than 0.4 mm long, 4 - 8 times as long as wide; petiole 1 - 2 times as long as the lamina. Lower cauline leaves very similar to basal but becoming progressively smaller up the stem. Upper cauline leaves narrowly ovate to oblong, acute, weakly cordate to cuneate, becoming progressively smaller up the stem; lamina glabrous above, less distinctly pubescent on veins below than lower or basal leaves; trichomes cylindrical; petiole much shorter than the lamina, glabrous to weakly pubescent. Inflorescence a narrow open panicle occupying the upper half of the stem, the branches subtended by leaves. Whorls commonly consisting of more than 20 flowers arranged on branches and portion of main stem above branches, the lower whorls remote, the upper almost contiguous, none of the whorls subtended by bract-like leaves. Pedicels filiform, 4.5 - 10.8 mm long, less than 0.3 mm wide at middle,

articulated in the lower two-fifths, commonly in the lower third; articulation distinctly swollen. Sepals oblong, obtuse, 1.5 - 3.1 mm long and about half as wide. Valves reniform, abruptly acute, weakly to distinctly cordate, indented 0.2 - 1.7 mm at base, 6.0 - 9.5 mm long, 6.6 - 11.0 mm wide, the margin below the apex weakly crenate to rarely weakly serrate; teeth less than 8 per side, the largest less than 0.3 mm long; veins diverging from the midrib at ca 90° with 2 - 3 anastomoses on each side of valve; distance to the first anastomosis from the midrib 1.3 - 2.3 mm. Tubercles 1, ovate, obtuse to acute at apex, obtuse at base, 1 - 2.3 mm long, 0.5 - 1.4 mm wide, 0.6 - 1.4 mm deep, widest at the middle; base of tubercle (0.2-) 0.5 (-1.0) mm above base of valve. Anthers usually less than 1 mm in length. Mature achene, ovate, red-brown, 3 - 3.4 mm long, 1.7 - 2.3 mm wide, 1.5 - 2.0 mm deep, widest at or near the middle, acute at the apex, cuneate to obtuse at the base.

Distribution: Native in eastern and east-central Europe (Rechinger 1964) to the Caucasus, and most of Russia (Lousley 1939). A rare introduction in North America.

Canada: AB, MB. United States: ND (Map 8)

Habitat: Occurring in roadside areas, mainly ruderal.

Discussion: The first known collection of this species in North America was made by Scoggan in 1953 at Ethelbert, Manitoba. It was then mistaken for R. petientia; the specimens were correctly identified in 1964, by Frankton (DAO). The first published report of its occurrence in North America was by Boivin (1968). The present study has revealed two additional North American sites for the

species: one in North Dakota and one in Alberta; the former dates from 1955 but the latter is a relatively recent collection made in 1971. At first glance, the species can be mistaken for R. alpinus or R. patientia, but closer examination of its basal leaves or fruits shows its distinctiveness. From R. alpinus, it can be separated by its ovate-triangular basal and lower leaves in which the length is slightly greater than the width, while in R. alpinus, these leaves are orbicular and the length and width are approximately equal. In addition, the valves of the fruit are reniform with an abruptly acute apex and one valve is tubercled, whereas in R. alpinus the valves are ovate to deltoid with an acute apex and lack tubercles. R. confertus can be distinguished from R. patientia as the latter has oblong-lanceolate basal and lower leaves which are rounded, broadly cuneate or rarely weakly cordate at the base, rather than deeply and widely cordate as is the leaf-base in R. confertus leaves. The abaxial leaf surfaces of R. confertus, are weakly to distinctly pubescent, whereas in R. patientia the abaxial surfaces are glabrous to indistinctly papillose. The valves of the fruit of R. patientia are orbicular to weakly reniform, commonly obtuse at the apex, whereas those of R. confertus, are more distinctly reniform and abruptly acute at apex. As R. patientia can have more than one tubercle, it would be readily-separable from R. confertus, which only has one tubercle.

12.

Rumex alpinus Linnaeus, Sp. Pl. 334, 1753.

Localities cited: "in Helvetia, Monspeli" (in Switzerland (and) at Montpellier (France))

Linnaean specimen: No. 464.35, labelled "alpinus" and "H.U," (LINN photo!) (See discussion)

Perennial. Rootstock creeping. Stems ca 1m tall, ca 1cm wide at base, erect, sulcate, brown to red-brown, leafy, branched alternately above middle, glabrous or weakly to distinctly pubescent; trichomes cylindrical, evident on upper stem and branches, less than 0.1 mm long, longer than to as long as wide; ochreae deciduous to usually weakly persistent on mature plants. Branches diverging at 20° - 45° to the main stem and often weakly arching. Basal leaves large, the largest observed 32.2 cm long, 27.2 cm wide, orbicular to broadly ovate, as long as or longer than wide, widest near middle to above base, obtuse, deeply and widely cordate, entire; middle veins of lamina diverging at 40° - 70° to the midrib; commonly at 50° ; lamina glabrous above, distinctly pubescent on veins below; trichomes cylindrical, less than 0.3 mm long, 2 to 5 times as long as wide; petiole longer than lamina, glabrous to distinctly pubescent toward the lamina. Lower cauline leaves very similar to basal but becoming progressively smaller up the stem; petiole shorter than lamina. Upper cauline leaves oblong-lanceolate, acute, broadly cuneate, entire, glabrous above, weakly to distinctly pubescent on veins below; petiole much shorter than lamina. Inflorescence a wide panicle, occupying the upper half of the stem, all but the very upper branches subtended by leaves. Whorls commonly consisting

of 10 - 20 flowers, present on branches and portion of main stem above branches, not subtended by leaf-like bracts, the lower whorls remote, the upper almost contiguous. Pedicels filiform, 3.3 - 9.6 mm long, less than 0.3 mm wide at middle, articulated in the lower quarter; articulation distinctly swollen. Sepals oblong, obtuse, at right angles to pedicel, 1 - 1.5 mm long and about half as wide. Valves ovate-triangular, acute, truncate to weakly cordate at base, 2.6 - 5.1 mm long, 2.5 - 5.0 mm wide, widest at middle to just above base, the margin below the apex weakly and irregular serrate; distinct teeth 1 - 4 per side, the longest typically less than 0.4 mm long; veins raised, diverging from the midrib at ca 45° with 1 - 2 anastomoses on each side of the valve; distance to first anastomosis from the midrib 0.5 - 1.0 mm. Tubercles lacking. Anthers 0.6 - 1.0 mm long. Mature achenes elliptic, brown, 2.0 - 3.3 mm long, 1.0 - 1.9 mm wide, 1.0 - 1.9 mm deep, widest at or near middle, acute at the apex and base; apex often narrower than base.

Distribution: Native to central and southern Europe (Rechinger 1964) Caucasasia and W. Iran (Cullen 1967). A rare introduction in north-eastern North America.

Canada: NS. United States: ME, VT. (Map 8)

Habitat: In Europe reported as being strongly nitrophile (Lousley 1944, Rechinger 1964); in North America reported from along roadsides and in old fields.

Discussion: The Linnaean specimen No. 464.35, labelled "alpinus" and "H.U.", in the hand-writing of Linnaeus, consists of one small

leaf and a portion of an inflorescence that is only in flower; the Linnaean photograph and observations made by Dr. J. McNeill in 1975 confirmed that it was R. alpinus. However, this specimen cannot provide any fruit characteristics - "valvulis integerrimis graniferis" as noted in the protologue (Linnaeus 1753): the protologue cannot have been drawn up from this specimen alone and therefore, it cannot be considered as the type as proposed by Losin-Losinskaya (1936). Whether or not a fruiting specimen of R. alpinus, seen by Linnaeus and in agreement with the protologue has survived, is not known.

In 1920, Fernald (1921) reported finding R. alpinus in the Rockville area of Yarmouth Co., Nova Scotia and this was noted by Rechinger (1937). Several confirmed Nova Scotian sites as well as one each from Vermont and Maine are now known (Map 8). One of the specimens from Maine, previously misidentified as R. patientia, represents a collection made in 1892 and it is possible that the species has had a slightly larger but still sparse distribution in the north-east. Lousley (1944) reported that in England, R. alpinus is often a relic of cultivation having been introduced for its herbal qualities; this may also apply in North America. A 1974 collecting trip in the Rockville area, Nova Scotia, failed to find the species and no information is available on the present status of R. alpinus. Its ability to persist vegetatively may allow it to remain in some localities. The species is included in most floristic treatments that cover Nova Scotia and the New England states (Fernald 1950, Gleason and Cronquist 1963, Gleason 1968, Roland and Smith 1969, Seymore 1969) although they do not indicate the full North American range.

Rumex longifolius de Candolle, in Lamarck and DeCandolle, Fl. Fr. ed. 3,5: 368. 1815.

Type: (France) Pyr. orient. M. Coder c. No. 183. 1814 (G-DC No. 231, microfiche!) (See discussion)

R. domesticus Hartman, Scand. Fl. ed 1:148. 1820

R. patientia auct. amer. prior, non L.

Perennial. Root fusiform. Stem 45 - 150 cm tall, usually around 1 cm wide at base, erect, sulcate, leafy, branched alternately above middle, red-brown to red-purple, glabrous to indistinctly papillose; trichomes most evident on branches, papilliform, less than 0.1 mm long, shorter than to as long as wide; ochreous deciduous to partially persistent on mature plants. Branches diverging at 25° - 40° to the main stem, most usually at 30°, but often weakly arching. Basal leaves lanceolate to oblong-lanceolate, acute to subacute, cuneate to rounded or very weakly cordate, 2 to 4 times as long as wide, slightly undulate to weakly crisped; middle veins of lamina diverging at (40°-) 50° - 70° to the midrib; lamina glabrous above, weakly to distinctly papillose on veins below; trichomes papilliform, less than 0.1 mm long, longer than wide, petiole usually longer than the lamina, glabrous. Lower cauline leaves very similar to basal but becoming progressively smaller up the stem; petiole shorter than lamina. Upper cauline leaves progressively more linear-lanceolate, narrowly acute, narrowly cuneate, becoming progressively smaller up the stem; lamina glabrous above, weakly to moderately papillose on veins below; trichomes papilliform, less than 0.1 mm long, shorter

to as long as wide; petiole much shorter than the lamina, glabrous, the leaf sometimes sessile. Inflorescence a narrow, dense panicle, occupying the upper half of the stem, the branches subtended by leaves. Whorls commonly consisting of 10 - 20 flowers, present on the branches and the main stem above the branches, the lower whorls remote and rarely subtended by leaf-like bracts, the upper contiguous and not subtended by leaf-like bracts. Pedicels filiform, (3-) 4.8 - 5.8 - 9.0 (-9.3) mm long, less than 0.2 mm wide at middle, articulated in the lower two-fifths of the pedicel, usually in the lower third; articulation distinctly swollen. Sepals oblong, obtuse (-0.8) 1.5 - 1.5 - 2.0 (-2.3) mm long, and about half as wide, angled slightly downwards. Valves orbicular to weakly reniform, obtuse, cordate, indented (0.3-) 0.6 - 1.0 - 1.5 mm at the base, (-3.5) 5.2 - 6.0 - 6.5 (-7.5) mm long, (4.0-) 6.0 - 6.8 - 7.5 (-9.5) mm wide, the margin entire to weakly serrate; teeth 2 - 6 per side, the longest less than 0.3 mm. Veins raised, diverging from the midrib at 45° - ca 90° with 2 - 3 anastomoses on each side of the valve; distance to the first anastomosis from the midrib (0.8-) 1.0 - 1.0 - 1.3 (-1.5) mm. Tubercles 0 or 1, variable even on same plant, oblong, obtuse at apex and base, (0.3-) 0.5 - 1.3 (-1.4) mm long, (1.3-) 0.4 - 1.0 mm wide, (0.1-) 0.3 - 0.8 (-1.0) mm deep, widest at or near middle; base of tubercle 0.3 - 0.5 mm above the valve base; surface smooth and faintly reticulate. Anthers (0.9-) 1.0 - 1.3 mm long. Mature achene ovate, brown (2.5-) 2.8 - 3.0 - 3.3 (-3.5) mm long, (1.3-) 1.5 - 1.6 - 1.8 (-2.0) mm wide, (1.0-) 1.3 - 1.4 - 1.6 (-1.8) mm deep, widest at or near middle, acute at the apex and base, apex narrower than the base.

Distribution: Native in the Pyrenees, northern Europe and most of Russia. (Losin-Losinskaya 1936, Rechinger 1964). Introduced in North America and most common in the northeast.

Canada: AB, BC, MB, NK, NF, NS, ON, PE, QU, SK. United States: AK, CT, ME, MA, MN, MH, ND, RI, VT, WI. (Map 10).

Habitat: Occurring in ruderal areas, roadsides, cultivated fields, usually on moist soil.

Discussion: De Candolle (1815) based his R. longifolius on a specimen sent to him by Coder from the "environs de Prades, in Roussillon". Lousley (1941) reported that in 1939, a request for a loan of this specimen from the de Candolle Herbarium in Geneva was met with the response that "we have no specimen of R. longifolius from France. De Candolle remarks that Coder has sent him that specimen, but it is likely that De Candolle sent the plant back without keeping a duplicate." However, the microfiche of the de Candolle Prodromus Herbarium contained a specimen of R. longifolius that had been annotated by Rechinger in 1946 as the type; the earlier confusion may have arisen from the fact that the collection site was given as "Pyr. orient", which may have been confused with Syr. orient or simply not recognized as part of France. Prades is located in the province of Pyrenees - Orientales while the historic name for the province used prior to 1790 (Hammond Atlas 1976) was Roussillon.

In North America, early authors confused R. longifolius with R. patientia. The R. patientia of Trelease (1892), Britton and Brown (1896) and Rydberg (1932) were all cited by Rechinger (1937).

under R. patientia, but these at least in part represent R. longifolius. Scoggan (1957) also applies the name R. patientia to plants of R. longifolius. In 1929, Rechinger placed R. domesticus Hartm. in synonymy under R. longifolius, but in 1937, he took up the name R. domesticus, again ignoring de Candolle's earlier R. longifolius. Still later, based on his annotation of the de Candolle specimen in 1946, (Rechinger 1948), he restored the name R. longifolius and this has been accepted in more recent taxonomic treatments of Rumex.

Since Rechinger's (1937) treatment, the geographic range of this species is larger, but whether this is due to an active expansion by the species, our increased knowledge of R. longifolius, or combination, both is uncertain.

Map 10 Distribution of R. longifolius

Map 10.

14.

Rumex pseudonatronatus Borbas, Ertek Term. Koreb. Magyar Tud.

Acad. 11:21 (see also p. 63) 1880.

Type: Hungary: Bekes Co., "in pseudonatronatus silvae Fas territorii Koros-Ladany passim copiosum fructiferumque legi Medio Augusto". Borbas (Holotype ? BP)

R. domesticus var. pseudonatronatus Borbas, Ertek Term.

Koreb. Magyar Tud. Acad. 11:63. 1880 (obligate synonym)

R. fennicus Murbeck, Bot. Not. 17:230. 1899.

Perennial. Root fusiform. Stem ca 1(-1.5) m tall, less than 1 cm wide at base, leafy, alternately branched above the middle, erect, sulcate, brown to red-brown, glabrous to very indistinctly papillose; trichomes most evident on branches, papilliform, less than 0.1 mm long, shorter than to as long as wide; ochreae deciduous to partially persistent on mature plants. Branches diverging at 20° - 30° - 40° (-50°) to the main stem but often weakly arching. Basal leaves lanceolate to narrowly lanceolate, narrowly acute, narrowly cuneate, narrowing almost equally at apex and base, greater than 4 times as long as wide, weakly crisped; middle veins of lamina diverging at 50° - 70° to the midrib; lamina glabrous above, indistinctly papillose on veins below; trichomes papilliform, less than 0.1 mm long, shorter than to as long as wide; petiole shorter than the lamina, glabrous to indistinctly papillose. Lower cauline leaves very similar to basal but becoming progressively smaller up the stem. Upper cauline leaves narrowly lanceolate, narrowly acute, narrowly cuneate, becoming progressively smaller up the stem; lamina glabrous above, indistinctly papillose

to glabrous on veins below; petiole much shorter than the lamina, the leaf sometimes subsessile. Inflorescence a narrow panicle, occupying the upper half of the stem, the branches subtended by leaves. Whorls commonly consisting of more than 20 flowers, the lower whorls remote and occasionally subtended by leaf-like bracts, the upper contiguous and not subtended by bracts. Bracts leaf-like, very similar to the uppermost cauline leaves but progressively smaller up the stem. Pedicels filiform, (2-) 3.0 - 6.3 - 7.0 (-8.5) mm long, less than 0.2 mm wide at middle, articulated in the lower two-fifths of the pedicel, commonly in the lower third; the articulation distinctly swollen. Sepals oblong, obtuse, (0.7-) 0.8 - 1.0 - 1.3 (-1.5) mm at the base, (2.0-) 3.8 - 4.0 - 4.9 (6.0) mm long, (2.3-) 4.0 - 4.5 - 5.3 (6.0) mm wide, the margin entire to weakly serrate; teeth 1 - 3 per side, the largest less than 0.1 mm long; veins raised, diverging from the midrib at ca 90° with (1-) 2 (-3) anastomoses on each side of the valve; distance to the first anastomosis from the midrib (0.5-) 0.8 - 1.0 - 1.3 mm. Tubercles 0 to 1, variable on a single plant, oblong, obtuse to subacute at apex, obtuse at base, 0.4 - 1.1 mm long, 0.2 - 0.8 mm wide, 0.2 - 0.6 mm deep, widest at middle, the base 0.2 - 0.5 mm above valve base, the surface smooth but faintly reticulate. Anthers ca 1 mm long. Mature achene ovate, red-brown (1.3-) 2.1 - 2.4 - 2.6 (-2.8) mm long, (0.8-) 1.2 - 1.3 - 1.4 (-1.5) mm wide, (0.8-) 1.0 - 1.2 - 1.3 mm deep, widest at or near the middle, acute at the apex and base, the apex narrower than the base.

Distribution: Native in eastern Europe, extending westward to eastern Sweden and eastern Austria (Rechinger 1964) and in parts of Siberia and Central Asia (Losin-Losinskaya 1936). Introduced in North America, predominantly in the prairie region.

Canada: AB, BC, MB, ON, SK, YT; United States: MN, ND, SD; to be expected in adjacent areas. (Map 11).

Habitat: Reported in Europe from riversides and seashores, more or less halophilous (Rechinger 1964). In North America, found in ruderal areas such as roadsides, low damp areas such as ditches and extending into cultivated fields.

Discussion: Although the species was collected in North America as early as 1908, R. pseudonatronatus is not included in Rechinger's (1937) account of the North American species as the earliest collections were misidentified as other species. The first American specimen collected in North Dakota in 1908, was misidentified as R. crispus and the earliest Canadian specimen collected in 1939 in Manitoba, was first called R. altissimus and then R. crispus.

Even after recognition of the species occurrence, it was generally known in Canada as R. fennicus (Budd and Best 1964; Bovin 1966, 1968), a name now regarded as a synonym of R. pseudonatronatus (Rechinger 1964). Although having occupied its present midwestern range for many years, R. pseudonatronatus, even under its synonym R. fennicus, is not included in any United States regional or state floras. (McNeill 1976). Most United States specimens have been incorrectly identified as R. domesticus Hartm.,

and this is the name used for example by McGregor, Barkley et al. (1977). R. domesticus Hartm., is a synonym of R. longifolius DC., (Rechinger 1929, 1964), from which R. pseudonatronatus can be distinguished by the features given below.

Specimens of this species can be confused with R. crispus, R. longifolius or even R. occidentalis. Cytologically, R. pseudonatronatus ($2n = 40$) can be separated from these species - R. crispus ($2n = 60$), R. longifolius ($2n = 60$) and R. occidentalis ($2n = 140$) (Table 10).

Morphologically, a number of characteristics can be used to distinguish this species from all of the four mentioned species by its narrowly lanceolate basal leaves, narrowing almost equally to the apex and base; from R. occidentalis by the distinctly swollen articulation and/or the presence of one small tubercle; from R. longifolius on the basis of its smaller valves and smaller achenes; from R. crispus by its orbicular to weakly reniform valves and its smaller single tubercle.

273

Map 11 Distribution of R. pseudonatronatus.

Map 11.

15.

Rumex conglomeratus Murray, Prodr. Stirp. Gotting. 52, 1770.

Protologue: Lapathum petiolis latescentibus foliis longe lanceolatis, floribus verticillatis verrucosis.

No type specimen indicated in the description, but according to Losinskaya (1936) described from Switzerland.

Perennial. Root fusiform. Stem usually ca 1.0 (-1.65) m tall, less than 1 cm wide at base, leafy, alternately branched in the upper two-thirds of the stem, erect, sulcate, brown or red-purple, glabrous to rarely indistinctly papillose; trichomes most evident on branches, papilliform, less than 0.1 mm long, shorter than to as long as wide; ochreous deciduous to partially persistent on mature plants. Branches diverging at 30° - 60° to the main stem, usually ca 40° but often weakly arching. Basal leaves oblong to oblong-lanceolate, obtuse to broadly acute, broadly cuneate to rounded, 3 to 5 times as long as wide, entire; middle veins of lamina diverging at 50° - 70° to the midrib; lamina glabrous above, glabrous to occasionally indistinctly papillose on veins below; trichomes papilliform less than 0.1 mm long, shorter than to as long as wide; petiole usually longer than the lamina, glabrous. Lower cauline leaves very similar to basal but becoming progressively smaller up the stem; the petiole shorter than the lamina. Upper cauline leaves lanceolate to linear-lanceolate, narrowly acute, narrowly cuneate, becoming progressively smaller up the stem, entire, glabrous above and below; petiole much shorter than the lamina, the leaf sometimes sessile. Inflorescence a wide, open

open, lax panicle, occupying as much as the upper two-thirds of the stem, the branches subtended by leaves. Whorls commonly consisting of (10-) ca 20 flowers, arranged on branches and main stem above branches, the lower whorls remote, the upper becoming less so, usually all to all but the uppermost whorls subtended by leaf-like bracts very similar to the uppermost cauline leaves. Pedicels filiform, (1.0-) 1.8 - 2.8 - 4.0 (-5.5) mm long, less than 0.3 mm wide at middle, articulated in lower third to lower two-fifths of pedicel, rarely at middle articulation distinctly swollen. Sepals oblong, obtuse (0.6-) 0.8 - 1.2 - 1.5 (-1.8) mm long and about half as wide, recurved upwards around base of valve. Valves oblong to ovate, obtuse, rounded at the base, (1.8-) 2.3 - 2.7 - 2.9 (-3.0) mm long, (1.0-) 1.2 - 1.3 - 1.5 (-1.8) mm wide, widest below middle, entire to rarely one indistinct serration near base; teeth less than 0.1 mm long; veins weak, diverging from the midrib at ca 90° with 1 - 2, usually 1 anastomoses on each side of the valve; distance to the first anastomosis from the midrib (0.3-) 0.4 - 0.5 - 0.7 mm. Tubercles 3, oblong to ovate, obtuse at apex and base, approximately equal in size, (0.9-) 1.3 - 1.7 - 1.8 (-2.0) mm long, (0.6-) 0.8 - 1.0 - 1.0 (-1.1) mm wide, (0.5-) 0.7 - 0.8 - 1.0 (-1.1) mm deep, widest near middle to near base, base of tubercle usually at the base of the valve to 0.5 mm above, the surface smooth but faintly reticulate. Anthers usually less than 1 mm long. Mature achene ovate, dark red-brown to black (1.3-) 1.5 - 1.8 - 1.9 (-2.1) mm long, (1.0-) 1.2 - 1.3 - 1.4 mm wide, 1.0 - 1.2 - 1.3 (-1.4) mm deep, widest in the lower two-fifths of the achene, distinctly acute at the apex, rounded at the base.

Distribution: Native in Europe from Scotland, southern Sweden and central Russia southwards (Rechinger 1964), the Caucasus (Losin-Losinskaya 1936), southwestern Asia, north Africa (Rechinger 1954). Reportedly naturalized in parts of the north and south temperate zones (Lawalree 1953).

North America: Canada: BC; United States: AL, AZ, CA, GA, MA, NC, OR, SC, TX, VA, WA. (Map 12)

Habitat: Occurring in moist areas such as waste places, ditches and stream-banks.

Discussion: Some of the early extensive ranges, incorrectly assigned to R. sanguineus in North America may have been based on misidentifications of R. conglomeratus. Since Rechinger's (1937) study, the species' range has apparently expanded, especially in the southeastern parts of the United States. Reports of the species in Ohio, Indiana and Michigan (Fernald 1950) appear doubtful, especially since Deam (1940) in the Flora of Indiana stated that he had seen no specimens of this species and assumed that its inclusion in Indiana may have been based on misidentified specimens of other Rumex, possibly R. obtusifolius. Steyermark (1963) reports it from Missouri and Strausbaugh and Core (1964) from West Virginia; no specimens from these areas have been seen however, the herbaria at which these floras were prepared, have not been consulted. Although a distinctive species, its superficial similarities with R. obtusifolius (Deam 1940) or with R. crispus (Correll and Correll)

1972) may have accounted for R. conglomeratus being included in the Missouri and West Virginia areas. As indicated in the key to the species, R. conglomeratus can be easily distinguished from R. obtusifolius and R. crispus.

R. conglomeratus is however, very similar in general habit and basal leaf characteristics with R. sanguineus, but a closer examination of the fruit and the inflorescence allows for separation of the two species. R. conglomeratus has fruit with three oblong to ovate tubercles that are approximately equal in size while R. sanguineus has fruit with 1 to 3 orbicular to oblong tubercles and if there are three present, they are distinctly unequal in size. In R. conglomeratus, all but the uppermost whorls are subtended by leaf-like bracts but in R. sanguineus, only the lowermost whorls are subtended by the bracts. The latter characteristic allows for separation of the species even when the fruit is immature and the tubercles are not fully developed.

Rechinger (1964) reports two subspecies for R. conglomeratus, subsp. conglomeratus and subsp. nevadensis H. Lindb., with the latter being restricted to southern Spain. The North American specimens are clearly referable to subsp. conglomeratus.

Map 12 Distribution of R. conglomeratus.

Map 12.

16.

Rumex sanguineus Linnaeus, Sp. Pl. 334, 1753.

Locality cited: "in Virginia".

Lectotype: Hb. LINN 464.2, "2 sanguineus" (LINN photo!) (See discussion)

Perennial. Root fusiform. Stems usually less than 1m tall, less than 8mm wide at base, erect, sulcate, red-brown, leafy, branched alternately, usually glabrous to rarely indistinctly papillose; trichomes papilliform, evident on upper branches, less than 0.1 mm long, shorter than to as long as wide; ochreose usually deciduous on mature plants. Branches diverging at 40° - 80° to the main stem and often weakly arching. Basal leaves oblong to ovate-lanceolate, subobtusate, weakly cordate to obtuse, 2 - 5 times as long as wide, widest below middle, entire; middle veins leaving the midrib at 50° - 65° ; lamina glabrous above, glabrous to very indistinctly papillose on veins below; trichomes papilliform, less than 0.1 mm long, shorter than to as long as wide; petiole usually shorter than the lamina. Lower cauline leaves similar to basal but becoming progressively smaller up the stem; petiole shorter than the lamina. Upper cauline leaves narrowly lanceolate, acute, cuneate, entire, glabrous above, commonly glabrous to indistinctly papillose on veins below; petiole glabrous, much shorter than the lamina, the leaf sometimes sessile. Inflorescence a wide, very open panicle, with most branches subtended by leaves. Whorls commonly consisting of less than 10 flowers, arranged on branches and on main stem above branches, all whorls remote, the lower subtended by leaf-like bracts very similar to uppermost leaves, the upper not

subtended. Pedicels filiform, usually longer than the valves, (1.0-) 1.8 - 4.3 mm long, less than 2mm wide at middle, articulated in lower third, rarely at middle; articulation distinctly swollen. Sepals oblong, obtuse, (0.8-) 1.0 - 1.5 mm long and about half as wide, curved upwards around base of valve. Valves oblong to ovate, obtuse, 2.0 - 3.0 mm long, (0.5-) 1.8 - 1.3 mm wide, widest at or near the middle, entire, rarely with one indistinct tooth near base; veins weakly raised, diverging the midrib at an angle of 45° - 90° with 1 anastomosis on each side of the valve; distance to the first anastomosis from the midrib 0.2 - 0.5 mm. Tubercles 1 - 3, if more than 1, then distinctly unequal in size, largest tubercle globular to oblong, obtuse at apex and base, (0.6-) 1.1 - 1.5 mm long, (0.5-) 0.8 - 1.0 mm wide, 0.5 - 1.0 mm deep, widest at or near the middle, base of tubercle at base of valve or up to 0.3 mm above, surface smooth but faintly reticulate. Anthers usually less than 1.0 mm long. Mature achene ovate, dark red-brown to black, 1.5 - 2.0 mm long, 1.0 - 1.3 mm wide, 0.9 - 1.3 mm deep, widest at or near the middle, abruptly contracted to an acute apex, obtuse to broadly cuneate at base.

Distribution: Native in most of Europe from c. 60° N. southwards but absent in Russia and rare in the Mediterranean region (Rechinger 1964), Turkey, N. Iran, Syria, N. Africa (Lawalree 1953). Introduced in North America but very rare.

Canada: BC; United States: AL, OR, PA. (Map 8)

Habitat: Damp and shady places in Europe but in North America apparently confined to waste places, commonly on ballast.

Discussion: Lousley (1938) designated Linn. Herb. 464.2, as the lectotype; however, an examination of the Linn. Herb. photograph of 464.2 and observations made by Dr. J. McNeill in 1975, indicate that this specimen is in flower and does not completely match the protologue (Linnaeus 1753). An examination of the microfiche of the Linnaeus material in the Riksmusset of Stockholm revealed that their specimen Sl49 15, although fitting the foliar characteristics of the protologue, also is either in bud or flower and likewise, could not have provided the fruit characteristics of the protologue. The location of a Linnaean specimen fitting all aspects of the protologue is not known.

Linnaeus (1753) reported that the native habitat of R. sanguineus was Virginia. The species does not occur there today and other than Linnaeus' report of it being native in North America, it is known only as an introduction. Lousley (1938), postulated that because of its demand as a pot-herb, R. sanguineus was grown by the Virginian colonists. It then became at least temporarily naturalized and a specimen was sent to Linnaeus by one of his North American correspondents. From his diagnosis and his synonymy, it is clear that the species at present called R. sanguineus was indeed the one that Linnaeus had in mind, even though the Virginian origin may be doubtful. The R. sanguineus of late authors (Pursh 1814; Nuttall 1818; Campdera 1819; Eaton 1824;

Hooker 1840; Gray 1848; Meisner 1856; Macoun 1883; McSwain and Bain 1891) was based almost entirely on misidentifications of immature plants of R. obtusifolius; this error was corrected by Trelease (1892). Rechner (1937) cited only one site for R. sanguineus in North America. The present study adds only an additional three sites and is in agreement with Gleason (1968) who noted that the species is probably not persistent and any report of its widespread occurrence in North America must be held to be in suspect. Interestingly enough, Hitchcock and Cronquist (1969) reported R. sanguineus from Washington, Oregon and as a weed of the eastern United States perpetuating the inaccurate reports of the distribution of the species.

17.

Rumex britannica Linnaeus, Sp. Pl. 334. 1753.

Lectotype: LINN No. 464.5 "verticillatus 4" (written by Linnaeus has been crossed out by Smith), "Lapathum folius longis latis etc. fl. virg p. 39" (written by Gronovius) and "brittanica Gron: Britannica certe" (written by J. E. Smith). (LINN, DAO photo!) (see discussion)

R. hydrolapathum auct. amer. non Hudson.

R. hydrolapathum var. americanum A. Gray, Man. ed. 3. 377 1856. No specimens cited.

R. orbiculatus A. Gray, Man. ed. 4. 420, 1867.

Lectotype: Connecticut: New Haven, D.C. Eaton (GH!, F272261)

R. britannica var. borealis Rechinger f., Publ. Field Mus. (Bot.) 17:127, 1937

Locality cited: "in the northeastern part of its area."
(NK, NF, NS, QU).

Perennial. Root fusiform. Stem to 2m tall, usually around 1cm wide at base, moderately leafy, alternately branched above middle, erect, sulcate, glabrous to indistinctly papillose; trichomes papilliform, most evident on branches, much less than 0.1 mm long, shorter than to as long as wide; ochreous partially persistent on mature plants. Branches diverging at (20°-) 30° - 40° - 50° (-55°) to the main stem and often weakly arching. Basal leaves large, largest observed 55.5 cm long, 14cm wide, lanceolate to oblong-lanceolate, acute, cuneate to rounded, 3 to 4 times as long as wide, crisped; middle veins of lamina diverging at 65° - 80° to the midrib; lamina glabrous

above, indistinctly papillose on veins below; trichomes papilliform, less than 0.1 mm long, shorter than to as long as wide; petiole shorter than the lamina, glabrous. Lower cauline leaves very similar to basal but progressively smaller up the stem. Upper cauline leaves narrowly lanceolate to linear-lanceolate, narrowly acute, narrowly cuneate, becoming progressively smaller up the stem, crisped; lamina glabrous above, indistinctly papillose on veins below; petiole much shorter than the lamina, the leaf sometimes subsessile. Inflorescence a wide, open panicle, occupying the upper half of the stem, only the lower branches subtended by leaves. Whorls commonly consisting of ca 20 flowers, subtended by leaf-like bracts, the lower whorls remote, the upper contiguous. Pedicels filiform, (2.8-) 5.8 - 10.3 - 12.5 (-14.0) mm long, less than 0.2 mm wide at middle, articulated in lower third; articulation barely evident, not swollen. Sepals oblong, obtuse, (1.1-) 1.5 - 1.8 - 2.1 (-2.5) mm long, at right angles to the pedicel. Valves orbicular, obtuse to narrowly obtuse, truncate to weakly cordate, indented 0.1 - 1.7 mm at base, (3.3-) 4.5 - 5.5 - 6.0 (-7.5) mm long, (2.8-) 4.0 - 5.0 - 6.9 (-7.2) mm wide, the margin entire to very weakly serrate; teeth 1 - 9 per side, the largest usually less than 0.4 mm long; veins raised, diverging at ca 90° to the midrib with 1 - 2 anastomoses on each side of the valve; distance to the first anastomosis from the midrib 1.0 - 1.3 - 1.5 (-2.0) mm. Tubercles 3, fusiform, acute at apex, obtuse at base, approximately equal in size, (1.6-) 2.5 - 3.0 - 3.5 (-4.8) mm long, (0.6-) 0.8 - 1.0 - 1.2 (-1.4) mm wide,

widest near base, base of tubercle (0.2-) 0.3 - 0.5 - 0.5 (-0.8) mm above valve base, surface smooth but faintly reticulate. Anthers usually less than 1mm long. Mature achene ovate, light red-brown, (2.8-) 3.2 - 3.7 - 4.0 (-4.5) mm long, (1.5-) 1.7 - 2.0 - 2.3 (-2.7) mm wide, (1.0-) 1.5 - 1.7 - 1.9 (-2.0) mm deep, widest at or near the middle, acute at apex and base, the former sometimes more sharply so, seed coat thin, easily removed.

Distribution: Native in North America, most common in the eastern portion of its range.

Canada: AB, NT, MB, NK, NF, NS, ON, PE, QU, SK; United States: CA, CT, DC, IL, IA, ME, MA, MI, MN, (MO), (NE), NH, NJ, NY, ND, OH, PA, RI, SD, VT, WI. () reported in these states, but no specimens seen. (Map 13)

Habitat: Damp areas such as marshes, meadows, shores.

Discussion: Until 1945, when Fernald reexamined its typification, the name R. britannica was applied to the native North American water dock that occurs from Newfoundland westward to Alberta (with a disjunct population in California) and southward to New Jersey and to Missouri. This was the usage of for example, Trelease (1892), Britton and Brown (1896), Robinson and Fernald (1908) and Rechinger (1937). Since 1945, most authors (eg. Fernald 1950; Gleason 1952; Gleason and Cronquist 1963) have accepted Fernald's treatment of R. britannica as a nomen confusum and have adopted the name R. orbicu-

latus A. Gray for this species. The present International Code of Botanical Nomenclature (Stafleu et al. 1978) does not however, recognize nomina confusa per se. Any name that can be typified comes under the provisions of the Code and can only be rejected "if it has been widely and persistently used for a taxon not including its type" (Art. 69) or if it is otherwise illegitimate. Consequently, the typification of R. britannica has been re-examined and it is concluded that the name has to be reinstated for the species recently known as R. orbiculatus.

Rumex britannica was published by Linnaeus in the first edition of Species Plantarum (1753). The phrase-name is attributed to his own Materia medica (Linnaeus 1749). The protologue also includes (1) two other synonyms, one referring to Colden (1743) and the other to Gronovius (1739), (2) the statement "Habitat in Virginia 4", (3) a comparison with R. verticillatus as to ochrea and inflorescence characters and, (4) the note that the specimen that he has from Gronovius does not match details of Gronovius' description in Flora Virginia (1739).

The phrase-name is always the most important element in a Linnaean protologue; in this case, it is particularly critical because despite the citation of Materia medica, it is in fact, a new diagnosis. The Species Plantarum phrase-name reads "RUMEX floribus hermaphroditis: valvulis integerrimis: omnibus graniferis, folius lanceolatis: vaginis obsoletis.", whereas that in Materia medica has "...valvulis integerrimis nudis, folius cordato-lanceolatis."

Almost all that is in common is that both are members of the hermaphrodite section Rumex, a much more substantial change in the diagnosis than Fernald (1945) implies in saying that it was "some-what altered"! Stearn (1957) notes that "Linnaeus based the phrase-name nearly always on a specimen or an illustration seen by him, for he distrusted the description of others and the phrase-name is thus likely to give a significant indication of his intent, certainly he would never have chosen an element disagreeing with the phrase-name to typify his species."

As noted by Fernald (1945), the Colden (1743) reference in the protologue may very well apply to the species presently known as R. orbiculatus. The description was from Colden's treatment of the plants growing around his residence (now near Newburgh) in New York, (Gray 1843), a region in which this North American native species is still known to occur. Colden's Plantae Coldenhamiae had been sent to Gronovius who in turn, sent it to Linnaeus; Linnaeus was responsible for its publication in 1749, which accounts for his familiarity with this work. As Gray (1872) and Fernald (1945) have suggested that the Gronovian synonym refers to a different species. In Flora Virginia, this description is marked by an asterisk and followed by "Clayt" (on) indicating that the description is Clayton's as Gronovius did not possess a specimen upon which he could base a description. The examination by Gray (1872) of the specimen retained in the Clayton herbarium and the illustrations given by Fernald (1945) confirm that it is referable to another species, namely R. obtusifolius. In this context, it is noteworthy that

Linnaeus had indicated in the protologue of R. britannica that his specimen did not agree with the Gronovian diagnosis. As there is no indication that either Gronovius nor Linnaeus saw the Clayton specimen, its identity does not affect the application of the name R. britannica. The Linnaean specimen No. 464.5 being proposed here as the lectotype of R. britannica bears part of the Gronovian citation and reference to Flora Virginia and would therefore, appear to have come from Gronovius; if so, it must have come into his possession subsequent to the publication of Flora Virginia in 1739 and then reached Linnaeus prior to 1753 but probably after 1749 when the Materia Medica was prepared.

As Fernald (1945) stressed, the species at present known as R. orbiculatus does not occur in Virginia. Linnaeus' choice of habitat "in Virginia" was probably based entire on the fact that the specimen in his possession came from Gronovius. Rather than being a Clayton specimen, it is quite possible that the specimen sent to Linnaeus by Gronovius was from Colden. Colden, in his correspondence with Gronovius (Gray 1849) indicated that he was sending North American plant specimens to him.

Believing that both R. britannica and R. verticillatus were native to Virginia (both protologues include a Gronovian (1739) synonym), Linnaeus (1753) included a brief comparison of the two species that must have been based on plants available to him. The comparison clearly separates the proposed type of R. britannica,

LINN 464.5 from the Linnsean specimens of R. verticillatus, LINN 464.3, 464.4 and 464.6

Gray (1872) during his examination of the material in the Linnsean herbarium in London, reported finding a specimen named R. britannica, referable to his R. orbiculatus; he wrongly discounted it as a possible type since the Clayton specimen had been referable to R. obtusifolius. Later, a search for the Linnsean specimen of R. britannica was carried out for Rechinger (1937) by H. W. Pugsley. Pugsley reported that the specimen's characters were not in agreement with those of R. britannica. Based on the description given to Rechinger, Pugsley appears to be describing a specimen of R. verticillatus, probably No. 464.4, as it is labelled "R. britannica HU". Apparently, Pugsley ignored or was unaware of the transposition in naming the specimens that was corrected by Smith (Gray 1872) and did not consider examining No. 464.5 which was labelled as a specimen of R. britannica. It is interesting to note that Fernald (1945) accepted Pugsley's observations and believed that a specimen of R. britannica referable to R. orbiculatus did not exist at LINN, thus ignoring Gray's (1872) earlier observations on this specimen. Observations made on LINN 464.5 by Dr. J. McNeill have indicated clearly that it is referable to the plant known as R. orbiculatus. A Gray, and agrees with the Linnsean (1753) diagnosis.

In summation, no aspect of the protologue of R. britannica

(Linnaeus 1753) has been found to preclude its employment for the North American native species, presently referred to as R. orbiculatus. Also, the morphological characteristics expressed in the diagnosis and in the comparison of R. verticillatus with R. britannica are evident in LINN 464.5. Based on these factors, LINN 464.5 is here proposed as the lectotype of R. britannica.

The only basis for rejecting the name R. britannica would be if it had been "widely and persistently used for a taxon not including its type". Hudson (1762) in his Flora Anglica, took up R. britannica for a European species of Rumex, but in the second edition (Hudson 1778), he replaces it by R. hydrolapathum, the name still used for the species, making no reference to R. britannica. Ascherson and Graebner (1906) include R. britannica Hudson, not Linnaeus as a synonym of R. hydrolapathum. In Flora Europaea, Rechinger (1964) does not mention R. britannica and so despite the epithet, the name has scarcely ever been used for a European species of Rumex. As indicated by Gray (1872), it is uncertain whether Michaux (1803) and Pursh (1816) were employing R. britannica in what I have established is the Linnaean sense or in Gray's sense, i.e. for the native species now known as R. altissimus (section Axillares). From the description that Meisner (1856) provided, his use of R. britannica is certainly not that of Linnaeus and would also appear to be referable to R. altissimus. However, there has been no widespread usage of the name R. britannica for R. altissimus.

In his treatment of the North American species of Rumex, Trelease (1892) employed the name R. britannica for the species to which its lectotype belongs placing R. orbiculatus A. Gray in synonymy. Moreover, all subsequent uses of R. britannica in North America prior to Fernald's (1945) declaring it a nomen confusum, were for this species. Because R. britannica can be typified and because it has scarcely ever been used for a taxon not including its type (Article 69), it must be reinstated, despite its inappropriateness for the North American species referred to in most recent publications as R. orbiculatus A. Gray.

The reason for the inappropriate epithet britannica is to be found in Linnaeus' (1749) Materia Medica. There Linnaeus noted that the root of an aquatic European Rumex was known in pharmacy as "Herbae Britannicae Radix". The species that he was dealing with in Materia Medica appears to have been the European R. aquaticus, and not R. britannica, or R. hydrolapathum as Fernald claims. Although Linnaeus, in totally altering his Materia Medica diagnosis dealt in the Species Plantarum with a North American rather than a European species, he did draw on the Materia Medica for his binomial. This fact alone upset Fernald (1945) who considered R. britannica as "ill-begotten" and "inconsiderately assigned." It is unfortunate that Linnaeus did not employ a more appropriate name for this North American taxon. However, no matter how inappropriate name, R. britannica cannot be dismissed on this

basis (Article 62). A further example of an inappropriate Linnaean name employed for a North American species is that of Asclepias syriaca which although not from Syria is the correct name for this native North American species.

In his Catalogue of Canadian Plants, Macoun (1883) placed R. acutus sensu Hooker and R. sanguineus sensu Hooker (Hooker 1840) in synonymy under R. orbiculatus; Reehinger (1937) included these as synonyms of R. britannica. Specimens cited by Hooker have not been seen; however, as the habitat of his "R. acutus" was described as "woody country" and that of "R. sanguineus" as "barren country beyond lat. 64", it is very doubtful if Hooker was referring to R. britannica as the habitats are atypical for that species. Moreover, R. britannica is not otherwise known north of 64° N. lat.

As pointed out by Reehinger (1937) R. britannica does not appear to be closely related to any other North American species and its delimitation is easily described (see the key to the species). R. britannica appears most closely related to the European R. hydrolapathum Huds. from which it can be distinguished by its valves being orbicular, obtuse rather than triangular, acute.

Mar 13 Distribution of R. britannica.

Map 13

16.

Rumex obtusifolius Linnaeus, Sp. Pl. 335. 1753.

Lectotype: LINN No. 464.18, "obtusifolius" (LINN, DAO photo!)

R. crispatus Michaux, Fl. Bor. Amer. 1: 217. 1803.

Type Specimen: "Kentucky" (Michaux Herbarium, DAO microfiche!)

R. sanguineus auct. amer. prior., non Linnaeus.

Perennial. Roots fusiform. Stems ca 1m tall, ca 1cm wide at base, erect, sulcate, brown to red-brown, often with red-purple areas, leafy, branched alternately above the middle, glabrous to distinctly pubescent or papillose, even on same plant; trichomes papilliform to commonly cylindrical, less than 0.3 mm long, shorter than to 3 times longer than wide; ochreae deciduous to partially persistent on mature plants. Branches diverging at (20°-) 25° - 30° - 40° to the main stem and often weakly arching. Basal leaves large, largest observed 37.0 cm long, 18.2 cm wide, broadly ovate to oblong-ovate, subacute to obtuse, cordate or rarely truncate, indented at base 0.8 - 3.0 cm, 1.7 - 2.5 times as long as wide, widest near middle to just above base; margin somewhat undulate, entire to crisped; middle veins of lamina diverging at 40° - 70°, commonly at 60° to the midrib; lamina glabrous above, distinctly pubescent to papillose on veins below, rarely glabrous; trichomes papilliform or more commonly cylindrical, less than 0.3 mm long, shorter than to 3 times as long as wide; petiole somewhat shorter than lamina, glabrous to moderately pubescent or papillose.

Lower cauline leaves similar to basal but becoming ovate-lanceolate, acute, weakly cordate to broadly cuneate or rounded, becoming progressively smaller up the stem; petiole shorter than the lamina. Upper cauline leaves ovate-lanceolate to narrowly lanceolate, acute to acuminate, cuneate to narrowly cuneate, entire to weakly crisped, glabrous above, distinctly pubescent to papillose or glabrous on veins below; petiole much shorter than lamina, the leaf sometimes subsessile. Inflorescence a narrow to wide panicle, occupying the upper half of the stem, all branches subtended by leaves. Whorls commonly consisting of 10 - 26 flowers, arranged on branches and portion of main stem above branches, only the lowermost whorls subtended by leaf-like bracts similar to uppermost cauline leaves, the lower remote, the upper becoming contiguous. Pedicels filiform, 2.3 - 5.8 - 8.5 (-10.0) mm long, ca 0.2 mm wide at middle, articulated in the lower two-fifths, commonly in the lower third; articulation distinctly swollen. Sepals oblong, obtuse, commonly at right angles to pedicel, occasionally reflexed slightly downwards, (1.2-) 1.5 - 1.7 - 2.0 (-2.8) mm long and about half as wide. Valves deltoid, acute to subacute, truncate to weakly emarginate, indented to 0.5mm, (3.0-) 3.5 - 4.3 - 5.0 (-6.0) mm long, (2.0-) 2.5 - 3.0 - 3.5 (-4.0) mm, widest near base, the margin below the apex dentate-spring; distinct teeth (1-) 3- 4 - 4 (-5) per side, the largest (0.6-) 1.0 - 1.5 - 1.8 (-2.0) mm long; veins raised, diverging from midrib at ca 45° with (1-) 2 anastomoses on each side of the valve; distance

to the first anastomosis from the midrib 0.5 - 0.8 - 1.0 mm.

Tubercles 1 (-3), somewhat variable even in the same inflorescence, ovate, obtuse to acute, obtuse at base, very unequal in size, surface smooth but faintly reticulate; the largest tubercle (1.1-) 1.5 - 2.0 - 2.5 (-2.9) mm long, (0.4-) 0.7 - 1.2 - 1.5 (-1.8) mm wide, (0.5-) 0.8 - 1.0 - 1.3 (-1.5) mm deep, widest at middle to lower fifth, base of tubercle over-arching base of valve or terminating up to 0.5 mm above the base; the smallest tubercle 0.5 - 1.5 mm long, 0.3 - 0.5 mm wide, 0.5 - 0.9 mm deep, widest at middle to near base, base of tubercle 0.3 - 0.5 mm above base of valve. Anthers 1.3 - 1.5 mm long. Mature achene ovate, red-brown, (1.8 -) 2.3 - 2.5 - 2.9 (-3.0) mm long, (1.2-) 1.3 - 1.5 - 1.7 (-1.8) mm wide, (1.0-) 1.2 - 1.4 - 1.5 (-1.7) mm deep, widest at middle to lower third, acute at the apex and base, the base somewhat broadly so.

Distribution: Native in Europe (excluding the arctic, subarctic and mediterranean regions) and western Asia; introduced in eastern Asia, southern Africa, Australia, North and South America (Rechinger, 1954). Subsp. obtusifolius is native in western Europe, whereas subsp. transiens is native in central Europe, the Balkan Peninsula and southern Scandinavia.

Widely distributed in Canada and the United States but absent in the Canadian Arctic and much of the prairie regions of both countries.

Canada: BC, NK, NF, NS, ON, PE, QU;

United States: AL, AK, AZ, AR, CA, CO, CT, DE, DC, FL, GA, ID, IL, IN, IA, KS, KY, LA, ME, MD, MA, MI, MN, MS, MO, MT, NB, NH, NJ, NM, NY, NC, OH, OK, OR, PA, RI, SC, TN, TX, UT, VT, VA, WA, WV, WI. (Map 14)

Habitat: Occurs in damp to arid soils of roadsides, fields, waste ground, about dwellings, woodlands and along streams.

Discussion: Lousley (1939) designated LINN No. 464.17 as the lectotype of R. obtusifolius, a specimen in bud with only cuneate upper leaves present. This specimen was selected rather than No. 464.18, a fruiting specimen that has a single cordate leaf, since Lousley regarded it as "an annual form of subsp. agrestis Rech. fil - perhaps from secondary growth". Photographs of the Linnaean specimens and observations made on these specimens by Dr. J. McNeill in 1975 showed that in addition to not being capable of providing the fruit characteristics of the Species Plantarum protologue (Linnaeus 1753), LINN 464.17 has cuneate leaves rather than the cordate ones referred to by Linnaeus. Consequently, it is not a suitable lectotype and I propose to reject Lousley's choice on the grounds that he misinterpreted the protologue. (Code Article 8; Staffeu et al. 1978). Although agreeing with Lousley that LINN 464.18 probably does represent secondary growth, this specimen agrees with the protologue in both the fruit and leaf character and is here designated as lectotype.

R. crispatus Michaux (Michaux 1803) was considered by

Trelease (1892) to be a synonym of R. persicarioides; Fernald's examination of the type in the Michaux herbarium indicated that it is referable to R. obtusifolius. This was confirmed by an examination of the DAO microfiche of the specimen.

Trelease (1892) noted that variant of R. obtusifolius with purple and red veined leaves (R. obtusifolius var. discolor Wallroth = f. purpureus (Poiret) Lousley) had often been misidentified as R. sanguineus. It was these misidentifications that led early authors to attribute a much greater geographic range to R. sanguineus that it actually possesses. This mistake must have been based entirely on the red colour of the stem and leaf veins as other morphological characteristics are quite distinct. The distinctness of the two species is evident in the following table (Table 24).

TABLE 24 COMPARISON OF R. sanguineus and R. obtusifolius

Characteristic	<u>R. sanguineus</u>	<u>R. obtusifolius</u>
Basal Leaves	Oblong to ovate-lanceolate; Abaxial veins glabrous to very indistinctly papillose; trichomes papilliform, less than 0.1 mm long.	Ovate or oblong-ovate; Abaxial veins usually distinctly pubescent, occasionally papillose; trichomes commonly cylindrical, greater than 0.1 mm long
Inflorescence	Branches diverging at 40°-80°	Branches diverging at (20°-) 25° - 40°
Valve Shape	Oblong	Deltoid
Valve Margin	Entire	Dentate-spiny

Comparison of R. sanguineus and R. obtusifolius, cont.

<u>Characteristic</u>	<u>R. sanguineus</u>	<u>R. obtusifolius</u>
Seed Colour	Dark red-brown to black	Light red-brown
Seed Size	1.5 - 2.0 mm long, 1.0 - 1.3 mm wide	(1.8-) 2.3 - 2.9 (-3.0) mm long, (1.2-) 1.3 - 1.7 (-1.8) mm wide
Chromosome No.	2n = 20	2n = 40

R. obtusifolius has also been confused with R. crispus, R. dentatus, R. pulcher and R. stenophyllus. The characteristics distinguishing R. obtusifolius from all but R. stenophyllus have been dealt with under the discussion of those species. R. obtusifolius (2n = 40) differs cytologically from R. stenophyllus (2n = 60) and the two species can readily be separated morphologically as indicated in the accompanying table (Table 25)

TABLE 25 COMPARISON OF R. obtusifolius and R. stenophyllus

<u>Characteristic</u>	<u>R. obtusifolius</u>	<u>R. stenophyllus</u>
Basal Leaves	Ovate to oblong-ovate, distinctly cordate	Narrowly lanceolate to oblong, narrowly to broadly cuneate
Leaf Pubescence	Usually distinctly pube- scent, occasionally papillose	Glabrous to indistinctly papillose
Valve Margin	(1-) 3 - 4 (-5) teeth per side	(4-) 5 - 8 (-10) teeth per side
Valve veins	Diverging from midrib at ca 45°	Diverging from midrib at ca 90°
Tubercles	1 - 3, if 3, unequal	3, approximately equal in size

Rechinger (1937) recognized that the majority of the North American specimens of R. obtusifolius belonged to subsp. obtusifolius (which he incorrectly referred to as subsp. agrestis (Fries) Danser), "characterized by leaves somewhat papillous on the under side along the nerves and by large (about 6mm) fruiting perigonia", he also recognized subsp. transiens (Simonkai) Rechinger f., Beih., Bot. Centr. 49 (2);52, 1932, "characterized by somewhat smaller fruiting perigonia with 3 usually unequal grains and shorter fruiting segments as occurring in North America. Fernald (1921) had reported var. sylvestris (Lam.) Koch treated by Rechinger (1932) as a synonym of subsp. sylvestris (Wallroth) Rechinger from Nova Scotia and Prince Edward Island; Rechinger (1937) makes no mention of subsp. sylvestris occurring in North America nor have I seen any North American specimens with its distinguishing features, namely: (a) valves entire or with a few short or indistinct teeth near the base, (b) all valves with equal to subequal tubercles occupying the whole width of the valve, and (c) glabrous throughout. (Rechinger 1932, 1964; Lousley 1939).

In trying to relate the North American specimens to the subspecies recognized by Rechinger, some confusion was apparent. The characteristics that are supposed to distinguish subsp. obtusifolius from subsp. transiens (Rechinger 1932, 1937, 1964; Lousley 1939) and those actually observed on North American specimens, are compared in Table 26. One North American specimen (collected by A. Ruth from the District of Columbia) is cited by Rechinger (1937) as belonging to subsp. transiens; the only feature that it possesses that is characteristic of that

subspecies is the absence of pubescence; the other characteristics of this specimen - valve size, teeth length, tubercle number and shape were well within the range shown by the other North American specimen of R. obtusifolius. The Ruth specimen lacked basal and lower leaves and the leaf L:W ratio could not be used to help in its identification. This specimen does not appear readily referable to subsp. transiens as that subspecies is presently delimited in Europe. From Table 26 it can be seen that the remaining North American specimens also do not exactly agree with the characteristics described for either subspecies in Europe. In five specimens annotated by Rechinger as subsp. obtusifolius, the length of the valves ranged from 3.5 - 4.8 mm and that of the largest tooth from two-fifths to half of the valve width. As seen in Table 26, these valves are closer to those reported for subsp. transiens. Two of these specimens had fruit with only one tubercle, although the midribs of the other two valves were often slightly enlarged while the remaining three had fruit with one to three tubercles of unequal size, the size of the tubercles appears to indicate subsp. obtusifolius but the variation in number of tubercles from one to three on the same inflorescence has not been reported in this subspecies. All five specimens were pubescent and the L:W Leaf Ratios were ca 2:1, both more characteristic of subsp. obtusifolius than subsp. transiens. These specimens annotated by Rechinger as subsp. obtusifolius are typical of almost all the North American specimens and as they appear to combine some of the characteristics

there are said to distinguish these two subspecies, a division into subspecies along the lines adopted in Europe will not be attempted in this study.

A more detailed examination of the range of variation that occurs within the native geographic range of R. obtusifolius would be necessary to assess whether it is the existing subspecific classification that is unsatisfactory or whether subspecific differentiation that is clear in Europe has become blurred in the introduced North American populations. An examination of available European specimens tend to indicate that the former is more likely than the latter.

TABLE 26

CHARACTERISTICS said to DISTINGUISH
R. obtusifolius subsp. *obtusifolius*
 FROM subsp. *transiens* COMPARED WITH
 those observed on the NORTH AMERICAN
 SPECIMENS of *R. obtusifolius*.

Characteristic	subsp. <i>obtusifolius</i>		subsp. <i>transiens</i>		North American Specimens	
	cs 6mm		cs 5mm		(3.0-) 3.5 - 5.0 (-6.0) mm	
Valve Length						
Teeth Length	Usually equal to or somewhat shorter than the valve width but in f. <i>subulatus</i> , longer than the valve width in f. <i>subdentatus</i> much shorter as the margin is almost entire.		Usually up to half the valve width but often much shorter (cs 0.5 mm)		cs half the valve width	
Tubercles	Usually only one tubercle, but in f. <i>frigoris</i> , three well developed tubercles.		Three tubercles but unequal in size.		One to three tubercles; if three, very unequal in size; variable even in same inflorescence.	
Pubescence	Petioles and leaves usually Pubescent		Plant usually glabrous, rarely the petioles and the leaves sparingly pubescent		Petioles and leaves usually pubescent	
Basal and Lower Leaves:						
Average Length:	2.6:1		1.2:1		2.1:1	
Width	(Housley (1939) gave no ranges for the two subspecies)				(range 1.7 - 2.5:1)	303

Map 14 Distribution of R. obtusifolius.

Map 74

Rumex obtusifolius L.

NORTH AMERICA

0 500

MILES

19.

Rumex pulcher Linnaeus, Sp. Pl. 336. 1753.

Localities cited: "in Gallia, Italia, Verona" (in France, (and) at Verona in Italy)

Lectotype: Hb. LINN 464.19, "pulcher 12", "H.U.", selected by Lousley (1938). (LINN DAO photo!) (See discussion)

Perennial. Root fusiform. Stem 38 - 111 cm tall, commonly less than 100 cm, less than 1 cm wide at base, leafy, alternately branched in the upper two-thirds, erect, but often somewhat zig-zag above, red-brown, glabrous to distinctly papillose or pubescent; trichomes commonly most evident on lower stem, papilliform to cylindrical, up to 0.9 mm long, shorter than to ca. 6 times as long as wide; ochreae deciduous to partially persistent on mature plants. Branches diverging at (40° -) 50° - 60° - 70° (-80°) to the main stem, often intricate and weakly arching. Basal Leaves small, largest observed 16.0 cm long, 5.0 cm wide, oblong to oblong-lanceolate or occasionally pandurate, obtuse to subacute, rounded to very weakly cordate at base, commonly 3 - 4 times as long as wide, widest between base and middle, weakly undulate, crisped to entire; middle veins of lamina diverging at 50°-75° to the midrib; lamina glabrous above, weakly to distinctly pubescent to papillose on veins below; trichomes papilliform to cylindrical, less than 0.9 mm long, shorter than to ca 6 times as long as wide; petiole shorter than to as long as lamina, glabrous to distinctly papillose or pubescent. Lower cauline leaves similar

to basal but becoming progressively smaller up the stem; the apex becoming more acute; petiole shorter than lamina. Upper cauline leaves, oblong-lanceolate to linear-lanceolate, commonly acute to obtuse, narrowly cuneate, becoming progressively smaller up the stem; lamina glabrous above, glabrous to distinctly pubescent to papillose; petiole much shorter than the lamina, the leaf sometimes sessile. Inflorescence a wide, open panicle, occupying the upper two-thirds of the stem, all branches subtended by leaves. Whorls commonly consisting of 10 - 20 flowers arranged on branches and portion of main stem above branches, the lower whorls remote and subtended by bracts, the upper somewhat more crowded, all but the uppermost subtended by bracts. Bracts leaf-like, similar to the uppermost cauline leaves but progressively reduced in size upward along the branches or main stem above the branches. Pedicels uniformly thickened, not filiform, (1.6-) 2.5 - 4.0 - 5.0 (-6.0) mm long, commonly 0.3 - 0.4 mm wide at middle, articulated just below middle to the lower third, occasionally at the middle, articulation distinctly swollen. Sepals oblong, obtuse, spreading at right angles to pedicel, (1.0-) 1.3 - 1.8 - 2.0 (-2.5) mm long and about half as wide. Valves ovate-deltoid to oblong-deltoid, acute, truncate, (2.7-) 3.5 - 4.2 - 4.8 (-5.5) mm long, 1.8 - 2.3 - 3.0 (-3.5) mm wide, widest at base to just above the middle, the margin below the apex dentate-spiny; teeth (0-) 5 - 7 - 9 (-11) per side, the largest (0.2-) 1.0 - 1.3 - 1.6 (-2.0) mm long; veins distinctly raised, diverging from the midrib

at ca 90°, with 1 - 2 (-3), commonly 2 anastomoses on each side of the valve; distance to the first anastomosis from midrib 0.5 - 1.0 - 1.0 (-1.3) mm. Tubercles 1 - 3, variable on same plant; ovate to oblong, obtuse to subacute at the apex, obtuse at the base, unequal in size; verrucose but faintly reticulate; the largest (1.5-) 1.8 - 2.3 - 2.5 (-3.0) mm long, (0.5-) 0.8 - 1.0 - 1.3 (-1.7) mm wide, (0.5-) 0.8 - 1.0 - 1.3 (-1.5) mm deep, widest at or near base, the base of the tubercle at the base of the valve or protruding somewhat below the base; the smallest 1.0 - 1.5 mm long, (0.3-) 0.5 - 0.6 mm wide, (0.3-) 0.5 - 0.6 mm deep, widest at or near base, the base of the tubercle at the base of the valve to ca 0.3 mm above. Anthers 1.0 - 2.0 mm long. Mature achene ovate, dark red-brown to almost black, (1.8-) 2.2 - 2.5 - 2.8 mm long, (1.4-) 1.5 - 1.6 - 2.0 mm wide, (1.1-) 1.3 - 1.4 - 1.5 mm deep, acute at the apex, obtuse to subtruncate at the base.

Distribution: Southern and western Europe (Rechinger 1964) and southwestern Asia (Asia Minor). Losin-Losinskaya (1936). Introduced in the United States, common in the southeast and southwest.

United States: AL, AR, CA, FL, GA, LA, MS, NV, NJ, NC, OK, OR, PA, SC, TN, TX, VA. (Map 15)

Habitat: Moist to dry ground; roadsides, waste ground, low fields, edges of lakes and ponds, bogs and muddy areas.

Discussion: The lectotype designated by Lousley (1939) agrees with the protologue for R. pulcher (Linnaeus 1753); this was confirmed by examination of the LINN photograph of 464.19, and by observations made by Dr. J. McNeill in 1975.

Rechinger (1932) recognized five subspecies in R. pulcher and later, (Rechinger 1937) reported that in North America, three of these subspecies had been introduced. The following key is adapted from that given most recently by Rechinger (1964).

Valves entire, or with teeth up to 0.5 mm near the base; branches arcuate-divaricate; leaves seldom panduriform. subsp. anodontus

Valves with several teeth on each side; teeth more 0.5 mm.

Valves suborbicular or ovate triangular, broadly acuminate at apex, teeth short, up to 8 on each side; usually 3 meshes of venation on each side of the tubercle; branches arcuate-divaricate; leaves seldom panduriform. subsp. divaricatus

Valves narrowly ovate or oblong with lingulate apex, 4 - 5 mm long; teeth 1 - 2 mm, 4 on each side; one or two meshes of venation on each side of the tubercle; branches angular-flexuous, often intricate, leaves usually panduriform. subsp. pulcher

In attempting to utilize Rechinger's key to the subspecies, many difficulties were recognized. Specimens from South Carolina

and Texas appeared relatable to subsp. anodontus (Hauskn.) Rech. f. The number of teeth varied from 0 - 2 (5) on each side of the valve but teeth up to 0.8 mm long were observed. Several specimens referred to this subspecies by Rechinger (1937) were not fully mature and the size of the teeth could not be adequately assessed. Even when using specimens cited by Rechinger (1937) and later annotations by him in 1953, of subsp. pulcher and subsp. divaricatus (L.) Murb., the reported distinctiveness of these subspecies was not apparent. The specimens were intermediate in their outline, the number of teeth on both; the size of the teeth did not indicate any separation nor did the meshes of venation. In annotating a number of the North American specimens, Rechinger used, ⁺ subsp. divaricatus as well as subsp. divaricatus non typicus; these specimens did not differ from those annotated as subsp. divaricatus. Apparently, Rechinger was himself uncertain of the delimitation of this subspecies at least in North America. As the specimens usually lacked basal or lower leaves, this characteristic could not be checked; however, Lousley (1944) reported that the panduriform leaves of subsp. pulcher are "best marked on dry sandy or chalky soils, and less evident when the plant grows in moister habitats" indicating that even this characteristic may not be as diagnostic as implied by Rechinger (1964). Due to the difficulties in recognizing these subspecies in North America, these taxa will not be used in this study.

Examination of early herbarium specimens suggests that prior

to the treatment of R. pulcher by Trelease (1892), the species was generally misidentified as R. obtusifolius. Correll and Johnston (1970) reported that references to R. obtusifolius in Texas were based on misidentified specimens of R. pulcher. R. pulcher can be distinguished from R. obtusifolius on both vegetative and fruit characteristics, as indicated in the following table (Table 27).

<u>Characteristic</u>	<u>R. pulcher</u>	<u>R. obtusifolius</u>
Basal leaves	Oblong to oblong-lanceolate	Broadly ovate to oblong-ovate
	Rounded to weakly cordate at base	Distinctly cordate at base
	3 to 4 times as long as wide	1.7 - 2.5 times as long as wide
Inflorescence	Branches diverging at (40° - 50° - 60° - 70° - 80°)	Branches diverging at (20° - 25° - 30° - 40°)
	Branches often intricate	Branches not intricate
Pedicel	Usually shorter than valve	Usually longer than valve
	Thick, 0.3 - 0.4 mm wide at middle	Thin, ca. 0.2 mm wide at middle
Valve	Thick and the veins very distinctly raised	More membranous and less distinctly veined
	teeth usually more than 5 per side	teeth usually less than 5 per side

Table 27.

Comparison of R. pulcher and R. obtusifolius.

Table 27, cont.

311

<u>Characteristic</u>	<u>R. pulcher</u>	<u>R. obtusifolius</u>
Tubercle	Verrucose	Smooth
Achenes	Almost black Base distinctly rounded to truncate	Red-brown Base cuneate

Mar 15 Distribution of R. pulcher.

Map 15

20.

Rumex dentatus Linnaeus, Mant. Alt. 226, 1771.

Type Locality: "in Aegypto" (In Egypt).

Lectotype: Hb. LINN 464.12, "dentatus" and "H.U." (LINN, DAO photo1)

Annual or biennial. Roots fusiform. Stem 17.4 - 88.0 cm tall, less than 1 cm wide at base, erect, weakly sulcate, red-brown, leafy, branched alternately above middle, glabrous to indistinctly papillose or pubescent; trichomes cylindrical to commonly papilliform, evident on upper stem and branches, less than 0.2 mm long, commonly less than 0.1 mm, shorter than to twice as long as wide; ochrace deciduous to partially persistent on mature plants. Branches diverging at 40° - 50° - 70° to the main stem and often weakly arching. Basal leaves small, oblong to oblong-lanceolate, obtuse to subacute, broadly cuneate to truncate, rarely subcordate, 2 - 7 times as long as wide, weakly undulate, entire to weakly crisped; middle veins of lamina diverging at ca 45° to the midrib; lamina glabrous above, glabrous to indistinctly or weakly papillose or pubescent on veins below; trichomes cylindrical to commonly papilliform, less than 0.2 mm long, commonly less than 0.1 mm long, shorter than to twice as long as wide; petiole shorter than to as long as lamina. Lower cauline leaves similar to basal, but becoming progressively smaller up the stem; petiole shorter than lamina. Upper cauline leaves narrowly oblong to narrowly lanceolate, acuminate to narrowly cuneate, entire to weakly crisped, glabrous above, glabrous to indistinctly or weakly papillose on veins below; petiole much shorter than the lamina. Inflorescence a wide, open panicle, occupying the upper half of the stem, all branches subtended by leaves. Whorls consisting

of 10 - 20 flowers arranged on branches and portion of main stem above branches, all or all but a few uppermost whorls subtended by leaf-like bracts similar to but progressively smaller than uppermost cauline leaves, the lower whorls remote, the upper becoming less so. Pedicels filiform, (1.8-) 2.8 - 3.8 - 5.0 mm long, less than 0.3 mm wide at middle, articulated in the lower third; the articulation distinctly swollen. Sepals oblong, obtuse, at right angles to pedicel to weakly curved upwards, (1.0-) 1.2 - 1.5 - 2.0 mm long and about half as wide. Valves deltoid, acute, broadly cuneate, (3.0-) 3.8 - 4.1 - 4.8 (-5.0) mm long, (1.8-) 2.0 - 2.3 - 2.5 mm wide, widest in lower third, the margin below the apex dentate to spiny; teeth (1-) 2 - 3 - 3 (-4) per side, the largest (1.0-) 1.4 - 2.0 - 2.5 mm long; veins raised, diverging from the midrib at 45° - 90° ; with 1 - 2 anastomoses on each side of the valve; distance to first anastomosis from midrib 0.5 - 0.8 - 0.8 mm. Tubercles 3, oblong-ovate, obtuse to subacute to apex, obtuse at base, approximately equal in size, (2.0-) 2.1 - 2.6 - 3.0 mm long, (0.8-) 1.1 - 1.5 - 1.8 mm wide, (0.7-) 0.8 - 1.0 - 1.0 (-1.3) mm deep, widest near the middle to above the base, the base of the tubercle usually at the valve base or projecting past the valve base, surface typically verrucose but faintly reticulate. Anthers 0.5 - 0.9 mm long. Mature achene ovate, red-brown, (1.8-) 2.0 - 2.4 - 2.5 mm long, (1.3-) 1.4 - 1.6 - 1.6 mm wide, (1.2-) 1.3 - 1.4 - 1.5 mm deep, widest just below middle, acute at apex, more broadly acute to obtuse at base.

55

OF / DE

R. dentatus is represented in Eurasia by several subspecies and most of the North American specimens are referable to R. dentatus L. ssp. Klotzschianus (Meisner) Rechinger f., Beih. Bot. Centralbl., 49 (2): 19.1932.

Distribution: The species as a whole is native from tropical and subtropical Asia to southeastern Europe and North Africa (Rechinger 1954). Ssp. klotzschianus is confined to southern and eastern Asia. Introduced in North America, but rare.

Canada: AT, ON; United States: AZ, CA, OR, TX. (Map 16)

Habitat: Occurring in damp soil; waste and cultivated fields, along streams or ditches.

Discussion: Observations made on the Linnaean photograph of LINN 464.12 and an examination of the specimen by Dr. J. McNeill in 1975, indicated that this specimen agrees with the protologue (Linnaeus 1753) and therefore, it has been selected as the lectotype.

Rechinger (1937) recognized R. dentatus from sites in Oregon and California. The present range, although somewhat greater appears to be based on further introductions of the species rather than an active spread from previous areas of introduction. The species is known from only two sites in Canada: Toronto, Ontario, from 1904 and Lethbridge, Alberta, from 1946. The Ontario specimen lacks mature fruit and it cannot be determined to subspecies. No recent

Canadian collections have been reported and its persistence is doubtful. Similarly, the specimens from Oregon represent old collections (late 1809's) and no recent collections have been reported from that state. However in Arizona, California and Texas, most of the collections are from the mid 1960's to the mid 1970's.

The majority of the specimens of R. dentatus had been misidentified as R. pulcher or R. obtusifolius. Cytologically, R. dentatus ($2n = 40$) can be separated from R. pulcher ($2n = 20$), but not from R. obtusifolius. Although superficially similar to the species, R. dentatus can be recognized by the following comparisons of morphological characteristics, (Table 28):

TABLE 28

COMPARISON OF R. dentatus WITH
R. pulcher and R. obtusifolius.

Characteristic	<u>R. dentatus</u>	<u>R. pulcher</u>	<u>R. obtusifolius</u>
Basal leaf Base	Broadly cuneate to truncate	Obtuse to weakly cordate	Distinctly cordate
Inflorescence Branches	Diverging at 40° - 50° - 70°	Diverging at (40°) 50° - 60° - 70° (-80°) often intricate	Diverging at (20°) 25° - 30° - 40°
Pedice	Usually shorter than valve. Filiform, less than 0.3 mm wide at middle	Usually shorter than valve. Not noticeably narrower, 0.3-0.4 mm wide at middle	Usually longer than valve. Filiform, ca 0.2 mm wide at middle
Valve Width	(1.8-)2.0-2.5mm	1.8-3.0(-3.5)mm	(2.0-)2.5-3.5(-4.0)mm

Table 26, cont.

<u>Characteristic</u>	<u>R. dentatus</u>	<u>R. pulcher</u>	<u>R. obtusifolius</u>
No. of Teeth per side	(1-)2-3(-4)	(0-)5-9(-11)	(1-)3-4(-5)
Tubercles	3, approximately equal in size and verrucose	1-3, if 3, une- qual in size and verrucose	1-3, if 3, unequal in size and smooth
Achenes	Red-brown. Acute at apex, broadly acute to obtuse at base.	Dark red-brown, to black, Acute at apex, obtuse to truncate at base.	Red-brown. Acute at the apex, more broadly so at base.

318

Map 16 Distribution of R. dentatus, R. brownii
and R. palustris.

21.

Rumex violascens Rechinger f., Repert. Spec. Nov. Regni Veg.

39:171, 1936.

Type: New Mexico: In the valley of the Rio Grande below Donana, Mexico Boundary Survey under Major W. H. Emory, C.C. Parry, J.M. Bigelow, C. Wright and A. Schott 1173 (US 62361).

R. berlandieri auct. amer. non Meisner.

Annual, biennial or perennial. Stem 24.5 - 75.0 cm tall; ca 1cm wide at base, erect, sulcate, leafy, alternately branched above middle, brown to red-brown, glabrous; ochreae deciduous to weakly persistent in part on mature plants. Branches diverging at 35° - 40° - 50° (-60°) to the main stem, often weakly arching. Basal leaves oblong-lanceolate to elongate-obovate, broadly acute to obtuse, cuneate, to obtuse or truncate, 2 - 4.5 times as long as wide, entire; middle veins of lamina diverging at 45° - 70° to the midrib; lamina glabrous above and below; petiole shorter than lamina. Lower cauline leaves similar to basal but becoming progressively smaller up the stem. Upper cauline leaves becoming linear-lanceolate, narrowly acute, narrowly cuneate, becoming progressively smaller up the stem; lamina glabrous above and below; petiole much shorter than the lamina, the leaf sometimes sessile. Inflorescence a wide, open panicle, occupying the upper half of the stem, the branches usually subtended by leaves. Whorls commonly consisting of 10 - 20 flowers, arranged on branches and portion of main stem above branches, the lower whorls remote, rarely subtended by leaf-like bracts very similar to uppermost leaves, the upper whorls becoming contiguous, not subtended by leaf-like bracts. Pedicels filiform,

2.9 - 4.0 - 8.0 mm long, less than 0.3 mm wide at middle; articulation distinctly swollen in the lower two-fifths to commonly in the lower third of the pedicel. Sepals oblong, obtuse to acute, (0.8-) 1.0 - 1.3 - 1.5 mm long and about half as wide, commonly angled slightly upwards. Valves deltoid, acute, broadly cuneate, (2.5-) 2.8 - 3.2 - 3.7 (-4.0) mm long, (2.0-) 2.5 - 2.8 (-3.0) mm wide, widest in lower third to near base, the margin below the apex serrate; teeth 3 - 5 - 6 per side, the longest 0.2 - 0.4 - 0.5 mm long; veins raised, diverging at ca 90° to the midrib with 1 - 2, usually 1, anastomoses on each side of the valve; distance to first anastomosis from the midrib 0.4 - 0.8 - 0.8 mm. Tubercles 3, ovate, obtuse to weakly acute at apex, obtuse at base, unequal in size, surface smooth but faintly reticulate; largest tubercle (1.6-) 1.8 - 2.0 - 2.3 mm long, 0.7 - 0.8 - 1.0 mm wide, 0.6 - 0.8 - 1.0 mm deep, widest near base, base of tubercle at the valve base or projecting beyond the valve base; smallest tubercle ovate, 1.0 - 1.3 - 1.5 mm long, 0.5 - 0.6 - 0.7 mm wide, (0.5-) 0.6 - 0.7 - 0.8 mm deep, widest near base, base of tubercle at the valve base. Anthems less than 1 mm long. Mature achene ovate, brown to red-brown, (1.8-) 1.9 - 2.3 - 2.3 mm long, (1.1-) 1.3 - 1.3 - 1.5 mm wide, 1.1 - 1.1 - 1.3 mm deep, widest at or near the middle, acute at the apex and base; apex often narrower than base.

Distribution: Native to the S.W. United States and Mexico.

U.S.A.: AZ, CA, NM, TX. (Map 17)

Habitat: Occurring in wet, low land and along ditch banks.

Discussion: Rechinger (1937) named R. violascens to represent the species previously misidentified by American Authors (Trelease 1892; Small 1903; Jepson 1923) as R. berlandieri Meisner; the latter is a species belonging to the section Axillares. This separation is justified. R. violascens differs from R. berlandieri in that (a) it lacks regular axillary branches, (b) the basal and lower leaves are oblong-lanceolate to elongate obovate rather than linear-lanceolate to oblong-linear, (c) the valves are smaller with the width less than 3.0 mm wide, (d) the margin below the apex is serrate rather than entire, (e) the tubercles are unequal in size rather than approximately equal in size and (f) the achenes are smaller with the length less than 2.5 mm. R. berlandieri was reported by Rechinger (1937) from Louisiana, Texas and Mexico.

Map 17 Distribution of R. violascens.

Map 17

22.

Rumex maritimus Linnaeus, Sp. Pl. 335. 1753.

Locality cited: "in Europae".

Lectotype: Hb. LINN No. 464.13, "maritimus 8" (written by Linnaeus but the specimen was collected by Hudson). (LINN, DAO Photo!)
(see discussion)

Annual or biennial. Root fusiform to fibrous. Stem (6-) 12.5 - 68 (-88) cm tall, less than 1 cm wide at base, usually erect or ascending; smooth to weakly sulcate, brown to red-brown, leafy, alternately branched in the upper two-thirds to half; weakly to distinctly pubescent, trichomes cylindrical, less than 0.4 mm long, 2 - 5 times longer than wide; ochreae deciduous to commonly persistent in part on mature plants. Branches diverging at (20°-) 30°-40° - 60°(-80°) to the main stem and often weakly arching. Basal leaves narrowly elliptic, narrowly lanceolate to narrowly oblong-lanceolate, acute to acuminate, narrowly to widely cuneate, truncate or weakly cordate, 3 - 7 times longer than wide, evident at or near middle, crisped to entire; middle veins leaving lamina at 45° - 70°; glabrous above, distinctly pubescent on veins below; trichomes cylindrical, less than 0.3 mm long, 3 - 6 times as long as wide; petiole shorter than the lamina, glabrous to distinctly pubescent. Lower cauline leaves similar to basal but progressively smaller up the stem. Upper cauline leaves narrowly elliptic to linear-lanceolate, acute to acuminate, narrowly to widely cuneate, crisped or entire, glabrous above, somewhat papillose on veins below; trichomes papilliform, less than 0.1 mm long, 1.5 times longer to as long as wide; petiole much shorter than lamina, the leaf

sometimes subsessile. Inflorescence a wide, open panicle occupying the upper two-thirds to half of the stem, all branches subtended by leaves. Whorls commonly consisting of more than 20 flowers, present on branches and portion of main stem above branches, all subtended by leaf-like bracts similar to uppermost cauline leaves but progressively smaller up the stem and branches, the lower whorls remote, the upper contiguous. Pedicels filiform (1.5-) 2.5 - 4.0 - 5.8 (-8.5) mm long, less than 0.2 mm wide at middle, articulated near base; the articulation at most weakly evident. Sepals oblong, obtuse, (0.5-) 0.8 - 1.0 - 1.5 (-1.8) mm long and about half as wide, angled upwards along valve base. Valves deltoid, narrowly acute, cuneate, (1.5-) 2.3 - 2.5 - 3.0 (-3.5) mm long, (0.8-) 1.1 - 1.3 - 1.5 (-1.6) mm wide, widest in lower two-fifths to lower fifth, the margin below the apex spiny; distinct spines 1 - 2 - 3 per side, the largest (1.0-) 1.5 - 2.3 - 2.8 (-3.5) mm long; veins weakly raised, leaving midrib at ca 90° with 1 - 1 - 2 anastomoses on each side of valve; distance to first anastomosis from midrib 0.3 - 0.4 - 0.5 (-0.7) mm. Tubercles 3, fusiform to oblong-ovate, acuminate to subacute or obtuse at apex, obtuse at base, approximately equal in size, (1.0-) 1.2 - 1.5 - 1.6 (-2.0) mm long, 0.3 - 0.4 - 0.5 (-1.0) mm wide, 0.3 - 0.5 - 0.5 (-0.9) mm deep, widest in lower third to near base; base of tubercle usually at the valve base or sometimes up to 0.1 mm above. Anthers 0.3 - 0.5 (-0.7) mm long. Mature achenes ovate, light brown, (1.1-) 1.3 - 1.5 - 1.7 (-2.0) mm long, (0.6-) 0.7 - 0.8 - 0.9 (-1.1) mm wide, (0.4-) 0.6 - 0.7 - 0.8 (-0.9) mm deep, widest at middle, acute at the apex, broader at the base than at the apex.

In North America, four varieties are distinguished as follows:

key:

Basal and lower to mid-cauline leaves narrowly elliptical, tapering nearly equally at both ends; tubercles fusiform, acute to subacute at apex. var. maritimus

Basal and lower to mid-cauline leaves narrowly oblong to narrowly lanceolate, acute to acuminate at apex, widely cuneate, truncate or weakly cordate at base; tubercles fusiform or oblong-ovate, acute to obtuse at apex.

Tubercles fusiform, acute or subacute at apex, 0.3 - 0.5 mm wide. var. fueginus

Tubercles oblong-ovate, commonly obtuse at apex, 0.4 - 1.0 mm wide.

Tubercles less than 1.9 mm long, 0.4 - 0.7 mm wide, eastern coastal areas. var. persicarioides

Tubercles greater than 1.9 mm long, 0.7 - 1.0 mm wide, western coastal areas. var. pacificus

Rumex maritimus Linnaeus var. maritimus

Distribution: native in most of Europe except the extreme north and south (Rechinger 1964), Asia Minor, Siberia, Mongolia, China and Japan. (Losin-Losinskaya 1936). Introduced very rarely in North America.

United States: NJ, MA. (Map 18)

Rumex maritimus Linnaeus var. fueginus (Philippi) Dusen, Svensk.

Exped. Magell. 3(5) : 194. 1900.

R. fueginus Philippi, Ann. Univ. Chile 91:493. 1895.

Type locality: in Fuegia orientali, Febrero 1879 lecta.

R. maritimus subsp. fueginus (Philippi) Hulten, Ark. Bot., n.s. 7:45. 1968.

R. maritimus var. stethrix St. John, Rhodora 17 (196):79. 1915.

Type specimen: Utah: Vermilion, clay, altitude 5,400 ft.

Aug. 20, 1894, M. E. Jones 5839 (Holotype, NO; GH!, US 259496!)

R. fueginus var. brachythrix Rechinger f., Publ. Field Mus.

Bot. 17:143. 1937.

Lectotype: California: Siskiyou Co., Butte Valley, altitude

4,200 ft., saline flat, 12 Sept. 1910. G. D. Butler 1877

(US 669588!, RM 669588!)

R. fueginus var. tanythrix Rechinger f., Publ. Field Mus.

Bot. 17:143. 1937.

Lectotype: Oregon: Willows Co., "The Meadows" near spring

or deserted range, 4250 ft., 18 Aug. 1897, E. P. Sheldon 8718

(US 528399!)

R. fueginus var. ovata-cordatus Reehinger f., Field Mus.

Bet. 17:143. 1937.

Lectotype: California; Ventura Co., Oxnard. B. Dary

7804 (UC).

R. persicarioides auct. plur. non Linnaeus (1753).

R. maritimus auct. amer. nonn, non Linnaeus (1753).

Distribution:

Native in Mexico, Ecuador, southern Chile and Argentina (Mitchell 1978) and throughout most of North America; reportedly introduced elsewhere.

Canada: AB, BC, NT, MB, NK, NS, ON, PE, PU, SK, YT; United

States: AK, AZ, AR, CA, CO, CT, DE, ID, IL, IA, KS, ME, MA, MI,

MO, MT, MS, NV, NH, NY, ND, OH, OR, RI, SD, (TX?), UT, VA, WI,

WY. (Map 16).

Rumex maritimus Linnaeus var. persicarioides (Linnaeus) Mitchell,
Brittonia 30 (2); 295. 1978.

R. persicarioides Linnaeus; Sp. Pl. 335. 1753.

Type Locality: in Virginia.

Lectotype: Hb. LINN No. 464.9, labelled "9 persicarioides
7 K" (alm) (pinned to sheet 10) (LINN, DAO photo!)

R. maritimus auct. amer. nonn., non Linnaeus (1753.)

Distribution:

native in coastal areas of eastern North America, not
common.

Canada: (NB), NS, PE, QU. Reported from New Brunswick (Rechinger
1937), but no specimens referable to this variety as here
delimited have been seen.

United States: CT, MA, NY. (Map 16).

Rumex maritimus Linnaeus var. pacificus J. E. Dawson, var. nov.

Differt ab aliis varietatibus magnis tuberculis, quae sunt maiora
1.9mm longa et 0.7 - 1.0 mm. lata.

Type Specimen: Oregon: Lincoln Co., Seal Rock, beach, 31 Aug.
1921. M. E. Peck 10695 (F 7583201)

R. persicarioides sensu Reehinger (1937) in part, Munz
and Keck (1959), Peck (1961), Hitchcock (1964), Hitchcock
and Cronquist (1973).

*Distribution:

Native in coastal areas of western North America, not
common.

Canada: BC; United States: CA, OR, (WA). Reported from
Washington (Hitchcock and Cronquist 1973), but no specimens
referrable to this variety have been seen from that state. (Map 16)

Habitat: wet places such as marshes, edge of ponds, sloughs,
lakes, irrigation ditches. var. maritimus has only been reported
on ballast but the habitats of the other varieties appear to be
very similar.

Discussion:

In Species Plantarum, Linnaeus (1753) named R. maritimus
(from Europe) and R. persicarioides (from Virginia), two species
with very similar characteristics. The most apparent differences
were the linear leaves of the former, versus the lanceolate leaves
of the latter and that R. persicarioides had large pale grains
(tubercles)

St. John (1915) reported that a small specimen collected by Kalm (LINN No. 464.9) (pinned to a larger garden grown (Hortus Uppsalensis), specimen (LINN No. 464.10) and labelled "canade" (nsis)) was the type of R. persicarioides. After examining the records of Kalm's travels, St. John (1915) believed that Kalm's specimen was collected not in Virginia, but at a site on the Richelieu or the lower St. Lawrence River in Canada, an area in which similar plants are known to occur. Although plants corresponding to the type specimen are not known to range south of Long Island, New York, numerous 19th century authors, e.g. Hooker (1840), Meisner (1856), Trelease (1892), etc., employed the binomial, R. persicarioides for most American specimens. Hooker (1840) in employing R. persicarioides for California specimens, noted, that they were not really distinct from R. maritimus.

Lousley (1939) designated LINN 464.13 as the lectotype of R. maritimus. Examination of the photograph of this specimen and observations made on this specimen by Dr. J. McNeill in 1975 confirmed that it is referable to the variety to which the epithet maritimus is currently applied. As the specimen was collected by Hudson, it would not be the type if it was found to be a replacement. Some authors, e.g. Meisner (1856), Gray (1868), Watson (1880), Macoun (1883) and Rydberg (1932) applied the name to most American specimens, excluding those to which R. persicarioides had been applied.

St. John (1915) restricted the use of R. persicarioides to plants with large tubercles and short spines, on the valves

that occur in salt marshes and along saline shores in the New England States and southeastern Canada. Plants with narrow tubercles and long spines were recognized as R. maritimus; within this species, St. John recognized three varieties, two natives and one introduced. Var. maritimus, a rarely introduced taxon was separated from the other natives on the basis of the mid-cauline leaves being cuneate, rather than cordate or truncate at the base. The more common wide-ranging plant was said by St. John to be identical with the South American R. fueginus Philippi and as it was very similar in vegetative and fruit characteristics to the Eurasian, R. maritimus, it was called R. maritimus var. fueginus (Philippi) Dusen. In the drier parts of the western United States, St. John recognized plants in which the spines were virtually lacking as var. athrix.

Rechinger (1937) accepted St. John's treatment of R. persicarioides; St. John informed him of an Oregon specimen referable to this species and in his treatment, Rechinger recognized California and Oregon specimens as being referable to R. persicarioides. The west coast plants were "distinguished from the eastern American specimens only by the valves and grains being somewhat longer and the teeth of the valve relatively shorter". (Rechinger 1937). Rechinger (1937) did not however, accept St. John's treatment of R. fueginus as a variety of R. maritimus, but considered it to be a distinct species. He, however concluded that only the tubercle character was a reliable feature for separating the two species. In addition to recognizing

St. John's var. athrix as a variety of R. fueginus, three further varieties also separable only on the basis of their spine size (var. brachythrix, var. typicus, var. tanythrix) were described by Rechinger; Rechinger stated that these varieties were very distinct in extreme cases, connected by transitional forms and of little taxonomic significance. An additional variety, var. ovato-cordatus with ovate-cordate cauline leaves was described by Rechinger on the basis of one California specimen.

Most manuals and popular treatments have tended to follow either St. John (1915) or Rechinger (1937). Recent west coast treatments (Munz and Keck 1958, Hitchcock and Cronquist 1973) recognize R. persicarioides from coastal habitats from northern California to British Columbia.

In addition to Hooker's (1840) recognition that the North American plants were not clearly distinct from R. maritimus, Hitchcock (1964) noted that R. persicarioides was "a species so similar to R. maritimus, that the recognition of both is perhaps more mechanical than natural". The first author to place one of the binomials in synonymy with the other was apparently Macoun (1883), who treated R. persicarioides as a synonym of R. maritimus. On this basis, R. maritimus is the correct name when a single species is recognized as is done by Mitchell (1973). Within R. maritimus, Mitchell recognizes two native varieties, var. persicarioides and var. fueginus. In his treatment, var. persicarioides is represented by east and west coast populations with three "usually prominent pale plump-ellipsoid tubercles" and

"spines 0.5 - 1.5 mm" long, while var. fueginus is more widespread and with "narrower lanceolate tubercles" and "usually longer spines (up to 2.7 mm)" (Mitchell 1978). No measurements were given for the tubercle, even though it is the main feature separating the two native varieties.

The present study agrees with Mitchell in that only one species, R. maritimus, is recognized but based on the numerical analysis of the North American specimens (Intraspecific Numerical Studies, p155) it is proposed to recognize three native varieties, var. fueginus, var. persicarioides and var. pacificus. Var. persicarioides represents the large-tubercled east coast plants that are most similar to the Linnaean type, whereas var. pacificus represents the large tubercled west coast specimens, previously included in var. persicarioides, while var. fueginus represented the narrower, usually acute, tubercled, wide ranging specimens. As noted in the numerical analysis, although morphologically similar to var. persicarioides, var. pacificus can be readily separated from it and from var. fueginus on the basis of the larger tubercles. The morphological differences and the geographic separation were considered to represent sufficient reasons for recognizing the western large tubercled plants as a separate variety.

The introduced European variety, var. maritimus at present known from only two North American sites, is separable from the native varieties by the shape of its basal and lower to middle

cauline leaves. This is in agreement with the findings of Jorgensen (1971) that the other features given by Rechinger (1935b, 1937) are of little use in separating var. maritimus from introduced var. fueginus in Norway.

Mitchell (1976) considers "the characteristics expressed in Western Hemisphere populations to represent just a few permutations from a much wider range of variability found in the Eurasian species complex" and that the "New World taxa of this group merely represent facies of the worldwide species complex that have been dispersed to the Western Hemisphere on three or more separate occasions". As noted by Mitchell, further study of the Eurasian portion of the species complex would help in understanding the variability expressed by R. maritimus in North America.

R. maritimus is not easily confused with other species of Rumex occurring in North America with the exception of the apparently rare introduction of R. palustris. In Europe, confusion often arises in separating these two species; the characteristics used in differentiating them are dealt with in the discussion of R. palustris.

335

Mar 18 Distribution of R. maritimus var. maritimus,
var. fueginus, var. pacificus and var.
persicarioides.

Map 18

23.

Rumex palustris Smith, Fl. Brit. 1:394, 1800.

Type: England: By Acle Dam, Norfolk, 1781, Mr. Pritchford. Hb. Smith (LINN) (See discussion).

Annual, biennial or perennial. Root fusiform. Stem less than 1m tall, less than 1cm wide at base, leafy, alternately branched, erect, sulcate, red-brown, weakly papillose; trichomes cylindrical, less than 0.1 mm long, ca twice as long as wide; ochreae deciduous. Branches diverging at ca 55° to the main stem and often weakly arching. Basal leaves lanceolate, narrowly acute, narrowly cuneate, ca six times as long as wide, weakly crenulate-crisped; middle veins of lamina diverging at ca 60° to the midrib; lamina glabrous above, indistinctly papillose on veins below; trichomes papilliform, less than 0.1 mm long, shorter than to somewhat longer than wide; petiole equal to or longer than the lamina, glabrous to indistinctly papillose. Lower cauline leaves similar to basal but becoming oblong-lanceolate, acute, cuneate; petiole shorter than lamina, glabrous to indistinctly papillose. Upper cauline leaves oblong-lanceolate to narrowly linear-lanceolate, acuminate, narrowly cuneate, weakly crisped; glabrous above, indistinctly papillose on veins below; petiole much shorter than the lamina, glabrous, the leaf sometimes subsessile. Inflorescence a wide, open panicle, all of the branches subtended by leaves. Whorls commonly consisting of ca 20 flowers, present on branches and portion of main stem above branches, usually all subtended by leaf-like bracts, the

lower whorls remote, the upper becoming somewhat contiguous.

Bracts leaf-like, similar to uppermost cauline leaves but progressively smaller up the branches or stem. Pedicels weakly filiform, 3.3 - 5.0 mm long, ca 0.2 mm wide at middle, commonly articulated in the lower third; articulation distinctly swollen. Sepals oblong, acute, curving upwards, 1.0 - 2.0 mm long and about half as wide. Valves deltoid, acute, cuneate, 2.5 - 3.5 mm long, 1.0 - 1.3 mm wide, widest near base, the margin spiny; spines 2 - 3 per side, the longest 0.5 - 1.5 mm long; veins weakly raised, diverging from the midrib at ca 90° with 1 anastomosis on each side of the valve; distance to the first anastomosis from the midrib 0.4 - 0.5 mm. Tubercles 3, ovate, subacute at apex, obtuse at base, approximately equal in size, surface faintly reticulate, 1.5 - 2.0 mm long, 0.5 - 0.6 mm wide, 0.5 - 0.7 mm deep, widest in lower third; base of tubercle at or near base of valve. Anthers 0.9 - 1.0 mm long. Mature achene ovate, 1.8 - 2.0 mm long, 1.0 - 1.1 mm wide, 0.8 - 1.0 mm deep, widest in the lower two-fifths, acute at apex and base, somewhat more broadly acute at base.

Distribution: Native to almost all of Europe, except the north, but rare in the Mediterranean region; North Africa but extremely rare in Morocco. (Rechinger 1954). Introduced in North America, extremely rare. United States: CA, NJ (Map 16)

Habitat: In Europe, reported from marshes and riversides, but in North America, found only in ballast and on moist soil along drainage ditches.

Discussion: Lousley (1939) improperly attempted to alter the type of this species to England: St. George's Fields, London, 1800, Mr. E. Foster. Hb. Smith; his sole reason was that the specimen cited by Smith was immature. The specimen designated by Lousley (1939) can not be considered as the type which remains that designated by E. Smith; only when the holotype has been lost or destroyed, can a lectotype be designated.

R. palustris has not hitherto been reported in North America. The specimens upon which this report is based, were previously misidentified as R. maritimus and R. pulcher, collected in 1877 and 1959 respectively. It appears likely however, that these collections represent only chance introductions and the present status of the species in North America is very uncertain.

R. palustris

New Jersey: Camden, ballast, July 1877, Martindale (NY!)

California: Santa Barbara Co., 1.5 mi W. of Guadalupe, occasional in moist good soil in drainage ditch by an artichoke field, 2 Nov. 1959, Twisselmann 5867 (CAS 522038!)

R. palustris ($2n = 60$) differs cytologically from R. maritimus ($2n = 40$) and R. pulcher ($2n = 20$). Morphologically, it can be separated from these species by means of the characteristics employed in the key to the species. In Europe where R. palustris is native, confusion with R. maritimus is common; the morphological

characteristics separating these species have been discussed by Murbeck (1913), Lousley (1944) and Jorgensen (1971).

Rumex stenophyllus Ledebour, Fl. Altaica 2:58, 1830.

Type Locality: U.S.S.R.: Kirghiz S. S.R.; "in humidis deserti
sqongoro-kirg... v. gr. prope Belenki, ad radices mont. Ku".

Type Specimen reportedly at LE (Losin-Losinskaya 1936).

R. odontocarpus Sendor ex Borbas, Oesterr. Bot. Z. 37:334, 1887.

R. alluvius Gates and McGregor, Trans. Kansas Acad. Sci.
53:186, 1950.

Type: Kansas: collected along the Kansas R. north of
Lawrence in Douglas Co., July 4, 1949, McGregor 3328 (KANU)

Topotype: Kansas: wet alluvial floodland along the
Kansas R., 1½ mi. N. of Lawrence, July 2, 1950. McGregor
4373 (NY!, COLO!, US!, CAS!, SDU!, DAO!)

Perennial. Root fusiform. Stem 44 - 138 cm tall, less than 1 cm
wide at base, leafy, alternately branched above middle, brown to
red-brown often with red-purple areas, glabrous to weakly papillose;
trichomes most widest on branches near whorls, papilliform less
than 0.1 mm long, shorter than to as long as wide; ochreous usually
glabrous on mature plants. Branches diverging at 20° - 30° - 40°
to the main stem but often weakly arching. Basal leaves narrowly
lanceolate to oblong, acute, narrowly to broadly cuneate, up to
6 times as long as wide, entire to weakly crisped; middle veins
of lamina diverging at 40° - 80° to the midrib, commonly around
60°; lamina glabrous above, glabrous to indistinctly papillose on
veins below; trichomes papilliform, less than 0.1 mm long, shorter
than to as long as wide; petiole shorter than the lamina, glabrous

to indistinctly papillose. Lower cauline leaves very similar to basal but becoming progressively smaller up the stem. Upper cauline leaves linear-lanceolate to linear, narrowly acute, narrowly cuneate, becoming progressively smaller up the stem; lamina glabrous above, indistinctly papillose to usually glabrous below; petiole much shorter than the lamina, the leaf sometimes subsessile. Inflorescence a narrow, open panicle, occupying the upper half of the stem, the branches subtended by leaves. Whorls commonly consisting of 20 or more flowers per whorl, arranged on branches and main stem above branches, the lower whorls remote and subtended by leaf-like bracts, the upper whorls almost contiguous and not subtended. Bracts leaf-like, very similar to uppermost cauline leaves but becoming progressively smaller up the stem. Pedicels filiform, (2.5-) 5.3 - 7.0 - 8.5 (-11.3) mm long, less than 0.2 mm wide at middle, articulated in the lower two-fifths of the pedicel, commonly in the lower third, the articulation distinctly swollen. Sepals oblong, obtuse, (0.8-) 1.3 - 1.7 - 1.8 (-2.3) mm long and about half as wide; commonly at right angles to the pedicel or reflexed slightly downwards. Valves broadly deltoid, acute, truncate to weakly emarginate at the base, (2.5-) 3.8 - 4.3 - 4.8 (-5.3) mm long, (2.5-) 3.8 - 4.0 - 4.8 (-5.0) mm wide, widest at or near base, the margin below the apex dentate; teeth (4-) 5 - 6 - 8 (-10) per side, the longest (0.1-) 0.4 - 0.8 - 1.1 (-1.6) mm long; veins raised, diverging from the midrib at ca 90° with (1-) 2 - 3 anastomoses on each side of the valve; distance to the first anastomosis from the midrib 0.5 -

0.6 - 0.8 (-1.0) mm. Tubercles 3, ovate, obtuse at apex and base, approximately equal in size, (1.4-) 1.7 - 2.0 - 2.3 (-2.5) mm long, (0.8-) 0.9 - 1.0 - 1.3 (-1.4) mm wide, (0.7-) 0.6 - 1.0 - 1.0 (-1.2) mm deep, widest below the middle, base of tubercle usually at the valve base or up to 0.2 mm above, surface smooth but faintly reticulate. Anthers 0.6 - 0.9 mm long. Mature achene ovate, red-brown, (1.9-) 2.2 - 2.5 - 2.5 (-3.0) mm long, (1.0-) 1.2 - 1.3 - 1.5 mm wide, (0.9-) 1.0 - 1.2 - 1.3 mm deep, widest at or near the middle, acute at the apex and base.

Distribution: Native in central and eastern Europe (Rechinger 1964), the Caucasus, Siberia and central Asia (Losin-Losinskaya 1936). Introduced in North America, predominantly in the prairies and plains regions of Canada and the United States.

Canada: AB, MB, QU, SK; United States: CA, CO, KS, MN, MS, NE, NM, ND, OK, SC, UT, WA, WY, (Map 19)

Habitat: Occurring in damp, slightly saline soil, roadside ditches, marshy areas about lakes and in alluvial soil along streams; often spreading to road shoulders, waste areas or even crop fields.

Discussion: The earliest Canadian collection of R. stenophyllus is a plant collected in Saskatchewan in 1950. It and its American counterpart, an 1892 collection from Nebraska, were both originally misidentified as R. obtusifolius. Love and Bernard (1958) report finding specimens of R. stenophyllus misidentified as R. brittanica and R. pulcher. In 1950, Gates and McGregor realized that a species of dock known in Kansas and Nebraska since the 1930's could not be correctly identified with existing species in the treatment of North American Rumex (Rechinger 1937). Believing it to be native, they described it as a new species - R. alluvius. Seed samples under this name from Swift Current, Saskatchewan were sent by A. Budd to the Department of Agriculture, Ottawa, in 1954 and determined as R. stenophyllus (Frankton 1955). This led to the discovery of the earlier misidentifications. Meanwhile, in 1957, more reported the Eurasiatic R. odontocarpus Sandor ex Borbas as occurring in Minnesota, noting that it was the same species as Gates and McGregor had described as R. alluvius. R. odontocarpus is recognized to be conspecific with R. stenophyllus (Love and Bernard 1958).

Discussion of other names that have been applied to R. stenophyllus in Europe can be found in Love and Bernard (1958). The first introduction of the species may have been from grain brought to the United States in the late 1800's and early 1900's, by settlers from central and Eastern Europe (Love and Bernard 1958).

However, a recent introduction (based on a 1959 collection) in South Carolina was probably the result of an introduction of the species with imported wool as it was found in the vicinity of a wool combing mill. An additional wool alien - R. brownii from Australia was found in the same general area.

345

Mar 19 - Distribution of R. stenophyllus.

Map 19

25.

Rumex brownii Campdera, Monogr. Rumex, 64, 81. 1819.

Type: Australia: New South Wales: "ad Portum Jackson" (near Sydney) Dunn. Isotypes: BM (ex herb. Banks; G-DC).

Perennial. Stems 80 cm long, 6mm wide at base, erect or ascending, sulcate, red-brown, leafy, alternately branched, glabrous; ochreous partially persistent on mature plants. Branches diverging at 30° - 50° to the main stem but often weakly arching. Basal leaves narrowly lanceolate, acute, broadly cuneate to weakly cordate, narrowed above the base, 4 - 6 times as long as wide, widest at or near the base, the largest observed leaf 14.2 cm long, 3.1 cm wide, crisped; veins of lamina leaving midrib at 40° - 60°; lamina glabrous above, glabrous to indistinctly papillose on veins below; the trichomes papilliform, less than 0.1 mm long, shorter than to as long as wide; petiole shorter than to about as long as lamina, glabrous to indistinctly papillose. Lower cauline leaves very similar to basal, but becoming progressively smaller up the stem; petiole shorter than the lamina. Upper cauline leaves very narrowly lanceolate to linear, narrowly acute, narrowly cuneate, crisped; glabrous above and below; petiole much shorter than the lamina. Inflorescence a wide, open panicle, all branches subtended by leaves. Whorls commonly consisting of less than 10 flowers arranged on branches and portion of main stem above branches, only a few lower whorls subtended by leaf-like bracts, very similar to uppermost cauline leaves. Pedicels filiform, 2.3 - 5.0 mm long, less than 0.4 mm wide at middle, articulated in the lower third;

articulation distinctly swollen. Sepals oblong, acute, reflexed at right angles to pedicel, 1.3 - 2.0 mm long and about half as wide. Valves deltoid, narrowly acute and hooked, broadly cuneate at base, 2.6 - 4.3 mm long, 1.5 - 2.3 mm wide, the margin spiny; teeth ~~3~~(4) per side, hooked, the longest 0.8 - 1.3 mm long; veins diverging from the midrib at ca 90°, with 1 - 2, commonly 1 anastomosis on each side of the valve; distance to the first anastomosis from the midrib 0.5 - 0.6 mm. Tubercles lacking. Anthers 0.7 - 0.9 mm long. Mature achene elliptic, red-brown, 2 - 2.3 mm long, 1.3 - 1.5 mm wide, 1.0 - 1.3 mm deep, widest at or near middle, apex and base equally acute or the apex narrower than the base.

Distribution: Native of Australia, Java, Timor, New Guinea (Rechinger 1949). Introduced in parts of Europe (Clapham et al. 1962). Rechinger (1935) and North America.

United States: SC (Map 16)

Habitat: Spreading on waste ground, initially near wool combing mills.

Discussion:

My examination of Rumex specimens from South Carolina that had been collected during the late 1950's, revealed several specimens that did not fit the characteristics of any species little to known to occur in North America.. These specimens resembled R. pulcher, superficially. However, two striking differences from R. pulcher were evident: 1) a total lack of

tubercles, 2) the marginal spines and the apex of the valve recurved at the tips. As the habitats for the collection were all waste fields beside wool combing mills, it seems likely that the source of introduction of the species may have been the arrival of seeds with foreign wool shipments. The recurved hooks on the valves would certainly adhere to the wool. An examination of lists of common "wood aliens" (Dony 1953; Salisbury 1961) showed that only one Rumex species - R. brownii Campdera, had been reported as being regularly introduced with wool shipments. Published descriptions (Rechinger 1935, Clapham et al. 1962) of this species made it clear that the plants from South Carolina are referable to R. brownii. This species has been included as an aberrant member of the section Axillares (Rechinger 1935, 1949) but in more recent literature (Clapham et al. 1962; Degraeve 1975) it is included in section Rumex. R. brownii is reported to persist in localities in Britain for several years after introduction (Clapham et al. 1962). Whether or not it is still present in the South Carolina region is uncertain; the climate of the Carolinas would certainly be mild enough to allow for its survival. Also, reintroduction via new wool shipments would also be likley.

Summary and Conclusions

Section Rumex in Canada and the United States is now recognized to consist of twenty-five species; fifteen of these and one variety of another represent introduced taxa while the remaining taxa are native. Preliminary observations on the species of section Rumex indicated that most could be readily discriminated from each other utilizing morphological and/or cytological characteristics. Because of the limited material of R. orthoneurus, R. pycnanthus, R. densiflorous and R. praecox these species have been retained as separate taxa but as Rechner (1937) noted, they could represent one collective species. Previous treatments had maintained R. occidentalis and R. fenestratus as separate species but the preliminary observations did not support this; a similar situation existed in the recognition of R. maritimus and R. persicarioides as separate species. Traditional taxonomic methods combined with numerical methods were employed to resolve these two problems. Confusion between the species was found to have been based on poor understandings of the variation within the species and especially of the characteristics needed to differentiate them. The present study has examined herbarium material and where possible field and cultivated specimens of the North American species of section Rumex in order to provide a better understanding of the variation possible within the species. The observations made during the study have been used in providing better taxonomic

descriptions and a key for the discrimination of the species.

Experiments were carried out that indicated that the species are self-fertile producing high percentages of fertile pollen and large numbers of viable achenes. Within section Rumex, the chromosome number is very variable ranging from $2n = 20$ to $2n = ca. 200$; pairing during meiosis proceeds normally. The features of the breeding biology and the cytology of the species can be used as guidelines in the recognition of plants of the species or in indicating hybrids.

In Europe, hybrids involving almost every possible pairwise combination of species have been reported while in North America, only 11 different hybrids have been identified. Although the number of hybrids is now greater than that reported by Rechinger (1937), only one hybrid, R. crispus x obtusifolius can be considered as being widespread and relatively common. Factors contributing to the paucity of the hybrids are their tendency to produce low percentages of viable pollen and few viable achenes; observations made during meiosis of R. crispus x obtusifolius and R. longifolius x obtusifolius have indicated that pairing during meiosis in the F₁'s is very irregular. However, as indicated by the study of Williams (1971), backcrosses to one of the parents can restore a degree of fertility and viability to subsequent hybrid generations. In the course of this study no hybrid sites were observed where introgression was taking place and the examination of the herbarium material obtained on loan from the major herbaria did not reveal any apparent confusion between the species brought about by intro-

gression. Generally, the F1 hybrids can be easily identified morphologically as their characteristics combine some attributes of each parent whereas hybrids involving backcrosses with one of the parents can become morphologically very similar to one of the parents being discernible by means of a hybrid index or an examination of pairing at meiosis. Because of the apparent lack of introgression, hybridization in section Rumex in North America does not represent a major area of difficulty in species recognition. Further experimental studies are needed to determine to what extent hybridization and introgression are possible in the species occurring in North America. Ecological studies on the requirements of the species are also needed to further understand the factors restricting the occurrence of the hybrids and the maintaining of separate species in nature.

To provide further assistance in the recognition of the species occurring in North America, comparisons of many of the species with those species with which they have been confused have been included in the taxonomic treatment of the species. Identification of any infraspecific taxa recognized within the species has been provided by the keys included with those species. The infraspecific units recognized in many of the introduced European species were generally not found to be of use in circumscribing the variation observed within the species in North America.

Within the native species R. occidentalis, two varieties have been recognized while in the species R. maritimus, three

native varieties and one introduced variety have been recognized.

The distribution maps provided for each species represent an up-to-date representation of the range of the species based on an examination of approximately 9,000 herbarium specimens from the major herbaria and herbaria chosen to ensure coverage of the species ranges. As many of the introduced species tend to be somewhat weedy in nature, their ranges will always be in a state of flux. The increased ranges of the native species recognized by Rechinger (1937) are probably due to more specimens being available for the present study rather than active expansion of the species.

Only one nomenclatural problem was found to exist within section Rumex, the use of R. britannica or R. orbiculatus for a native species. This problem was resolved as a type specimen of R. britannica was found to exist in the Linnaean herbarium thus placing R. orbiculatus in synonymy.

The present study has provided a better understanding of the species and of the hybridization occurring within section Rumex in Canada and the United States which will help to eliminate the previous confusion observed within the section.

BIBLIOGRAPHY

- Ascherson, P. and Graebner, M. 1908. Synopsis der Mitteleuropaischen Flora. Wilhem Engelmann. Leipzig.
- Boivin, B. 1966. Énumération des plants du Canada. III - Herbidiées, 1^{re} partie: Digitatae: Dimerae, Liberae. Naturaliste Canadien 93:583-646.
- Boivin, B. 1968. Flora of the Prairie Provinces. Part II Provaqcheria 3. Université Laval, Quebec.
- Borodina, A.E. 1979. Main directions of evolution in the genus Rumex L. (Polygonaceae) and its system. Botanicheskii Zhurnal 64(4):541-553.
- Britton, N. L. and Brown, A. 1896. Illustrated Flora of the Northern United States, Canada and the British Possessions. Vol. I. Charles Scribner's Sons. New York.
- Budd, A.C. and Best, K.F. 1964. Wild Plants of the Canadian Prairies. Canada Dept. of Agriculture. Publication 983.
- Campdera, F. 1819. Monographie des Rumex précédée de quelque vues générales sur la famille de Polygonacées. Paris.
- Cavers, P.B. and Harper, J.L. 1964. Rumex obtusifolius L. and R. crispus L. Journal of Ecology 52(3): 737-766.
- Cavers, P.B. and Harper, J.L. 1966. Germination polymorphism in Rumex crispus and Rumex obtusifolius. Journal of Ecology 54(2): 367-382.
- Cavers, P.B. 1974. Germination polymorphism in Rumex crispus. The effects of different storage conditions on germination responses of seeds collected from individual plants. Can. J. Bot. 52(3): 575-583.
- Calder, J.A. and Taylor, R.L. 1968. Flora of the Queen Charlotte Islands. Pt. 1. Systematics of the Vascular Plants. Can. Dept. Agric. Res. Branch Monogr. 4(1): 1-659.
- Chancellor, P. 1956. Studies on the ecology of some species of the genus Rumex in Proceedings of Third British Weed Control Conference 1:197-203.

- Clapham, A.R., Tutin, T.J. and Warburg, E.F. 1962. Flora of the British Isles. University Press, Cambridge.
- Colden, C. 1743. Plantae Coldenhamiae, in Provincia Novaboracensis Americes. Acta Soc. Reg. Upsal.
- Correll, D.S. and Johnston, M.C. 1970. Manual of the Vascular Plants of Texas. Texas Research Foundation, Renner.
- Correll, D.S. and Correll, H.B. 1972. Aquatic and Wetland Plants of Southwestern United States. Environmental Protection Agency.
- Cullen, J. 1967. Rumex. in Davis, P.H. ed. Flora of Turkey 2:281-293. University Press, Edinburgh.
- Deam, C.C. 1940. Flora of Indiana. Department of Conservation, Indianapolis.
- Degraeve, N. 1975. Contribution a l'etude cytotaxonomique des Rumex - I. Le genre Rumex L. sensu stricto. Caryologia 28(2): 187-201.
- Degraeve, N. 1975b. Contribution a l'etude cytotaxonomique des Rumex II. Le genre Bucephalophora Pau. Caryologia 28(2):203-206.
- Dony, J.G. 1953. Wool aliens in Bedfordshire, in Lousley, J.E. (ed.) The Changing Flora of Britain. T. Buncle and Co. Arbroath. p. 160-163.
- Eaton, A. 1824. Manual of Botany for the Northern and Middle States of America. Websters and Skinner. Albany.
- Fernald, M.L. 1921. The Gray Herbarium expedition to Nova Scotia. Rhodora 23:257-278.
- Fernald, M.L. 1945. So-called Rumex britannica not a Virginian. Rhodora 47:133-137.
- Fernald, M.L. 1950. Gray's Manual of Botany. ed. 8. American Book Co. New York.
- Gates, F.C. and McGregor, R.L. 1950. Rumex alluvius, a new species of dock from Kansas. Trans. Kans. Acad. Sci. 53: 186-189.
- Gleason, H.A. and Cronquist, A. 1963. Manual of Vascular Plants of Northeastern United States and Adjacent Canada. Van Nostrand. Princeton, N.J.

UNITED STATES, cont.

MI	Michigan	OR	Oregon
MN	Minnesota	PA	Pennsylvania
MO	Missouri	RI	Rhode Island
MT	Montana	SC	South Carolina
NB	Nebraska	SD	South Dakota
NV	Nevada	TN	Tennessee
NH	New Hampshire	TX	Texas
NJ	New Jersey	UT	Utah
NM	New Mexico	VT	Vermont
NY	New York	VA	Virginia
NC	North Carolina	WA	Washington
ND	North Dakota	WV	West Virginia
OH	Ohio	WI	Wisconsin
OK	Oklahoma	WY	Wyoming

- Hultén, E. 1968. Flora of Alaska and Neighbouring Territories. Stanford University Press. Stanford.
- Hultén, E. 1971. The Circumpolar Plants. II Dicotyledons. Almquist and Niksell. Stockholm.
- Ichikawa, S., Sparrow, A.H., Frankton, C., Nauman, A.F., Smith, E.B. and Pond, V. 1971. 'Chromosome number, volume and nuclear volume relationships in a polyploid series (2X-20X) of the genus Rumex. Can. J. Genet. Cytol. 13: 842-863.
- Jackowska, I. 1964. See Listowski, A. and Jackowska, I.
- Jalas, J. and Lindholm, R.K. 1975. Biosystematics of Rumex longifolius D.C. of Fennoscandia and the Pyrenees. Anal. Inst. Bot. Cavanilles 32(2): 197-202.
- Jardine, N. and Sibson, R. 1970. Quantitative attributes in taxonomic descriptions. Taxon 19: 862-870.
- Jensen, H.V. 1936. Meiosis in Rumex. I. Polyploidy and the origin of new species. Cytologia 7: 1-22.
- Jepson, N.L. 1923. A Manual of the Flowering Plants of California, Berkeley.
- Johnson, A.W. and Packer, J.G. 1965. Chromosome numbers in the flora of Ogotoruk Creek, N. Alaska. Botan. Notiser 121: 403-456.
- Jorgensen, Per Magnus. 1971. Rumex maritimus L. and R. palustris Sm. in Norway. Blyttia 29(3): 133-139.
- Johnson, A.W. and Packer, J.G. 1968. Chromosome numbers in the flora of Ogotoruk Creek, N.N. Alaska. Botan. Notiser 121: 403-456.
- Kihara, H. 1927. Über die vorbehandlung einiger pflanzlicher objekte bei der fixierung der pollen-mutterzellen. Bot. Mag.
- Knuth, P. 1909. Handbook of Pollination. Vol. III. Clarendon Press. Oxford.
- Lance, G.N. and Williams, W.T. 1971. 1967. A general theory of classificatory sorting strategies - 1. Hierarchical systems. Comput. J. 9: 373-380.
- Lawalree, A. 1953. Flore Générale de Belgique. Spermatophytes. Ministère de l'Agriculture Bruxelles.
- Lawrence, G.H. 1959. Taxonomy of Vascular Plants. The MacMillan Co., New York.

- Lepage, E. 1954. Études sur quelques plantes américaines. III
Naturaliste Canadien 81(3-4): 59-68.
- Lepage E. 1954. Nouveautés dans la flore de la baie James.
Naturaliste Canadien 81(12): 235-261.
- Linnaeus, C. 1749. *Materia Medica*. Stockholm
- Linnaeus, C. 1737. *Genera Plantarum*. Leyden.
- Linnaeus, C. 1753. *Species Plantarum*. Laurentii Salvii, Holmiae.
- Listowski, A. and Jackowska, I. 1964. Observations on plant
development (VIII). The development of Rumex
obtusifolium. Acta Societatis Botanicorum Poloniae
33(4): 705-717. (in text as Jackowska 1964)
- Losin - Losinskaya, A. 1936. Rumex. In Flora of the U.S.S.R.
Vol. V. Moscow and Leningrad pp. 444-482.
- Lousley, J.E. 1939. Notes on British Rumices. I. Bot. Exch.
Club. Rep. 12(1): 118-157.
- Lousley, J.E. 1944. Notes on British Rumices. II. Bot. Exch.
Club Rep. 12(5): 547-585.
- Lousley, J.E. and Williams, J.T. 1975. Rumex in Stace, A.
(ed.) Hybridization and the Flora of the British
Isles. Part 2. Hybridization. No. 325. Academic
Press. London.
- Löve, A. and Bernard, F.J.P. 1958. Rumex stenophyllus in
North America. Rhodora 60: 54-57.
- Löve, A. and Kapoor, B.M. 1967. A chromosome atlas of the
collective genus Rumex. Cytologia 32: 328-342.
- Löve, A. and Love, D. 1961. Chromosome numbers of central and
northwest European plant species. Opera Bot. 5:581.
- Löve, A. and Love, D. 1967. In IOPB chromosome numbers report
XIII. Taxon 16: 191-192.
- Löve, A. and Patil, S. 1967. In IOPB chromosome number reports
XIII. Taxon 16: 445-461.
- Löve, A. and Ritchie, J.C. 1966. Chromosome numbers from central
northern Canada. Can. J. Botany 44: 429-439.
- Löve, A. 1967. In IOPB chromosome numbers reports XIII.
Taxon 16: 423-425.

- Love A. 1975. In IOPB chromosome numbers report LIV
Taxon 25: 631-649.
- Macoun, J. 1883. Geological Survey of Canada. Part 1-
Polypetalae. Dawson Brothers. Montreal.
- Marek, S. 1954. Morphological and anatomical characteristics
of fruits of genera of Polygonum L. and Rumex L.
and keys for their identification. Polskie Towarzystwo
Botaniczne, Monographiae Botanicae 11: 77-161.
- Marriott, F. 1974. See Addenda.
- Martin, A.C. Barkley, N.D. 1931. Seed Identification Manual.
Univ. of California Press. Berkeley.
- Maun, M.A. and Cavers, P.B. 1969. Influence of Photoperiod on
flowering of Rumex crispus L. Agron. J. 61: 823
- Maun, M.A. and Cavers, P.B. 1970. Influences of soil tempera-
ture on reproduction of Curley Dock. Weed Sci.
18(2): 202-204.
- McGregor, R.L. and Barkley, T.M. et al. 1977. Atlas of the
Flora of the Great Plains. The Iowa State University
Press. Ames.
- McNeill, J. 1975. A generic revision of Portulacaceae tribe
Montieae using techniques of numerical taxonomy. Cana-
dian Journal of Botany 53(8): 789-809.
- McNeill, J. 1976. The taxonomy and evolution of weeds.
Weed Research 16: 398-413.
- McSwain, J. and Bain, T. 1891. List of Prince Edward Island
Plants. Natural History of Prince Edward Island.
- Meisner, K.F. 1856. Polygonaceae. in De Candolle, A.
Prodromus Systematis Naturalis Regni Vegetabilis. Volume XIV.
Victoris Masson. Paris.
- Michaux, A. 1803. Flora Boreali Americana. Hafner Press.
New York. (1974 facsimile).
- Mitchell, R.S. 1978. Rumex maritimus L. versus R. persicarioides
L. (Polygonaceae) in the western hemisphere. Brittonia
30(2): 293-298.
- Montgomery, F.H. 1977. Seeds and Fruits of Eastern Canada and
Northeastern United States. University of Toronto
Press. Toronto.
- Moore, J.W. 1957. Notes on flowering plants in Minnesota.
Rhodora 59: 608.

- Moore, R.J. ed. 1973. Index to Plant Chromosome Numbers 1967-1971. Oosthoek's Uitgeversmaatschappij B.V. Utrecht.
- Mulligan, G.A. 1957. Chromosome numbers of Canadian weeds. I. Canadian Journal of Botany 35: 779-789.
- Mulligan, G.A. 1959. Chromosome numbers of Canadian weeds. II Can. J. Botany 37: 81-92.
- Mulligan, G.A. 1965. Recent colonization by plants in Canada. pp. 138-139. in Bakker, H.G. and Stebbins, G.L. ed. The Genetics of Colonizing Species. Academic Press. New York.
- Mulligan, G.A. 1969. In IOPB chromosome reports XX. Taxon 18: 213-221.
- Mulligan, G.A., and Porsild, A.E. 1969. Chromosome numbers of some plants from the unglaciated central Yukon plateau. Can. J. Botany 47: 655-662.
- Mulligan, G.A. and Frankton, C. 1972. Chromosome races in Rumex arcticus (Polygonaceae) Can. Bot. 50(2): 378-380.
- Munz, P.A. and Keck, D.D. 1959. A California Flora, University of California Press. Berkeley.
- Murbeck, S. 1899. Die nordeuropaischen Formen der Gattung Rumex. Bot. Not. 1899. Rumex fennicus. 17.
- Murbeck, S. 1913. Beitrag zur Kenntnis der Gattung Rumex. Bot. Not. 1913: 201-237.
- Musil, A.F. 1963. Identification of Crop and Weed Seeds. Agriculture Handbook No. 219. U.S. Dept. of Agriculture. Washington.
- Nuttall, J. 1818. Genera of North American Plants. D. Heartt Philadelphia.
- Osterhout, G.E. 1898. A new species of Rumex from Colorado: R. densiflorus Erythea 6: 13.
- Porter, C.L. 1967. Taxonomy of Flowering Plants. W.H. Freeman and Company. San Francisco.
- Pursh, F. 1814. Flora Americanae Septentrionalis. White, Cochrane and Co. London.
- Pursh, F. 1816. Flora Americanae Septentrionalis. Black, London.

- Radford, A.C., Alles, H.E. and Bell, C.R. 1968. Manual of the Vascular Flora of the Carolinas. The University of North Carolina Press. Chapel Hill.
- Rechinger, K.H. 1929. Beitrage zur Kenntnis von Rumex, I, Sekt. Lapathum XXI. Repert. Spec. Nov. 26:12.
- Rechinger, K.H. 1932. Vorarbeiten zu einer Monographie der Gattung Rumex I. Beih. Bot. Centralbl. 49: 1-132.
- Rechinger, K.H. 1933. See Addenda.
- Rechinger, K.H. 1935. Vorarbeiten zu einer Monographie der Gattung Rumex IV. Österreichische Botanische Zeitschrift 84: 31-52.
- Rechinger, K.H. 1935b. Die sud- und zentralamerikanischen Arten der Gattung Rumex. Ark. Bot. 26. A 3.
- Rechinger, K.H. 1937. Vorarbeiten zu einer Monographie der Gattung Rumex V. The North American Species of Rumex. Publ. Field Mus. (Bot.) 17(1): 1-151.
- Rechinger, K.H. 1948. See Addenda.
- Rechinger, K.H. 1949. Vorarbeiten zu einer Monographie der Gattung Rumex VII. Rumices Asiatici. Candollea 12: 9-152.
- Rechinger, K.H. 1954. Vorarbeiten zu einer Monographie der Gattung Rumex VIII. Monograph of the genus Rumex in Africa. Bot. Notis. Suppl. 3(3): 1-114.
- Rechinger, K.H. 1961. Beitrag zur Kenntnis von Rumex. No. XV. Watsonia 5: 64-66.
- Rechinger, K.H. 1964. Rumex in Tutin, T.G., Heywood, V.H., Burges, N.A., Valentine, D.H., Walters, S.M., and Webb, D.A. (ed.) Flora Europaea. Volume I. University Press, Cambridge.
- Robinson, B.L. and Fernald, M.L. 1908. Gray's New Manual of Botany 7th ed. American Book Co., New York.
- Roland, A.E., and Smith, E.C., 1969. The Flora of Nova Scotia. Nova Scotia Museum. Halifax.
- Rydberg, Per Axel, 1932. Flora of Prairies and Plains of Central North America. New York Botanical Gardens. New York.
- ~~St. John, W. 1915. Rumex persicarioides and its allies in North America. Rhodora 17: 73-83.~~
- Salisbury, E.J. 1942. The Reproductive Capacity of Plants, Studies in Quantitative Biology. G. Bell. London.

- Salisbury, E.J. 1961. Weeds and Aliens. Collins. London.
- Sarkar, N. 1958. See Addenda.
- Scoggan, H.J. 1978. The Flora of Canada. Part 3, Dicotyledoneae. National Museums of Canada. Ottawa.
- Seymour, F.C. 1969. The Flora of New England. The Charles E. Tuttle Co. Rutland.
- Small, J.K. 1903. Flora of the Southeastern United States, New York.
- Small, J.K. 1933. Manual of the Southeastern Flora. University of North Carolina Press. Chapel Hill.
- Sneath, P.A., and Sokal, R.R. 1973. Numerical Taxonomy. W.H. Freeman and Co. San Francisco.
- Snow, R. 1963. Alcoholic Hydrochloric Acid-Carmine as a stain for chromosomes in squash preparations. Stain Technology 38: 9-13.
- Sokal, R.R. and Sneath, P.A. 1963. See Addenda.
- Stafleu, F.A. et al. 1978. International Code of Botanical Nomenclature. Bohn, Schetema and Holkema and Int. Assoc. Plant Taxon. Utrecht.
- Stearn, W.T. 1957. An Introduction to the Species Plantarum and Cognate Botanical Works of Linnaeus in Species Plantarum, a Facsimile of the First Edition. Ray Society. London.
- Stevens, O.A. 1950. Handbook of North Dakota Plants. North Dakota Agricultural College. Fargo.
- Steyermark, J.A. 1963. Flora of Missouri. The Iowa State University Press. Ames.
- Strausbaugh, P.D., and Core, E.L., 1964. Flora of West Virginia. West Virginia University. Morgantown.
- Taylor, R.L., and Mulligan, G.A., 1968. Flora of the Queen Charlotte Islands, Part 2. Cytological aspects of the vascular plants. Queen's Printer, Ottawa.
- Thompson, K., Grime, J.P., and Mason, G. 1977. Seed germination in response to diurnal fluctuations of temperature. Nature 267: 147-149.
- Trelease, W. 1892. Revision of the American species of Rumex occurring north of Mexico. Missouri Botanical Garden, Third Ann. Rept. St. Louis.
- Watkins, A.E. 1925. Cotton blue - lacto-phenol method. Journ. Genet. IV: 340.

- Watson, S. 1877. Rumex occidentalis. Proc. Amer. Acad. 12: 253
- Watson, S. 1880. Rumex. In: Geological Survey of California. vol. 2, Botany. pp 7-10. J. Wilson and Son, Univ. Press. Cambridge.
- Welsh, S.L. 1974. Anderson's Flora of Alaska and Adjacent Parts of Canada. Brigham Young University Press. Provo.
- Williams, J.T. 1971. Seed polymorphism and germination. II The role of hybridization in the germination polymorphism of Rumex crispus and R. obtusifolius. Weed Res. 11: 12-21.
- Addenda
- Gower, J.C. 1966. Some distance properties of latent root and vector methods used in multivariate analysis. Biometrika 53: 325-328.
- Rechinger, K.H. 1933. Vorarbeiten zu einer Monographie der Gattung Rumex. Die Arten der Subsektion Patientiae. Repert. Sp. Nov. 31: 225-283.
- Rechinger, K.H. 1948. Beitrage zur Kenntnis von Rumex IX. Candollea 11: 229-241.
- Sarkar, N.M. 1958. Cytotaxonomic studies on Rumex section Axillares. Can. J. Bot. 36(6): 947-996.
- Sokal, R.R. and Sneath, P.A. 1963. Principles of Numerical Taxonomy. W.H. Freeman and Co. San Francisco.
- Marriott, F. 1974. The Interpretation of Multiple Observations. Academic Press. London.

Herbaria-Consulted in Study

Loans from the following herbaria were obtained and utilized in the morphological and distributional aspects of the study:

1. University of Colorado, Boulder, Colorado, U.S.A. (COLO).
2. University of Alberta, Calgary, Alberta, Canada (UAC).
3. Gray Herbarium of Harvard University, Cambridge, Massachusetts, U.S.A. (GH).
4. Herbarium of University of North Carolina, Chapel Hill, North Carolina, U.S.A. (NCU).
5. Chicago Natural History Museum, Chicago, Illinois, U.S.A. (F).
6. University of Alberta, Edmonton, Alberta, Canada (ALTA).
7. North Dakota State University and Experiment Station, Fargo, North Dakota, U.S.A. (NDA).
8. Rocky Mountain Herbarium, University of Wyoming, Laramie, Wyoming, U.S.A. (RM).
9. University of Nebraska State Museum, Lincoln, Nebraska U.S.A. (NEB).
10. Herbarium of Montana State University, Missoula, Montana, U.S.A. (MONTU).
11. McGill College Herbarium, McGill University, Montreal, Canada (MTTG).
12. The New York Botanical Garden, New York, New York, U.S.A. (NY).
13. Museum of Natural Sciences, National Museums of Canada, Ottawa, Canada (CAN).
14. Vascular Plant Herbarium, Biosystematic Research Institute, Dept. of Agriculture, Ottawa, Canada (DAO).

15. Saskatchewan Museum of Natural History, Regina, Saskatchewan, Canada. (USAS).
16. California Academy of Science, San Francisco, California, U.S.A. (CAS).
17. The W. P. Fraser Memorial Herbarium, University of Saskatchewan, Saskatoon, Saskatchewan, Canada. (SASK).
18. Department of Botany, University of Toronto, Toronto, Ontario, Canada. (TRT).
19. University of Arizona, Tucson, Arizona, U.S.A. (ARIZ).
20. The University of British Columbia, Vancouver, British Columbia, Canada. (UBC).
21. Provincial Museum, Victoria, British Columbia, Canada. (V).
22. U.S. National Museum, Smithsonian Institution, Washington, DC. U.S.A. (US).
23. Herbarium of the Department of Botany, University of Manitoba, Winnipeg, Manitoba, Canada. (MB).
24. Acadia University, Wolfville, Nova Scotia, Canada (ACAD).

DISTRIBUTION ABBREVIATIONS

The following abbreviations have been employed in this study.

CANADA

AB	Alberta	NF	Newfoundland excl. Labrador
AA	Arctic Archipelago	NS	Nova Scotia
BC	British Columbia	ON	Ontario
NT	Northwest Territories	PE	Prince Edward Island
LB	Labrador	QU	Quebec
MB	Manitoba	SK	Saskatchewan
NK	New Brunswick	YT	Yukon Territory

UNITED STATES

AL	Alabama	GA	Georgia
AK	Alaska, excl. Aleut I.	ID	Idaho
AI	Aleutian Islands	IL	Illinois
AZ	Arizona	IN	Indiana
AR	Arkansas	IA	Iowa
CA	California	KS	Kansas
CO	Colorado	KY	Kentucky
CT	Connecticut	LA	Louisiana
DE	Delaware	ME	Maine
DC	District of Columbia	MD	Maryland
FL	Florida	MA	Massachusetts

UNITED STATES, cont.

MI	Michigan	OR	Oregon
MN	Minnesota	PA	Pennsylvania
MO	Missouri	RI	Rhode Island
MT	Montana	SC	South Carolina
NB	Nebraska	SD	South Dakota
NV	Nevada	TN	Tennessee
NH	New Hampshire	TX	Texas
NJ	New Jersey	UT	Utah
NM	New Mexico	VT	Vermont
NY	New York	VA	Virginia
NC	North Carolina	WA	Washington
ND	North Dakota	WV	West Virginia
OH	Ohio	WI	Wisconsin
OK	Oklahoma	WY	Wyoming

END

1 6 1 2 8 1

FIN