**PM-1 3½"x4" PHOTOGRAPHIC MICROCOPY TARGET
NBS 1010a ANSI/ISO #2 EQUIVALENT**

PRECISIONSM RESOLUTION TARGETS

National Library
of Canada

Acquisitions and
Bibliographic Services Branch

395 Wellington Street
Ottawa, Ontario
K1A 0N4

Bibliothèque nationale
du Canada

Direction des acquisitions et
des services bibliographiques

395, rue Wellington
Ottawa (Ontario)
K1A 0N4

Your file: Votre référence

Our file: Notre référence

NOTICE

The quality of this microform is heavily dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to ensure the highest quality of reproduction possible.

If pages are missing, contact the university which granted the degree.

Some pages may have indistinct print especially if the original pages were typed with a poor typewriter ribbon or if the university sent us an inferior photocopy.

Reproduction in full or in part of this microform is governed by the Canadian Copyright Act, R.S.C. 1970, c. C-30, and subsequent amendments.

AVIS

La qualité de cette microforme dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

S'il manque des pages, veuillez communiquer avec l'université qui a conféré le grade.

La qualité d'impression de certaines pages peut laisser à désirer, surtout si les pages originales ont été dactylographiées à l'aide d'un ruban usé ou si l'université nous a fait parvenir une photocopie de qualité inférieure.

La reproduction, même partielle, de cette microforme est soumise à la Loi canadienne sur le droit d'auteur, SRC 1970, c. C-30, et ses amendements subséquents.

Canada

**SIGNAL TRANSDUCTION AND THE FUNCTION OF SECOND MESSENGERS
AND PROTEIN KINASES IN THE CONTROL OF GLYCOGENOLYSIS AND
CRYOPROTECTANT PRODUCTION IN FREEZE TOLERANT
VERTEBRATES**

by

CLARK PETER HOLDEN

M.Sc. Carleton University, 1991

**A thesis submitted to the Faculty of
Graduate Studies and Research in partial
fulfillment of the requirements
for the degree of**

Doctor of Philosophy

Department of Biology

Carleton University

Ottawa, Ontario

October, 1994

©copyright

1994, Clark Holden

National Library
of Canada

Acquisitions and
Bibliographic Services Branch

385 Wellington Street
Ottawa, Ontario
K1A 0N4

Bibliothèque nationale
du Canada

Direction des acquisitions et
des services bibliographiques

385, rue Wellington
Ottawa (Ontario)
K1A 0N4

Your file Votre référence

Our file Notre référence

THE AUTHOR HAS GRANTED AN
IRREVOCABLE NON-EXCLUSIVE
LICENCE ALLOWING THE NATIONAL
LIBRARY OF CANADA TO
REPRODUCE, LOAN, DISTRIBUTE OR
SELL COPIES OF HIS/HER THESIS BY
ANY MEANS AND IN ANY FORM OR
FORMAT, MAKING THIS THESIS
AVAILABLE TO INTERESTED
PERSONS.

L'AUTEUR A ACCORDE UNE LICENCE
IRREVOCABLE ET NON EXCLUSIVE
PERMETTANT A LA BIBLIOTHEQUE
NATIONALE DU CANADA DE
REPRODUIRE, PRETER, DISTRIBUER
OU VENDRE DES COPIES DE SA
THESE DE QUELQUE MANIERE ET
SOUS QUELQUE FORME QUE CE SOIT
POUR METTRE DES EXEMPLAIRES DE
CETTE THESE A LA DISPOSITION DES
PERSONNE INTERESSEES.

THE AUTHOR RETAINS OWNERSHIP
OF THE COPYRIGHT IN HIS/HER
THESIS. NEITHER THE THESIS NOR
SUBSTANTIAL EXTRACTS FROM IT
MAY BE PRINTED OR OTHERWISE
REPRODUCED WITHOUT HIS/HER
PERMISSION.

L'AUTEUR CONSERVE LA PROPRIETE
DU DROIT D'AUTEUR QUI PROTEGE
SA THESE. NI LA THESE NI DES
EXTRAITS SUBSTANTIELS DE CELLE-
CI NE DOIVENT ETRE IMPRIMES OU
AUTREMENT REPRODUITS SANS SON
AUTORISATION.

ISBN 0-315-98475-9

Name

Dissertation Abstracts International is arranged by broad, general subject categories. Please select the one subject which most nearly describes the content of your dissertation. Enter the corresponding four-digit code in the spaces provided.

Biology, General

SUBJECT TERM

0306

U·M·I

SUBJECT CODE

Subject Categories

THE HUMANITIES AND SOCIAL SCIENCES

COMMUNICATIONS AND THE ARTS

Architecture	0729
Art History	0377
Cinema	0900
Dance	0378
Fine Arts	0357
Information Science	0723
Journalism	0391
Library Science	0399
Mass Communications	0708
Music	0413
Speech Communication	0459
Theater	0465

EDUCATION

General	0515
Administration	0514
Adult and Continuing	0516
Agricultural	0517
Art	0273
Bilingual and Multicultural	0282
Business	0688
Community College	0275
Curriculum and Instruction	0727
Early Childhood	0518
Elementary	0524
Finance	0277
Guidance and Counseling	0519
Health	0680
Higher	0745
History of	0520
Home Economics	0278
Industrial	0521
Language and Literature	0279
Mathematics	0280
Music	0522
Philosophy of	0998
Physical	0523

Psychology	0525
Reading	0535
Religious	0527
Sciences	0714
Secondary	0533
Social Sciences	0534
Sociology of	0340
Special	0529
Teacher Training	0530
Technology	0710
Tests and Measurements	0288
Vocational	0747

LANGUAGE, LITERATURE AND LINGUISTICS

Language	
General	0679
Ancient	0289
Linguistics	0290
Modern	0291
Literature	
General	0401
Classical	0294
Comparative	0295
Medieval	0297
Modern	0298
African	0316
American	0591
Asian	0305
Canadian (English)	0352
Canadian (French)	0355
English	0593
Germanic	0311
Latin American	0312
Middle Eastern	0315
Romance	0313
Slavic and East European	0314

PHILOSOPHY, RELIGION AND THEOLOGY

Philosophy	0422
Religion	
General	0318
Biblical Studies	0321
Clergy	0319
History of	0320
Philosophy of	0322
Theology	0469

SOCIAL SCIENCES

American Studies	0323
Anthropology	
Archaeology	0324
Cultural	0326
Physical	0327
Business Administration	
General	0310
Accounting	0272
Banking	0770
Management	0454
Marketing	0338
Canadian Studies	0385
Economics	
General	0501
Agricultural	0503
Commerce-Business	0505
Finance	0508
History	0509
Labor	0510
Theory	0511
Folklore	0358
Geography	0366
Gerontology	0351
History	
General	0578

Ancient	0579
Medieval	0581
Modern	0582
Black	0328
African	0331
Asia, Australia and Oceania	0332
Canadian	0334
European	0335
Latin American	0336
Middle Eastern	0333
United States	0337
History of Science	0585
Law	0398
Political Science	
General	0615
International Law and Relations	0616
Public Administration	0617
Recreation	0814
Social Work	0452
Sociology	
General	0626
Criminology and Penology	0627
Demography	0938
Ethnic and Racial Studies	0631
Individual and Family Studies	0628
Industrial and Labor Relations	0629
Public and Social Welfare	0630
Social Structure and Development	0700
Theory and Methods	0344
Transportation	0709
Urban and Regional Planning	0999
Women's Studies	0453

THE SCIENCES AND ENGINEERING

BIOLOGICAL SCIENCES

Agriculture	
General	0473
Agronomy	0285
Animal Culture and Nutrition	0475
Animal Pathology	0476
Food Science and Technology	0359
Forestry and Wildlife	0478
Plant Culture	0479
Plant Pathology	0480
Plant Physiology	0817
Range Management	0777
Wood Technology	0746

Biology

General	0306
Anatomy	0287
Biostatistics	0308
Botany	0309
Cell	0379
Ecology	0329
Entomology	0353
Genetics	0369
Limnology	0793
Microbiology	0410
Molecular	0307
Neuroscience	0317
Oceanography	0416
Physiology	0433
Radiation	0821
Veterinary Science	0778
Zoology	0472

Biophysics

General	0786
Medical	0760

EARTH SCIENCES

Biogeochemistry	0425
Geochemistry	0996

Geodesy	0370
Geology	0372
Geophysics	0373
Hydrology	0388
Mineralogy	0411
Paleobotany	0345
Paleoecology	0426
Paleontology	0418
Paleozoology	0985
Polynology	0427
Physical Geography	0368
Physical Oceanography	0415

HEALTH AND ENVIRONMENTAL SCIENCES

Environmental Sciences	0768
Health Sciences	
General	0566
Audiology	0300
Chemotherapy	0992
Dentistry	0567
Education	0350
Hospital Management	0769
Human Development	0758
Immunology	0982
Medicine and Surgery	0564
Mental Health	0347
Nursing	0569
Nutrition	0570
Obstetrics and Gynecology	0380
Occupational Health and Therapy	0354
Ophthalmology	0381
Pathology	0571
Pharmacology	0419
Pharmacy	0572
Physical Therapy	0382
Public Health	0573
Radiology	0574
Recreation	0575

Speech Pathology	0460
Toxicology	0383
Home Economics	0386

PHYSICAL SCIENCES

Pure Sciences

Chemistry	
General	0485
Agricultural	0749
Analytical	0486
Biochemistry	0487
Inorganic	0488
Nuclear	0738
Organic	0490
Pharmaceutical	0491
Physical	0494
Polymer	0495
Radiation	0754
Mathematics	0405
Physics	
General	0605
Acoustics	0986
Astronomy and Astrophysics	0606
Atmospheric Science	0608
Atomic	0748
Electronics and Electricity	0607
Elementary Particles and High Energy	0798
Fluid and Plasma	0759
Molecular	0609
Nuclear	0610
Optics	0752
Radiation	0756
Solid State	0611
Statistics	0463

Applied Sciences

Applied Mechanics	0346
Computer Science	0984

Engineering	
General	0537
Aerospace	0538
Agricultural	0539
Automotive	0540
Biomedical	0541
Chemical	0542
Civil	0543
Electronics and Electrical	0544
Heat and Thermodynamics	0348
Hydraulic	0545
Industrial	0546
Marine	0547
Materials Science	0794
Mechanical	0548
Metallurgy	0743
Mining	0551
Nuclear	0552
Packaging	0549
Petroleum	0765
Sanitary and Municipal	0554
System Science	0790
Geotechnology	0428
Operations Research	0796
Plastics Technology	0795
Textile Technology	0994

PSYCHOLOGY

General	0621
Behavioral	0384
Clinical	0622
Developmental	0620
Experimental	0623
Industrial	0624
Personality	0625
Physiological	0989
Psychobiology	0349
Psychometrics	0632
Social	0451

AGREEMENT

This is an agreement between the author (a) and University Microfilms International, Ann Arbor, Michigan, providing for the publication of an abstract of not more than 350 words of the dissertation (d) in *Dissertation Abstracts International*, which is distributed on a subscription basis. If more than 350 words are contained in the abstract, publication is delayed. *Mathematical formulas, diagrams, and other illustrative materials are not recommended for the printed abstract.*

INSTRUCTIONS

Please read the Agreement form carefully. It authorizes the publication by University Microfilms International of a 350-word abstract in *Dissertation Abstracts International*. Indicate your approval of the Agreement by signing item 9 below.

DO NOT WRITE IN THIS SPACE

Abstract Length _____

Supervisor _____

Personal Data

1. Full Legal Name Holden
Clark (last name) Peter (first name)
(first name) (middle name)

2. Present mailing address R.R. #3
Carp, Ontario
Canada K0A 1L0

Future mailing address _____

Telephone No. 831-4040 (613)Effective Date January, 1995

Doctoral Degree Data

3. Abbreviation for Degree Authorized by University Ph.D.4. Degree date authorized by University 19955. Full name of university conferring degree
Carleton University

Title Data

6. Title of Dissertation exactly as it appears on Title Page and Abstract.
(Please attach a copy of your title page.)

7. Enter Subject Category selected for your dissertation.

(Choose from list on overleaf) Main Field Biology
Sub Field _____

Reprints

8. Please send me _____ reprints of my abstract.
(Minimum order 100 copies.)

First set of 100 \$20.00

Additional sets of 100 @ \$5.00 each set _____

Enclosed please find my check for _____

Please Note: Reprints are not mailed until the abstract appears in *Dissertation Abstracts International*, which is generally four months after we receive the material.

9. I agree to the provisions of the Agreement.

Signature [Signature]Date JANUARY 24, 1995**IMPORTANT!** Please select subject matter on overleaf.

The undersigned hereby recommend to
the Faculty of Graduate Studies and Research
acceptance of the thesis,

**"SIGNAL TRANSDUCTION AND THE FUNCTION OF SECOND MESSENGERS
AND PROTEIN KINASES IN THE CONTROL OF GLYCOGENOLYSIS AND
CRYOPROTECTANT PRODUCTION IN FREEZE TOLERANT VERTEBRATES"**

submitted by

CLARK P. HOLDEN, MSc., B.Sc. (Hons)

in partial fulfilment of the requirements
for the degree of Doctor of Philosophy

Chair, Department of Biology

Thesis Supervisor

External Examiner

Carleton University

January 1995

ABSTRACT

A unique group of lower vertebrates, termed freeze tolerant, have developed complex biochemical mechanisms that allow them to survive whole body freezing. Perhaps the most important of these mechanisms is the production of cryoprotectant, usually in the form of glucose, via an activation of glycogenolysis as a response to freezing. This thesis investigates: (1) whether protein kinase enzymes are responsible for the early activation of glycolysis that stimulates cryoprotectant production in freeze tolerant vertebrates and/or the subsequent inhibition of the pathway in the later stages of a freezing episode, (2) the comparative aspects of the responses of protein kinases elicited by the environmental stresses of dehydration and anoxia, to those of freezing, and (3) the role of the free catalytic subunit of cyclic AMP-dependent protein kinase (PKAc) and protein kinase C (PKC) in cryoprotectant production in *Rana sylvatica* liver.

Short freezing episodes caused a dramatic increase in the percentage of free catalytic PKA (PKAc). This response diminished with increased time frozen in tissues of freeze tolerant vertebrates. The measurement of second messengers over a freezing time course revealed significant increases in adenosine 3'-5' cyclic monophosphate (cAMP) and inositol 1,4,5-trisphosphate (IP₃) in liver and muscle of 4 freeze tolerant animals. An activation of PKA and PKC by these molecules, in response to freezing, is likely. Dehydration stress, not anoxia stress, elicited a remarkably similar protein kinase response to that seen during freezing. Levels of PKAc and its second messenger cAMP, as well as the second messenger for PKC, IP₃, rose in dehydrated liver tissue whereas anoxia did not activate PKAc and only increased liver IP₃. Purified *R. sylvatica* liver PKAc was not inhibited by high concentrations of salt or glucose and showed an increased affinity for substrates at low temperatures. Additional evidence showed that a freezing stress did not initiate a translocation of PKC to the plasma

membrane in *R. syivatica* liver and brain tissues. Freeze tolerant vertebrates possess an adaptive strategy to combat freezing which links second messengers and protein kinases to the post-translational modification of glycolytic enzymes conducting the biosynthesis of cryoprotectant.

ACKNOWLEDGMENTS

First and foremost, I wish to thank Dr. Ken Storey for all his guidance, helpful suggestions, and encouragement throughout the course of my studies here at Carleton University. I would like to thank Jan Storey whose help, calm demeanor, and critical writing skills are a great benefit both to the lab and to me. Thanks goes out to my committee members Dr. Stuart Jacobson and Dr. Tom Moon for their valued suggestions, advice, and guidance both in and out of the laboratory.

I would also like to thank my fellow lab-mates (you know who you are) that lent an ear and provided a smile or a joke in a time of need. Thank you all for providing me with a topnotch scientific working environment. Finally, thanks goes out to my family and to my girlfriend Nadia for providing me with love, understanding, support, and late night meals; all of which supplied me with the mental and metabolic fuel that made my studies just that much easier. Thanks all!

TABLE OF CONTENTS

	Page
Acceptance Sheet	ii
Abstract	iii
Acknowledgments	v
Table of Contents	vi
List of Abbreviations	viii
List of Tables	x
List of Figures	xiii
Chapter One: General Introduction	1
Chapter Two: General Materials and Methods	16
Chapter Three: Signal transduction, second messenger, and protein kinase responses during freezing in four freeze tolerant vertebrates.	32
Introduction	33
Materials and Methods	35
Results	36
Discussion	54
Chapter Four: Signal transduction, second messenger, and protein kinase responses during dehydration in two anurans	63
Introduction	64
Materials and Methods	66
Results	67
Discussion	74

	Page
Chapter Five: Signal transduction, second messenger, and protein kinase responses during anoxia stress in two anurans	83
Introduction	84
Materials and Methods	86
Results	87
Discussion	93
Chapter Six: Purification of the free catalytic subunit of cyclic AMP-dependent protein kinase from <i>Rana sylvatica</i> liver	100
Introduction	101
Materials and Methods	102
Results	105
Discussion	123
Chapter Seven: Activation and translocation of protein kinase C during freezing stress	130
Introduction	131
Materials and Methods	133
Results	139
Discussion	145
Chapter Eight: General Discussion	151
References	159
Publication List	166

LIST OF ABBREVIATIONS

32 -P ATP - gamma labelled ATP

ANOVA - analysis of variance

antiPKC- α - the primary antibody of α PKC from rabbit brain

antiPKC- β - the primary antibody of β PKC from rabbit brain

ATP - adenosine triphosphate

BCIP - 5-bromo-4-chloro-3-indolyl-phosphate

cAMP - adenosine 3'-5' cyclic monophosphate

cGMP - guanosine 3'-5' cyclic monophosphate

DAG -diacylglycerol

DMF - dimethylformamide

EDTA - ethylenediamine tetraacetate

EGTA - ethyleneglycol bis(β -aminoethyl ether) tetraacetate

G6PDH - Glucose-6-phosphate Dehydrogenase

GP - glycogen phosphorylase

GS - glycogen synthase

h - Hill coefficient

H89 - protein inhibitor of mammalian PKA

HEPES - (N-[2-hydroxyethyl] piperazine-N'-[2-ethanesulphonic acid])

I₅₀ - concentration of inhibitor that reduces V_{max} activity by one half

IgG - the secondary antibody, goat antirabbit IgG (H+L)

IP₃ - inositol 1,4,5-trisphosphate

K_a - concentration of activator that produces one half of maximal enzyme activation

K_m - Michaelis-Menten constant

K_p - partition coefficient

KP_i - a mixture of KH_2PO_4 and K_2HPO_4 , Mono- and di-potassium phosphate

M.W. - molecular weight

Mg(OAc)₂ - magnesium acetate

Mg-ATP - Mg^{2+} and ATP complex

NAD⁺, NADH - oxidized and reduced forms of nicotinamide adenine dinucleotide

NaP_i - a mixture of $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$ and Na_2HPO_4 , Mono- and di-sodium phosphate

M.W. - molecular weight

NBT - nitro blue tetrazolium

PBS - phosphate-buffered saline

PCA - perchloric acid, HClO_4 (specific gravity = 1.54)

PFK - phosphofructo-1-kinase

PFKII - 6-phosphofructo-2-kinase

PK - pyruvate kinase

PKA - holoenzyme of cyclic AMP-dependent protein kinase

PKAc - free catalytic subunit of cyclic AMP-dependent protein kinase

PKA_i - protein inhibitor of mammalian PKA

PKC - calcium and phospholipid dependent protein kinase

PKG - cyclic GMP-dependent protein kinase

PKG - glycogen phosphorylase kinase

PMSF - phenylmethylsulfonyl fluoride

PS - phosphatidylserine

RIA - radioimmune assay

SDS - sodium dodecyl sulphate

SDS-PAGE - sodium dodecyl sulphate polyacrylamide gel electrophoresis

SNK - Student-Newman-Keuls test

TRIS - (Tris[hydroxymethyl]aminomethane)

V_{max} - maximal enzyme velocity

LIST OF TABLES

Table	Page
1. Percentage of the free catalytic subunit of PKA, total PKA, and total activity from control (5°C), frozen (24 h, -2.5°C), and thawed (24 h, 5°C) <i>Rana sylvatica</i> tissues.	37
2. Percentage of the free catalytic subunit of PKA, total PKA, and total activity from control (5°C), frozen (4 h, -2.5°C), and thawed (24 h, 5°C) <i>Trachemys scripta elegans</i> tissues.	38
3. Percentage of the free catalytic subunit of PKA from control (5°C), frozen (4 h, -2.5°C), and thawed (24 h, 5°C) <i>Trachemys scripta elegans</i> , and <i>Chrysemys picta marginata</i> and <i>Chrysemys picta bellii</i> control (5°C), frozen (24 h, -2.5°C), and thawed (24 h, 5°C) liver and muscle tissue.	40
4. Second messenger concentrations (cAMP, IP ₃ , and cGMP) in control (5°C), frozen (24 h, -2.5°C), and thawed (24 h, 5°C) <i>Rana sylvatica</i> tissues.	42
5. Second messenger concentrations (cAMP, IP ₃ , and cGMP) in control (5°C), frozen (4 h, -2.5°C), and thawed (24 h, 5°C) <i>Trachemys scripta elegans</i> tissues.	43
6. Second messenger concentrations (cAMP, IP ₃ , and cGMP) in control (5°C), frozen (24 h, -2.5°C), and thawed (24 h, 5°C) midland painted turtle <i>Chrysemys picta marginata</i> tissues.	44
7. Second messenger concentrations (cAMP, IP ₃ , and cGMP) in control (5°C), frozen (24 h, -2.5°C), and thawed (24 h, 5°C) western painted turtle <i>Chrysemys picta bellii</i> tissues.	45
8. Effect of freezing and thawing on glucose concentrations in <i>Rana sylvatica</i> and <i>Trachemys scripta elegans</i> tissues (spring, 1992).	48
9. Changes in the percentage of the free catalytic subunit of cAMP-dependent protein kinase over a time course of freezing at -2.5°C, followed by thawing for 24 h at 5°C in <i>Rana sylvatica</i> tissues.	50
10. Effect of freezing exposure on cAMP and IP ₃ concentrations over a freezing time course at -2.5°C, followed by thawing for 24 h at 5°C in <i>Rana sylvatica</i> tissues.	51
11. Changes in tissue glucose concentrations over a time course of freezing exposure at -2.5°C, followed by thawing at 5°C in three <i>Rana sylvatica</i> tissues.	53
12. Total protein concentrations in tissue samples from <i>Rana sylvatica</i> sampled over a course of whole body dehydration and rehydration.	69

Table	Page
13. Effect of changes in total body water content on the percentage of protein kinase A present as the free catalytic subunit in <i>Rana sylvatica</i> liver and muscle.	70
14. Effect of changes in total body water content on the percentage of protein kinase A present as the free catalytic subunit in <i>Rana pipiens</i> liver and muscle.	71
15. Effect of changes in total body water content on the concentrations of cyclic AMP (cAMP) and inositol trisphosphate (IP ₃) in <i>Rana sylvatica</i> tissues.	73
16. Effect of anoxia exposure at 5°C on the percentage of protein kinase A present as the free catalytic subunit in <i>Rana sylvatica</i> tissues.	88
17. Effect of anoxia exposure at 5°C on the percentage of protein kinase A present as the free catalytic subunit in <i>Rana pipiens</i> tissues.	89
18. Effect of anoxia exposure at 5°C on the concentrations of IP ₃ in <i>Rana sylvatica</i> tissues.	91
19. Effect of anoxia exposure at 5°C on the concentrations of IP ₃ in <i>Rana pipiens</i> tissues.	92
20. Purification of the free catalytic subunit of protein kinase A from <i>Rana sylvatica</i> liver using various chromatographic techniques.	106
21. Stability of the purified free catalytic subunit of protein kinase A from <i>Rana sylvatica</i> liver over time stored in the presence versus the absence of 0.25 % (v/v) Triton X-100 detergent.	111
22. Stability of the purified free catalytic subunit of protein kinase A from <i>Rana sylvatica</i> liver when stored at different subzero temperatures in the presence of 0.25 % (v/v) Triton X-100 detergent.	112
23. Substrate specificity of the purified free catalytic subunit of protein kinase A from <i>Rana sylvatica</i> liver; V _{max} activities with several different protein kinase A substrates at 22°C and 5°C.	114
24. Substrate affinity coefficients for the purified free catalytic subunit of protein kinase A from <i>Rana sylvatica</i> liver.	115
25. Inhibitor coefficients, I ₅₀ values, for the purified free catalytic subunit of protein kinase A from <i>Rana sylvatica</i> liver.	117

Table**Page**

26. Effect of freezing exposure (24 h, at -2.5°C) and subsequent thawing (24 h, at 5°C) on the percentage of bound PKC and the levels of inositol 1,4,5-trisphosphate (IP_3) in *Rana sylvatica* liver and brain.

140

LIST OF FIGURES

Figure		Page
1.	Effect of increasing amounts of free cyclic AMP on the activity of crude <i>Rana sylvatica</i> liver cAMP-dependent protein kinase.	28
2.	Three elution profiles showing the purification of the free catalytic subunit of cyclic AMP-dependent protein kinase from <i>Rana sylvatica</i> liver: A) hydroxylapatite column, B) protamine agarose column, C) Bio-Gel P-200 gel filtration column.	107
3.	Sodium dodecyl sulphate polyacrylamide gel indicating the subunit molecular weight determination for the free catalytic subunit of cAMP-dependent protein kinase purified to homogeneity from <i>Rana sylvatica</i> liver.	109
4.	A plot of the molecular weight versus the partition coefficient (K_p) of standard proteins from a Bio-Gel P-200 column in order to calculate the native molecular weight of the free catalytic subunit of cAMP-dependent protein kinase purified from <i>Rana sylvatica</i> liver.	110
5.	Effect of increasing amounts of free magnesium on the activity of the free catalytic subunit of cAMP-dependent protein kinase purified from <i>Rana sylvatica</i> liver.	116
6.	An Arrhenius plot for the free catalytic subunit of cAMP-dependent protein kinase purified from <i>Rana sylvatica</i> liver.	119
7.	The effect of pH on the activity of the free catalytic subunit of cAMP-dependent protein kinase at 22°C (A) and 5°C (B).	120
8.	The tissue specific isozymes of the free catalytic subunit of cyclic-AMP dependent protein kinase from <i>Rana sylvatica</i> liver, muscle, and brain tissue as expressed using hydroxylapatite column chromatography.	122
9.	An elution profile showing the single isozyme of <i>Rana sylvatica</i> liver PKC determined using hydroxylapatite column chromatography.	141
10.	Two nitrocellulose immunoblots showing the single PKC isozyme found in <i>Rana sylvatica</i> liver; detected with antiPKC antibodies, (a) antiPKC- α (negative stain), and (b) antiPKC- β (positive stain).	142
11.	An elution profile showing the isozymes of <i>Rana sylvatica</i> brain PKC determined using hydroxylapatite column chromatography.	144

CHAPTER #1

GENERAL INTRODUCTION

Hypotheses: The following hypotheses were the ideas tested and examined in this thesis. Signal transduction, second messengers, and protein kinases, specifically cyclic AMP-dependent protein kinase (PKA) and Ca^{2+} -phospholipid dependent protein kinase (PKC), are responsible for initiating the metabolic adjustments seen in vertebrates that are able to survive freezing, dehydration, and anoxia. In addition, protein kinases initiate and mediate the biosynthesis of the cryoprotectant glucose throughout a freezing stress in *R. sylvatica*, the freeze tolerant wood frog. The phenomenon of vertebrate freeze tolerance developed from similar metabolic capacities, particularly the second messenger-protein kinase signal transduction responses, seen in amphibians that readily survive dehydration and anoxia stresses. Through the examination of the metabolic responses to dehydration and anoxia, the probable origin of vertebrate freeze tolerance adaptations can be implied.

Vertebrate Freeze Tolerance

Terrestrial anurans and reptiles that live at high latitudes have developed elaborate physiological and biochemical mechanisms for surviving cold winter temperatures. These vertebrates must be able to survive subzero temperatures which are far below the equilibrium melting point of body fluids. There are three main behavioral strategies that animals can choose from to deal with the subzero temperatures of winter. The first strategy is to avoid the possibility of body fluids freezing by either: (1) burrowing into the forest floor like many toad species, (2) spending the winter at the bottom of fresh water ponds like the leopard frog *Rana pipiens* and many turtle species, or (3) migrating to warmer climates like many bird species (Storey and Storey, 1988). The second strategy, employed by many terrestrial arthropods, is that of freeze-avoidance (Storey and Storey, 1992a). These animals endure subzero temperatures without freezing. By using adaptations including the production of thermal hysteresis proteins and polyhydric alcohols (such as glycerol and sorbitol) these animals

are able to stabilize the liquid state of their body fluids (Storey and Storey, 1992a). The third strategy, the main focus of this thesis, is freeze tolerance or the ability to endure the freezing of extracellular body fluids. Adaptive strategies involved in freeze tolerance include those that: (1) regulate the place and rate of ice formation in the extracellular spaces of all tissues, (2) limit the extent of cell volume decrease during extracellular ice formation, and (3) stabilize macromolecules at low temperature in cells that, as a consequence of freezing, contain low free water content and high solute concentrations. The molecular mechanisms involved in these strategies have been widely studied in insects (Zachariassen, 1985; Storey and Storey, 1992a) and, more recently, in freeze tolerant frogs and turtles (Storey and Storey, 1988, 1992b). The wood frog, *R. sylvatica*, has been the primary model for studies of freeze tolerance in vertebrates. Known adaptations identified in this species include: (1) the occurrence of ice nucleating proteins in plasma that stimulate and control the rate of ice formation, and (2) the production of high concentrations of glucose as a colligative cryoprotectant to help limit cell volume reduction and stabilize macromolecules in the frozen state.

The definition of vertebrate freeze tolerance includes the survival of whole body freezing, with 50-60 % of total body water sequestered as ice, for lengths of time longer than 4 h at -2.5°C . To date, there are only 11 known species of freeze tolerant vertebrates, all from terrestrial habitats; 6 from the class Amphibia and 5 from the class Reptilia. Species that meet these criteria are: the wood frog, *R. sylvatica*; the grey tree frog, *Hyla versicolor*; Cope's tree frog, *Hyla chrysoscelis*; the chorus frog, *Pseudacris triseriata*; the spring peeper, *Pseudacris crucifer*; the Siberian salamander, *Hynobius keyserlingi*; the box turtle, *Terrapene carolina carolina*; the western box turtle, *Terrapene ornata*; the western and midland subspecies of the painted turtle, *Chrysemys picta bellii*, and *Chrysemys picta marginata*; the red-eared slider turtle, *Trachemys scripta elegans*; and the common garter snake, *Thamnophis sirtalis parietalis* (Storey, 1984; Berman et al, 1984; Storey and Storey, 1985a; Storey, 1987a; Costanzo et al.,

1990; Costanzo and Claussen, 1990; Churchill and Storey, 1991; Churchill and Storey, 1992a,b).

Ice-Formation and Cryoprotectant Production During Freezing

The formation of intracellular ice is lethal to all animals due to the destruction of membrane integrity resulting from the penetration of ice crystals into cellular membranes (McGann et al., 1988). Limiting and controlling the formation of extracellular ice and preventing the formation of intracellular ice is essential to the overall survival of a freeze tolerant organism. Typical cytoplasmic osmolalities in terrestrial and freshwater organisms range from 200 to 400 mM, contributed by salts, total metabolite pools, and proteins. The net colligative effect of these compounds are not enough to reduce the freezing and/or supercooling point of the intracellular matrix. Adaptive pressures and seasonal variations in temperature in temperate zones have resulted in the ability of some terrestrial organisms to synthesize additional polyhydrated carbon compounds within the cytoplasm. These 3 to 6 carbon sugars or sugar alcohols are commonly called cryoprotectants. Freeze tolerant insects such as *Eurosta solidaginis* synthesize glycerol and sorbitol whereas freeze tolerant vertebrates accumulate glucose produced from liver glycogen stores (Zachariassen, 1985; Storey and Storey, 1988).

The two types of protective measures which all cryoprotectants provide for the cells of freeze tolerant animals are: (1) membrane stabilization, and (2) colligative regulation of cell volume. Membrane stabilization is achieved through the interactions between the cryoprotectant and the polar hydrophilic head groups of membrane phospholipids (Carpenter and Crowe, 1988). Cryoprotectant colligative effects are due to the molar levels of these carbon compounds, 200 to 300 mM glucose in *R. sylvatica*. By virtually doubling cellular osmolality through cryoprotectant production, the normal osmolality of *R. sylvatica* cells being 300 mosmol/l (Storey and Storey, 1986), the

presence of cryoprotectants reduces the amount of ice that can form at any given subzero temperature and also reduces the consequent decrease in cell volume that occurs with freezing.

Another functional quality of polyhydroxylated compounds is their structural, and concurrently functional similarity to water. Due to the presence of multihydroxyl groups, intercellular water undergoes hydrogen bonding with these cryoprotectants. This hydrogen bonding reduces the degree of intercellular dehydration that accompanies a freezing episode (Carpenter and Crowe, 1988). Polyols are also capable of stabilizing protein structure, by acting as water does under normal conditions, through the formation of a hydration shell around the hydrophilic region of the protein. This in turn prevents the denaturation of proteins by stabilizing their tertiary structure.

All naturally synthesized cryoprotectants are non-toxic to the cells of freeze tolerant vertebrates; even at high concentrations. Chemical inertness is an essential property of all cryoprotectants. These molecules do not participate in any irreversible chemical reactions with cellular compounds or structures (Zachariassen, 1985). Most cryoprotectants are not directly exposed to the regulatory effects, such as reversible phosphorylation reactions, that are observed in many biochemical pathways. It follows then that most cryoprotectants, once synthesized, are not readily catabolized; a necessary characteristic that allows for the conservation of the cryoprotectant pool throughout the winter months when the animal is frozen.

In addition to carbohydrate cryoprotectant synthesis, many freeze tolerant animals, such as the midland painted turtle *C.p. marginata*, rely on the build up of amino acid pools to stabilize membrane structure while in the frozen state (Churchill and Storey, 1991). Freeze tolerant turtles increase total amino acid pools as well as the concentrations of glutamate and taurine to survive freezing (Storey et al., 1988).

Ice Nucleation and Thermal Hysteresis in Freeze Tolerant Animals

In addition to the biosynthesis of cryoprotectants, many freeze tolerant animals utilize ice nucleating proteins (INP) and thermal hysteresis proteins (THP) to help control the freezing process (Duman, 1982). These proteins nucleate extracellular ice formation (INP) or bind to the growing outer edges of the ice crystals (THP), halting their expansion before any damage can occur in the cell. Only two freeze tolerant vertebrates are known to accumulate these proteins *R. sylvatica* and *C.p. marginata* (Wolanczyk et al., 1990; Storey et al., 1991). However, studies are ongoing in other freeze tolerant vertebrates since high supercooling points are found in some and others only synthesize small quantities of protective sugars, but all can survive freezing episodes for prolonged periods of time.

Freeze Tolerance in *R. sylvatica*

The development of a freeze tolerance strategy is the major adaptation that enables many terrestrial vertebrates to survive the cold winter environment of northern latitudes. One of the most successful freeze tolerant vertebrate species is the wood frog *R. sylvatica*. This species can supercool down to -2 to -3°C and can survive for weeks in the frozen state at temperatures as low as -6 to -8°C with over 50% of its total body water as ice (Storey, 1984). Under laboratory conditions the time elapsed before ice nucleation on the skin of *R. sylvatica* is approximately 30 min at -4°C. This anuran always freezes from the outside in; the last organs to freeze being the liver, heart, and brain. Ice formation continues in this manner under these conditions for 24-48 h until such a time as equilibrium ice (65 % of body water frozen as extracellular ice) is formed.

The synthesis of glucose from liver glycogen reserves is a well known biochemical process in many vertebrate organisms (Hers, 1976). *R. sylvatica* has refined this process so that it is able to mobilize its glycogen stores almost immediately

after an external inoculative freezing stress. Glucose is the cryoprotectant of choice for *R. sylvatica* and is found in high concentrations in every major tissue and organ in the frozen animal. There is an active production of glucose (up to 200-300 $\mu\text{mol/g}$ wet weight) from liver glycogen within the early minutes to hours of a freezing stress (Storey and Storey, 1986). The glucose that is produced from liver glycogen stores is transported to other organs by the circulatory system (Storey, 1987b). *R. sylvatica* does not synthesize cryoprotectant prior to freezing, a common metabolic strategy utilized by cold hardy insects (Storey and Storey, 1992a). Instead, glucose biosynthesis is initiated as a response to ice crystallization on the body surface and begins less than 5 min after nucleation (freezing) (Storey, 1987b). The immediate breakdown of glycogen to glucose via glycogenolysis allows for a fast and efficient synthesis and distribution of cryoprotectant to protect the animal in its frozen state.

The rapid output of glucose from *R. sylvatica* liver is facilitated by two freezing-stimulated events (1) an increase in the conversion of glycogen phosphorylase to its active α form, and (2) an increase in the total activity of glycogen phosphorylase measurable in liver (Mommensen and Storey, 1992). In addition, large glucose concentrations can be synthesized due to the presence of very high concentrations of liver glycogen accumulated during summer/autumn feeding (equivalent to as much as 1000 glucose units/g wet weight). Cryoprotectant synthesis may be controlled by a β -adrenergic hormone signal (Storey and Storey, 1992b), and the protein phosphorylation-mediated change in liver glycogen phosphorylase properties during freezing exposure (Mommensen and Storey, 1992) suggests a probable role for cyclic AMP-dependent protein kinase (PKA) and/or protein kinase C (PKC) in the phosphorylation of this glycolytic enzyme. The enzyme that activates glycogen phosphorylase is phosphorylase kinase, an enzyme which must first be converted to its active form by cAMP-dependent protein kinase (PKA). PKA activation then, must be one of the critical rate limiting steps leading to the activation of glycogen

phosphorylase, and the immediate production of glucose that is seen in *R. sylvatica* during a freezing stress.

Although many biochemical events during cryoprotectant biosynthesis in *R. sylvatica* have been well documented (Storey and Storey, 1992b) the role of PKA and PKC in this process has not. The coordination of this biological process with respect to the levels of second messengers, their relationship to the activity of PKA and PKC, and whether or not PKA and/or PKC controls glucose production in this amphibian are investigated in this study. It has been determined that PKA and PKC may play very critical roles in the production of glucose from liver glycogen stores in this freeze tolerant frog.

Additionally, this study relates information on a new and novel purification of the free catalytic subunit of cAMP-dependent protein kinase from *R. sylvatica* liver. Kinetic studies will provide details on how PKAc acts to ensure production of the cryoprotectant glucose via liver glycogenolysis and its concurrent distribution throughout the tissues and organs of this freeze tolerant anuran. PKC and IP₃ distribution in liver and brain tissue of *R. sylvatica* is investigated in this study in order to determine: (1) whether this kinase is activated by a freezing stress, and (2) whether PKC is involved in the control of glucose biosynthesis and cell volume regulation in this freeze tolerant vertebrate.

Metabolic Control of Glycolytic Enzymes by Protein Kinase Enzymes in Natural Freeze Tolerant Animals

Maintenance of glycolytic control during freezing is essential in effectively coordinating cryoprotectant synthesis. This may be achieved through the highly regulated actions of protein kinase enzymes that are responsible for the phosphorylation and subsequent activation/inhibition of many rate limiting enzymes. Through the covalent modification of specific enzymes the flux of carbon through different

metabolic pathways can be tightly regulated. Regulation of the glycolytic pathway during cryoprotectant synthesis is carried out by several key enzyme control points; glycogen phosphorylase (GP), phosphofructokinase (PFK), and pyruvate kinase (PK). In order to successfully achieve cryoprotectant synthesis, efficient activation/inhibition of the controlling mechanisms (kinetics) of these enzymes must be in place. During the freezing process in *R. sylvatica* an activation of glycogen phosphorylase (GP) increases the flux of carbon down the glycolytic pathway. Concurrent to this, PFK is inhibited halting carbon flux into the lower half of glycolysis (Storey and Storey, 1992b). This carbon is then diverted into the enzyme reactions that are responsible for glucose production (Storey and Storey, 1986). Inhibitory control on PK may further strengthen this block of carbon flow down the glycolytic pathway. Protein kinase cascade systems are known to operate on all three of these regulatory enzymes (Stryer, 1981). The resulting phosphorylation/dephosphorylation of these target enzymes undoubtedly ensures that protein kinase enzymes aid in the control of cryoprotectant production in many freeze tolerant animals. Glucose output from mammalian liver is hormonally regulated by glucagon and catecholamine stimulation (Hers, 1976). Hormones activate protein kinase enzymes that, in turn, phosphorylate and activate GP kinase and then GP itself, while simultaneously inactivating PFK, PK, and 6-phosphofructo-2-kinase (PFKII) by phosphorylation. This system has been studied extensively in mammalian liver and comparable regulatory controls may also direct cryoprotectant production in freeze tolerant vertebrates.

Metabolic Responses to Dehydration in Anurans

Stresses put on cells due to dehydration have been a powerful selective agent in the colonization of terrestrial habitats by many amphibians (Thorson, 1955).

Dehydration puts a large osmotic stress on the cells of many anuran species (Hillman, 1980). Virtually the same osmotic stress on cells is observed in *R. sylvatica* when the

wood frog is in a dehydrated or frozen state (Churchill and Storey, 1993, 1994). Dehydration and freezing, although seemingly unrelated, have the same consequences on the regulation of cell volume in *R. sylvatica*. Both stresses result in a reduction of cell volume, in one case caused by a net loss of water from the body (dehydration) and in the other by sequestering up to 65 % of total body water as extracellular and extra-organ ice (freezing) (Lee et al. 1992; Churchill and Storey 1993; Storey and Storey, 1993). I propose, therefore, that the initiation of cryoprotectant production through the activation of glycogenolysis by protein kinase enzymes, specifically PKA and PKC, in response to freezing in *R. sylvatica* evolved out of a pre-existing hyperglycemic response by amphibians to dehydration. Indeed, new studies of the effects of dehydration on the semi-aquatic leopard frog *R. pipiens* support this proposal; for this freeze intolerant frog developed a rapid hyperglycemia, based on liver glycogenolysis, in response to dehydration (Churchill and Storey, 1994). However, the magnitude of the response is much lower in *R. pipiens* than in *R. sylvatica*; maximum glucose levels in liver of autumn-collected frogs, dehydrated to 50 % of total body water, were only 20 $\mu\text{mol/g}$ wet weight in the leopard frog versus 127 $\mu\text{mol/gww}$ in the wood frog (Churchill and Storey, 1993).

Several concepts were investigated in this study that addressed the influences of dehydration on the activities of protein kinases and the levels of second messengers in tissues of a freeze tolerant frog, *R. sylvatica*, versus a non-freeze tolerant frog, *R. pipiens*. In particular, I analyzed the regulatory control exerted by protein kinase enzymes and their second messengers over liver glycogenolysis, the pathway that is responsible for dehydration-induced glucose export. To what extent do the metabolic responses observed for protein kinase enzymes and their respective second messengers during dehydration mimic those seen during a freezing episode in *R. sylvatica*? Do these same stress vectors appear in *R. pipiens*, a species of frog that is known to be desiccation-tolerant but not freeze tolerant?

Metabolic Responses to Anoxia in *R. pipiens* and to Whole Body Ice Formation in *R. sylvatica*

All cardiac activity ceases in freeze tolerant vertebrates once they are frozen (Storey and Storey, 1988). Blood flow to tissues and organs is shut down, creating ischemic conditions and starving tissues for nutrients and oxygen. A key to the survival of these animals in a frozen state must be their ability to survive anoxic conditions (Storey and Storey, 1988). Tissues of frozen *R. sylvatica*, including brain, are able to generate ATP anaerobically and can withstand the metabolic consequences associated with a large increase in the anaerobic endproduct lactate (Storey, 1986). The freeze intolerant leopard frog *R. pipiens* also shows a well developed anoxia tolerance that includes a depletion of liver glycogen reserves and an accumulation of lactic acid in many tissues (Christiansen and Penny, 1973). Therefore, the metabolic responses to anoxia in anoxia tolerant frogs, such as *R. pipiens*, are remarkably similar to those occurring during prolonged freezing-induced ischemia in *R. sylvatica*.

What type of protein kinase response is responsible for the control of this anaerobic energy metabolism and subsequent ATP production that is observed during these two very different environmental stresses? To what extent do the metabolic responses observed for protein kinase enzymes and their respective second messengers in these two anurans under anoxic stress mimic those seen during a freezing episode in *R. sylvatica*? Furthermore, do the metabolic events controlled by protein kinase enzymes during dehydration reflect those that occur during an anoxic stress in the wood frog? Finally, do these same stress related events appear in *R. pipiens*, a species of frog that possesses tolerance to anoxia but not to freezing?

Second messengers and Protein Kinases

Cellular second messengers play a critical role in all signal transduction pathways present within a cell. The main second messengers present in cells are: (1) the cyclic nucleotides, adenosine 3',5'-cyclic monophosphate (cAMP) and guanosine 3',5'-cyclic monophosphate (cGMP), (2) inositol 1,4,5-trisphosphate (IP₃), and (3) calcium ions (Ca²⁺). Hormone triggers, such as adrenaline, bind to receptors on the plasma membrane of cells causing the release or an increased production of second messenger. In almost all cases second messenger levels rise just prior to the activation of many protein kinase enzymes (Hunter, 1991). Second messengers mediate the activation of their respective protein kinases by binding to an active site on a kinase enzyme; a biochemical event that leads to the release of a catalytic or activated form of the kinase. Adenosine 3',5'-cyclic monophosphate (cAMP) activates PKA, inositol 1,4,5, trisphosphate (IP₃) activates PKC, and guanosine 3',5'-cyclic monophosphate (cGMP) activates PKG.

The release of hormones in response to a metabolic stress causes an increase in the amount of cellular phosphorylation of regulatory enzymes found in the metabolic pathways of many animals. Increases in the levels of second messenger usually reflect increases in the levels of active protein kinase enzymes within a tissue (Hunter, 1991). Therefore, experiments were designed to investigate whether the levels of second messengers such as cAMP, IP₃, and cGMP changed from control values in experimental animals that were exposed to either whole body freezing, dehydration, or anoxia.

Protein kinase enzymes are defined as catalysts that transfer a phosphate group from a phosphate donor (usually ATP) onto an acceptor amino acid, such as serine, threonine, or tyrosine on a substrate protein (Hunter, 1991). The coordination and mediation of many cellular processes are controlled by different types of protein kinases. The most widely studied of these enzymes are cyclic AMP-dependent protein

kinase (PKA), calcium and phospholipid dependent protein kinase (PKC), and cyclic GMP-dependent protein kinase (PKG). These enzymes are responsible for phosphorylating regulatory enzymes that are important in numerous aspects of cellular metabolism including: (1) cell growth and differentiation (Hashimoto and Yamamura, 1991), (2) control of ion flux through cellular membrane channel proteins (Bourinet et al., 1992), and (3) regulation of metabolic pathways (Storey, 1987b; Brooks and Storey, 1991). This indicates that protein kinase enzymes may have critical functions in controlling the regulation of cellular metabolism through the reversible phosphorylation of enzymes during periods of environmental stress.

Protein kinases have been purified from many animal species and characterized both kinetically and structurally. All protein kinases have a regulatory domain and a catalytic domain. However, there are a variety of structural differences between each protein kinase. For instance, 3',5'-cyclic AMP-dependent protein kinase (PKA) is a tetrameric enzyme with two regulatory and two catalytic subunits (Shenolikar, 1988), and cyclic GMP-dependent protein kinase (PKG) is a holoenzyme with the regulatory and catalytic sites on the same domain (Tremblay et al., 1988).

Cyclic-AMP dependent protein kinase (PKA) is one of the most widely distributed kinase enzymes in the cells of mammalian tissues. Increased levels of intracellular cAMP leads to the activation of this tetrameric enzyme. Cyclic-AMP binds to the two regulatory subunits of PKA causing the release of the two catalytic subunits. The catalytic subunits are then free to phosphorylate a number of regulatory enzymes that are important in the control of many metabolic pathways (Shenolikar, 1988).

The binding of a hormone signal to the cell membrane activates the PKC signal transduction pathway by initiating the hydrolysis of polyphosphoinositides. This in turn, results in the formation of inositol 1,4,5-trisphosphate, IP₃, and diacylglycerol, DAG (Berridge, 1987). These two molecules along with a release of calcium ions cause

the translocation of PKC to the plasma membrane, activating the kinase in the process (Rasmussen, 1989).

Signal transduction pathways are highly regulated within cells, particularly through the actions of protein kinases. The level of phosphorylation at any instant reflects the state of activation or inhibition of metabolic pathways. Many metabolic processes are regulated by the actions of protein kinases including glycogenolysis, glycolysis, and gluconeogenesis. Protein kinases, especially PKA and PKC, may play critical synergistic roles in the regulation of glucose synthesis via glycogen phosphorylase phosphorylation in the freeze tolerant wood frog *R. sylvatica* during freezing and dehydration, and may be responsible for the control of anaerobic metabolism in anoxia stressed anurans.

Summary of Aims and Objectives

The first aim of this thesis is to determine if signal transduction cascades particularly at the second messenger and protein kinase level are responsible for initiating and controlling the metabolic adjustments seen in vertebrates that are able to survive one or more of the following stresses; freezing, dehydration, and anoxia.

The second aim is to elucidate whether these same protein kinases initiate and mediate the biosynthesis of the cryoprotectant glucose throughout a freezing stress in the freeze tolerant wood frog *R. sylvatica*.

The third aim of this study is to determine whether the development of vertebrate freeze tolerance, specifically that observed in *R. sylvatica*, stemmed from this anuran's natural ability to survive dehydration, anoxia, or a combination of both these environmental stresses. This will be accomplished by relating the actions and functions of protein kinases, with particular focus on purified PKAc and semi-purified PKC, and their respective second messengers in response to vertebrate freezing, dehydration, and anoxia stresses. Data collected on the responses of signal transduction

pathways to these three stresses will yield a clearer overall picture of the vertebrate freeze tolerance phenomenon.

CHAPTER #2

GENERAL MATERIALS AND METHODS

MATERIALS AND METHODS

Chemicals

Chemicals were purchased from Boehringer-Mannheim Corp., Montreal, PQ or Sigma Chemical Company, St. Louis, MO. Bio-Gel P-200 was from Bio-Rad Laboratories, Richmond, CA. Radiolabelled gamma ^{32}P -ATP was purchased from New England Nuclear (Dupont) at a concentration of approximately 3000 Ci/mmol (0.5 mCi/ml) and stored at -20°C .

Animal Collection, Acclimatization, and Storage

Wood frogs, *R. sylvatica*, and leopard frogs, *R. pipiens*, were collected from ponds in the Ottawa area in mid-April of the years 1992-1994 by a local supplier. Frogs were held in temperature controlled incubators at $5 \pm 1^{\circ}\text{C}$ for at least three days prior to experimentation. Hatchling red-eared slider turtles, *Trachemys scripta elegans*, were purchased from the commercial breeder, Reptiles Unlimited. Eggs of midland painted turtles, *Chrysemys picta marginata*, were obtained from Algonquin Provincial Park, Ontario in mid-July 1989 and were then buried in artificial nests in Ottawa. Hatchling *C.p. marginata* were excavated in November 1989 and transferred to a laboratory incubator set at 5°C . Eggs of the western subspecies of the painted turtle, *C.p. bellii*, were collected from Turtle Mountain Provincial Park, Manitoba in July 1989, and were incubated in Winnipeg in artificial nests. Hatchling *C.p. bellii* were excavated and shipped to Ottawa in September 1989 and then held in an incubator at 5°C . All turtles were held for two weeks prior to their use in experiments. In the incubators, all species were held in plastic boxes containing damp sphagnum moss. Animals were used in different experiments involving exposure to either freezing, dehydration, or anoxia stresses in order to elucidate whether the development of freeze tolerance by *R. sylvatica* was related to (or might have evolved from) pre-existing

amphibian abilities to survive dehydration or anoxia exposure. These experiments are described in the following sections.

General Freezing Protocol for Frogs and Turtles

A typical freezing exposure was conducted in a similar fashion to that described in Storey (1987a), with only slight modifications to the protocol. Each animal was placed on a pad of dry paper toweling with the ventral side (the plastron of turtles) centered over a thermistor. The animal and thermistor were secured in place with several strips of masking tape to ensure that no movement was possible during the freezing process. Previous experiments have shown that the taping of *R. sylvatica* produced no activation of glycogenolysis or glucose production in this anuran. Further, glycogenolysis was activated only after ice nucleation occurred on the skin of these animals (Storey and Storey, 1985b). This unit was then placed in an incubator set at a given subzero temperature, usually $-4 \pm 0.1^{\circ}\text{C}$, and cooling was monitored until a freezing exotherm was observed. Thermistors were connected to a Yellow Springs Instrument telethermometer, which in turn was connected to a Kipp and Zonen linear chart recorder.

Body surface temperature was recorded over time as animals cooled to their supercooling points over 30-40 min. Temperature recordings for each individual were made every 10-15 s as the body temperature of the animal neared the supercooling point. If the body temperature of individual animals reached the temperature of the incubator without nucleation occurring, then the incubator temperature was lowered in 0.5°C increments. After each adjustment body temperature was allowed to fall until it again equalled incubator temperature or until nucleation was seen.

Nucleation was observed by an exotherm after the animals had reached their supercooling points. The initiation of freezing was monitored as a brief exotherm (seconds) as the animals' body temperatures warmed from their supercooling point of

-2 to -3°C to a temperature of -1 to -0.5°C. Body temperatures 1 min after exotherms appeared were taken as the rebound temperature (closely approximates the freezing point of body fluids) and was $-0.5 \pm 0.1^\circ\text{C}$ for both frogs and turtles. Incubator temperature was raised to -2.5°C after exotherms were recorded for all animals being monitored and was held there for the remainder of the freezing exposure. The lengths of all freezing periods were timed from the appearance of the exotherm.

Two freezing protocols were designed for *R. sylvatica*. The first, a bulk freeze, involved freezing for 24 h at -2.5°C. Sixty frogs were placed in boxes on damp toweling, 10 frogs per box. Two frogs in each box were attached to thermistors (as described above) for monitoring of body surface temperature and the mean time to exotherm for these two was taken as the time zero for measuring the length of freezing exposure for all 10 frogs in the box. The overall mean time to exotherm for all 6 boxes was 30 ± 3 min and agrees with previous reports (Storey and Storey, 1986). Boxes were placed at -4°C until all exotherms were observed and then at -2.5°C for 24 h. Thirty frogs were sampled (5 from each box) in the frozen state at 24 h post-exotherm. The 30 remaining frogs were allowed to recover in an incubator set at 5°C for 24 h and were then sampled as thawed animals.

The second freezing protocol, a time course of freezing, was completed using a total of 70 frogs in the following manner. Frogs were individually taped to thermistors as described above and placed in incubators set at -4°C until each individual exotherm was recorded; then the incubator temperature was quickly readjusted to -2.5°C for the remainder of the freeze. Animals were sampled after 2, 5, or 20 min, or 1, 4, 8, or 24 h of freezing exposure (timed from the exotherm). For the 4, 8, and 24 h freezing time points, frogs were frozen in groups of ten, in bulk, in plastic boxes containing damp paper towelling with two animals taped to thermistors as described previously; the mean time to exotherm for these two was taken as time zero for determining the length of freezing exposure. However, frogs sampled at the 2, 5, and 20 min and 1 h time

points were monitored individually with freezing time calculated from the appearance of the individual exotherm.

C.p. marginata, *C.p. bellii*, *T.s. elegans* were subjected to a similar bulk freezing protocol as was used for *R. sylvatica*. Sixty individuals of each species were frozen for either 24 h (*C.picta* subspecies) or 4 h (*T.s. elegans*). Groups of 10 turtles were placed into plastic boxes lined with damp paper towelling (6 boxes per species). In each box two individuals were taped to thermistors as described above for monitoring exotherms. Boxes were then placed in an incubator set to -4°C until all exotherms were recorded, and then turtles were frozen at -2.5°C for 4 or 24 h. Half of the animals from each species were sampled at the end of the freezing time while still frozen, whereas the other half were allowed to thaw and recover at 5°C for 24 h before sampling.

Control animals for all freezing experiments were sampled directly from the 5°C incubator. These and experimental animals were all killed by double pithing (frogs) or decapitation (turtles); tissues were then excised and immediately frozen in liquid nitrogen. The tissues sampled from frogs were liver, hind leg skeletal muscle, brain, heart, and kidney. These, as well as gut (stomach and intestines combined), were collected from hatchling turtles. Frozen tissues were removed from liquid nitrogen, placed in appropriately marked vials, and stored at -80°C until required.

General Dehydration Protocol for Frogs

Dehydration experiments were begun by placing *R. sylvatica* or *R. pipiens* in large glass vacuum desiccators with 300 g of desiccant (silica gel) on the bottom. Desiccant was separated from the frogs by a 2 cm thick pad of sponge. Five frogs were put in each desiccator; the ratio of frog weight to silica gel was chosen to result in a rate of body water loss of 0.5-1.0 % of total body water per hour in both species. The tops of the desiccators were sealed with petroleum jelly to prevent any excess moisture

from entering the desiccating chambers but the stopcock on each desiccator was left open so that oxygen availability would not be impeded. Desiccators were placed in an incubator set at 5°C. Seventy-five *R. sylvatica* and 75 *R. pipiens* frogs were used in five experimental groups ($n = 15$ per group). In addition, 15 control frogs from each species were used in the following dehydration experiment for a total of 90 frogs per species. Animals were sampled when they had lost 10, 20, or 40% of total body water. In addition, two other groups were first dehydrated to 40 % of total body water lost and were then allowed to rehydrate either to 20 % of total body water lost or to the fully rehydrated state. Tissues used from 5% dehydrated frogs were taken from dehydration experiments conducted a year earlier. For rehydration, frogs were placed in plastic containers with 1 cm distilled water on the bottom so that the water was not covering the heads of the animals. Complete rehydration took 1 day for both species. The percentage of total body water lost (or regained) was determined by the following equation:

$$\begin{aligned} \text{Equation 1: } \% \text{ Dehydrated} &= \% \text{ of total body water lost due to dehydration} \\ &= [W_i - W_d] / [W_i \times \% \text{ H}_2\text{O}] \end{aligned}$$

where; W_i is the initial weight of the frog before dehydration, W_d is the weight of the frog at any given time during the experiment, and % H_2O is the percentage of total body weight of control frogs that is water. This last parameter was determined from initial and final weight measurements of control frogs that were killed and then dried to a constant weight for 72 h at 80°C. This value was $80.8 \pm 1.2 \%$ for *R. sylvatica* and $81.4 \pm 1.9 \%$ for *R. pipiens*.

A typical dehydration experiment was completed in the following manner. Bladders of all frogs were evacuated by gently squeezing frogs between thumb and forefinger before beginning. This allowed for the elimination of all variable water reserves from influencing the water loss pattern of individuals. Frogs were weighed on

top loading balance and the initial body weights were recorded. The weight of water in each animal was then calculated assuming an 80 % body water content and recorded; for example, a 5 g frog would have 4 g of water. Then a target body weight was calculated for each frog according to the target percent dehydration required. For example, if the 5 g frog above was to be sampled at the 10 % dehydrated point, then the target weight for sampling would be $5 \text{ g} - (10 \% \times 4 \text{ g})$ or 4.6 g. To minimize handling during experimental dehydration, all animals were routinely weighed every 4 h for the first day and twice a day subsequently. This level of animal handling resulted in constant replenishment of oxygen to the animals, therefore no hypoxic conditions were encountered. When each frog neared its calculated target weight it was weighed more frequently, usually every 30 min. Once each frog reached its target weight, it was sacrificed by double pithing. Tissues were immediately excised, frozen in liquid nitrogen, placed in properly labelled containers, and stored at -80°C until needed.

General Anoxia Protocol for Frogs

R. sylvatica and *R. pipiens* were given anoxic exposure in the following manner. Several 700 ml dark brown glass bottles were covered with aluminum foil and placed on ice so that only the tops were exposed. To each bottle was added 50 ml of distilled water and this was bubbled with a 2% CO_2 / 98% N_2 gas mixture for 20 min. The hose was always placed in the distilled water to ensure that oxygen was removed from the water and the air inside each bottle. Five frogs were placed in each bottle and the bottles were then sealed with parafilm except for a small hole. Bottles were flushed with N_2 gas through this hole for a further 30 min to ensure an essentially anoxic environment. This N_2 gassing procedure results in a drop of O_2 content to less than 2 % of normal air (15 Torr compared to 760 Torr). Although oxygen is present, at this very low level, oxygen limitation is severe and presents an anoxic metabolic challenge to the animals. Henceforth the term "anoxia" (or anoxic), which appears throughout the

thesis, will refer to the above metabolic stress condition. Bottles were then completely sealed with double parafilm and each bottle top was tightly secured. All bottles were then placed in an incubator equilibrated at 5°C. Time of anoxia exposure was taken from the point at which each bottle was sealed and placed in the incubator. Times for anoxia exposure were ½ h, 1 h, 4 h, and 2 d. Control frogs were sampled directly from the holding boxes that were stored in 5°C incubators. Individuals from both species were all alive and seemed quite healthy after the 2 d anoxia exposure. Once each frog reached its target time of anoxia exposure, it was sacrificed by double pithing. Tissues were immediately excised, frozen in liquid nitrogen, placed in properly labelled containers, and stored at -80°C until needed.

Preparation of Tissue Extracts for Metabolite and Second Messenger Measurements

Perchloric acid extracts were prepared from frozen tissue samples of all species for the measurement of cyclic AMP (cAMP), cyclic GMP (cGMP), inositol 1,4,5-trisphosphate (IP₃), and glucose concentrations. Frozen tissue samples were ground to a powder under liquid nitrogen using a mortar and pestle. The frozen tissue powder was then added to a preweighed polypropylene tube which was then quickly weighed a second time (to give tissue weight by subtraction). Five volumes (1:5 w/v) of ice-cold 6 % perchloric acid (PCA) containing 1 mM EDTA were then added and the sample was immediately homogenized on ice using a sonicating Polytron homogenizer. Samples were centrifuged at 27,000 g at 5°C in a Sorvall RC-5B refrigerated centrifuge for 20 min to precipitate denatured protein. A measured aliquot of the supernatant was removed and a volume of KOH solution (3.0 N KOH, 0.4 M KCl, 0.3 M Tris base) that was 0.25 x the measured volume of the supernatant aliquot was added and mixed for 1 min. Further adjustments of the extract to a pH between 7-7.5 was achieved with subsequent additions of small volumes (1-3 µl) of the KOH

solution. Neutralized solutions were then centrifuged for 20 min at 5°C. The supernatant was saved and stored at -20°C until needed. There was no degradation of cyclic nucleotides, IP₃, or glucose in the extracts as long as they were kept frozen.

Cyclic Nucleotide and Inositol 1,4,5-Trisphosphate Assays

Cyclic AMP, cGMP, and IP₃ were assayed using Amersham radioimmune assay (RIA) kits that are specific for each of these second messengers. Competitive binding assays for cAMP and IP₃ are based on the competition between unlabelled second messenger (in the sample extract) and tritiated second messenger for binding to a protein that has a high specificity for the second messenger. For cGMP competition is for binding to an antiserum. The amount of radiolabelled second messenger bound is inversely proportional to the amount of second messenger in the tissue extract. All measurements were of bound radioactive second messenger so that, for example, a low amount of radioactivity in a sample indicated a large concentration of second messenger in the original tissue extract. All samples were counted in a water soluble scintillation fluid (Ready GelTM) on a Packard 1900 CA liquid scintillation analyzer. Second messenger concentrations in tissue extracts were then determined by comparison with a standard curve prepared with known amounts of second messenger. Data for second messenger standards were treated by linear regression to calculate the slope and intercept, and then the picomoles of second messenger in each sample was calculated as follows:

Equation 2: $(C_0/C_{\text{unknown}} - \text{intercept})/\text{slope}$

where C_0 = number of counts/min (cpm) in the zero standard (with no added IP₃,

C_{unknown} = number of counts/min (cpm) in a sample. Final second messenger

concentrations in tissues, in pmol per gram wet weight (gww), were determined from the following equation:

Equation 3: $\text{pmol/gww} = (\text{pmol}/V_4) \times \text{D.F.}$

The dilution factor (D.F.) is:

Equation 4: $[(V_1 + W)/W] \times [(V_2 + V_3)/V_2]$

where W = weight of tissue sample in grams, V_1 = volume of PCA used, V_2 = volume of acid supernatant removed, V_3 = volume of KOH solution added, and V_4 = volume of neutralized extract added to the binding assay, all volumes in milliliters (1 ml).

Glucose Assay

Glucose concentrations in PCA extracts of *R. sylvatica* and *T.s. elegans* tissues were measured using a fluorometric coupled enzyme assay. The excitation wavelength used was 340 nm and the emission wavelength was 460 nm. A standard curve of NADH at concentrations of 0, 1, 2, 5, 10, 20, and 50 μM was completed so that the fluorescence change seen for each extract could be converted to μmol of NADH produced; glucose consumption in the reaction occurs in a 1:1 ratio with NADH produced. The reaction mixture contained 0.1 mM NAD^+ , 0.1 mM ATP, 5 mM MgSO_4 , 1 IU NAD^+ -dependent glucose-6-phosphate dehydrogenase, G6PDH, in 50 mM Tris buffer, pH 8.0, in a final volume of 1 ml. Sample extract (10 - 100 μl) was added to each cuvette and an initial fluorescence (F_1) reading was taken. G6PDH was added first to remove any endogenous G6P in the sample. After a stable baseline was achieved (approximately 30 min) 1 IU of hexokinase was added to measure total glucose in the sample. The reaction was allowed to proceed to completion (typically 1 h). Final fluorescence readings (F_2) were then taken. A high concentration of ATP (5

mM) was then added to the cuvette after F_2 was measured to ensure that the reaction end point had been determined by the consumption of all glucose and not by a limitation of the cosubstrate. The difference between the two fluorescence measurements ($F_2 - F_1$) for each cuvette was then converted into $\mu\text{moles NADH}$ produced for each sample. The equation for the line of the NADH standard curve was calculated in the form of $y = mx + b$, where m = the slope, x = the concentration of NADH produced, and b = the y intercept. The change in fluorescence for each sample was substituted into the rearranged equation $x = (y - b)/m$, which gave $\mu\text{moles NADH}$ produced in each sample. For all experiments there was no difference between tissue and cellular glucose concentrations. Tissue glucose concentrations, in $\mu\text{mol/g wet weight}$, were then calculated as follows, using the dilution factor as defined above:

Equation 5: $\mu\text{mol/gww} = \mu\text{mol NADH} \times \text{D.F.}$

Cyclic AMP-Dependent Protein Kinase Assay Protocol

The activity of cAMP-dependent protein kinase (PKA) was monitored by a radioactive assay that measured the incorporation of radiolabelled ^{32}P -ATP onto the phosphate-accepting artificial substrate "kemptide" (Jiang and Corbin, 1991). Tissue samples previously frozen at -80°C were quickly weighed and then immediately homogenized 1:30 w/v in PEM buffer; 10 mM KPi , 2 mM β -mercaptoethanol, and 1 mM EDTA, pH 6.8, with 1 mM phenylmethylsulfonylfluoride (PMSF) added immediately prior to homogenization. Samples were homogenized with a Polytron PT 10 homogenizer. This homogenization protocol was chosen to ensure that the *in vitro* concentration of cAMP in each homogenate was extremely low so that endogenous cAMP concentrations would not interfere with the measurements of total versus free catalytic forms of PKA. Homogenates were centrifuged at 27,000 g for 30 s in a Baxter Canlab Biofuge 15 microcentrifuge and the supernatants were removed and used to assay for PKA and PKAc activities.

Various stock solutions were prepared fresh daily as follows: Solution I: 120 mM KP_i , 0.9 mM kemptide peptide, 6 mM EDTA; Solution II: 1.42 mM ATP, 12.48 mM magnesium acetate ($MgOAc$), 60 $\mu Ci/ml$ ^{32}P -ATP; Solution III: 6 μM cAMP, 300 mM KCl; Solution IV: 300 mM KCl. For each sample determination, 3 conditions were run: (1) reaction mixture with no added enzyme (control), (2) reaction mixture with enzyme but minus cAMP (measures catalytic subunit, PKAc activity), and (3) reaction mixture with enzyme and 1 μM cAMP (total PKA activity). The cAMP concentration used to measure total PKA was chosen based on preliminary optimization tests that showed that frog liver PKA was saturated at a cAMP concentration of 1 μM (Fig. 1). Reactions were carried out in disposable glass tubes with standard assay conditions which gave the following concentrations of reactants in a final volume of 60 μl : 20 mM phosphate, 150 μM kemptide, 237 μM cold ATP, 2.1 mM Mg^{2+} , 1.0 mM EGTA, 2.2×10^6 dpm/assay (1.0 $\mu Ci/assay$) ^{32}P -ATP, and plus/minus 1 μM cAMP. The final pH of the assay mix was always 6.8. Additions were made in the following order: distilled water, solution III (+ cAMP) or solution IV (-cAMP), solution I, enzyme preparation, and finally solution II. Reactions were run for 5 min at 22°C or 15 minutes at 5°C. Initial trials determined the optimum amount of tissue extract (homogenate) used in each assay and demonstrated that reaction rates were linear for longer than the standard 5 min reaction time. Reactions were not limited by protein (enzyme) concentration.

Reactions were stopped using the P81 filter paper method of Roskoski (1983). Briefly, 40 μl aliquots of each reaction mixture were spotted onto individual 2 x 2 cm pieces of Whatman P81 filter paper. The papers were washed with four changes (50 ml each) of 75 mM H_3PO_4 for 5 min. Then they were placed in 1.5 ml microcentrifuge tubes held in 20 ml scintillation vials and radioactivity was counted using Cerenkov methodology on a Packard Model 1900CA scintillation counter. One unit of PKAc

Fig. 1. Effect of increasing amounts of free cyclic AMP on the activity of crude liver PKA. Data are means $\pm$ S.E.M. for $n = 3$ separate enzyme preparations of crude homogenate. The S.E.M. bars are contained within the symbols.

activity is defined as the amount of enzyme which catalyzes the incorporation of 1 nmol of ^{32}P from ATP onto kemptide in 1 min at 22°C .

The percentage of free catalytic subunit of PKA and the total PKA activity in all tissue extracts was determined in the following manner. For each tissue type at each sampling point four separate ($n = 4$) homogenates were prepared. A total of 48 reaction mixtures were made prior to the addition of the PKA enzyme source, 12 reaction tubes for each extract. All reaction tubes were prepared as described above with the following changes. These included four control reactions; the first two run in the absence of added enzyme for 5 min and with or without added cAMP and then stopped, and the second two in the presence of added enzyme with or without cAMP but stopped immediately. The remaining 8 reaction tubes contained four tubes with no cAMP, which measured PKAc, and four tubes containing $1\ \mu\text{M}$ cAMP, which measured total PKA.

Unless otherwise stated, a $10\ \mu\text{l}$ aliquot of a 1:30 (w/v) tissue homogenate was added to each reaction tube. The addition of enzyme source was done as quickly as possible, approximately 2 min after homogenization, to ensure that endogenous phosphatases and kinases had no time to affect the overall distribution of PKA and PKAc. Radioactive ^{32}P -ATP was added last and the reactions were run for 5 min at 22°C to allow for the full incorporation of ^{32}P into the kemptide substrate. Preliminary trials altered time and protein amount to ensure that the standard reaction conditions chosen were not limited by protein (enzyme) concentration.

Determination of Protein Concentrations

Protein concentrations for purified enzymes and PCA-precipitated protein fractions of tissue homogenates were determined using the Coomassie blue G-250 dye-binding method (Bradford, 1976). The Bio-Rad protein assay kit was used with bovine serum albumin as the standard. Before assaying, acid precipitates from

perchloric acid extractions were dissolved in 0.2 N NaOH. Absorbance was monitored at 595 nm using a Pye Unicam 8610 UV/Vis spectrophotometer.

Data Calculations and Statistical Analyses

Data are reported as means $\pm$ S.E.M. (standard error of the mean). Statistical significance was determined using either the Student's t-test or a one way analysis of variance (ANOVA) with a Student-Newman-Keuls test. Correlation coefficients for lines fit to data were determined using linear regression analysis (1st order regression). Statistical analysis was completed using a simple computer program with statistical tests for the analysis of enzyme kinetic data (Brooks, 1992).

All kinetic data for substrate binding was initially analyzed through Hill plots. When $h = 1$ (the Hill coefficient) the data was described using Michaelis-Menten hyperbolic kinetics. I_{50} for inhibitors were determined by Job plots (V_0/V_i versus $[I]$); where V_0 = enzyme velocity with no inhibitor present, V_i = enzyme velocity with inhibitor present, and $[I]$ = concentration of inhibitor used (Brooks, 1992). All Job plots were linear.

For purified PKAc all kinetic parameters at 22 and 5°C are presented as means $\pm$ S.E.M. for $n = 3$ separate preparations. All other data are presented as means $\pm$ S.E.M. for $n = 3$ to $n = 6$ separate homogenates or acid precipitations. Kinetic parameters were defined as the following: (1) K_m (Michaelis constant), the concentration of substrate at which the enzyme reaction rate proceeds at 50% of its maximal velocity, and (2) I_{50} , the concentration of inhibitor at which the enzyme reaction rate is inhibited by 50%. All kinetic values were determined by fitting the data to Michaelis-Menten and Job plots using a simple computer kinetics program (Brooks, 1992). The concentrations for free Mg^{2+} , ATP, and all ionic species such as K^+ , Cl^- , PO_4^{3-} , EDTA needed to determine the correct pH-temperature relationship for the pH

curves of *R. sylvatica* PKAc were calculated using the BAD program developed by Brooks and Storey (1992).

CHAPTER #3

SIGNAL TRANSDUCTION, SECOND MESSENGER, AND PROTEIN KINASE RESPONSES DURING FREEZING IN FOUR FREEZE TOLERANT VERTEBRATES

INTRODUCTION

A full understanding of the regulation of glucose output as a cryoprotectant in response to freezing in *R. sylvatica* requires a study of the basic signal transduction molecules that may be responsible for this phenomenon. The primary enzyme that is important in glucose production is glycogen phosphorylase. The enzyme in *R. sylvatica* liver is activated within seconds after ice formation begins at peripheral body sites and initiates the rapid breakdown of glycogen that leads to the export of large quantities of glucose into the blood for cryoprotection (Storey, 1987b). Glycogen phosphorylase regulation has been well studied in mammals. The enzyme is controlled by reversible protein phosphorylation and activation of the enzyme is catalyzed by phosphorylase kinase, which converts the inactive, dephosphorylated β form of the enzyme into the active, phosphorylated α form. In turn, phosphorylase kinase must first be converted to its active form through phosphorylation by cAMP-dependent protein kinase (PKA). The activation of PKA, then, could be the critical initiating step that leads to the activation of glycogen phosphorylase, and to the rapid production of cryoprotectant by *R. sylvatica* liver in response to a freezing stress.

The coordination and mediation of many cellular processes is controlled by a family of enzymes known as protein kinases. The most widely studied of these enzymes are cyclic AMP-dependent protein kinase (PKA), calcium and phospholipid dependent protein kinase (PKC), and cyclic GMP-dependent protein kinase (PKG). The second messengers for these enzymes activate their respective protein kinase by binding to the kinase and causing either a release of an activated form (subunit) of the kinase as in PKA and PKC, or as in the case of PKG, an activated form of the holoenzyme. Modulation of the levels of second messengers is regulated by changes in the levels of hormones and other extracellular signals and these second messenger signals then

activate the appropriate protein kinases which, in turn, phosphorylate one or more proteins/enzymes to achieve the desired metabolic response.

Although many of the biochemical events accompanying cryoprotectant biosynthesis in *R. sylvatica* have been well documented (Storey and Storey, 1992b), the roles of protein kinases in the process, particularly protein kinase A, have not. Although it has been presumed that the freezing-stimulated activation of PKA is the initiating event in activating liver cryoprotectant synthesis, and the inhibitory effects of the β -adrenergic blocker, propranolol, on cryoprotectant output tend to support this, there has been no direct evidence of PKA involvement. The coordination of freezing-induced liver glycogenolysis with respect to changes in the levels of second messengers, their relationship to the activity of PKA, and even whether or not PKA controls glucose production in this amphibian have not previously been investigated and these topics became the initial goals of this study.

Furthermore, despite a primary interest in the second messenger and protein kinase controls of cryoprotectant synthesis, the overall involvement of protein kinase mediation in regulating the many other metabolic adaptations required for freezing survival has never received attention. Thus, by identifying both the types and patterns of second messengers produced and protein kinases activated during freezing, insights could be gained into the regulation of various cellular adjustments that support freezing survival. The studies described in this chapter document the pattern of changes in cellular cAMP levels and PKA activation in *R. sylvatica* tissues as well as the distinctly different effects of freezing on the levels of cGMP and IP₃, the second messengers of other cellular protein kinases. In addition, the analysis of second messengers and PKA activity is extended to the tissues of freeze tolerant turtle species. These produce little, if any, cryoprotectant and a comparison of the second messenger and PKA responses between species could further highlight and differentiate between events that are linked

solely to cryoprotectant synthesis and those that stimulate/regulate other aspects of cellular adaptation for freezing survival.

MATERIALS AND METHODS

Preparation of Experimental Animals

Freezing protocols were carried out as described in chapter 2 with frogs sampled from two different experiments: (1) control (5°C), 24 h frozen at -2.5°C, and 24 h frozen followed by 24 h thaw at 5°C, or (2) control (5°C) versus 2, 5, or 20 min, and 1, 4, 8, or 24 h freezing exposure at -2.5°C. Freezing protocols for turtles were conducted as described in chapter 2 with sampling under the following conditions: (1) control (5°C), (2) 4 h or 24 h frozen at -2.5°, and (3) 4 h or 24 h freezing followed by 24 h thaw at 5°C.

PKA Assay and Determination of Percent PKAc

The activity of cAMP-dependent protein kinase (PKA) was monitored with the use of a radioactive assay as in described in chapter 2. Reactions measuring the amount of free catalytic subunit (PKAc) were completed in the absence of cAMP whereas the total PKA activity was measured in assays run in the presence of 1 μ M cAMP to fully dissociate the holoenzyme; the percentage of total activity present as PKAc was then calculated.

Second Messenger and Glucose Concentrations

The second messengers cAMP, cGMP, and IP₃ were measured in all tissues sampled from *R. sylvatica* and *T.s. elegans*, but only liver, muscle, and brain of *C.p. marginata* and *C.p. bellii*. Glucose levels were assayed in *R. sylvatica* and *T.s. elegans* tissues. Methods for metabolite assay were as described in chapter 2.

RESULTS

Effect of Freezing Exposure on Total PKA Activity and the Percent PKAc in *R. sylvatica* Tissues

Initial studies centered around characterizing changes in the activity of cAMP-dependent protein kinase in tissues of *R. sylvatica* as a result of freezing (24 h at -2.5°C) and thawing (24 h returned to 5°C after freezing). Table 1 shows that the percentage of total PKA that was present as the active free catalytic subunit (PKAc) changed significantly with freezing and thawing in liver, skeletal muscle, and heart, but not in brain and kidney. The highest percent PKAc was 33.4 % in skeletal muscle of frozen frogs. PKAc content doubled in muscle in response to freezing stress and remained elevated in thawed tissue. Liver and heart showed different patterns of response. In both tissues, the percentage of PKAc decreased slightly during freezing but then increased to a value significantly above the control level after thawing. Liver showed a 4-fold increase in the percent PKAc between the frozen and thawed states whereas the corresponding change in heart was 1.4-fold.

Total activity of PKA was not altered by freezing or thawing in any of the *R. sylvatica* tissues tested (Table 1). Skeletal muscle exhibited the lowest total PKA activity of 54 pmol P_i transferred/min/g wet weight; PKA activities in liver, heart, and kidney were 18-22 fold higher than this and brain showed the highest total PKA activity, 44-fold greater than in muscle. A similar pattern was observed for total PKA activity in turtle, *T.s. elegans*, tissues. Freezing (4 h at -2.5°C) and thawing (24 h at 5°) had no effect on the total activity of the enzyme in any of the six tissues tested (Table 2). Skeletal muscle again showed the lowest total PKA activity of 162 pmol P_i transferred/min/gww with values in liver, gut, brain, heart, and kidney being 5, 9, 10, 11, and 17-fold higher, respectively. Comparisons of total PKA activity in the tissues

Table 1. Changes in the amount of active PKA (expressed as a percentage) present as the free catalytic subunit and total PKA activities in tissues from control (5°C), frozen (24 h, -2.5°C), and thawed (24 h, 5°C) *Rana sylvatica*.

Tissue	Condition	% Free PKAc	Total Activity
Liver	Control	7.1 ± 0.3	1031 ± 39
	Frozen	5.1 ± 0.3 ^a	1114 ± 101
	Thawed	20.6 ± 0.6 ^{a,b}	1232 ± 77
Muscle	Control	15.3 ± 0.9	54 ± 5
	Frozen	33.4 ± 1.1 ^a	70 ± 6
	Thawed	28.2 ± 1.3 ^{a,b}	59 ± 2
Brain	Control	12.3 ± 0.6	2373 ± 168
	Frozen	13.7 ± 0.6	2078 ± 41
	Thawed	13.0 ± 0.3	2275 ± 71
Heart	Control	11.3 ± 0.6	984 ± 63
	Frozen	9.6 ± 0.4 ^a	974 ± 12
	Thawed	13.4 ± 0.5 ^{a,b}	1003 ± 26
Kidney	Control	7.1 ± 0.3	1210 ± 153
	Frozen	6.2 ± 0.3	1331 ± 34
	Thawed	7.1 ± 0.4	1400 ± 66

Data are means ± S.E.M., n = 4 animals. Total PKA activity in crude tissue homogenates is reported as pmol P_i transferred/min/g wet weight.

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, P < 0.05.

^b - Significantly different from the corresponding 24 h frozen value, P < 0.05.

Table 2. Changes in the amount of active PKA (expressed as a percentage) present as the free catalytic subunit and total PKA activities in tissues from control (5°C), frozen (4 h, -2.5°C), and thawed (24 h, 5°C) *Trachemys scripta elegans*.

Tissue	Condition	% Free PKAc	Total Activity
Liver	Control	6.0 ± 0.3	886 ± 74
	Frozen	3.0 ± 0.1 ^a	1056 ± 68
	Thawed	4.1 ± 0.1 ^a	1167 ± 57
Muscle	Control	32.2 ± 0.8	162 ± 16
	Frozen	34.2 ± 1.4	216 ± 22
	Thawed	20.1 ± 1.3 ^{a,b}	227 ± 34
Brain	Control	13.2 ± 0.2	1571 ± 42
	Frozen	25.2 ± 0.8 ^a	1418 ± 37
	Thawed	14.4 ± 0.4 ^b	1533 ± 47
Heart	Control	4.8 ± 0.2	1814 ± 204
	Frozen	5.2 ± 0.2	1471 ± 10
	Thawed	6.0 ± 0.2 ^{a,b}	1490 ± 34
Kidney	Control	4.2 ± 0.1	2826 ± 144
	Frozen	4.6 ± 0.2	2725 ± 181
	Thawed	7.4 ± 0.5 ^{a,b}	3124 ± 557
Gut	Control	5.9 ± 0.1	1400 ± 219
	Frozen	5.6 ± 0.2	1231 ± 144
	Thawed	5.7 ± 0.2	930 ± 93

Data are means ± S.E.M., n = 4 animals. Total PKA activity in crude tissue homogenates is reported as pmol P_i transferred/min/g wet weight. Gut includes stomach and intestine.

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, P < 0.05.

^b - Significantly different from the corresponding 4 hr frozen value, P < 0.05.

of the two freeze tolerant species showed the following: (1) no significant difference in total liver PKA activity between species, (2) a 1.5-fold higher total PKA activity in brain of the wood frog, and (3) 2-3 fold higher PKA activities in turtle muscle, heart, and kidney (Table 1 and Table 2).

Table 2 also shows the effect of freezing and thawing on the distribution of PKAc in red-eared slider tissues. This turtle can survive short term freezing episodes of up to 4 h at -4°C (Churchill and Storey, 1992a). Freezing significantly altered the % PKAc only in liver (a 50 % decrease) and in brain (a 2-fold increase). The percent PKAc remained reduced in liver during thawing and also decreased significantly in skeletal muscle of thawed turtles. In brain PKAc content returned to control values during thawing whereas in heart and kidney PKAc content increased significantly during thawing. Neither freezing nor thawing had an effect on PKAc content in gut (stomach + intestines combined).

Table 3 shows the effects of freezing (24 h at -2.5°C) on the percent PKAc in liver and skeletal muscle of two turtle species with well-developed freeze tolerance, the Midland *C.p. marginata* and Western *C.p. bellii* subspecies of painted turtle (data for *T.s. elegans* are repeated for comparison). Freezing exposure stimulated a significant decrease in the percent PKAc in liver of *C.p. marginata*, similar to the results for *T.s. elegans* liver. During thawing, liver PKAc content rose significantly above the value in liver of frozen turtles, but did not regain the control value. Neither freezing nor thawing altered PKAc content in liver of the Western subspecies. Skeletal muscle PKAc showed opposite responses in the two painted turtle subspecies. In *C.p. marginata* muscle, PKAc content increased significantly during freezing by 1.5-fold and remained elevated during thawing. In *C.p. bellii* muscle, however, PKAc content decreased by nearly half during freezing and remained significantly suppressed in the thawed tissue.

Table 3. Changes in the amount of active PKA (expressed as a percentage) present as the free catalytic subunit in three species of turtles under control, frozen, and thawed conditions.

Tissue	Condition	% Free Catalytic PKA		
		<i>T.s. elegans</i>	<i>C.p. marginata</i>	<i>C.p. bellii</i>
Liver	Control	6.0 ± 0.3	7.8 ± 0.5	2.7 ± 0.2
	Frozen	3.0 ± 0.1 ^a	2.9 ± 0.5 ^a	3.2 ± 0.2
	Thawed	4.1 ± 0.1 ^a	4.4 ± 0.4 ^b	3.4 ± 0.3
Muscle	Control	32.2 ± 0.8	19.5 ± 1.6	13.7 ± 0.8
	Frozen	34.2 ± 1.4	29.4 ± 1.2 ^a	8.8 ± 0.4 ^a
	Thawed	20.1 ± 1.3 ^{a,b}	30.4 ± 1.1 ^a	8.1 ± 0.3 ^a

Conditions are: controls (5°C), frozen at -2.5°C (4 h for *T.s. elegans* and 24 h for *C. picta* subspecies), and thawed (24 h, 5°C). Data are means ± S.E.M., n = 4.

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, P < 0.05.

^b - Significantly different from the corresponding frozen value, P < 0.05.

Effects of Freezing and Thawing on Tissue Second Messenger Concentrations

Tables 4 through 7 show the effects of freezing and thawing on the levels of cAMP, cGMP and IP₃ in tissues of *R. sylvatica*, *T.s. elegans*, *C.p. marginata* and *C.p. bellii*, respectively.

Adenosine 3'-5' monophosphate: In the freeze tolerant wood frog *R. sylvatica*, freezing and thawing *in vivo* affected cAMP concentration only in the liver (Table 4); cAMP rose by 1.9-fold during freezing and remained elevated in liver from thawed animals. Levels of the second messenger were unchanged in the four other organs but notably, skeletal muscle cAMP concentrations were substantially lower (by 5-10 fold) than the amounts of cAMP in other organs. During freezing in *T.s. elegans* cAMP content increased by 54 % in brain but decreased significantly in both heart and gut to values 63 and 34 %, respectively, of the corresponding controls (Table 5). Levels of cAMP returned to control values during thawing in brain and heart but remained suppressed in gut. In *C.p. marginata* freezing exposure led to significant increases in cAMP levels in skeletal muscle and brain and in both organs, cAMP was sharply reduced again upon thawing (Table 6). By contrast, cAMP in *C.p. marginata* liver was unaffected by freezing exposure but cAMP levels increased after thawing. The same pattern was seen in liver of *C.p. bellii* (Table 7). Skeletal muscle cAMP in the Western subspecies also responded similarly to the pattern seen in the Midland painted turtle with a significant increase in cAMP during freezing and a strong decrease in the second messenger content in the tissue after thawing. However, freezing and thawing had no effect on cAMP levels in brain of *C.p. bellii*.

Guanosine 3'-5' cyclic monophosphate: Overall, levels of cGMP in tissues of both frogs and turtles were much lower than the amounts of cAMP. Freezing and thawing stimulated changes in this second messenger in some tissues although the patterns were distinctly different than those seen for cAMP. In *R. sylvatica* freezing exposure resulted in a significant decrease in cGMP content in skeletal muscle and

Table 4. Second messenger concentrations in tissues of control (5°C), frozen (24 h, -2.5°C), and thawed (24 h, 5°C) *Rana sylvatica*.

Tissue	Condition	[Second Messenger] (pmol/gww)		
		cAMP	cGMP	IP ₃
Liver	Control	615 ± 57	50 ± 7	1406 ± 164
	Frozen	1138 ± 60 ^a	62 ± 4	16525 ± 2983 ^a
	Thawed	1147 ± 430 ^a	106 ± 13 ^{a,b}	3214 ± 737 ^b
Muscle	Control	246 ± 42	76 ± 8	2131 ± 150
	Frozen	118 ± 20	52 ± 5 ^a	3310 ± 341 ^a
	Thawed	242 ± 138	80 ± 1 ^b	1518 ± 212 ^{a,b}
Brain	Control	1387 ± 186	101 ± 3	20133 ± 1689
	Frozen	921 ± 204	81 ± 2	22807 ± 2465
	Thawed	1109 ± 210	88 ± 9	21345 ± 1922
Kidney	Control	1323 ± 127	190 ± 5	10335 ± 1944
	Frozen	1538 ± 186	96 ± 5 ^a	13203 ± 1335
	Thawed	1894 ± 257	94 ± 5 ^a	11660 ± 3132
Heart	Control	1522 ± 69	94 ± 8	5724 ± 879
	Frozen	1122 ± 197	116 ± 12	7773 ± 1316
	Thawed	1261 ± 98	85 ± 6	4490 ± 811

Data are pmol/g wet weight, means ± S.E.M., n = 4 separate acid precipitates for each tissue.

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, P < 0.05.

^b - Significantly different from the corresponding 24 h frozen value, P < 0.05.

Table 5. Second messenger concentrations in tissues of control (5°C), frozen (4 h, -2.5°C), and thawed (24 h, 5°C) *Trachemys scripta elegans*.

Tissue	Condition	[Second Messenger] pmol/gww)		
		cAMP	cGMP	IP ₃
Liver	Control	1325 ± 215	53 ± 4	3487 ± 624
	Frozen	771 ± 145	120 ± 11 ^a	17945 ± 3124 ^a
	Thawed	1296 ± 167	96 ± 7 ^a	2915 ± 277 ^b
Muscle	Control	1080 ± 257	116 ± 12	4570 ± 359
	Frozen	662 ± 92	92 ± 2	3902 ± 80
	Thawed	544 ± 98	57 ± 8 ^{a,b}	4739 ± 673
Brain	Control	1623 ± 64	70 ± 7	17938 ± 3245
	Frozen	2504 ± 112 ^a	72 ± 8	6233 ± 426 ^a
	Thawed	1832 ± 69 ^b	58 ± 3	3287 ± 330 ^{a,b}
Kidney	Control	2025 ± 330	88 ± 7	8446 ± 1297
	Frozen	1343 ± 170	102 ± 9	12034 ± 2567 ^a
	Thawed	1606 ± 105	78 ± 7	5753 ± 534 ^{a,b}
Heart	Control	1579 ± 128	65 ± 5	11298 ± 639
	Frozen	1007 ± 58 ^a	45 ± 3 ^a	14756 ± 1643 ^a
	Thawed	1403 ± 59	86 ± 10 ^b	9286 ± 609 ^{a,b}
Gut	Control	3892 ± 607	515 ± 40	7816 ± 1371
	Frozen	1313 ± 89 ^a	58 ± 9 ^a	22193 ± 2490 ^a
	Thawed	1205 ± 264 ^a	76 ± 7 ^a	16542 ± 3175 ^{a,b}

Data are pmol/g wet weight, means ± S.E.M., n = 4 separate acid precipitates for each tissue. Gut includes stomach and intestine.

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, P < 0.05.

^b - Significantly different from the corresponding 4 h frozen value, P < 0.05.

Table 6. Second messenger concentrations in tissues of control (5°C), frozen (24 h, -2.5°C), and thawed (24 h, 5°C) midland painted turtles *Chrysemys picta marginata*.

Tissue	Condition	[Second Messenger] pmol/gww		
		cAMP	cGMP	IP ₃
Liver	Control	575 ± 17	25 ± 1	3439 ± 248
	Frozen	518 ± 19	38 ± 4	15094 ± 946 ^a
	Thawed	658 ± 25 ^{a,b}	26 ± 3	1770 ± 113 ^{a,b}
Muscle	Control	548 ± 23	30 ± 2	618 ± 51
	Frozen	2097 ± 56 ^a	46 ± 4	1379 ± 140
	Thawed	1265 ± 50 ^{a,b}	34 ± 2	730 ± 88
Brain	Control	955 ± 33	40 ± 2	17868 ± 1873
	Frozen	1213 ± 96 ^a	49 ± 3 ^a	16091 ± 2437
	Thawed	591 ± 24 ^{a,b}	36 ± 3 ^b	19892 ± 1288

Data are pmol/g wet weight, means ± S.E.M., n = 4 separate acid precipitates for each tissue.

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, P < 0.05.

^b - Significantly different from the corresponding 24 h frozen value, P < 0.05.

Table 7. Second messenger concentrations in tissues of control (5°C), frozen (24 h, -2.5°C), and thawed (24 h, 5°C) western painted turtles *Chrysemys picta bellii*.

Tissue	Condition	[Second Messenger] pmol/gww		
		cAMP	cGMP	IP ₃
Liver	Control	523 ± 21	33 ± 2	3008 ± 273
	Frozen	427 ± 36	34 ± 1	13249 ± 1999 ^a
	Thawed	708 ± 50 ^b	28 ± 1	4917 ± 324 ^{a,b}
Muscle	Control	1121 ± 94	34 ± 2	1025 ± 86
	Frozen	1788 ± 283 ^a	33 ± 2	1092 ± 239
	Thawed	555 ± 44 ^{a,b}	27 ± 2 ^{a,b}	895 ± 116
Brain	Control	788 ± 22	45 ± 3	16505 ± 936
	Frozen	727 ± 40	41 ± 2	16587 ± 3196
	Thawed	762 ± 30	44 ± 3	11642 ± 1196

Data are pmol/g wet weight, means ± S.E.M., n = 4 separate acid precipitates for each tissue.

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, P < 0.05.

^b - Significantly different from the corresponding 24 h frozen value, P < 0.05.

kidney by 32 and 50 %, respectively (Table 4). Muscle cGMP rebounded upon thawing but kidney cGMP remained reduced. By contrast, liver cGMP levels, which were unaffected during freezing, doubled during thawing. Freeze/thaw exposure altered cGMP levels in four tissues of *T.s. elegans* (Table 5). In liver, cGMP increased by 2.3-fold during freezing and remained elevated in thawed animals. However, freezing exposure reduced cGMP in heart and gut, levels in frozen gut being only 11 % of control values. cGMP remained low in gut after thawing but levels increased above control values in heart. Skeletal muscle showed a 50 % decrease in cGMP during thawing. The contents of cGMP in liver and muscle of *C.p. marginata* were unaffected by freeze/thaw but cGMP increased significantly during freezing in brain with levels reverting to control values after thawing (Table 6). In *C.p. bellii* cGMP levels were unaffected by freeze/thaw in liver and brain but levels of the second messenger decreased slightly in thawed skeletal muscle (Table 7).

Inositol 1,4,5-trisphosphate: Levels of IP₃, one of the second messengers for protein kinase C activation, also changed in response to *in vivo* freezing and thawing exposures of frogs and turtles. In the wood frog, levels of this second messenger increased dramatically in liver, by 11-fold, during freezing and also rose to a lesser extent, by 55 %, in skeletal muscle (Table 4). IP₃ in both tissues was reduced again after thawing. Contents of the second messenger were unaffected by freeze/thaw in other tissues of the frog. Freezing stimulated changes in IP₃ levels in all tissues of *T.s. elegans* except skeletal muscle (Table 5). IP₃ increased during freezing by 5-fold in liver, 2.8-fold in gut, and by 42 and 31 % in kidney and heart, respectively; oppositely, freezing reduced IP₃ levels in turtle brain by 65%. After thawing, IP₃ content was reduced even further in brain and IP₃ also fell in all other tissues during thawing, as compared with the frozen state (although only in liver were control values reestablished). Painted turtles also showed dramatic increases in IP₃ content in liver of frozen animals, amounts increasing by 4.4-fold in both subspecies (Table 6,7). Levels

were again reduced upon thawing but remained higher than control values in *C.p. marginata* and fell to 50 % of the control value in *C.p. bellii*. Freeze/thaw had no effect on the IP₃ content of either skeletal muscle or brain of painted turtles.

Tissue Distribution of Glucose

In freeze tolerant frogs, freezing exposure stimulates a rapid biosynthesis and export of glucose by liver that leads to the accumulation of high levels of this sugar in all organs of the frozen animal where it acts as a cryoprotectant (Storey and Storey, 1986). Since changes in second messenger levels and protein kinase activities may be intimately involved in triggering and regulating cryoprotectant synthesis, it is important to correlate changes in these parameters with changes in organ glucose contents. Table 8 shows the effects of freezing and thawing on tissue glucose levels in *R. sylvatica*. For comparison, values for *T.s. elegans* are also shown; turtle species typically show only minimal increments in glucose content during freezing and do not, apparently, make any substantial amounts of colligative cryoprotectants derived from glycogen (Churchill and Storey, 1992a). Table 8 shows that 24 h of freezing at -2.5°C stimulated an enormous increase in glucose concentration in *R. sylvatica* liver and heart, over 200-fold in both cases. Glucose also rose to high levels in brain and kidney, increasing by 76-fold in both cases, and skeletal muscle glucose increased by 16-fold. Similar freezing-induced tissue-specific distributions of glucose have been reported previously; the progressive freezing of the frog from the outside inwards results in a gradation of glucose values with the lowest values in peripheral organs and the highest in core tissues (Storey, 1987b). After 24 h of thawing, glucose levels were significantly reduced in four organs (except for skeletal muscle which showed an increase) and glucose content was much more uniform throughout the frog as would be expected once circulation was restored.

Table 8. Effect of freezing and thawing on glucose concentrations in tissues of spring *Rana sylvatica* and *Trachemys scripta elegans*.

<i>Rana sylvatica</i>		$\mu\text{mol glucose/gww}$	
Tissue	Control	Frozen	Thawed
Liver	1.3 ± 0.1	261.0 ± 29.9^a	$42.2 \pm 1.0^{a,b}$
Muscle	0.9 ± 0.04	14.4 ± 0.7^a	$21.6 \pm 1.8^{a,b}$
Brain	0.8 ± 0.16	60.8 ± 3.2^a	$31.1 \pm 2.1^{a,b}$
Kidney	1.2 ± 0.1	90.8 ± 5.8^a	$45.6 \pm 2.1^{a,b}$
Heart	1.0 ± 0.1	241.9 ± 15.8^a	$49.5 \pm 5.8^{a,b}$
<i>Trachemys scripta elegans</i>		$\mu\text{mol glucose/gww}$	
Tissue	Control	Frozen	Thawed
Liver	1.5 ± 0.4	7.4 ± 0.7^a	$15.8 \pm 2.4^{a,b}$
Muscle	1.2 ± 0.1	1.8 ± 0.3	$7.9 \pm 0.7^{a,b}$
Brain	1.5 ± 0.1	1.5 ± 0.1	$0.5 \pm 0.1^{a,b}$
Kidney	1.3 ± 0.1	2.1 ± 0.3^a	$5.2 \pm 0.5^{a,b}$
Heart	1.4 ± 0.1	1.1 ± 0.1	$8.3 \pm 0.6^{a,b}$
Gut	1.1 ± 0.1	1.9 ± 0.4^a	3.1 ± 0.2^a

Data are $\mu\text{mol/g}$ wet weight, means $\pm$ S.E.M., $n = 4$ animals. Conditions are: control (5°C), frozen at -2.5°C (24 h for frogs, 4 h for turtles), and thawed (24 h, 5°C).

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, $P < 0.05$.

^b - Significantly different from the corresponding frozen value, $P < 0.05$.

Freezing had much less of an effect on tissue glucose levels in *T.s. elegans*. Glucose rose by 5-fold in liver of frozen turtles and increased slightly in kidney and gut but was unchanged in skeletal muscle, brain, and heart. After thawing, however, glucose levels had risen in five tissues (but were reduced in brain), reaching a maximum of 15.8 $\mu\text{mol/g}$ wet weight in liver and about 8 $\mu\text{mol/gww}$ in heart and skeletal muscle.

Changes in PKAc, cAMP, and Glucose in *R. sylvatica* Tissues Over a Time Course of Freezing.

The percentage of active PKA rose quickly in *R. sylvatica* liver after nucleation, reaching a maximum of $62.3\% \pm 2.5$ after 5 min of freezing exposure at -2.5°C (Table 9). The free catalytic (active) form of liver PKA remained elevated well above control levels up to 8 h post-exotherm but had declined to control values after 24 h. Interestingly, however, when the frogs were thawed PKAc was again elevated; PKAc content was 2.5 fold higher in liver from thawed frogs when compared with control PKAc levels for the same organ. PKAc content also increased quickly in muscle during freezing, rising 3.5-fold within the first 2 min of freezing. Subsequently, muscle PKAc content remained well above control levels throughout the freezing time course (Table 9). Freezing also stimulated small increases in PKAc content in brain and heart with significant changes noted in as little as 2 min post-exotherm. Subsequently, PKAc gradually returned to control levels as the time of freezing exposure approached 24 h and remained similar to control values in thawed frogs. In kidney, PKAc content showed only a minor increase after 1 h freezing but otherwise was unaltered over the freeze/thaw time course.

The levels of two second messengers, cAMP and IP_3 , were also measured in selected *R. sylvatica* tissues over the same time course of freezing exposure. The data are presented in Table 10. The concentration of cAMP in *R. sylvatica* liver increased

Table 9. Changes in the amount of active PKA (expressed as a percentage) present as the free catalytic subunit in *Rana sylvatica* tissue over a time course of freezing at -2.5°C, followed by thawing for 24 h at 5°C.

Tissue	Liver	Muscle	Brain	Heart	Kidney
Control	7.1 ± 0.3	15.3 ± 0.9	12.3 ± 0.6	11.3 ± 0.6	7.1 ± 0.4
Frozen					
2 min	43.4 ± 2.3 ^a	55.5 ± 3.9 ^a	16.9 ± 0.7 ^a	9.2 ± 0.4 ^a	5.4 ± 0.3
5 min	62.3 ± 2.5 ^a	44.9 ± 1.8 ^a	16.4 ± 0.5 ^a	16.2 ± 0.5 ^a	6.2 ± 0.3
20 min	60.6 ± 2.3 ^a	42.5 ± 1.8 ^a	16.5 ± 0.7 ^a	13.1 ± 0.5 ^a	5.3 ± 0.2
1 h	58.0 ± 0.6 ^a	41.8 ± 1.3 ^a	13.5 ± 0.5	8.9 ± 0.5 ^a	9.4 ± 0.7 ^a
4 h	29.5 ± 0.6 ^a	45.4 ± 2.7 ^a	14.3 ± 0.5	11.6 ± 0.5	7.4 ± 0.4
8h	33.1 ± 3.0 ^a	24.4 ± 1.7 ^a	17.4 ± 0.6 ^a	13.3 ± 0.5 ^a	6.2 ± 0.3
24 h	5.1 ± 0.3	33.4 ± 1.1 ^a	13.7 ± 0.6	9.6 ± 0.4	6.2 ± 0.3
Thawed					
24 h	20.6 ± 0.6 ^a	28.2 ± 1.3 ^a	13.0 ± 0.3	13.4 ± 0.5 ^a	7.1 ± 0.4

Data are means ± S.E.M., n = 4 animals. The data for control, 24 h frozen, and 24 h thawed animals are taken from Table 1.

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, P < 0.05.

Table 10. Effect of freezing exposure on cAMP and IP₃ concentrations in *Rana sylvatica* tissues over a freezing time course at -2.5°C, followed by thawing for 24 h at 5°C.

Condition	Liver pmol cAMP/gww	Muscle pmol cAMP/gww	Brain pmol IP ₃ /gww
Control	615 ± 57	214 ± 60	20133 ± 1689
Frozen			
2 min	1125 ± 68 ^a	238 ± 4	15264 ± 1452
5 min	1322 ± 214 ^a	232 ± 12	21609 ± 1186
20 min	1512 ± 109 ^a	290 ± 10	19272 ± 1187
1 h	1414 ± 72 ^a	130 ± 5 ^a	20689 ± 3086
4 h	1020 ± 201 ^a	130 ± 24 ^a	34569 ± 1444 ^a
8 h	1042 ± 130 ^a	124 ± 8 ^a	35157 ± 1132 ^a
24 h	1138 ± 60 ^a	118 ± 3 ^a	22807 ± 2465
Thawed			
24 h	1147 ± 97 ^a	242 ± 57	21345 ± 1922

Data are means ± S.E.M., n = 4 animals. All second messenger values for control liver, muscle, and brain tissues are taken from the bulk freezing experiment, Table 4.

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, P < 0.05.

quickly after nucleation, doubling after only 5 min and remaining elevated at 20 min and 1 h post-exotherm. Subsequently, cAMP levels decreased somewhat but remained elevated, compared with control values, throughout the freezing time course. Wood frog skeletal muscle cAMP content showed a different response; cAMP levels were unaffected during the early minutes of freezing. With longer periods of freezing, however, cAMP concentrations fell to values about 60 % of control. Muscle cAMP content rose again during thawing.

IP₃ concentrations in *R. sylvatica* brain are also shown in Table 10. IP₃ concentration was unaltered over the early minutes of freezing exposure but amounts had risen significantly after 4 and 8 h of freezing. Levels of this second messenger returned to near-control values after 24 h of thawing.

Changes in tissue glucose concentrations over the course of freezing are presented in Table 11. Glucose levels rose over time during freezing in each tissue. In liver, a significant increase in glucose was noted at as little as 2 min post-exotherm and glucose content had risen by 224-fold after 24 h of freezing. Muscle and brain showed significant increases in tissue glucose within 2 and 5 min post-exotherm, respectively, and in both tissues glucose continued to increase over time during freezing although total accumulation in muscle was low. After 24 h thawing, glucose content was decreased substantially in liver and brain but glucose in muscle was actually higher after 24 h thawing than in frozen frogs.

Table 11. Changes in tissue glucose concentrations over a time course of freezing exposure at -2.5°C , followed by thawing at 5°C in *Rana sylvatica*.

Condition	Liver	Muscle	Brain
Control	1.2 ± 0.01	1.0 ± 0.09	0.4 ± 0.02
Frozen			
2 min	3.1 ± 0.2^a	3.2 ± 0.1^a	0.5 ± 0.04
5 min	4.4 ± 0.3^a	1.9 ± 0.2^a	0.8 ± 0.2^a
20 min	4.8 ± 0.4^a	2.2 ± 0.1^a	0.8 ± 0.1^a
1 h	4.1 ± 0.1^a	6.0 ± 0.3^a	1.9 ± 0.2^a
4 h	12.9 ± 1.4^a	6.2 ± 0.9^a	3.5 ± 0.2^a
8 h	32.9 ± 9.7^a	6.4 ± 0.3^a	2.7 ± 0.2^a
24 h	261.1 ± 29.9^a	7.2 ± 0.4^a	60.8 ± 3.2^a
Thawed			
24 h	42.2 ± 1.0^a	21.6 ± 1.8^a	31.1 ± 2.1^a

Data are means $\pm$ S.E.M., $n = 4$ animals.

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, $P < 0.05$.

DISCUSSION

The synthesis of cryoprotectants from carbohydrate reserves, liver glycogen in freeze tolerant frogs and fat body glycogen in insects, is a well known biochemical process in many cold hardy vertebrates and invertebrates (Storey and Storey, 1988; Zachariassen, 1985). All these freeze tolerant animals have adopted a similar metabolic strategy. This strategy involves directing carbon flow from glycogen stores into the production of carbohydrate cryoprotectant; for example, glucose in *R. sylvatica* and glycerol in *Epiblema scudderiana* (Storey and Storey, 1988; Churchill and Storey, 1989). In preparation for winter freezing episodes, glycogen stores in *R. sylvatica* liver are increased to 1 M levels and the contents of enzymes involved in the catabolism of glycogen must be maximized. This occurs during the autumn cold-hardening period and, in fact, glycogen levels in fall frogs have been measured at levels as high as 1000 $\mu\text{mol/g}$ (amounting to 18 % of liver fresh weight) (Storey and Storey, 1984, 1986), a more than adequate supply of carbon for the biosynthesis of cryoprotectant glucose. Typically, the glucose levels measured in tissues of frozen wood frogs have ranged from 200 to 400 mM in previous studies (Storey, 1987b; Storey and Storey, 1984).

My initial investigations aimed at identifying the molecular mechanism of protein kinase action involved in generating the mobilization of carbon stores in *R. sylvatica* upon the initiation of an inoculative freeze. Previous studies on this freeze tolerant anuran have shown overwhelming evidence that freezing induces: (1) an increase in total activity of glycogen phosphorylase (GP), and (2) an increase in the conversion of GP into its active a form from its inactive b form (Mommensen and Storey, 1992). What then is the biochemical mechanism that allows glycogenolysis and GP to become active in order to produce glucose during freezing? The answer may involve the activation of the glycogenolytic cascade via one or more protein kinases. The relationship between PKAc activity and that of glycogen phosphorylase activation has

been well documented in mammals (Hers, 1976). However, information on this relationship in freeze tolerant vertebrates is limited. Therefore these relationships were investigated in the wood frog *R. sylvatica*, the red eared slider turtle hatchling *T.s. elegans*, and the midland and western painted turtle hatchlings *C.p. marginata* and *C.p. bellii*, all species known to be freeze tolerant to varying degrees (Churchill and Storey, 1992a; Churchill and Storey, 1992b; Storey, 1984).

The wood frog has refined the glycolytic process so that it is able to mobilize its liver glycogen stores almost immediately after an external inoculative freezing stress. There is an active output of glucose in liver up to 200 $\mu\text{mol/gww}$ within the early minutes to hours of a freezing stress (Storey and Storey, 1986). Glucose is then shunted to other tissues via the circulatory system (Storey and Storey, 1992b). Once glucose reaches these tissues a concomitant increase in tissue glucose concentration is observed (Storey and Storey, 1986; this study). Tissue glucose concentrations measured in tissues of spring collected *R. sylvatica* (Table 8 and 11) are certainly within these values. Thus, the metabolic machinery responsible for cryoprotectant production in the fall must still be in place in *R. sylvatica* during the spring months. The wood frog then is an excellent model to use in the study of second messenger/protein kinase interrelationships in conjunction with vertebrate cyroprotectant production.

The fact that PKAc, cAMP, and cGMP levels did not show a significant increase in any tissue except muscle in *R. sylvatica* (Table 1 and 4) during the 24 h freeze would lead to the conclusion that PKA plays no role in the phosphorylation of the proteins responsible for glucose production in *R. sylvatica*. However, this is simply not the case. Upon freezing the wood frog for short periods of time, major increases in the level of the free catalytic subunit of PKA (PKAc) were seen in liver tissue (Table 9). The total concentration of cellular cAMP in liver also rose in the early minutes to hours of freezing (Table 10). Mommsen and Storey (1992) showed that a freezing stress increased both the total activity of phosphorylase expressed and the percentage of

the enzyme in the active a form in isolated hepatocytes from *R. sylvatica*. The results of the present experiment confirmed that freezing induced an increase in the levels of cAMP and an elevation of the free catalytic subunit of PKA. Elevated cAMP is well known to increase PKAc content by dissociating the holoenzyme PKA after binding to its regulatory subunit. The subsequent release of PKAc observed following an increase in cellular cAMP could, and in fact did, occur within minutes of a freezing stress. The release of PKAc may be responsible for the activation of glycogen phosphorylase within the early minutes of a freezing stress. This guarantees that the production and distribution of glucose from the liver to the major organs will occur well in advance of the animal reaching a fully frozen state.

Mommsen and Storey (1992) have shown that *R. sylvatica* liver phosphorylase, like the typical vertebrate form of the enzyme, responded to hormones (glucagon, epinephrine) that are known to increase cellular cAMP concentrations and initiate the activation of PKA. Additionally, the cAMP analogue, dibutyryl cAMP, effectively activated frog liver GP and caused a concurrent glucose output from the organ. The results of this thesis indicate that cAMP concentrations always increase prior to the release of PKAc during the early stages of a freezing stress. Thus, the response shown by cAMP and PKAc to freezing must be linked to the hormone response that activates glucose output from *R. sylvatica* liver. The cAMP and PKAc responses in liver occurred very quickly, within 2 min, after freezing began (Tables 9 and 10), indicating the potential for a rapid activation of GP by this protein kinase cascade system in *R. sylvatica* liver.

Only when PKA is activated by increased levels of cAMP and phosphorylates GPK (glycogen phosphorylase kinase), which in turn phosphorylates GP, is the glycolytic cascade primed to begin synthesizing glucose in large quantities from *R. sylvatica* liver glycogen. The data collected above indicate that there is a direct relationship between PKAc activity, second messenger levels and glucose production in

R. sylvatica. This also shows that freeze tolerant frogs have coordinated their cellular metabolic pathways, specifically the PKA/glycolytic cascade, prior to, and so that they may survive, a freezing stress.

Evidence is presented that indicates that organs other than liver may produce glucose in response to whole body freezing in the wood frog. Increases in the free catalytic subunit of PKA were observed in muscle, brain, heart, and kidney within 1 h after freezing began in *R. sylvatica* (Table 9). Glucose measurements in *R. sylvatica* muscle and brain revealed that this carbohydrate does in fact increase steadily during the early stages of a freezing stress (Table 11). This suggests that these organs may be involved in synthesizing at least some of their own glucose as a metabolic response to freezing. Glucose pools synthesized in these organs may provide protection until such time that cryoprotectant is delivered to these organs from the liver.

The three turtle species in this study are known to be freeze tolerant vertebrates that do not produce any substantial amounts of glucose as a cryoprotectant (Churchill and Storey, 1991; Churchill and Storey, 1992a). A study was completed in order to determine whether a similar pattern of second messenger production/release and protein kinase response is observed in these vertebrates as is seen in the glucose producing freeze tolerant vertebrate, *R. sylvatica*.

Initial results showed a similar distribution pattern of total PKA activity in *R. sylvatica* and *T.s. elegans* tissues (Table 1 and Table 2). These data showed that total enzyme activity did not change significantly from control levels in frozen and thawed tissues of either species. It seems reasonable to conclude then, that the total levels of cAMP-dependent protein kinase do not change upon exposure to whole body freezing in all freeze tolerant vertebrates. The main difference between glucose-producing freeze tolerant vertebrates, *R. sylvatica*, and those that are not, *T.s. elegans*, seems to lie in the activities of the free (active) catalytic subunit of PKA (PKAc). PKAc levels were

much higher in all organs, especially liver, of glucose-producing freeze tolerant vertebrates.

The only significant increase in the content of the free catalytic subunit of PKA in response to freezing was observed in *T.s. elegans* brain, frozen for 4 h (Table 2). A concurrent increase in the cAMP concentration was seen in frozen brain (Table 5), but no increase in the level of glucose output was observed in this organ during freezing (Table 8). Most notable, however, was the lack of an increase in the levels of active catalytic PKA in the liver of the three turtles in response to a freezing stress (Table 3). There was either no change (*C.p. bellii*) or a decrease (*C.p. marginata* and *T.s. elegans*) in the percentage of active PKA in this organ. The metabolic response to freezing for the PKA phosphorylation cascade system is the opposite to that seen in *R. sylvatica*. There seems to be no indication of a protein kinase A response in freeze tolerant turtles.

Glucose data showed that only minimal amounts of cryoprotectant biosynthesis occurred in response to freezing in the 3 turtle species (Table 8). Significant glucose elevations were observed in frozen liver, kidney, and heart tissue, the other increases were seen under the thawed tissue condition. These data were consistent with previous results that showed no significant glucose accumulation in the tissues of freeze tolerant turtles including, *C.p. marginata* and *C.p. bellii*, and the box turtle *Terrapene carolina* (Costanzo et al. 1988; Churchill and Storey 1991; Churchill and Storey 1992b; Storey et al. 1993). The concentrations of glucose in the organs of these turtles were clearly much less than the few hundred millimolar amounts produced by freeze tolerant frogs (Storey, 1990). Without the activation of glycogenolysis via the PKA cascade mechanism it is impossible to produce large quantities of glucose for the sole purpose of cryoprotection, as is the case in *R. sylvatica*. It seems that, at least in liver, these reptiles do not bother with the activation of their respective glycolytic enzyme pathways in response to a freezing stress. Instead, these freeze tolerant reptiles may have

developed an alternative strategy to survive for long periods of time in a frozen state. It appears then, that the response of the protein kinase A signal transduction cascade during freezing is entirely different in freeze tolerant vertebrates that produce cryoprotectant, *R. sylvatica*, and those that do not, *T.s. elegans* and *C. picta* subspecies.

The levels of glucose in these reptiles are not nearly enough to serve as a colligative cryoprotectant against the physical stresses imposed by the reduction of cell volume seen during a long term freezing episode. This is supported by the evidence that *T.s. elegans* is only able to survive brief episodes of freezing, a maximum of up to 4 h with 30-40% of its body water in the form of ice (Storey and Storey 1992b). The fact that this animal does not produce any large amount of cryoprotectant prior to freezing and does not depress its metabolism as it begins to freeze suggests that long term survival of this animal in the frozen state is unlikely. However, tissue glucose concentrations were elevated in all tissues (except brain) of the red-eared slider after thawing. Increased tissue glucose at this time may provide the needed energy currency to aid the turtle during post-freeze recovery.

The primary metabolic fuel for the vertebrate brain is glucose. It would seem critical for *T.s. elegans* and *R. sylvatica* to increase the supply of glucose to their brains prior to freezing. This was indeed the case for the wood frog but was not observed in red-eared sliders. Glucose and other putative cryoprotectants such as glycerol and lactate did not increase significantly in frozen tissues of *T.s. elegans* in a previous study (Churchill and Storey; 1992a). However as was shown in that study, levels of these metabolites are very similar to those found in anoxic tissues of *T.s. elegans* (Kelly and Storey, 1988). Increases seen in the levels of second messengers and PKAc for the red-eared slider turtle may reflect the animal's natural ability to survive anoxic, hypoxic, and ischemic stresses. Studies on the levels of second messengers and PKA

activation under these particular stresses must first be completed in order to test this hypothesis.

Red-eared sliders and painted turtles, unlike *R. sylvatica*, do not have the capacity for a rapid build up of cryoprotectant when exposed to a freezing stress, which is consistent with previous studies involving other reptiles. Many reptiles which exhibit limited freeze tolerance such as hatchling painted turtles, garter snakes, and box turtles also exhibit this slow production of low concentrations of glucose (Churchill and Storey 1991; Churchill and Storey, 1992b; Storey et al., 1993). Nevertheless a truly freeze tolerant animal must have the metabolic machinery in place so that a rapid build up and distribution of cryoprotectant occurs just prior to, and during a freezing episode. This is extremely critical for an organism if it is to survive in a frozen state for a prolonged period of time.

The second messengers cAMP, cGMP, and IP₃ act as allosteric effectors to directly activate protein kinase enzymes (Shenolikar, 1988). The mechanisms of action for many second messengers are quite well documented, perhaps the best known being that of cyclic AMP. The production of cAMP is triggered by the binding of a hormone to a cell surface receptor. This hormone stimulates adenylate cyclase, a membrane bound enzyme, to produce cAMP from ATP (Shenolikar, 1988). Free cytosolic cAMP covalently modifies PKA releasing the catalytic subunits of this protein kinase throughout the cell.

The target protein kinase enzymes for all second messengers analyzed in this study are known; PKA for cAMP, PKC for IP₃ and Ca²⁺, and PKG for cGMP. Evidence for the kinds of protein kinases involved in the response to freezing in vertebrates can be derived from the analysis of changes in second messenger levels during freezing and thawing. An increase in the level of a second messenger usually reflects an increase in the activity of the corresponding protein kinase enzymes in a

tissue. Hence, levels of the second messengers cAMP, cGMP, and IP₃ were quantified in this manner in the tissues of control, frozen, and thawed frogs and turtles.

The increased IP₃ levels seen in frozen *R. sylvatica* liver and brain (Table 4) may be indicative of protein kinase C activation. There was also an increased level of IP₃ in the frozen liver tissues obtained from all 3 turtle species (Tables 5, 6 and 7). An increase in the second messenger for protein kinase C (IP₃) during freezing seems to be a universal phenomenon in the liver of all 4 freeze tolerant vertebrates involved in this study. Therefore, the activation of PKC by IP₃ may have an important major function in both species that produce glucose and those that do not, and in good, *R. sylvatica*, and poor, *T.s. elegans*, freeze tolerant species. How then could the activation of liver PKC benefit these freeze tolerant vertebrates during a freezing episode? The answer may lie in the control of cell volume, specifically cell shrinkage due to water loss (Watson, 1991).

A recent study looking at the control of cell volume by protein kinases and phosphatases clearly resolved that the control of cell volume was not accomplished by PKA (Grinstein, et al. 1992). However the evidence for cell volume control by PKC remains inconclusive. Further studies are needed before PKC is ruled out as a potential regulator of cell volume, specifically by controlling the influx/efflux of ions through the plasma membrane. Liver PKC may play a critical role in these freeze tolerant vertebrates in controlling the influx of ions, particularly K⁺ and Na⁺, into the cell during the freezing process in *R. sylvatica*. During freezing, water is drawn out of cells down its concentration gradient where it is sequestered as ice. Extracellular ice formation elevates ion concentrations in the remaining extracellular fluid and this causes ion inflow into the cells and elevates intracellular ions in the frozen state. A strict control over the influx of these ions in the cells may be facilitated by the phosphorylation and subsequent inhibition of ion channel proteins by PKC. Thus, the activation of PKC may not only be controlled by the release of hormones in response to

freezing, as is the case for PKAc, but may in addition, be linked to changes in cell volume (Grinstein et al., 1992). Cell volume changes are known to occur in frozen and dehydrated *R. sylvatica* (Lee et al. 1992; Churchill and Storey, 1994). The fact that IP₃ levels, an activator of PKC, increased during freezing suggests that PKC may have a significant role in controlling the rate of freezing in all freeze tolerant vertebrates. By phosphorylating membrane channel proteins PKC may restrict the rate of water loss out of, and the influx of ions into, cells during a freezing episode.

A second potential mode of action for PKC activation must not be overlooked, and that is in glycogen phosphorylase control. Although the results of this study clearly show that PKA is very important in phosphorylase activation, PKC could also play a complementary role in regulating glycogenolysis during cryoprotectant production. Hence, both the extent of phosphorylase activation and the magnitude of the hyperglycemia stimulated by freezing are far beyond the "normal" vertebrate response to stimuli that activate liver glycogenolysis (such as the catecholamine-mediated fight-or-flight response). It can be proposed then, that the control of *R. sylvatica* glycogenolysis might need two regulatory mechanisms, one to allow responses to normal hyperglycemic stimuli (e.g. muscle exercise, starvation) and one to permit the massive glycogenolysis in response to freezing. Furthermore, the homeostatic control of vertebrate liver glycogen phosphorylase activity that is normally exerted by glucose (rising glucose inhibits phosphorylase α activity by making the enzyme more susceptible to phosphorylase phosphatase) must be overridden somehow to allow freezing-induced glucose accumulation to occur. The job of sustaining phosphorylase activation until several-hundred millimolar amounts of glucose have been accumulated may be a function of PKC.

CHAPTER #4

SIGNAL TRANSDUCTION, SECOND MESSENGER, AND PROTEIN KINASE RESPONSES DURING DEHYDRATION IN TWO ANURANS

INTRODUCTION

Stresses put on cells due to dehydration appear to have been a powerful selective agent in the colonization of terrestrial habitats by many amphibians (Thorson, 1955). Because their skins are highly water-permeable (unlike those of higher vertebrates), amphibian species have generally chosen relatively moist terrestrial habitats, and when they are faced with dry environments, one of two general strategies for avoiding dehydration are used. These are: (1) the development of water-proofing mechanisms, such as cocoons, that form a protective barrier to water loss, or (2) enduring the large variations in total body water that are a part of a dehydration stress (Katz, 1989). Proteins and macromolecular structures inside cells are still surrounded by large quantities of water when the whole animal is in a dehydrated state, even at 40 % of total body water lost. The two anurans in this study, *R. sylvatica* and *R. pipiens*, utilize the latter method to survive the stress of dehydration.

Severely dehydrated amphibians die due to the loss of critical water stores found in the blood and internal organs. As plasma water content drops, blood viscosity increases dramatically, retarding blood flow to tissues and compromising gas exchange. Death due to dehydration in animals occurs long before cellular water contents fall to a level that could destroy macromolecular structures by disrupting their bound water. The critical element for dehydrated whole animals is circulatory collapse due to the extreme viscosity and low volume of the blood as it loses water, stresses the heart, and eventually causes cardiac arrest (Hillman, 1987).

In a previous study it was determined that the wood frog *R. sylvatica*, a freeze tolerant species (Schmid 1982), also exhibited excellent dehydration tolerance. Autumn-collected animals readily tolerated the loss of 50 % of total body water whereas spring-collected frogs tolerated 60 % dehydration in experiments conducted at 5°C (Churchill and Storey, 1993). Furthermore, one of the major metabolic responses

to dehydration by *R. sylvatica* was an activation of liver glycogenolysis leading to a rapid elevation of glucose in liver, blood, and other organs (Churchill and Storey, 1993). This response to dehydration at 5°C was virtually identical to the well-described accumulation of glucose as a cryoprotectant in response to freezing in this species (Storey and Storey, 1984, 1992b) where the colligative actions of high glucose concentrations serve to lessen the extent of cellular and organ water loss during freezing (Costanzo et al., 1993). High glucose may also stabilize the structure and/or activities of enzymes that may be inhibited under the high ionic strength conditions that exist in cells of desiccated amphibians (Katz, 1989; Shoemaker, 1992). Dehydration and freezing both have the same effect in reducing cell volume, the one being caused by a net loss of water from the body whereas the other is caused by the sequestration of up to 65 % of total body water as extracellular and extra-organ ice (Lee et al. 1992; Churchill and Storey 1993; Storey and Storey 1993). Recent studies have indicated that changes in cell volume can have a wide range of important metabolic effects on cells (Watson, 1991).

I proposed, therefore, that the initiation of cryoprotectant production through the activation of glycogenolysis by protein kinase enzymes in response to freezing in *R. sylvatica*, evolved out of a pre-existing hyperglycemic response by amphibians to dehydration. Recent studies of the effects of dehydration on the semi-aquatic leopard frog *R. pipiens* support this proposal for this freeze intolerant frog also showed a rapid hyperglycemia, based on liver glycogenolysis, in response to dehydration (Churchill and Storey, 1994). However, the magnitude of the response was much lower in *R. pipiens* than in *R. sylvatica*; maximum glucose levels in liver of autumn-collected frogs, dehydrated to 50 % of total body water lost, were only 20 $\mu\text{mol/g}$ wet weight in the leopard frog versus 127 $\mu\text{mol/gww}$ in the wood frog (Churchill and Storey, 1993).

Several concepts were investigated in this study that addressed the influences of dehydration on the levels of protein kinases and second messengers in a freeze-tolerant

frog, *R. sylvatica*, versus a non-freeze-tolerant frog, *R. pipiens*. Dehydration puts a large osmotic stress on the cells of many anuran species (Hillman, 1980). Virtually the same osmotic stress on cells is observed in *R. sylvatica* when the wood frog is dehydrated or frozen (Churchill and Storey, 1993). Thus, which protein kinases and second messengers are involved in mediating the responses to dehydration stress and, if more than one protein kinase is involved, can different functions for each protein kinase be identified? To what extent do the metabolic responses observed for protein kinase enzymes and their respective second messengers during dehydration mimic those seen during a freezing episode in *R. sylvatica*? Do these same responses appear in *R. pipiens*, a species of frog that is known to be desiccation-tolerant but not freeze-tolerant?

The present study examines the metabolic responses to dehydration stress by the organs of the freeze-tolerant wood frog *R. sylvatica* and the freeze-intolerant leopard frog *R. pipiens*. Organ-specific metabolic responses to dehydration are investigated including changes in protein kinase A activity and the concentrations of its second messenger, cAMP, as well as effects on IP₃ concentrations with its implications for PKC activity. These are correlated with measurements of body water loss and tissue protein content, the latter providing an index of individual tissue water loss. Furthermore, the protein kinase responses to dehydration by *R. sylvatica* liver indicate the mode of regulatory control of liver glycogenolysis during dehydration-induced glucose export.

MATERIALS AND METHODS

Preparation of Experimental Animals

R. sylvatica and *R. pipiens* were collected, held, and dehydration experiments were conducted as stated in chapter 2.

Protein Measurements in Dehydrated *R. sylvatica* Tissues.

Protein concentrations in acid precipitates from *R. sylvatica* dehydrated tissues were determined as described in chapter 2. Protein content is expressed as mg protein/gram wet weight of tissue.

Protein Kinase A and Second Messenger Assay Protocols

The activity of cAMP-dependent protein kinase in tissues from dehydrated *R. sylvatica* and *R. pipiens* was monitored by a radioactive assay that measured the incorporation of radiolabelled ^{32}P -ATP onto the phosphate-accepting artificial substrate 'kemptide' as described in chapter 2. Perchloric acid extracts of frozen tissues samples from dehydrated frogs were prepared and levels of cAMP and IP_3 were determined using competitive binding radioimmune assay kits as described in chapter 2.

RESULTS

Protein Contents of Tissues from Dehydrated *R. sylvatica*

The water contents of individual tissues can change during whole body dehydration, although generally by low amounts, as tissue water content seems to be defended at the expense of extra-organ water (Churchill and Storey, 1993). However, if tissue water contents change significantly over the course of dehydration then values for metabolite concentrations, measured per g wet weight, would be inaccurate. Hence, to avoid the introduction of errors due to changes in organ wet mass during dehydration, metabolite levels in these experiments were expressed per mg protein. Previous studies have shown that protein content of tissues remains constant during dehydration as tissue protein, measured per g dry weight, did not change over the course of dehydration in either species (Churchill and Storey 1993, 1994).

Table 12 shows the effect of whole body dehydration and rehydration on the protein concentrations in 5 tissues in *R. sylvatica*. All tissues and organs showed an increase in the amount of protein, measured as mg/g wet weight, when dehydration reached high values. These increases were significant at whole body water loss values of 20 % in kidney and 40 % in other organs. However, protein concentrations fell again when frogs were rehydrated to the 20 % dehydrated or the fully rehydrated states. These data support the idea that despite whole body water loss, organ water contents are largely undisturbed until total body water loss reaches high values.

Effect of Dehydration on the Percent of PKAc in Frog Liver and Muscle

Changes in the percentage of PKA present as the free catalytic subunit were monitored over the course of dehydration in liver and muscle of both *R. sylvatica* and *R. pipiens* (Tables 13 and 14). Dehydration to 20 % of total body water lost resulted in a significant 3-fold increase in PKAc content in *R. sylvatica* liver. PKAc fell somewhat in liver of 40 % dehydrated frogs but remained above control values. When frogs were rehydrated, the % PKAc dropped and after full rehydration, PKAc was actually significantly lower than the original control values. In skeletal muscle, dehydration also resulted in a significant increase in PKAc content that was sustained in muscle of 40 % dehydrated frogs. However, PKAc content again decreased during rehydration, dropping below the original control value (Table 13).

Dehydration in leopard frogs *R. pipiens* gave different results (Table 14). The percent PKAc was unchanged by dehydration in both liver and muscle of leopard frogs and also remained constant during rehydration in muscle. However, as also occurred in *R. sylvatica*, PKAc content was sharply reduced in liver of rehydrated frogs.

Tables 13 and 14 also show the actual values for the percentage of total body water lost by frogs in each experimental group; these were calculated from measurements of total body weight lost during dehydration, using a value of 80 % for

Table 12. Total protein concentrations in tissue samples from *Rana sylvatica* sampled over a course of whole body dehydration and rehydration.

Condition	Protein (mg/gww)				
	Liver	Muscle	Brain	Heart	Kidney
Control	164 ± 2.6	182 ± 15.3	64 ± 3.0	113 ± 4.3	82 ± 7.3
Dehydrated	5 %	159 ± 2.5	55 ± 3.3	101 ± 5.9	100 ± 1.8
	10 %	170 ± 7.9	71 ± 5.1	115 ± 1.7	93 ± 5.5
	20 %	171 ± 20.3	66 ± 4.0	119 ± 10.4	118 ± 2.7 ^a
Rehydrated	40 %	257 ± 22.5 ^a	86 ± 3.5 ^a	157 ± 8.1 ^a	123 ± 11.6 ^a
	20 %	167 ± 4.5	79 ± 5.5	110 ± 3.4	84 ± 2.2
	0 %	165 ± 5.9	74 ± 1.3	113 ± 4.4	81 ± 2.8

Data are means ± S.E.M. for n = 4 animals.

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, $P < 0.05$.

Dehydration: frogs were sampled when the percentage of body water lost reached 5, 10, 20 or 40 % of total body water.

Rehydrated frogs were first dehydrated to 40 % of total body water lost and then allowed to regain water to return to a value equivalent to 20 % or 0 % total body water lost as compared to controls. Fully rehydrated (0 %) frogs were allowed to rehydrate for 24 h at 5 °C in a container filled with water to a depth of 1 cm; final water regained was within ± 5 % of the original weight.

Table 13. Effect of changes in total body water content on the amount of active PKA (expressed as a percentage) present as the free catalytic subunit in *Rana sylvatica* liver and skeletal muscle.

Condition		Liver PKAc (% active)	Muscle PKAc (% active)	Calculated % Body Water Lost
Control		9.9 ± 1.1	58.3 ± 3.2	--
Dehyd.	20 %	31.1 ± 1.6 ^a	67.5 ± 2.1 ^a	19.8 ± 0.1
	40 %	16.4 ± 0.8 ^a	65.7 ± 1.7 ^a	40.4 ± 0.2
Rehyd.	20 %	10.2 ± 0.7	48.3 ± 2.7 ^a	18.1 ± 1.0
	0 %	6.6 ± 0.4 ^a	41.0 ± 2.3 ^a	--

Data are means ± S.E.M. for n = 4 animals.

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, P < 0.05.

Dehydration: frogs were sampled when the percentage of body water lost reached 20 or 40 % of total body water. Rehydrated frogs were first dehydrated to 40 % of total body water lost and then allowed to regain water to return to a value equivalent to 20 % or 0 % total body water lost when compared to controls. Fully rehydrated (0 %) frogs were allowed to rehydrate for 24 h at 5°C in a container filled with water to a depth of 1 cm; final water regained was within ± 5 % of the original weight.

Table 14. Effect of changes in total body water content on the amount of active PKA (expressed as a percentage) present as the free catalytic subunit in *Rana pipiens* tissues.

Condition	Liver PKAc (%a)	Muscle PKAc (%a)	Calculated % Body Water Lost
Control	21.9 $\pm$ 1.4	41.0 $\pm$ 2.3	--
Dehy. 10 %	26.3 $\pm$ 1.4	45.8 $\pm$ 3.2	9.9 $\pm$ 0.2
20 %	28.3 $\pm$ 1.6	48.0 $\pm$ 2.6	19.7 $\pm$ 0.6
40 %	26.7 $\pm$ 1.2	43.6 $\pm$ 0.4	40.3 $\pm$ 0.3
0 %	11.9 $\pm$ 0.5 ^a	44.3 $\pm$ 1.9	--

Data are means $\pm$ S.E.M., n = 4 animals. The symbol %a represents the percent active catalytic subunit of PKA. Other information as in Table 13, except frogs were allowed to dehydrate to 10 % as well as 20 or 40 % of total body water lost and then sampled.

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, P < 0.05.

the water content of control frogs and assuming that all weight loss is due to water loss. The data show that actual measured values were virtually identical to target values.

Tissue Distribution of Second Messengers

Increases in the tissue concentrations of second messengers usually indicate increased activities of the associated protein kinases in the tissue. The effects of dehydration and rehydration of the levels of cAMP and IP₃ in five tissues of *R. sylvatica* are shown in Table 15. Liver and kidney were the only tissues that showed significant increases in cellular cAMP concentration in response to whole body dehydration; cAMP levels were unaltered over the dehydration/rehydration course in the other tissues (Table 15). In liver the response by cAMP was very sensitive to dehydration with cAMP increasing rapidly by 3-fold in frogs that had lost only 5 % of total body water. Levels remained 2-fold higher than control values in 10 % dehydrated frogs but cAMP levels fell at higher levels of dehydration and were not significantly different from control values at these points or when frogs were rehydrated. Kidney showed a different pattern of response; cAMP levels did not change at the lower levels of dehydration but at 40 % dehydration, the kidney concentration of cAMP increased 5-fold. With rehydration, however, kidney cAMP returned to values close to control (Table 15). Tissue-specific differences in cAMP levels were again noted (as were seen in Chapter 3). Levels of cAMP were far lower in skeletal muscle than in any other tissue and were about 10-fold higher in liver and progressively higher still in heart, kidney, and brain.

Dehydration also stimulated changes in liver IP₃ concentrations. Levels were significantly elevated by 5 % dehydration and rose progressively to reach values 4-fold higher than controls in the 40 % dehydrated animals (Table 15). In partially rehydrated frogs, IP₃ levels fell somewhat but remained 3-fold higher than controls. As also occurred for cAMP, IP₃ levels in kidney increased significantly in 40 % dehydrated

Table 15. Effect of changes in total body water content on the concentrations of cyclic AMP (cAMP) and inositol triphosphate (IP₃) in *Rana sylvatica* tissues.

Condition		Tissue levels of cAMP (pmol/mg protein)			
		Liver	Muscle	Brain	Kidney
Control		38 ± 3.4	5 ± 0.9	381 ± 23.3	265 ± 22.6
Dehydrated	5 %	128 ± 21.3 ^a	6 ± 0.3	345 ± 23.3	340 ± 22.1
	10 %	80 ± 5.4 ^a	4 ± 0.5	309 ± 8.5	224 ± 13.1
	20 %	28 ± 2.1	4 ± 0.4	390 ± 13.3	268 ± 9.5
Rehydrated	40 %	33 ± 1.9	5 ± 0.2	324 ± 15.1	1431 ± 77.9 ^a
	20 %	41 ± 5.8	4 ± 0.2	317 ± 18.8	330 ± 21.7
	0 %	67 ± 4.3	7 ± 0.3	414 ± 12.6	411 ± 98.5
Condition		Tissue levels of IP ₃ (pmol/mg protein)			
		Liver	Muscle	Brain	Kidney
Control		41 ± 2.9	7 ± 0.2	2442 ± 145.4	698 ± 25.0
Dehydrated	5 %	68 ± 7.9 ^a	6 ± 0.3	2218 ± 227.9	899 ± 49.0
	10 %	131 ± 32 ^a	7 ± 0.3	2794 ± 482.1	758 ± 33.8
	20 %	114 ± 35 ^a	8 ± 0.4	2173 ± 152.5	752 ± 69.2
Rehydrated	40 %	166 ± 14.6 ^a	4 ± 0.5 ^a	2076 ± 148.2	1110 ± 45.2 ^a
	20 %	113 ± 7.7 ^a	8 ± 1.0	1807 ± 410.0	680 ± 4.3
	0 %	89 ± 6.6 ^a	6 ± 0.2	3164 ± 67.4	606 ± 120.0

Data are means ± S.E.M., n = 4 animals. Other information is as in Table 12. For reference purposes 38 pmol cAMP/mg protein corresponds to a value of 623 ± 36 pmol/gwt; 41 pmol IP₃/mg protein corresponds to a value of 1358 ± 364 pmol/gwt. ^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, P < 0.05.

animals but returned to control levels during rehydration. IP₃ content was not altered by dehydration or rehydration in either brain or heart and fell slightly in skeletal muscle from 40 % dehydrated frogs. Tissue specific differences in IP₃ levels were again seen with very low values in skeletal muscle, about 10-fold higher values in liver, even higher amounts in heart and kidney, and the highest IP₃ content was found in brain.

Survival Following Dehydration

The mortality of both frog species was monitored throughout the experimental dehydration courses. Three of 75 *R. sylvatica* died over the course of dehydration, a survival rate of over 95 %; whereas 7 of 75 *R. pipiens* died under the same experimental conditions, a survival rate of 90 %. Death was indicated by no righting response and no heart beat, even after rehydration. Smaller frogs from both species lost water at a faster rate than larger animals, but dehydration tolerance did not vary with the size of the frog. Frogs that died during dehydration exposure were not used for experimentation; however, an $n = 4$ was retained for all dehydration points.

DISCUSSION

Both species of frogs readily endured the loss of up to 40 % of total body water during experimental dehydration. This agrees with previous studies on these two species (Churchill and Storey, 1993, 1994) and is also generally within the vital limits for various terrestrial and semi-terrestrial anurans (Shoemaker, 1992). Water loss during dehydration appears to come first from extra-organ fluids, such as abdominal fluid and plasma (bladder water was not a factor in these studies as frogs were induced to void bladders before experimentation began) since Table 12 shows that organ protein contents showed no significant change until 20 % (for kidney) or 40 % (for other organs) of total body water had been lost. Because total organ protein (mg/g dry

weight) remained constant under dehydration stresses (Churchill and Storey 1993, 1994), the changes in protein concentration reported as mg/g wet weight in Table 12 must be a reflection of a change in water content of the organs. Thus, it is clear that the water contents of individual organs were strongly defended during whole body dehydration so that only at high levels of dehydration did disruptions of organ water content occur. The data also show that upon rehydration, organ water contents were fully restored by the time that frogs were rehydrated to 20 %. The data in Table 12 also show that all organs lost proportionally about the same amount of water; the ratio of protein concentrations for 40 % dehydrated frogs versus controls spanned a narrow range from 1.4 for brain to 1.6 for liver. These values demonstrate that the rate of water loss from each tissue is remarkably similar, a good indication that this process is highly regulated in *R. sylvatica*. This finding, plus the fact that significant changes in individual organ water contents (that were not calculated in this study) only appeared at high dehydration values, are important to remember in interpreting the remainder of the data on the metabolic effects of whole body dehydration which showed some very specific effects on protein kinases and second messengers only in selected organs and initiated at very low percent dehydration values.

Both this and a previous study (Churchill and Storey, 1993) showed that the percentage of tissue water lost due to dehydration was only a minor fraction of the total tissue weight in all *R. sylvatica* tissues. Liver, brain, heart, and kidney perform specific functions that are crucial for the overall survival of *R. sylvatica*; hence, it is not surprising that the water contents of key internal organs are strongly defended when the animal faces dehydration stress.

In order to minimize the overall tissue water loss, and the potential for tissue damage during dehydration, both frog species must make metabolic compromises. The major metabolic compromise made in anuran species is to reduce the amount of water in the blood and extra organ spaces such as the fluids found in the abdomen and the

bladder (Hillman, 1980). This leads to an increase in blood viscosity, placing a greater burden on the cardiovascular system to deliver oxygen and nutrients to the tissues and organs. Furthermore, as the content of water in the blood is reduced there is a concurrent increase in the concentration of solutes in the blood (Hillman, 1980; Hillman, 1987). The hyperosmolality observed during anuran dehydration can reach levels which are comparable to those observed in frozen *R. sylvatica* (Storey, 1990). Most amphibians can tolerate plasma sodium ion concentrations up to 200 mM, well above the range tolerated by other vertebrates (Hillman, 1988). This is excellent evidence which supports the hypothesis that the development of the freeze-tolerance phenomenon in *R. sylvatica* arose out of the capability of anurans to survive prolonged episodes of dehydration in which elevated solute concentrations are a common occurrence.

Glucose biosynthesis via glycolysis is a major metabolic response during dehydration stress in *R. sylvatica* (Churchill and Storey, 1993). Glycogen breakdown resulted in a rise of 127 $\mu\text{mol/g}$ wet weight glucose in the liver of wood frogs dehydrated to 50 % of total body water lost. Dehydration stimulated a strong increase in glucose content in all other organs that resulted in maximal levels that were up to 300-fold higher than in organs of control frogs (Churchill and Storey, 1993). Glucose levels rose progressively as a function of total body water loss in all *R. sylvatica* organs. These data indicated that the glucose accumulation response to dehydration is virtually the same as the cryoprotectant response of wood frogs to freezing (Storey and Storey, 1984, 1988).

Dehydration stimulated a 3-fold increase in the percentage of free catalytic subunit of PKA in *R. sylvatica* liver (Table 13). A comparable effect was not observed in the liver of the freeze intolerant *R. pipiens* (Table 14). The elevation of liver PKAc in 20 % dehydrated *R. sylvatica* was preceded by a 3.4-fold increase in liver cAMP that occurred by 5 % dehydration and high cAMP was sustained at 10 % dehydration

(Table 15). I hypothesize that this rise in cAMP, along with the rise in the free catalytic subunit of PKA, stimulated the activation of glycogenolysis that is responsible for the production of glucose in the wood frog in response to dehydration. Even though the rise in percent PKAc was not as large during dehydration as was observed for the freezing stress, the response by PKAc to the two stresses were remarkably similar. Thus the accumulation of glucose during dehydration is probably a result of the activation of glycogen phosphorylase as was also seen in *R. sylvatica* in response to a freezing stress (Storey and Mommsen, 1992). The metabolic machinery is in place that allows *R. sylvatica* to respond immediately to produce enough glucose to substantially contribute to the colligative regulation of cell volume during dehydration. However, another potential use for dehydration-induced hyperglycemia could be proposed and this is to provide fermentative fuel to organs when dehydration begins to compromise aerobic metabolism. This advances the argument that key biochemical events that occur in freezing may have evolved from preexisting mechanisms for dealing with the variations in body water content that occur in dehydration in the wood frog. Because *R. sylvatica* responds with a cAMP elevation in liver by 5 % dehydration, well before liver shows any indication of being dehydrated itself, the freeze tolerant frog may have a very sensitive detection system for picking up small changes in water content or cell volume in particular cells. *R. sylvatica* may be able to directly detect minute changes in water content or cell volume in liver tissue itself, but perhaps more likely, probably detects signals coming from a more peripheral site on the body, for example, the skin. The freeze intolerant species, *R. pipiens*, seems to lack this response. The reason may be that the leopard frog likely responds to any dehydration stress by simply seeking water and corrects its water balance by behavioral means. For *R. sylvatica*, on the other hand, water stress is typically associated with freezing and hence immobility, and therefore the species has had to develop a metabolic response to conserve organ water instead.

An interesting observation was the change in *R. sylvatica* muscle PKAc and the lack of change in PKAc content in *R. pipiens* muscle. This suggests that some other function of PKAc activation exists in the muscle of the freeze tolerant frog. Notably in both dehydration (Table 13, 15) and freezing (Table 9,10), *R. sylvatica* muscle PKAc increased under the stress but cAMP did not. Thus, something else must be activating the kinase, perhaps cGMP. In addition, the function of this PKAc activation may not be related to glycogenolysis but might act to achieve some other metabolic response to dehydration/freezing in muscle tissue, perhaps promoting the uptake of blood glucose into muscle tissue under these stresses.

Dehydration also resulted in a large increase in the levels of the second messenger for protein kinase C, IP₃, in the liver of *R. sylvatica* (Table 15). The concentration of this second messenger increased progressively with the rising level of dehydration stress reaching a net 4-fold increase at 40 % dehydration. Kidney was the only other organ to show a dehydration-induced increase in IP₃, a rise of 1.59 fold at 40 % dehydration. Again, the response by IP₃ in the liver to dehydration was similar to that observed in wood frogs as a response to freezing. The concentrations of this second messenger increased significantly over control levels in both cases. This evidence points to a specific role for PKC in the liver response to freezing or dehydration. This role could be in the control of glycogenolysis through regulatory effects on GP activity, or in the regulation of some other aspect of metabolism.

Several interesting patterns emerged in the examination of second messenger levels that suggested a coordinated control of glycogenolysis by both PKA and PKC during dehydration. The pattern of IP₃ accumulation was entirely different from that for cAMP. In *R. sylvatica* liver a rapid increase in cAMP was seen which peaked at 5 % body water lost and then declined (Table 15). This is most likely linked to the rapid initial activation of phosphorylase in order to initiate glucose output from the liver during the early stages of dehydration stress. By contrast, IP₃ levels increased

progressively over the course of dehydration giving the impression that the target of this second messenger was also something that was stimulated progressively as dehydration progressed. Hence IP_3 , and by extension PKC, may be linked with changes in cell volume, either directly stimulated by small changes in liver cell volume as dehydration progresses, or activated by external signals that stimulate liver to respond in an anticipatory or adaptational way to the coming dehydration stress. The effects of IP_3 stimulation could include the regulation of plasma membrane ion channel proteins and actions that are related to the PKAc stimulated activation of glycogenolysis such as enhancing membrane glucose transporter activity to aid in the export of glucose.

PKC might also be directly involved in phosphorylase control. PKAc clearly controls the immediate activation of phosphorylase via the phosphorylation of GP kinase, as it does in all vertebrates that respond to epinephrine or glucagon stimulation while under stress. However, given the pattern of IP_3 accumulation, PKC might be involved in sustaining high levels of glucose output during both freezing and dehydration, and overriding the homeostatic controls on liver glycogen metabolism that should normally keep blood and tissue glucose below about 5 mM. Thus, in mammalian liver, high glucose leads to a reduction in GP_a activity; this is because a glucose- GP_a complex is the preferred substrate for phosphorylase phosphatase which then dephosphorylates the enzyme to convert it back to the inactive b form (Hers, 1976). Thus, high liver glucose concentrations effectively turn off further phosphorylase activity. In freeze-tolerant frogs, however, phosphorylase clearly remains active while cellular glucose concentrations climb to 200 mM or more. Perhaps the role of PKC could be to help keep net GP activity high, even in the presence of high levels of its allosteric inhibitor, glucose. This might be accomplished by making one or more additional phosphorylation modifications to the enzymes involved, perhaps to change GP_a affinity for glucose so that GP_a -glucose complex formation is reduced

2

PM-7 3 1/4"x4" PHOTOGRAPHIC MICROCOPY TARGET
NBS 1010a ANSI/ISO #2 EQUIVALENT

PRECISIONSM RESOLUTION TARGETS

(despite the very high glucose concentrations) or to change the affinity of phosphorylase phosphatase for the GP_a-glucose complex.

Thus, if two protein kinases, PKA and PKC, are involved in liver glycogenolysis regulation then the mechanisms would be in place to allow differential responses by liver glycogenolysis to the glucose output demands of normal life (e.g. in response to exercise, hypoxia, or starvation) and the extreme hyperglycemia demanded for freezing-induced cryoprotectant synthesis. The normal glycogenolytic responses by vertebrate liver to catecholamine or glucagon stimulation would be mediated via PKA (which also appears to stimulate the initial phosphorylase activation during freezing) whereas the sustained activation of glycogenolysis needed to accumulate high cryoprotectant levels could be maintained by PKC mediation.

Interestingly, data collected over a time course of dehydration showed that liver glucose output was low up to about 20 % dehydration, and then glucose (and glucose-6-phosphate) levels rose rapidly with more severe dehydration (Churchill and Storey, 1994). This pattern closely matches the pattern of IP₃ changes seen in the present study. Further studies involving the distribution/activation of PKC in liver over a freezing and dehydration time course could give great insights into how cryoprotectant metabolism is controlled by PKAc and PKC, a dual protein kinase cascade mechanism.

No increases in PKAc content occurred throughout the entire dehydration course in *R. pipiens* liver or muscle (Table 14). It seems reasonable to suggest then, that no significant contribution was made by the PKA enzyme cascade system to the dehydration-induced output of glucose via liver glycogenolysis that occurred in this freeze intolerant anuran (Churchill and Storey, 1994). A different protein kinase cascade system, perhaps that of PKC, may be responsible for the rapid hyperglycemia, based on liver glycogenolysis, in response to dehydration in this semi-aquatic frog. Again, PKC activation could stem from direct and minute changes in liver cell volume which cause a translocation of the enzyme to the cell membrane or from external

signals (hormones) that come in from peripheral sites that pick up dehydration volume changes. However, further studies are needed in order to test these hypotheses.

There may be more than one cellular signal in response to dehydration that causes a release of hormones along with a simultaneous biochemical activation of signal transduction cascades in *R. sylvatica* liver. The first signal may be a biochemical response to an increase in evaporative water loss across the skin. The increase in cAMP and PKAc observed in the early stages of dehydration (Table 13 and Table 15) was most likely a response to a β -adrenergic signal that was released due to evaporative water loss. The PKA signal transduction cascade must be responsible for the initial mobilization of liver glycogen into glucose under these conditions. Hence, it appears very likely that the increases in wood frog liver cAMP and PKAc are linked to the activation of glycogenolysis in this organ. Dehydration may be detected at some peripheral site, possibly as a cell volume change in skin cells, and this may, in turn, send a signal that triggers liver glycogenolysis during dehydration.

The second signal could be a biochemical response to cell volume changes, or actual tissue water loss from liver itself. The increase in IP₃ in the later stages of dehydration may be an indication that the PKC signal transduction cascade is responding to a hormone signal that is released in response to a decrease in cell volume. Changes in liver cell volumes during freezing in *R. sylvatica* have been well documented (Storey et al., 1992), and may take place in this organ in response to a dehydration stress as well. The stimulus for the activation of glycogenolysis in liver during dehydration may be a direct change in liver cell volume. This causes an activation of the PKC signal transduction cascade, sustaining glucose biosynthesis during periods of severe dehydration (greater than 20 % tissue water loss).

These data indicate a synergistic cooperation between the PKA and PKC signal transduction cascades which allows for a continuous production of glucose throughout a dehydrational stress. During short term dehydration PKA is responsible for the

activation of glycolysis; if dehydration conditions persist it is the PKC signal transduction cascade that takes over the phosphorylation of GP, resulting in a sustained activation of glycolysis and glucose biosynthesis in cases of severe dehydration.

Data from this study show that the mechanism of glucose synthesis and clearance could be hormone-stimulated changes in the activity of cAMP-dependent protein kinase along with protein kinase C. Further investigations of the actions of protein kinase C are required to fully elucidate its role in glucose biosynthesis in response to freezing and/or dehydration stresses in wood frogs.

In summary, freeze tolerance and dehydration tolerance are intimately linked in the wood frog *R. sylvatica*. Key metabolic events that occur in a freezing episode such as increases in second messenger concentrations (cAMP and IP₃), the mobilization of the cAMP-dependent protein kinase cascade (and perhaps that of PKC), along with the production of glucose as a cryoprotectant, are also elicited as a response to dehydration stress. Therefore, it is very likely that the metabolic events that occur in natural freeze tolerance in amphibians like *R. sylvatica* may have evolved from the pre-existing capabilities that these anurans possessed to survive the wide variations in body water content that occur in a dehydration stress. Initiation of cryoprotectant production in *R. sylvatica* by PKAc and/or PKC must be under direct hormonal control, a classic metabolic response that is known to occur with changes in cell volume.

CHAPTER #5

SIGNAL TRANSDUCTION, SECOND MESSENGER, AND PROTEIN KINASE RESPONSES DURING ANOXIA STRESS IN TWO ANURANS

INTRODUCTION

Extracellular freezing imposes an ischemic and anoxic state on the cells of freeze tolerant animals. Thus, all cells must have sufficient endogenous fuel and energy reserves to allow them to endure a long freezing episode. Cells must also tolerate the accumulation of anaerobic metabolic end products, and any acidification of the cytosol that these cause, until such time as thawing permits a re-establishment of circulation (Storey and Storey, 1986). Energy metabolism in the frozen state relies on anaerobic metabolism, specifically on the glycolytic pathway. In freeze tolerant frogs, freezing activates glycogen phosphorylase in liver for the purpose of cryoprotectant production in the form of glucose (Mommensen and Storey, 1992). This sugar is delivered to other organs, where in addition to its use as a cryoprotectant, some small percentage may also be used as a fermentative substrate, along with endogenous organ reserves of glycogen, to sustain energy metabolism in the frozen state (net metabolic rate is very low due to the subzero temperatures). Other responses to the ischemic/anoxic state of freezing may include an overall metabolic rate depression in organs which, along with subzero body temperatures, may ensure animal survival over long periods of time in the frozen state (Storey and Storey, 1988).

While frozen, the wood frog undergoes cardiac arrest and all blood flow to the organs and tissues ceases, causing anaerobic conditions to prevail within the organism. Thus, the tolerance to anoxic and ischemic conditions may be another key factor that enables the wood frog to survive in the frozen state. Anurans are able to survive 2 d of anoxia under laboratory conditions (this study) and are able to survive for months in nature in environments where severely hypoxic and sometimes anoxic conditions prevail (Christiansen and Penny, 1973). Semi-terrestrial frogs, such as *R. pipiens*, encounter periods of prolonged anoxia only during the winter months when the body temperature of these animals is just a few degrees above freezing, approximately 3-5°C

(Christiansen and Penny, 1973). At this time leopard frogs remain in a lethargic state on or in the mud at the bottom of ponds for periods of up to five months. (Poczopko, 1959). The oxygen tension in the environment surrounding the animal during this period can fall to very low levels, for once the surface is frozen over, there is no aeration of the water to replenish oxygen consumed by living organisms (Hutchinson and Dady, 1964).

It is likely that the anoxia tolerance displayed by *R. sylvatica* may be a key component of its ability to survive freezing. Thus, the development of the freeze tolerant strategy in *R. sylvatica* may be linked to a natural tolerance of anoxia. The study described in this chapter addressed the influences of anoxia on the activities of protein kinases and the levels of second messengers in a freeze tolerant frog, *R. sylvatica*, compared with a freeze intolerant species, *R. pipiens*. The two species would typically experience anoxia stress in nature under entirely different physiological scenarios. *R. sylvatica* undergoes anoxia stress as part of the ischemia imposed by freezing and would be unlikely to encounter oxygen limitation in other situations. *R. pipiens*, on the other hand, overwinters at the bottom of ponds and as winter progresses must endure an increasingly hypoxic environment, becoming even fully anoxic, as oxygen is gradually consumed in ice-locked waters. Aquatic-hibernating frogs are well adapted for oxygen uptake across their skin and maintaining aerobic metabolism even under severely hypoxic conditions but, unlike the freeze tolerant species, do not endure extended anoxia well. This has been linked to the rapid depletion of liver glycogen supply and accumulation of lactic acid in their tissues in anoxia (Christiansen and Penny, 1973).

The effects of whole animal anoxia exposure on the activities and the percentage of free catalytic subunit of PKA and the concentration of the second messenger for PKC, IP₃, were quantified in several tissues of *R. sylvatica* and *R. pipiens*. The present study aimed to analyze the protein kinase and second messenger responses to anoxia

exposure in these two ranid species. The goals included determination of: (1) the similarities and/or differences in the metabolic responses to anoxia between these two ranid species, (2) whether these differences represent adaptive strategies that could promote anoxia/ischemia resistance during freezing in *R. sylvatica*, and (3) whether various known metabolic responses/adaptation observed during freezing exposure in *R. sylvatica* are, in fact, linked to the survival of the anoxia/ischemia imposed by freezing.

MATERIALS AND METHODS

Preparation of Experimental Animals

R. sylvatica and *R. pipiens* were collected, held, and given anoxia exposure as stated in chapter 2. Timing of anoxia exposure was taken from the point at which each bottle was sealed and placed in the incubator at 5°C. Times for anoxia exposure were ½ h, 1 h, 4 h, and 2 days. Control animals were taken from incubators held at 5°C and sacrificed by double pithing. Tissues used included liver, leg skeletal muscle, brain, heart, and kidney.

PKA and IP₃ Assays

The activity of cAMP-dependent protein kinase and the percentage of the free catalytic subunit in anoxia-exposed *R. sylvatica* and *R. pipiens* were assayed by a radioactive assay that measured the incorporation of radiolabelled ³²P-ATP onto the phosphate-accepting artificial substrate 'kemptide' as described in chapter 2. IP₃ was measured in frog tissue samples as described in chapter 2.

RESULTS

Effect of Anoxia Exposure on the Percent of PKAc in *R. sylvatica* and *R. pipiens* Tissues

Both *R. sylvatica* and *R. pipiens* showed a good ability to tolerate anoxia (exposure to an N₂ gas atmosphere) at 5°C. Initial studies showed that frogs readily endured 5 d under these conditions and all animals used in the present 2 d time course experiment were alive at all sampling times.

The total activities of PKA in *R. sylvatica* tissues were not significantly different from those measured for control tissues in the experimental freezing studies (see Chapter 3). Furthermore, total PKA activity was not affected by anoxia exposure in any *R. sylvatica* and *R. pipiens* tissue tested.

Table 16 shows the effect of anoxia exposure time on the percentage of PKA present as the free catalytic subunit in five tissues of *R. sylvatica*. The amount of PKAc was never higher than 30% of the total in any of the tissues. PKAc contents of four organs, including liver, were unaffected by anoxia exposure. In brain, however, PKAc rose significantly after the longer anoxia exposure times, 4 h and 2 d; PKAc in these animals was 1.6-fold higher than in controls.

Table 17 shows the effect of different lengths of anoxia exposure on the percentage of PKA present as the free catalytic subunit in four tissues of *R. pipiens*. Each tissue showed a different response. PKAc content of *R. pipiens* liver was unaffected by short anoxic exposures but increased nearly 2-fold after 2 d anoxic exposure. The highest overall percentages of PKAc were in leg muscle with values of about 32 % in 1 h, 4 h, and 2 d anoxic frogs representing about a 1.45-fold increase in PKAc content in anoxia. By contrast the percent PKAc in brain decreased during short anoxia exposures, falling to 8.7, 12.6, and 11.3% after ½ h, 1 h, and 4 h of anoxia, respectively. After 2 d of anoxia exposure, however, brain PKAc content had returned

Table 16. Effect of anoxia exposure at 5°C on the amount of active PKA (expressed as a percentage) present as the free catalytic subunit in *Rana sylvatica* tissues.

Condition		% Free Catalytic PKA			
		Liver	Muscle	Brain	Heart
Control		6.8 ± 0.5	24.4 ± 3.2	13.0 ± 0.7	20.5 ± 1.4
Anoxic	½ h	5.2 ± 0.6	28.2 ± 4.0	14.0 ± 0.7	18.7 ± 0.9
	1 h	4.9 ± 1.0	28.8 ± 2.8	17.5 ± 0.9	19.7 ± 0.7
	4 h	7.9 ± 1.7	28.7 ± 3.8	20.6 ± 1.2 ^a	20.9 ± 1.1
	2 d	8.4 ± 0.3	26.8 ± 3.0	21.6 ± 0.4 ^a	19.8 ± 1.7
					9.9 ± 0.5
					11.7 ± 0.5
					10.9 ± 0.3
					10.9 ± 0.7
					10.4 ± 0.6

Data are means ± S.E.M., n = 4 animals.

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, P < 0.05.

Table 17. Effect of anoxia exposure at 5 °C on the amount of active PKA (expressed as a percentage) present as the free catalytic subunit in *Rana pipiens* tissues.

Condition	% Free Catalytic PKA			
	Liver	Muscle	Brain	Heart
Control	8.1 ± 1.9	22.5 ± 2.0	16.5 ± 0.9	5.5 ± 1.2
Anoxic	½ h	11.1 ± 2.1	21.5 ± 1.2	8.7 ± 0.5 ^a
	1 h	7.9 ± 2.0	31.6 ± 4.7 ^a	12.6 ± 3.2 ^a
	4 h	6.7 ± 0.5	32.1 ± 3.1 ^a	11.3 ± 0.7 ^a
	2 d	15.6 ± 1.7 ^a	32.0 ± 4.2 ^a	15.3 ± 0.8
				8.2 ± 1.4

Data are means ± S.E.M., n = 4 animals. No PKA activity was found in *R. pipiens* kidney.

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, P<0.05.

to control values. Heart PKAc content was unaffected by experimental anoxia exposure.

Effect of Anoxia Exposure on Tissue IP₃ Concentrations in *R. sylvatica* and *R. pipiens*

Because PKAc content was unaffected by anoxia exposure in all but one tissue of *R. sylvatica*, it was decided that cAMP levels would not be measured and that tissue samples would be used, instead, to analyze the effects of anoxia on protein kinase C via analysis of IP₃ levels. Direct evidence of an activation of PKC under anoxia can be derived from measured increases in the concentrations of its second messenger IP₃.

Table 18 shows the effect of anoxia exposure on IP₃ concentration in four tissues of *R. sylvatica*. In liver IP₃ concentration rose very rapidly, increasing by 7-fold after only ½ h of anoxic stress. Subsequently, liver IP₃ levels declined progressively after 1 and 4 h but remained about 2-fold higher than control values even after 2 d of anoxia. By contrast, IP₃ concentration in skeletal muscle did not change over the short term but fell by 40 % after 2 d of anoxia exposure. Anoxia exposure had no effect on IP₃ in *R. sylvatica* brain and in heart IP₃ increased by 30 % after 1 h anoxia and remained elevated over the remainder of the anoxia exposure.

Table 19 shows the comparable effects of anoxia exposure on the concentrations of IP₃ in *R. pipiens* tissues. As in the wood frog, anoxia stimulated a rapid increase in liver IP₃ concentration, levels rising by 3-fold after only ½ h of anoxic exposure. Liver IP₃ remained at this level through 4 h of anoxia exposure but had declined slightly after 2 d of anoxia. By contrast, and again like in *R. sylvatica*, skeletal muscle IP₃ was unaltered by short term anoxia exposure but dropped by 55 % in the 2 d anoxic animals. Again, there was no change in brain IP₃ content over the course of anoxia and in heart, IP₃ showed a transient increase of 30 % after ½ h anoxia but then IP₃ returned to control levels.

Table 18. Effect of anoxia exposure at 5°C on IP₃ concentrations in *Rana sylvatica* tissues.

Condition		IP ₃ , pmol/g wet weight		
		Liver	Muscle	Brain
Control		1406 ± 164	715 ± 32	12816 ± 1205
Anoxia	½ h	2376 ± 99 ^a	775 ± 4	13888 ± 2688
	1 h	1250 ± 111	712 ± 37	9766 ± 1507
	4 h	680 ± 122 ^a	737 ± 37	12716 ± 1473
	2 d	667 ± 172 ^a	460 ± 17 ^a	6642 ± 1480
				1944 ± 86
				2084 ± 114
				2520 ± 174 ^a
				2473 ± 182 ^a
				2794 ± 70 ^a

Data are means ± S.E.M., n = 4 animals. The control value for liver IP₃ was taken from Table 4.

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, P < 0.05.

Table 19. Effect of anoxia exposure at 5°C on IP₃ concentrations in *Rana pipiens* tissues.

Condition		IP ₃ , pmol/g wet weight		
		Liver	Muscle	Brain
Control		375 ± 24.6	1405 ± 20.4	7493 ± 441
Anoxic	½ h	1136 ± 21.2 ^a	1478 ± 28.8	6985 ± 326
	1 h	1179 ± 57.3 ^a	1581 ± 60.7	9147 ± 199
	4 h	1013 ± 43.1 ^a	1525 ± 15.5	7053 ± 252
	2 d	911 ± 51.1 ^a	630 ± 43.3 ^a	5824 ± 1067
				1980 ± 55.2
				2586 ± 95.3 ^a
				1847 ± 98.7
				2092 ± 73.8
				1568 ± 117

Data are means ± S.E.M., n = 4 animals.

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, P < 0.05.

DISCUSSION

Metabolic rate depression and accumulation of anaerobic end products are well known characteristics that many animal species exhibit as responses to severe hypoxic or anoxic conditions (Storey and Storey, 1990). Anoxia tolerant species greatly suppress the rates of energy-utilizing reactions in order to extend the time that they can survive based on anaerobically-generated ATP alone. Freshwater turtles are a prime example of ectotherms that can survive extended periods without oxygen (Belkin, 1963). Species from the genera *Chrysemys* and *Pseudemys* can survive anoxia utilizing anaerobic glycolysis for several months at 3°C, reducing their metabolism far more than would be expected by the effects of temperature alone (Keiver and Hochachka, 1991; Lutz et al, 1984). Metabolic depression is the key to extended life when oxygen is limiting; this common element probably supports the survival of *R. sylvatica* throughout a freezing stress and *R. pipiens* throughout the severely hypoxic conditions that are prevalent at the bottom of ice covered ponds.

In freeze tolerant frogs metabolism becomes restricted by the environmental constraints themselves, low water availability and low temperature. Therefore the cessation of metabolism in the frozen state is imposed on the organism as a consequence of the stress of freezing itself. Conversely, the metabolic depression seen in *R. pipiens* during anoxia stress, is used as a survival strategy to endure periods in which oxygen is limiting at the bottom of ponds during winter hibernation (Storey and Storey, 1990). The strategy of anoxia tolerance exhibited in *R. pipiens* saves energy in the form of ATP equivalents, carbohydrates, and lipids, until such a time as normoxic environmental conditions return that allow for normal cell functions and metabolism to be restored (Christiansen and Penny, 1973).

Research to this point has identified three major mechanisms that control the entrance into, and sustaining of, the reduced metabolism seen in organisms under

anoxic stress. These are: (1) the post translational covalent modification of enzymes by means of protein phosphorylation or dephosphorylation reactions, controlled by protein kinases and protein phosphatases, (2) fructose-2,6-bisphosphate regulation of the rate-limiting glycolytic enzyme PFK, and (3) the reduction of membrane ion transport and ion channel function via protein phosphorylation (Storey and Storey, 1990; Storey, 1994).

The key enzymes that are controlled by protein kinases during post-translational modification reactions are glycogen phosphorylase (GP), phosphofructo-1-kinase (PFK), 6-phosphofructo-2-kinase (PFK-II), pyruvate kinase (PK), and pyruvate dehydrogenase (PDH). In the most well studied anoxia tolerant invertebrate the channeled whelk *Busycotypus canaliculatum*, there is an active dephosphorylation of GP and an active phosphorylation of PFK, PFK-II, and PK under anoxic conditions (Plaxton and Storey 1984; Storey 1988). These enzymes are all known to be strongly inhibited by covalent modification during many environmental stresses, including anoxia. Covalent modification is an important biochemical phenomenon that all anoxia tolerant animals utilize to combat this stress (Storey and Storey, 1990). The results of this study indicate that both PKA and PKC must be involved in the regulation of glycogenolysis via covalent modification. These results warrant further investigations in order to elucidate the exact target protein substrates for PKAc and PKC phosphorylation and to determine their specific function(s) in each organ during anoxia exposure in *R. pipiens* and *R. sylvatica*.

Table 16 shows that there was an activation of PKA by cAMP that increased the levels of PKAc in the brain of *R. sylvatica* during anoxia. Anoxic stress seemed to mimic the stress of freezing in this regard. However, there was no increase in the amount of active catalytic subunit of PKA in any other tissue, including liver. This was in direct contrast to the marked increase in PKAc observed in *R. sylvatica* liver, that was responsible for the initiation of cryoprotectant production in this tissue under both

freezing and dehydration stress. No large volume changes can occur in the cells of frogs under hypoxic or anoxic stress; therefore, it seems reasonable to conclude from the above data, that only environmental stresses that induce changes in cell volume, such as dehydration, invoke a similar protein kinase A response to that witnessed in *R. sylvatica* during freezing.

Furthermore, the percentage of PKAc in *R. sylvatica* brain increased in the later stages of anoxia, at times longer than 4 h (Table 16). This may be indicative of an activation of GP via the PKA cascade in this organ, rather than the suppression of this enzyme, via dephosphorylation, that is known to occur in other vertebrates that exhibit anoxia tolerance (Brooks and Storey, 1988). It is critical that the glucose used by the brain as an energy source to survive the early stages of anoxic exposure (and derived from endogenous glycogen) be replaced if the animal is to survive an anoxic stress for extended periods of time. By maintaining elevated glucose concentrations anoxia stressed frogs, like similarly stressed turtles, can maintain ionic, electrical, and mitochondrial integrity in the brain in order to survive prolonged anoxia (Lutz et al., 1984).

In contrast to *R. sylvatica*, there was a very large increase in liver and muscle PKAc throughout the entire period of anoxia exposure in *R. pipiens* (Table 17). The increase in the percentage of active catalytic subunit of *R. pipiens* liver PKA early in anoxic exposure suggests that PKAc may initiate the limited production of glucose from liver glycogen seen in leopard frogs as a metabolic response to this stress. This supports the fact that there is an active depletion of liver glycogen, followed by a rise in blood and lymph glucose, and finally a build up the anaerobic end product lactate in the tissues of *R. pipiens* upon exposure to anoxia (Christiansen and Penny, 1973). Glucose mobilization from the liver in *R. pipiens* permits organs lacking this metabolic substrate to take up glucose from the blood, once plasma concentrations of this sugar have increased. Thus, unlike *R. sylvatica*, this anuran may promote the typical covalent

modification of biochemical pathways via phosphorylation/dephosphorylation reactions in response to anoxia. These same enzyme modifications are known to occur in turtles and in the channeled whelk, two premier facultative anaerobic animals (Kelly and Storey, 1988; Plaxton and Storey, 1984; Kiever and Hochachka, 1991).

There may be several reasons for the increased levels in *R. pipiens* liver PKAc after 2 d of anoxia exposure (Table 17). If PKAc did mediate the phosphorylations of target proteins that help create metabolic depression, perhaps sustained increases in PKAc levels were required to maintain these conditions. Furthermore, given the low body temperature of these anurans it may have taken considerable time for the frogs to use up endogenous oxygen and go through the stages of hypoxia and then anoxia.

A general state of metabolic arrest can prevail even though liver glycogen breakdown is relatively activated, compared with the aerobic situation, due to the need to supply blood glucose as an anaerobic substrate to all tissues. The inactivation via phosphorylation of glycolytic, TCA cycle, and electron transport enzymes is critical in maintaining a depressed metabolic state under environmental conditions where oxygen is limiting. The rise in the levels of liver and muscle PKAc indicate that the activation of the PKA phosphorylation cascade is a component of the anoxia tolerance strategy of *R. pipiens*.

Table 17 displays the fact that there was a large decrease in the level of brain PKAc in leopard frogs after 1 h of anoxia exposure, the opposite response to that observed in *R. sylvatica*. This may indicate that there is a decrease in the rate of brain glycogenolysis in order to save fuel in the form of glucose and maintain ionic, electrical, and mitochondrial integrity in this organ while the animal depresses its metabolism throughout an entire anoxic exposure. Turtles, as well as mammals, show a rapid depletion of brain glucose in response to an anoxic exposure (Lutz et al., 1984). Export of glucose from the liver to other organs during anoxia restores the concentrations of glucose in the brain of these organisms. Export of glucose to the

brain tissue of *R. pipiens* is critical for the proper functioning of this organ, maximizing the survival time of individual frogs when environmental oxygen conditions are unfavorable.

Biochemical responses surrounding the activation of PKC by IP₃ that occurred during anoxia exposure were identical in these two ranid species (Tables 18 and 19). Anoxia exposure seemed to initiate the mobilization of the second messenger for protein kinase C, IP₃, in both frogs. Additionally, the rise in liver IP₃ concentrations during anoxia was a similar response to that observed during freezing, or dehydration exposure in *R. sylvatica*. The increased levels of this second messenger is a firm indication of cellular PKC activation. An 8-fold and 3-fold increase in IP₃ concentration was observed in the liver of *R. sylvatica* and *R. pipiens*, respectively. In fact, liver IP₃ levels remained elevated throughout the entire period of anoxia exposure (Tables 18 and 19). This preliminary evidence points to a role for the activation of PKC that may support glucose production by inhibiting the enzyme responsible for the synthesis of glycogen from glucose, glycogen synthase (GS). Recent accumulated evidence has shown that PKC, along with a number of other protein kinases, inhibits GS by phosphorylating this gluconeogenic enzyme at multiple phosphorylation sites (Roach, 1990). Accumulated liver glucose could then be shunted to other tissues and organs to sustain anaerobic glycolysis during metabolic arrest, as amphibians encounter prolonged anoxia during the winter months.

Cardiac IP₃ levels increased in response to an anoxic stress in both species (Table 18 and 19). The release of this second messenger, as in liver, may denote a stimulation of PKC in this organ in response to declining pO₂ levels in the blood. Endogenous glycogen reserves in heart are catabolized to produce glucose via the activation of the glycolytic pathway in cardiac muscle. Evidence for this was gathered from *R. pipiens* exposed to 96 h of anoxia. Cardiac glycogen reserves in this anuran fell dramatically from 186.9 $\mu\text{mol/g}$ wet wt to 3.5 $\mu\text{mol/g}$ wet wt, or 1.9% of control

glycogen levels, over this time period (Christiansen and Penny, 1973). Therefore, the activation of cardiac glycolysis may be critical for the maintenance of proper heart function during anoxia in all anurans. Studies involving the enzymatic analysis of PKC over a time course of anoxia exposure both in liver and cardiac tissues from several more anuran species are necessary to confirm this hypothesis.

The present data was in agreement with other studies involving the activation of glycolysis as a first initial response to declining oxygen tension (Storey, 1993). There is always an activation of protein kinase enzymes followed by an increase in glycolytic metabolism as an initial response to environmental stresses such as freezing exposure, dehydration, and anoxia. This strategy is used in an attempt to maintain a constant rate of ATP turnover (maintain energy charge) by an increase in the output of glycolytic ATP. When any of these stresses persist, for example if hypoxia becomes severe, the increased output of glycolytic ATP, or activation of the glycolytic pathway, is suspended. Anoxia tolerant animals opt for a strategy of metabolic arrest in which an inhibition of glycolysis is implemented as anoxic conditions persist in the environment. In this study there was a decline in the levels of IP₃ after 4 h of anoxia in all tissues of the 2 anurans, except in *R. sylvatica* heart (Table 18 and 19). This evidence along with the fact that PKAc did not increase during anoxia, reveals a general pattern of glycolytic inhibition in the later stages of anoxia, greater than 4 h, after an initial activation of this pathway by PKAc and/or PKC signal transduction cascade systems.

It appears from the results of this study that: (1) the stress of anoxia on *R. sylvatica* and *R. pipiens* caused a rise in liver IP₃ concentration in both anurans, a similar response to that observed in the liver of the wood frog during dehydration and freezing, (2) PKC may have initiated an activation of liver glycogenolysis during the early stages of anoxia in these two frog species, (3) the PKAc response to anoxia in *R. sylvatica* was the opposite to that of *R. pipiens*, and (4) in *R. sylvatica*, anoxic stress did not cause a stimulation of PKAc release (activation), especially in liver tissue, as

was observed during a freezing event or under conditions that promoted the active dehydration of wood frogs. Data obtained for anoxic *R. sylvatica* indicate that freezing induced anoxia was not responsible for initiating the major metabolic effects observed during a freezing episode, specifically the responses of PKAc. Only environmental stresses that cause a volume regulated response (freezing and dehydration) elicit the activation of protein kinase enzymes that support the controlled and highly regulated metabolic event of cryoprotectant production in freeze stressed *R. sylvatica*.

CHAPTER #6

PURIFICATION OF THE FREE CATALYTIC SUBUNIT OF CYCLIC AMP-DEPENDENT PROTEIN KINASE A FROM *Rana sylvatica* LIVER

INTRODUCTION

One of the most successful freeze-tolerant vertebrate species is the wood frog *R. sylvatica*. The synthesis of glucose from liver glycogen reserves is a well known biochemical process in many vertebrate organisms. *R. sylvatica* has refined this process so that it is able to mobilize its glycogen stores almost immediately after an external inoculative freezing stress. Glucose is the cryoprotectant of choice for *R. sylvatica* and is found in every major tissue and organ in the frozen animal.

Glucose output from *R. sylvatica* liver is due to an increase in (1) the active form of glycogen phosphorylase, (2) the total amount of glycogen phosphorylase activity, and (3) the total amount of liver glycogen. Cryoprotectant synthesis may be controlled by a hormone signal but although glycogenolysis is well known to be under hormonal control in vertebrates it is not yet known which hormone(s) may play a role in stimulating cryoprotectant output (Mommensen and Storey, 1992). However, β -adrenergic receptors (leading to changes in cAMP) in liver appear to be important since administration of the inhibitor, propranolol, reduces glucose production during freezing whereas the α -adrenergic antagonist does not (Storey and Storey, 1992b). Hence, catecholamines may be the involved in the regulation of cryoprotectant synthesis. The inhibitory effects of propranolol on cryoprotectant synthesis, the phosphorylation activation of glycogen phosphorylase (typically mediated by cAMP-dependent protein kinase in vertebrate liver) in response to freezing exposure (Storey and Storey, 1992), and the changes in PKAc content and cAMP levels during freezing documented in this thesis (chapter 3), all support the important role of cyclic AMP-dependent protein kinase in mediating cryoprotectant synthesis. Based on this information, a study of the properties of the catalytic subunit of cAMP-dependent protein kinase from frog liver was warranted.

The studies reported in this chapter relate information on a novel purification procedure for cAMP-dependent protein kinase from *R. sylvatica* liver. Kinetic studies provide details of how the wood frog liver enzyme functions and are related to the specific role of PKAc in activating liver glycogenolysis for cryoprotectant synthesis.

MATERIALS AND METHODS

Preparation of Experimental Animals and PKAc Assay

R. sylvatica and *R. pipiens* were collected as stated in Chapter 2 and were held on damp sphagnum moss at 5°C. Studies were conducted with tissue from control animals only; PKAc enzyme purification used liver from *R. sylvatica* whereas tissue specific isozyme studies were completed using liver, brain, and muscle from both *R. sylvatica* and *R. pipiens*. PKAc activity was assayed as described in chapter 2.

Purification of *R. sylvatica* Liver PKA Catalytic Subunit

A novel purification scheme was developed for the active free catalytic subunit of *R. sylvatica* liver PKA that involved the following steps.

Step 1. Frozen liver (2 g) was homogenized 1:3 w/v in Buffer A (10 mM KP_i , 2 mM 2-mercaptoethanol, 1 mM EDTA, 1 mM cAMP, 10 % v/v glycerol, pH 6.8) with 1 mM PMSF added immediately prior to homogenization with a Polytron PT 10 homogenizer. The addition of cAMP to the homogenization buffer was chosen to ensure that the *in vitro* concentration of cAMP in each homogenate reached a saturating level. This allowed 100 % of the free catalytic subunit, PKAc, to be released from the holoenzyme, PKA, for purification purposes. The incubated homogenate was centrifuged at 27,000 g for 30 min at 5°C in a Sorval RC-5B refrigerated centrifuge and the supernatant was saved.

Step 2. The supernatant was then loaded onto a hydroxylapatite column (4 cm x 1 cm) equilibrated in buffer B (10 mM KP_i , 2 mM 2-mercaptoethanol, 1 mM EDTA, 10 % v/v glycerol, pH 6.8). The column was washed twice with several volumes (10 ml) of buffer B. PKAc was then eluted in a single peak using a linear gradient of 0-500 mM potassium phosphate in buffer B. One ml fractions were collected and assayed as described in chapter 2 and active fractions (5 ml) were pooled and dialyzed for 2 h against buffer B.

Step 3. Dialyzed PKAc was then layered onto a protamine agarose column (4 x 1 cm), pre-equilibrated in buffer B. After two washes of 10 ml buffer B each, liver PKC was eluted using a linear gradient of 0-500 mM phosphate in buffer B. The enzyme eluted in a single peak and fractions were collected and assayed for PKAc activity. Fractions containing PKAc activity were pooled and concentrated using Centricons (Amicon) to a final volume of 1 ml.

Step 4. The concentrated enzyme was then loaded onto a Bio-Gel P-200 molecular weight column (40 x 0.5 cm) which was previously equilibrated in buffer B containing 0.25 % v/v Triton X-100 at pH 6.8. Fractions (0.5 ml) were collected and assayed for activity. Fractions containing PKAc activity were pooled (final volume 5 ml) and used as the enzyme source for kinetic studies.

Subunit and Native Molecular Weight Determinations

The subunit molecular weight of purified PKAc was determined by SDS polyacrylamide gel electrophoresis on a 10 % acrylamide gel (modified from Laemmli, 1970) and using a plot of K_p vs. log M.W. of the protein standards. Gels were stained with silver nitrate based on a procedure developed by Nielson and Brown, 1984. The native molecular weight of *R. sylvatica* PKAc was determined by gel filtration on a Bio-Rad P-200 (0.8 x 15 cm) column. The column was equilibrated and eluted in buffer B containing 0.25 % v/v Triton X-100. Purified PKAc (100 μ l) was loaded onto

the column, and 630 μ l fractions were collected and assayed for activity. Molecular weight markers were detected with spectrophotometric enzyme activity assays at 340 nm (hexokinase, enolase, α -glycerophosphate dehydrogenase, triose phosphate isomerase) or by absorbance at 280 nm (ovalbumin, chymotrypsinogen A).

Kinetic Characterization of PKAc

All kinetic studies involving frog liver PKAc were performed at 22°C and 5°C to determine whether temperature had effects on enzyme function. Substrate affinity constants (K_m values) were determined for kemptide and Mg-ATP. A magnesium curve was completed to determine whether the PKAc reaction was dependent on the free concentration of this divalent cation. Inhibitor constants (I_{50} values) were determined for various salts including KCl, KNO₃, NaCl, NaF, NH₄Cl, and (NH₄)₂SO₄. Additionally, the effects of two specific inhibitors of mammalian protein kinase A (PKA_i, and H89) were determined.

An Arrhenius plot of log relative activity versus $1/T \times 10^{-3}$ (°K) was completed under V_{max} assay conditions. Reaction rates were monitored from 1°C to 35°C at intervals of 5°C for an $n = 4$ for each temperature point on three separate purified preparations of frog liver PKAc.

For pH studies, the substrate concentrations for free Mg^{2+} , ATP, and all ionic species such as K^+ , Cl^- , PO_4^{3-} , EDTA needed to determine the correct pH-temperature relationship for pH curves were calculated using the BAD program developed by Brooks and Storey (1992). The program allows one to enter the desired initial concentrations of reactants and substrates for any particular enzyme assay and set the desired temperature and pH values needed for the reaction. The program then calculates the actual concentrations of stock reactants (in mM values) that must be added to the reaction in order to attain the desired set reaction pH and reaction conditions. This procedure was followed to monitor the effects of pH on *R. sylvatica*

liver PKAc activity at both 22°C and 5°C. In this case, 2.08 mM Mg^{2+} , 0.237 mM ATP, 80 mM potassium ion (K^+), 50 mM Cl^- , 20 mM PO_4^{3-} , 1 mM EDTA, and the extra ionic strength of the two buffers 20 mM TRIS and 20 mM Imidazole were the initial starting values entered into the program. The 20 mM phosphate, 20 mM TRIS, and 20 mM Imidazole buffer had a large combined range of pKa values (combined buffering capacity), which permitted the testing of a series of set pH values, ranging from 5-8.5.

Classification of PKAc Isozymes Using Hydroxylapatite Column Chromatography

To determine whether different tissue specific isozymes of the catalytic subunit of PKA existed, control tissue samples of liver, muscle, and brain from both *R. sylvatica* and *R. pipiens* were homogenized 1:6 (w/v) in PEM buffer (10 mM KP_i , 1 mM EDTA, 2 mM β -mercaptoethanol) plus 1 mM cAMP. An aliquot (2 ml) of each homogenate was then loaded onto separate 0.5 cm x 2 cm hydroxylapatite columns pre-equilibrated in PEM buffer without cAMP. Columns were washed with 10 ml of PEM buffer, and PKAc activity was then eluted with a salt gradient of 0.01-1.0 M KP_i , pH 6.8.

RESULTS

Purification of *R. sylvatica* Liver PKAc

The purification scheme developed for frog liver PKAc is summarized in Table 20. The purification involved an initial hydroxylapatite chromatography step, the enzyme eluted in a single peak at approximately 180 mM phosphate (Fig 2a). Chromatography on protamine agarose also yielded a single peak of enzyme activity that eluted at a mean phosphate concentration of 200 mM (Fig 2b). The final step utilized a Bio-Rad P-200 gel filtration (Fig 2c). Overall, an 86-fold purification was

Table 20. Purification of *Rana sylvatica* liver protein kinase A free catalytic subunit using various chromatographic techniques.

	Total Protein (mg)	Total Activity (nmol P _i /min)	% Yield	Fold Purification	Specific Activity (nmolP _i /min/mg)
Crude Homogenate	137	113.3	--	--	0.83
Hydroxyl-apatite	23.1	67.9	60	3.6	2.9
Protamine Agarose	7.5	25.1	22	4.0	3.3
S-300 Gel Filtration	0.047	3.3	3	85.7	70.9

Fig. 2. Elution profiles showing the chromatography steps in the purification of the free catalytic subunit of cyclic AMP-dependent protein kinase A from *Rana sylvatica* liver: A) hydroxylapatite, B) protamine agarose, C) Bio-Gel P-200 gel filtration column. The gradient used to elute the enzyme from the hydroxylapatite and protamine agarose columns was 10-500 mM KP_i , containing 2 mM 2-mercaptoethanol, 10% (v/v) glycerol, 1 mM EDTA, pH 6.8. One ml fractions were collected for the first two columns and 0.5 ml fractions were collected for the P-200 column. Enzyme assays were run at a pH of 6.8 under V_{max} conditions (see chapter 2). The pooled fractions collected from the Bio-Gel P-200 gel filtration column were used as the purified PKAc source for all kinetic parameter assays. The S.E.M. bars are contained within the symbols for $n = 3$ separate preparations.

achieved with a yield of 3 % and a final specific activity of 71 nmol Pi/min/mg protein. The novel purification scheme developed here for the isolation of *R. sylvatica* liver PKAc was extremely reproducible and resulted in highly purified enzyme.

Molecular Weight Determinations of *R. sylvatica* Liver PKAc

The subunit molecular weight of PKAc was determined to be 47 kDa by SDS polyacrylamide gel electrophoresis of the purified enzyme preparation (Fig. 3). The value obtained corresponded well with the molecular weight of 47.6 ± 1.1 kDa determined by Bio-Gel P-200 gel filtration (Fig. 4), confirming that the purified catalytic subunit was a monomer.

Stability of Purified Free Catalytic Subunit of PKA from *R. sylvatica* Liver

Purified PKAc lost activity quite rapidly over time. Therefore, tests were conducted with a number of different stabilizers to find one that would increase enzyme stability sufficiently to allow kinetic studies to be conducted over a number of days. Addition of various concentrations of glycerol (10%, 20%, 40% v/v) or inorganic phosphate (2, 5, 10, 20 mM) gave no improvement of stability. However, the detergent Triton X-100 was remarkably effective in stabilizing the enzyme, particularly when added to the buffer at the P-200 chromatography step of the purification. Enzyme stability over storage time at 4°C was tested in the presence versus absence of 0.25 % v/v Triton X-100 (Table 21). In the presence of Triton X-100, PKAc retained nearly full activity for 21 days whereas activity was rapidly lost during storage without the detergent. The enzyme preparation was also stable to freezing at low temperatures in the presence of Triton X-100 (Table 22). Full activity was retained after 2 h freezing at -196 or -80°C whereas about 28 % of activity was lost if storage was at -20°C. Preliminary kinetic studies were conducted in order to determine if the presence of Triton X-100 would alter enzyme kinetic properties. However, the detergent had no

Fig. 3. Subunit molecular weight determination for *Rana sylvatica* liver PKAc. SDS-PAGE used the following standards: (1) glycogen phosphorylase α (97.4 kDa), (2) bovine serum albumin (66.0 kDa), (3) ovalbumin (45.0 kDa), (4) bovine RBC carbonic anhydrase (29.0 kDa). Lane A contained crude supernatant; lane B, the enzyme preparation following hydroxylapatite column chromatography; lane C, the enzyme preparation following protamine agarose column chromatography; lanes D to I contained various peak fractions of PKAc activity from the P-200 gel filtration column.

1

2

3

4

A B C D E F G H I

Fig. 4. Native molecular weight of the free catalytic subunit of *Rana sylvatica* liver PKA as determined by P-200 gel filtration chromatography. An 100 μ l aliquot of purified PKAc was loaded onto a 10 ml (0.8 x 15 cm) Bio-Rad P-200 column equilibrated in buffer B plus 0.25 % v/v Triton X-100. The molecular weight standards were: (1) hexokinase (105 kDa), (2) enolase (82 kDa), (3) glycerol-3-phosphate dehydrogenase (68 kDa), (4) triose phosphate isomerase (53 kDa), (5) ovalbumin (43 kDa), and (6) chymotrypsinogen A (26 kDa). The void volume was determined with Blue Dextran and the total column volume was measured with KCl. Values are plotted as log molecular weight versus partition coefficient (K_p) according to Suelter (1985). The open circle shows the elution position of liver PKAc. Data presented is the mean $\pm$ S.E.M. for n = 3 separate preparations of purified liver PKAc.

Table 21. Stability of purified free catalytic subunit of frog liver PKA over time stored in the presence versus the absence of 0.25 % (v/v) Triton X-100 detergent at room temperature (22°C).

Days Stored	1	5	12	21
+ Triton X-100,(0.25 %)	70	70	64	64
- Triton X-100	70	40	27	12

Data are activity in units of nmol P_i/min/mg protein, n = 1.

Table 22. Stability of purified free catalytic subunit of frog liver PKA when stored at different subzero temperatures in the presence of 0.25 % (v/v) Triton X-100 detergent.

Temperature (°C)	Activity
+22	70.0 $\pm$ 4.1
-20	50.1 $\pm$ 3.2
-80	70.4 $\pm$ 2.4
-196	72.1 $\pm$ 1.7

Data are activity in units of nmol P_i/min/mg protein, n = 3.

effect on any of the kinetic parameters tested including K_m and I_{50} values, as well as Arrhenius plots and pH profiles of *R. sylvatica* liver PKAc.

Kinetic Characterization of *R. sylvatica* Liver PKA Free Catalytic Subunit

Table 23 shows the maximal activity of PKAc with five different phosphate-accepting substrates that have been used as PKAc substrates in other animal systems. Kemptide was by far the best substrate for the enzyme, particularly at the lower assay temperature. At 22°C the relative activity of the enzyme was 100:38:21:22:11 with kemptide, histone IIA, histone IIIS, histone VIS, and protamine used as substrates. The corresponding ratios at 5°C were 100:9:4:4:2. Based on these results, kemptide was chosen as the substrate for all subsequent kinetic analysis of the enzyme.

Substrate affinity constants for kemptide and Mg-ATP are shown in Table 24. Affinity for both substrates increased with decreasing assay temperature. The K_m value for Mg-ATP decreased by 50 % at 5°C, compared with 22°C, whereas the K_m for kemptide decreased by one-third at 5°C.

The effect of varying free magnesium concentration on PKAc activity is shown in Fig. 5. Enzyme activity was absolutely dependent on the presence of free magnesium (that is, a concentration of magnesium over and above that complexed with ATP). Optimal PKAc activity was reached at a concentration of 2.0 mM free magnesium. No enzyme activity was observed in assays lacking this divalent cation. Based on these results, a standard concentration of 2.1 mM Mg^{2+} was used for optimal activity in most PKAc assays.

Table 25 shows the inhibitory effects of various salts and metabolites on frog liver PKAc. Chloride salts had relatively limited inhibitory effects on the enzyme with high I_{50} values (495-562 mM) for KCl and NaCl at 22°C and a somewhat lower value for NH_4Cl (396 mM). Inhibition by both KCl and NaCl was reduced significantly (I_{50} increased) at 5°C. KNO_3 was also only a weak inhibitor but ammonium sulfate has a

Table 24. Substrate affinity coefficients for the purified free catalytic subunit of PKA from *Rana sylvatica* liver.

Temperature	22°C	5°C
K_m Mg-ATP, (μ M)	51.8 ± 1.0^a	24.8 ± 1.4
K_m Kemptide, (μ M)	9.0 ± 0.1^a	6.4 ± 0.3

Data are means $\pm$ S.E.M., $n = 3$ separate preparations of purified frog liver PKAc enzyme.

^a - Kinetic parameters are significantly different from the corresponding value at 5°C by Student's t-test, $P < 0.005$.

Fig. 5. Effect of increasing amounts of free magnesium on the activity of purified liver PKAc catalytic subunit. The enzyme showed an absolute dependence on magnesium ion for activity. Mg-ATP concentration was held constant at 2.08 mM throughout the experiment. Magnesium acetate was used and amounts of free and bound Mg^{2+} were calculated using a program developed for this purpose (Brooks and Storey, 1992). Data are means $\pm$ S.E.M. for $n = 3$ separate enzyme preparations.

Table 25. Inhibitor coefficients (I_{50} values) for the purified free catalytic subunit of PKA from *Rana sylvatica* liver.

Inhibitor	I_{50} Value	
	22°C	5°C
KCl (mM)	495 ± 10	720 ± 10 ^b
KNO ₃ (mM)	426 ± 12	--
NaCl (mM)	562 ± 16	700 ± 12 ^b
NaF (mM)	19.3 ± 1 ^a	--
NH ₄ Cl (mM)	396 ± 8 ^a	--
(NH ₄) ₂ SO ₄ (mM)	150 ± 12 ^a	--
Glucose (M)	2.1 ± 0.2	1.7 ± 0.2
PKA _i (nM)	3.3 ± 0.15	--
H89 (nM)	25.1 ± 0.6 ^c	--

Data are means ± S.E.M., n = 3 separate preparations of purified frog liver PKAc enzyme.

^a - All I_{50} values are significantly different from the corresponding value for the KCl salt at 22°C by Student's t-test, $P < 0.005$.

^b - All I_{50} values are significantly different from their corresponding salt at 5°C by Student's t-test, $P < 0.005$.

^c - I_{50} value is significantly different from the corresponding PKA_i value at 22°C by Student's t-test, $P < 0.005$.

substantial inhibitory effect on the enzyme. Fluoride anion was a powerful and specific inhibitor of PKAc with an I_{50} value at 22°C of only 19.3 ± 0.5 mM, a value nearly 30-fold lower than the corresponding value for NaCl.

Due to the fact PKA appeared to be important in the activation cascade of glycogenolysis that lead to the accumulation of glucose cryoprotectant in frogs and because glucose accumulated to high concentrations (as much as $350 \mu\text{mol/gww}$) in *R. sylvatica* liver during freezing, the effect of glucose itself on PKAc activity was tested. However, I_{50} values for glucose were very high, 2.1 ± 0.2 M at 22°C and 1.7 ± 0.2 M at 5°C (Table 25), and well above physiological levels, suggesting that PKAc activity would not be affected by the high glucose levels during freezing *in vivo*.

The effects of specific inhibitors of mammalian protein kinases on frog liver PKAc were also tested. PKA_i and H89 are known inhibitors of mammalian PKA and both also strongly inhibited the frog enzyme with I_{50} values in the nanomolar range (Table 25) (Walsh et al., 1971; Chijiwa et al., 1990). These data confirm the identity of the frog liver enzyme as a cAMP-dependent protein kinase.

An Arrhenius plot of log relative activity versus $1/T \times 10^{-3}$ (°K) was completed under V_{max} assay conditions (Fig. 6). A distinct break in the plot was seen at 10°C. The calculated activation energy, E_a , for the reaction at temperatures above 10°C was 51 ± 4 kJ/mol whereas below 10°C E_a was 110 ± 9 kJ/mol.

The effect of pH on frog liver PKAc activity is shown in Fig. 7. The pH optimum was strongly influenced by assay temperature. A sharp optimum at pH 7.4 was seen for the enzyme assayed at 22°C but at 5°C the optimum dropped to pH 6.0.

Assessment of Tissue Specific Isozymic Forms of PKAc in *R. sylvatica* and *R. pipiens*

Hydroxylapatite column chromatography was used to determine whether multiple isozymic forms of PKAc might exist in the tissues of *R. sylvatica*. A single

Fig. 6. Arrhenius plot for *Rana sylvatica* liver PKAc. The E_a for the reaction above 10°C was 51 ± 4 kJ/mol and below 10°C was 110 ± 9 kJ/mol. The distinct break in the plot at 10°C may be indicative of a conformational change in the enzyme below this temperature. Activities (dpm) were measured at 5°C intervals between 1°C to 35°C under V_{max} assay conditions. Data are means $\pm$ S.E.M. for $n = 4$ separate enzyme preparations.

Fig. 7. The effect of pH on PKAc activity at 22°C (A) and 5°C (B). Total Mg^{2+} and ATP concentrations were varied to maintain constant concentrations of Mg-ATP and free Mg^{2+} at each temperature and pH. Data are means $\pm$ S.E.M. for $n = 3$ separate enzyme preparations. The concentrations for free Mg^{2+} , ATP, and all ionic species such as K^+ , Cl^- , PO_4^{3-} , EDTA needed to determine the correct pH-temperature relationship for the pH curves were calculated using the BAD program developed by Brooks and Storey (1992). The pH was monitored at each temperature before and after every assay was run. In all cases it was found the pH values calculated using the BAD program remained constant throughout the entire pH range at both temperatures after the assays were completed.

sharp peak of PKAc activity eluted at exactly the same KP_i concentration of 420 mM for extracts from liver, muscle, and brain when each tissue was applied to separate hydroxylapatite columns and eluted with a salt gradient running from 0.01-1.0 M KP_i . Furthermore, when samples of the 3 tissues were homogenized together and applied to a single hydroxylapatite column, PKAc activity again eluted in a single peak at 420 mM KP_i (Fig. 8). An identical result was found for PKAc from extracts of liver, muscle, and brain of the leopard frog, *R. pipiens*. Hence, at least in these 2 anurans, only a single isoform of the free catalytic subunit of PKA appeared to reside in the 3 tissues examined.

Fig. 8. Assessment of the possible presence of tissue specific isozymes of PKAc in *Rana sylvatica* liver, muscle, and brain tissue were analyzed using hydroxylapatite chromatography. Individual tissues were homogenized 1:6 (w/v) in 10 mM KP_i , 1 mM EDTA, 2 mM β -mercaptoethanol, and 1 mM cAMP. Two ml of each homogenate was then loaded onto separate 0.5 cm x 2 cm hydroxylapatite columns. PKAc activity eluted in a single peak at the same salt concentration for all 3 tissues using a 10 mM - 1 M KP_i salt gradient. Only 1 form of PKAc was found in these 3 tissues after initial data analysis. Therefore, in order to get an $n = 3$ for these tissues, 1 liver, 1 piece of leg muscle, and 1 brain were homogenized together and run out on an hydroxylapatite column, 3 separate times. Accumulated data are means $\pm$ S.E.M. for $n = 4$ separate tissue preparations. The same procedure was followed using *R. pipiens* liver, muscle, and brain (data not shown).

DISCUSSION

In the present study a novel purification of the catalytic subunit of cAMP dependent protein kinase from *R. sylvatica* liver is presented. The homogenization procedure using cAMP ensured the dissociation of the catalytic from the regulatory subunits of this tetrameric protein. In fact all homogenates obtained with this procedure contained 100 % of the catalytic form of the enzyme as determined by assaying crude homogenates in the absence of cAMP.

PKAc from *R. sylvatica* had a final specific activity of 71 nmol P_i /min/mg protein (Table 20) which was lower than that reported for other purified catalytic subunits of the enzyme including *Drosophila melanogaster* and bovine heart (Foster and Hall, 1988; Haracska and Udvardy, 1992; Okuno and Fujisawa, 1990). However, the specific activity for frog liver PKAc was 2-fold higher than that for PKAc from the snail *Otala lactea* (Brooks and Storey, 1993). Most purifications of PKAc to date have been from mammalian tissue sources whose specific activity measurements are calculated at 37°C rather than at 22°C as in the present study (Sugden et al., 1976). When this temperature difference was factored into the calculation the resulting specific activity for wood frog PKAc was very similar to mammalian values.

The purification procedure for this enzyme was highly reproducible (Fig. 2). Although the overall yield was low, a highly stable purified enzyme preparation was easily obtained (Fig. 3). According to all recent literature on the methods of purification of PKAc from mammalian tissues, the enzyme should be highly purified after an initial step utilizing DEAE⁺-Sephadex and CM⁻-Sephadex chromatography (Livesey and Martin, 1988) or with a single DEAE⁺-Sephadex chromatographic step (Okuno and Fujisawa, 1990). The principle behind this dual column purification is to bind the inactive PKA (the regulatory subunit) to the DEAE⁺ column and the active PKA enzyme (catalytic subunit) to the CM⁻ column. In mammalian systems these 2

chromatographic steps yield a purified homogeneous PKAc enzyme. Despite numerous attempts, however, the yield of *R. sylvatica* PKAc from DEAE⁺-Sephadex was never higher than 30 %, and therefore, this step was not useful for purification of the frog enzyme. Therefore, other chromatographic techniques were required to purify the enzyme. The novel combination of hydroxylapatite, protamine agarose, and P-200 gel filtration proved highly effective for this purpose.

The native and subunit molecular weights of *R. sylvatica* liver PKAc, 47.6 kDa and 47 kDa, were very similar to those obtained for mammalian, reptilian, and invertebrate forms of the enzyme (Bechtel et al., 1977; Zoller et al., 1979; Mehrani and Storey, 1994; Brooks and Storey, 1993; Haracska and Udvardy, 1992). This is further evidence that the structure and function of this kinase has been highly conserved throughout evolution.

Storage of the enzyme in detergent was necessary for long term stability of PKAc (Table 21). Purified PKAc was stable even at -196°C as long as Triton X-100 was present (Table 22). Other PKAc enzymes such as bovine heart muscle and *Otala lactea* foot muscle also exhibited this characteristic (Okuna and Fujisawa, 1990; Brooks and Storey, 1993). However, unlike *R. sylvatica* PKAc, those enzymes were stable for several weeks in solutions containing glycerol. It was not known why the activity of PKAc was reduced at -20°C and not at the lower temperatures. However, this result did mimic the data of Reimann and Beham (1983) and Okuno and Fujisawa (1990). Kinetic studies using purified PKAc from *R. sylvatica* were conducted using enzyme stored at 4°C which ensured optimal activity.

The kinetic constants for *R. sylvatica* liver PKAc were similar to those reported for the mammalian catalytic subunit of PKA. Frog liver PKAc exhibited a much higher affinity for the synthetic substrate kemptide than for several different phosphate accepting histone substrates at both the temperatures tested (Table 23). Phosphate incorporation catalyzed by mammalian PKAc enzymes was also much

greater with kemptide as the substrate than with histones and the same was true for PKAc from turtle liver (Mehrani and Storey, 1994). Therefore, it seems reasonable to conclude that the catalytic site of PKAc is highly conserved throughout the phylogeny of vertebrates.

R. sylvatica liver PKAc had an absolute requirement for magnesium ions for its catalytic activity (Fig. 5). This was observed for both mammalian and invertebrate forms of the free catalytic subunit of PKA (Hofmann et al., 1975; Bishoff et al., 1990). The pH optimum for the enzyme was very broad at room temperature, as was also seen for *T.s. elegans* liver PKAc (Mehrani and Storey, 1994). However, this pH optimum was not conserved at lower temperatures, at 5°C the pH optimum fell to 6.0 (Fig. 7). This was not the case for other vertebrate forms of PKAc. *R. sylvatica* liver PKAc is unique in that at 5°C, the pH optimum is similar to that found for an insect form of the enzyme at warm temperatures (Haracska and Udvardy, 1992). The break in the Arrhenius plot at 10°C implies that there may have been a conformational change in the enzyme at lower temperatures (Fig. 6). This conformational change, as well as the acidic shift in pH optimum at a time when low body temperatures cause a basic shift in intracellular pH, may indicate that the enzyme is functioning under these "restricted" conditions both before freezing, and before cryoprotectant synthesis is activated. Thus, during all normal winter hibernation, (whether in the frozen state or at low temperatures) the wood frog could be dealing with a PKAc that is "subactive" as compared with what would be expected by a straight extrapolation of activity at temperatures above 10°C.

Other reasons for the break in the Arrhenius plot seen in Fig. 6 may be due to: (1) an increase in the viscosity of cell cytosol in the frozen condition which would lead to a change (reduction) in PKAc activity, and (2) PKAc and other multistep enzymes may change one "step" of their multiple catalytic steps of substrate to product

conversion as a result of a change in temperature of the PKAc reaction. The consequence would be an abrupt change in enzymatic activity with temperature.

Perhaps this confers an overall reduced response by PKAc activity to stimuli that would normally activate the enzyme and could be part of a general quiescence or metabolic depression in *R. sylvatica* during the winter. However, the stimulus of freezing must be powerful enough to override this inhibition, as the substrate kinetic data seem to indicate.

The K_m values for both substrates were within the range of mammalian values and much lower than those found for *Otala lactea* (Brooks and Storey, 1993). The lower K_m values for kemptide and Mg-ATP at 5°C suggests that at colder temperatures the enzyme shows a much higher affinity for both substrates (Table 24). All of the above characteristics show that a remarkable similarity exists between the mammalian and anuran enzymes. It appears that many kinetic characteristics are conserved among vertebrate forms of PKAc.

Frog liver PKAc was susceptible to inhibition by many different inorganic salts. As Table 25 shows, changing the cation had no effect on the inhibition of PKAc, and hence, inhibition must be linked to the anion effects. At lower temperatures, not only did the inhibition of PKAc activity by these dissociated salts decrease, but the concentrations needed for inhibition were 10 fold higher than those found *in vivo*! The inhibitory effect of all the tested salts on enzyme activity were similar to those found for mammalian, reptilian, and invertebrate forms of this kinase (Mehrani and Storey, 1994; Brooks and Storey, 1993). A large inhibition was observed when the anion was altered as was exemplified by sodium fluoride (Table 25). However, fluoride occurs in only trace amounts *in vivo*, and therefore, cannot be considered a physiological inhibitor of the enzyme. NaF is often used *in vitro* in PKAc purifications to inhibit the activity of endogenous protein phosphatases. Therefore, a careful balance must be struck in the design of *in vitro* experiments involving PKA in order to avoid the

inhibition of the enzyme while maintaining enough NaF in the assay to inhibit phosphatase activities.

The fact that inhibition by KCl and NaCl decreased (Table 25) with a decrease in temperature supports the observation that liver PKAc can be fully activated at subzero temperatures under which high intracellular ion concentrations can exist due to the freeze concentration phenomenon. This ensures that the production of cryoprotectant (glucose) in *R. sylvatica* liver can occur even at low temperatures and supports an activation of PKAc at these low temperatures since the intracellular ion concentrations are not high enough to inhibit liver PKAc activity.

The product of the activation of glycogenolysis in *R. sylvatica* liver during a freezing episode, glucose, had no inhibitory effect on frog liver PKAc (Table 25). The endogenous concentration of this cryoprotectant under normal physiological conditions is 2-5 mM. As the frog begins to freeze liver glucose concentrations rise dramatically reaching values ranging from 200-300 mM in fully frozen animals (Storey and Storey, 1986, this study). The I_{50} values for glucose were 10-fold higher than even the highest endogenous concentrations of this molecule, suggesting that this enzyme is not regulated by a negative feedback mechanism involving glucose.

The artificial inhibitors of mammalian PKAc, PKA_i and H89, also inhibited *R. sylvatica* PKAc (Table 25). These two inhibitors were useful as screening tools to determine whether or not PKAc from a lower vertebrate responded in a similar manner, as all mammalian PKAc enzymes do, to these two inhibitors. In fact, the I_{50} values for PKA_i and H89 were as low as those found for many mammalian forms of the enzyme indicating a similar type of inhibition for *R. sylvatica* PKAc, and a conservation of the inhibitor site of this enzyme in vertebrate organisms.

Hydroxylapatite chromatography revealed that only 1 isozyme of PKAc existed in liver, brain, and muscle of two ranid species, *R. sylvatica* and *R. pipiens*. This evidence suggests that the catalytic subunit of PKA is conserved in these 2

anurans. However, recent antibody studies have provided increasing evidence that there are multiple forms of the free catalytic subunit of mammalian PKA in variety of different tissue types (Chivia, et al., 1988; Hotz et al., 1989; Takhar and Munday, 1992). Thus, further studies are required using antibodies and reverse phase HPLC to confirm this hypothesis in relation to anuran species.

Three principles can be drawn from the kinetic data for *R. sylvatica* liver PKAc that indicate that the enzyme is capable of supporting cryoprotectant synthesis in this freeze-tolerant frog at very low temperatures, $<5^{\circ}\text{C}$. These are: (1) the low sensitivity of the enzyme to inhibition by cations and anions at high or low temperatures, (2) the lack of inhibition by glucose, and (3) an increase in the affinity of the enzyme for its substrates at low temperatures. *R. sylvatica* liver PKAc appears to function quite well in a cellular environment that contains high solute (glucose) and salt concentrations, both of which are known to occur in this animal during a freezing episode. The lack of any inhibitory effect on PKAc activity by the compounds tested *in vitro*, may signify that as the cell cytosol becomes more concentrated during freezing *in vivo*, PKAc can function to retain the activation of liver glycogenolysis, despite the rising levels of these ions and glucose.

It is clear from the above kinetic data that the free catalytic subunit of PKA in *R. sylvatica* liver is very similar to other vertebrate forms of the enzyme. In addition, the enzyme shares various kinetic and physical properties with invertebrate forms of PKAc. It appears then, that the structure and function of PKAc has been highly conserved throughout evolution (Taylor et al, 1990). Protein phosphorylation by PKAc is an important biochemical mechanism that cells utilize to regulate many different metabolic pathways, the primary one being liver glycogenolysis. *R. sylvatica* liver PKAc must be responsible for the phosphorylation and subsequent activation of liver GP in response to a freezing stress. The post-translational modification of GP is

critical for the production of glucose as a cryoprotectant, and thus, the winter survival of this freeze tolerant anuran.

CHAPTER #7**ACTIVATION AND TRANSLOCATION OF PROTEIN KINASE C DURING
FREEZING STRESS**

INTRODUCTION

Protein kinase C, PKC, and its signal transduction cascade is arguably the most widely studied mechanism of cellular post-translation modification. Once activated, PKC plays a pivotal role in the phosphorylation of serine and threonine residues on a myriad of target proteins in a variety of tissues (Qu et al., 1991). For example, PKC is responsible for phosphorylating regulatory enzymes that are important in cell growth and differentiation (Hashimoto and Yamamura, 1991) and in modulating ion channel flux (Nishizuka, 1986; Bourinet et al., 1992). Thus, PKC stimulated phosphorylation of target proteins is one of the most important mechanisms for regulating many different biochemical pathways in all types of cells. Perhaps then, PKC has a synergistic role along with PKAc in the direct activation of GP via phosphorylation that initiates the breakdown of liver glycogen stores to glucose as a response to freezing stress in *R. sylvatica*.

Initial kinetic studies involving PKC characterized the enzyme as a calcium and phospholipid dependent enzyme that required, along with a translocation to the plasma membrane, diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP₃) for its activation (Nishizuka, 1986). It has now been confirmed that there are 9 distinct isoforms of this kinase, subdivided into two groups (Bell and Burns, 1991). Group one includes the calcium-dependent isozymes (α , β I, β II, and γ) and group two the calcium-independent isozymes (δ , ϵ , ζ , θ , ι , κ , λ , μ , and ν) (Bell and Burns, 1991). One of the aims of the present study was to determine the specific PKC isozyme forms that exist in *R. sylvatica* liver and brain tissue with the use of hydroxylapatite chromatography and PKC antibody and immunoblotting techniques.

Many extracellular stimuli, in this instance external ice formation on the skin of *R. sylvatica*, must transmit their signals to the interior of cells by binding to specific receptors found in or on the plasma membrane of the cell (Qu et al., 1991). The

binding of these hormone signals induces an activation of the PKC signal transduction pathway beginning with the hydrolysis of polyphosphoinositides. This in turn, results in the formation of IP₃ and DAG (Berridge, 1987). An increase in inositol 1,4,5-trisphosphate, IP₃, causes a rise in the level of cellular calcium by releasing this cation from intracellular stores (Rasmussen, 1989). PKC then translocates to the membrane in a calcium dependent manner where it binds to phosphatidylserine producing an inactive but primed kinase enzyme. Concurrent to the release of calcium, elevations in diacylglycerol production result in the activation of primed PKC by increasing the enzyme's affinity for intracellular calcium. It seems that these two second messengers have a synergistic role in the activation of PKC in many cellular systems (Epand and Lester, 1991).

It has been shown that diacylglycerol is only transiently produced in the membrane of cells due to phosphoinositide breakdown (Nishizuka, 1984). Therefore, it is thought that PKC is a unique protein kinase enzyme, in that it must be bound to the membrane of the cell in the presence of DAG to be physiologically active. Without the correct complement of membrane phospholipids and calcium the enzyme is inactive.

The objectives of this study are three-fold: (1) to determine if PKC translocates to the membrane in liver and brain tissue during a 24 h whole animal freezing exposure of *R. sylvatica*, (2) to ascertain which PKC isozyme(s) are present in these two organs of the freeze-tolerant frog, and (3) to determine if the PKC signal transduction cascade system is involved in the activation of glycogenolysis and the freezing-induced activation of cryoprotectant synthesis by wood frog liver.

MATERIALS AND METHODS

Preparation of Experimental Animals and PKC assay.

R. sylvatica were collected, held, and experimental bulk freezing exposures were undertaken as described in chapter 2. Freezing time was 24 h at -2.5°C and frogs were thawed at 5°C for 24 h after freezing.

Protein Kinase C Assay Protocol

Frozen tissue samples were quickly weighed and then homogenized 1:30 w/v in buffer A (20 mM HEPES, 10 mM β -mercaptoethanol, 5 mM EDTA, 5 mM EGTA, 250 mM sucrose, pH 7.4) with 10 $\mu\text{g}/\text{ml}$ leupeptin, 1 $\mu\text{g}/\text{ml}$ aprotinin, and 1 mM PMSF added to buffer A just prior to homogenization of tissues with a Polytron PT 10 homogenizer. Homogenates were centrifuged at 24,000 g for 15 min in a Sorvall RC-5B centrifuge and then supernatants were removed. To remove inhibitors and ATP that were present in the supernatant, extracts were then loaded onto 0.5 cm x 2.0 cm columns of Sephadex DE-52 that were equilibrated in fresh buffer A, with 10 $\mu\text{g}/\text{ml}$ leupeptin, 1 $\mu\text{g}/\text{ml}$ aprotinin, and 1 mM PMSF. Columns were washed with 50 ml of buffer B (20 mM HEPES, 10 mM β -mercaptoethanol, 20 mM KCl, 1 mM EDTA, 1 mM EGTA, pH 7.4) and then PKC was eluted in 2 ml of buffer C (20 mM HEPES, 10 mM β -mercaptoethanol, 250 mM KCl, 0.1 mM EDTA, 0.1 mM EGTA, pH 7.4). This 2 ml eluate was used as the source of PKC activity for all assays.

The activity of protein kinase C was monitored by a radioactive assay that measured the incorporation of radiolabelled ^{32}P -ATP onto the phosphate-accepting artificial substrate 'Durkin peptide' (Chakravarthy et al., 1989). All solutions needed to monitor PKC activity were prepared fresh daily in distilled water as follows. Solution I: 500 μM Durkin Peptide; Solution II: 40 $\mu\text{Ci}/\text{ml}$ ^{32}P -ATP, 50 mM $\text{Mg}(\text{OAc})_2$, 200

μM ATP; Solution III: 100 mM EGTA; and Solution IV: 8 mM CaCl_2 . Solution V was the lipid preparation and was prepared just prior to PKC assay in the following manner. Stock phosphatidylserine (PS) and diacylglycerol (DAG), 40 μl and 10 μl respectively, were placed in a 1.5 ml microcentrifuge tube on ice. The mixture was evaporated under a constant stream of N_2 gas and then 1 ml of assay buffer containing 60 mM HEPES, 30 mM β -mercaptoethanol pH 7.4, was added. The solution was sonicated for 3 min, under a constant stream of N_2 gas, until the liquid turned a milky white color. The final concentrations of PS and DAG in the preparation were 20 $\mu\text{g/ml}$ and 2 $\mu\text{g/ml}$, respectively. The lipid preparation was kept on ice away from a light source. PKC assays were run in disposable glass tubes with 50 μl of a 1:20 w/v tissue homogenate plus 50 μl of reactants (Solutions I-V), plus water to make up a final volume of 100 μl . The standard assay had the following concentrations of reactants in a final volume of 100 μl : 100 μM Durkin peptide, 0.4 μCi ^{32}P -ATP/assay, 5 mM $\text{Mg}(\text{OAc})_2$, 20 μM ATP, 10 mM EGTA, 0.8 mM CaCl_2 , 4 μg PS, and 0.4 μg DAG. Control tubes which measured calcium-independent PKC activity contained no calcium and control tubes that measured background activity contained no phospholipid; however, each control had EGTA present. Total PKC activity was always assayed in the presence of both calcium and phospholipid. Solutions were added in the following order for a typical assay: solution I, solution IV, PKC enzyme source, solution V, and finally solution II, for a final reaction volume of 100 μl . Reactions were run for 20 min at 22°C and then stopped using the P81 filter paper method of Roskoski (1983), as described in chapter 2 for PKA, except that 60 μl aliquots were sampled from each reaction tube. One unit of PKC activity is defined as the amount of enzyme that catalyzes the incorporation of 1 nmol of ^{32}P from ATP onto Durkin peptide in one minute at 22°C.

Effect of Freezing and Thawing on PKC Distribution in Membrane versus Cytosolic Fractions

PKC exists in both free cytosolic (non-active) and membrane bound (active) forms *in vivo*. The effect of experimental freezing on the distribution of PKC in *R. sylvatica* liver and brain homogenates was determined using the protocol described below. PKC must be separated from its endogenous inhibitors in order to obtain the correct measurements of membrane bound versus free PKC. The use of Sephadex DE-52 column chromatography not only accomplished this but also separated the membrane bound (active) versus free cytosolic (non-active) fractions of PKC.

Frozen liver or brain tissue was homogenized (1:20 w/v liver or 1:30 w/v brain) in buffer A with 10 μ g/ml leupeptin, 1 μ g/ml aprotinin, and 1 mM PMSF added to the buffer just prior to homogenization. Homogenates were centrifuged at 100,000 g for 30 min at 5°C in a Beckman L2-65B Ultracentrifuge to separate the cytosolic (supernatant) and membrane (pellet) PKC fractions. Supernatants were removed and stored on ice. Pellets were placed in buffer A containing 0.1% (v/v) Triton X-100, and stored on ice for 30 min. Pellets were then resuspended by homogenization using an Ultra Turrax tissue homogenizer and centrifuged at 27,000 g at 5°C for 15 min using a Sorvall RC-5B centrifuge. The supernatant from this spin was removed and saved as the "resuspended pellet supernatant".

Both the original supernatant and the resuspended pellet supernatant were loaded onto 0.5 cm x 2.0 cm DE-52 columns that were pre-equilibrated in fresh buffer A. Columns were washed with 50 ml of buffer B and PKC activity was eluted in 2 ml of buffer C which contained 20 mM HEPES, 10 mM β -mercaptoethanol, 250 mM KCl, 0.1 mM EDTA, 0.1 mM EGTA at pH 7.4. PKC activity was assayed in each fraction as above. Percent bound PKC activity was calculated as the quotient of PKC activity in the resuspended pellet divided by the total PKC activity (PKC activity in the supernatant plus the PKC activity in the resuspended pellet).

Classification of PKC Isozymes Using Hydroxylapatite Column Chromatography and Immunoblotting

Isozymes of PKC in liver and brain tissues of control *R. sylvatica* were separated using hydroxylapatite column chromatography and their identity was confirmed using the modified immunoblotting procedure.

Samples of liver (0.5 g) were homogenized 1:20 (w/v) in isozyme buffer: 10 mM β -mercaptoethanol, 20 mM NaP_i , 0.1 mM EGTA, 0.1 mM EDTA, 20% glycerol (v/v), pH 7.0. Crude homogenate was loaded onto an 0.5 x 2.0 cm hydroxylapatite column pre-equilibrated in isozyme buffer. The column was washed with 40 ml of isozyme buffer and then PKC activity was eluted in a single peak using a 0.02-2 M NaP_i salt gradient. Peak fractions containing PKC activity were pooled and dialyzed for 2 h against isozyme buffer without glycerol. The pooled peak was then subjected to SDS-PAGE.

A similar protocol was used to analyze the PKC isozymes in *R. sylvatica* brain tissue except that brain tissue (0.2 g) was homogenized 1:30 (w/v) in isozyme buffer. Pooled brain isozymes were not subjected to the SDS-PAGE, immunoblotting, or PKC antibody staining procedures.

Gel Electrophoresis and Electoblotting Protocol

The single liver PKC isozyme contained in the pooled peak fractions from the hydroxylapatite column was run out on a 10 % acrylamide gel, a procedure modified from Laemmli (1970). Briefly, gels were run at 150 V for 1 h at room temperature. Prestained molecular weight standards from Bio-Rad, as well as a sample of rabbit brain extract (1:30 w/v crude homogenate in isozyme buffer) were run along with the PKC pooled peak fraction from frog liver on two identically prepared SDS gels. The standards served as markers during the transfer procedure to ensure the proper transfer

of proteins from the SDS gels to the nitrocellulose paper. The rabbit brain PKC source served as a marker to ensure that the antibody stains were, in fact, staining for PKC. After electrophoresis, gels were removed from the apparatus and placed in transfer buffer to equilibrate at 4°C for 1 h.

Several solutions were prepared prior to the electroblotting procedure. Transfer buffer contained 100 mM glycine, 50 mM TRIS base, 10 % methanol (v/v), pH 8.2. A 10x phosphate-buffered saline (PBS) solution contained 137 mM NaCl, 8 mM Na_2HPO_4 , 1.5 mM KH_2PO_4 , pH 7.4. Blocking solution consisted of 10x PBS plus 1% (w/v) bovine serum albumin (BSA). PBS-Tween 20 solution contained 100 ml of 10x PBS solution plus 1 ml of Tween 20 detergent with the volume then adjusted to 1 L with double distilled water. Alkaline phosphatase buffer contained 100 mM sodium bicarbonate with 5 mM MgCl_2 , pH 9.8. The nitro blue tetrazolium (NBT) solution was prepared by dissolving 5 mg of NBT in a 1 ml solution of 70% dimethylformamide (DMF), 30% H_2O . The 5-bromo-4-chloro-3-indolyl-phosphate (BCIP) solution was prepared by dissolving 5 mg of BCIP disodium salt, in 1 ml of 100% DMF. The latter two solutions were not prepared until the nitrocellulose papers were ready to be stained.

Two 10 x 6 cm nitrocellulose membranes were cut, wet with water by capillary action, and then equilibrated in 100 ml of transfer buffer plus 1 ml of 10 % w/v SDS at 4°C for 30 min. Six pieces of 10 x 21 cm Whatman filter paper were also equilibrated in cold transfer buffer. A gel-membrane sandwich was arranged inside the Owl-TM Semi Dry Electrophoretic apparatus in the following order; 3 pieces of saturated filter paper on the cathode base, two equilibrated gels, two nitrocellulose membranes (one over each gel), and finally the remaining pieces of saturated filter paper. All air bubbles were removed from the gel sandwich prior to the attachment of the anode. Proteins were transferred to the nitrocellulose membranes for 1.5 h with a current of 400 mA and a voltage setting between 5-10 V. The two nitrocellulose membranes were

then rinsed in several washes of distilled water. Gels were stained in Coomassie stain to check the efficiency of protein transfer onto the nitrocellulose membranes.

Immunodetection of PKC Isozymes

Membranes were placed protein side up in 20 ml of blocking solution for 2 h on a rotating shaker. Blocked membranes were then washed 3 times (each wash was 5 min) in PBS-Tween 20. Antibodies were purchased from Gibco-BRL; the primary antibody was either rabbit antiPKC- α or antiPKC- β whereas the secondary antibody was goat antirabbit IgG (H+L) attached to an alkaline phosphatase. Primary antibody was prepared by placing 10 μ l of either antiPKC- α or antiPKC- β and 12 ml of PBS-Tween 20 in a glass tube; these were then inverted 10 times to ensure good mixing. One membrane was placed in a dish with antiPKC- α and the other in antiPKC- β and both were incubated overnight on a rotating shaker. Following incubation, membranes were rinsed 3 times with 20 ml cold PBS-Tween 20 (for 10 min each). Two solutions of secondary antibody were prepared, each containing 20 μ l goat antirabbit IgG in 15 ml of PBS-Tween 20. Each membrane was then incubated in secondary antibody for 1 h followed by washing 3 times in 20 ml PBS-Tween 20 (10 min per wash).

Alkaline Phosphatase Staining of Nitrocellulose Membranes with Incorporated antiPKC- α and antiPKC- β Antibodies

Nitrocellulose membranes with incorporated antibody were then stained as follows. For the staining solution, 660 μ l of NBT solution was mixed with 10 ml of alkaline phosphatase buffer and then 330 μ l of BCIP solution was added, mixed, and the solution was placed in a staining dish. The nitrocellulose membranes were then added and the dish was placed in the dark for 5 min at room temperature. When the

stained protein bands reached a desired intensity the reaction was stopped by adding a solution of 200 μ l 0.5 mM EDTA (pH 8.0) in 50 ml of PBS-Tween.

RESULTS

Effect of Freezing and Thawing on the Distribution of Protein Kinase C in Frog Organs.

The effects of whole animal freezing exposure (24 h at -2.5°C) and subsequent thawing (24 h returned to 5°C) on the distribution of PKC between membrane-bound and free cytosolic fractions of liver and brain are shown in Table 26. Freezing exposure led to a significant reduction in the % bound for PKC in liver and this was maintained after thawing. By contrast, freezing had no effect on PKC distribution in brain but the percent membrane bound fell significantly from about 22 % in control or frozen frogs to 13.8 % in thawed animals. For comparison, Table 26 also shows the concentrations of IP_3 (from Table 4) in these same tissues. By contrast with the significant decrease in membrane-associated PKC activity in liver of frozen frogs, IP_3 levels were 11-fold higher in frozen, versus control, frogs. IP_3 levels fell again after thawing, although remained significantly above initial control values. Freeze/thaw had no effect on IP_3 content in brain.

Classification of Frog Liver PKC Isozyme(s) Using Hydroxylapatite Column Chromatography and Immunoblotting Techniques

A search for isozymes of PKC in *R. sylvatica* liver was conducted using hydroxylapatite column chromatography, a technique that has been shown to separate isozymes of PKC in mammalian tissues. The identity of the isozyme(s) was verified by a modified immunoblotting procedure. Results for these procedures are shown in Fig. 9 and 10. PKC eluted in a single large peak from the hydroxylapatite column at a NaP_i

Table 26. Effect of freezing exposure (24 h, at -2.5°C) and subsequent thawing (24 h, at 5°C) on the percentage of bound PKC and the levels of inositol triphosphate (IP_3) in two tissues of spring-collected *Rana sylvatica*.

Tissue/Condition	Bound PKC % of total activity	$[\text{IP}_3]$ (pmol/gww)
Liver		
Control	11.9 ± 1.0	1406 ± 164
Frozen (24 h, -2.5°C)	$8.5 \pm 1.2^{\text{a}}$	$16525 \pm 2983^{\text{a}}$
Thawed (24 h, 5°C)	$6.3 \pm 0.7^{\text{a}}$	$3214 \pm 737^{\text{a,b}}$
Brain		
Control	22.5 ± 0.9	20133 ± 1689
Frozen (24 h, -2.5°C)	21.7 ± 1.0	22807 ± 2465
Thawed (24 h, 5°C)	$13.8 \pm 0.4^{\text{a,b}}$	21345 ± 1922

Data are means $\pm$ S.E.M., $n = 4$ animals for bound PKC and IP_3 numbers. IP_3 values for both liver and brain tissue are taken from the bulk freezing experiment, Table 4.

^a - Significantly different from the corresponding control value using a one way ANOVA with an SNK test, $P < 0.05$.

^b - Significantly different from the corresponding 24 h frozen value using a one way ANOVA with an SNK test, $P < 0.05$.

Fig. 9. Assessment of the presence of isozymes of PKC in *R. sylvatica* liver using hydroxylapatite column chromatography. Liver (0.5 g) was homogenized 1:20 (w/v) in isozyme buffer pH 7.0. Isozyme buffer contained 10 mM β -mercaptoethanol, 20 mM NaP_i , 0.1 mM EGTA, 0.1 mM EDTA, and 20% glycerol (v/v). Crude homogenate was loaded onto hydroxylapatite columns. PKC activity eluted in a single peak using a 20 mM - 2 M NaP_i salt gradient. PKC activity was monitored using the assay protocol described in chapter 7. Data are means $\pm$ S.E.M. for $n = 4$ separate enzyme preparations.

Fig. 10. PKC isozymes from *R. sylvatica* liver (peak fractions with the highest PKC activity off of the hydroxylapatite column) were separated using SDS polyacrylamide gel electrophoresis on a 10% acrylamide gel (Laemmli, 1970). Prestained standards from Bio-Rad, along with a 1:30 (w/v) crude homogenate of rabbit brain were run together with the PKC fractions from frog liver on two identically prepared SDS gels. PKC isozymes were then transferred to nitrocellulose membrane (a) and (b) as described in chapter 7. Membranes were placed protein side up in 20 ml of blocking solution for 2 h, washed 3 times in PBS-Tween 20, and incubated with either antiPKC- α (membrane a) or antiPKC- β (membrane b) antirabbit primary polyclonal antibody. The secondary antibody was goat antirabbit IgG (H+L) attached to an alkaline phosphatase. Both nitrocellulose membrane a and b containing incorporated antibody were stained using the alkaline phosphatase staining procedure described in chapter 7. Membrane (a) contains lane E, the α -isozyme of rabbit brain PKC. Membrane (b) contains the stained lanes: A to D, the M-subunit of *R. sylvatica* PKC- β ; and E, rabbit brain PKC- β . The standards shown for both membranes are: (1) serum albumin (66.2 kDa); and (2) ovalbumin (45.0 kDa).

a

1

2

b

1

2

A B C D E

concentration of 120 mM (Fig. 9). This elution profile on hydroxylapatite is virtually the same as the profile for the β -isoform of PKC from mammalian tissues and suggests, therefore, that the single isoform of frog liver may be the β -isoforms of PKC. The fractions from this peak were pooled and used in the immunoblotting procedure to confirm the PKC isozyme type. Fig. 10a shows that staining with the antiPKC- α antibody was negative for frog liver PKC and positive for rabbit brain PKC; this indicates that frog liver PKC isozyme was not the α -isoform. Fig. 10b shows that staining with the antiPKC- β antibody gave a positive result for the frog liver PKC isozyme and a positive result for rabbit brain PKC. This strongly suggests that the single PKC isozyme found in frog liver was the β -isoform of PKC. However, since the liver isozyme was not tested for crossreactivity with a rabbit antiPKC- antibody the possibility that a portion of this single liver PKC peak may be the gamma PKC isoform cannot be ruled out.

The isozyme composition of *R. sylvatica* brain PKC was also determined by hydroxylapatite column chromatography. A typical chromatograph is shown in Fig. 11. Three peaks of enzyme activity were eluted at mean salt (NaP_i) concentrations of 60 mM, 100 mM, and 170 mM. The peaks were identified as the gamma (γ), beta (β), and alpha (α) isoforms of PKC, respectively, based on the nomenclature for mammalian brain PKC.

Fig. 11. Analysis of PKC isozymes of *R. sylvatica* brain using hydroxylapatite column chromatography. Brain (0.3 g) was homogenized 1:20 (w/v) in isozyme buffer pH 7.0. Isozyme buffer contained 10 mM β -mercaptoethanol, 20 mM NaP_i , 0.1 mM EGTA, 0.1 mM EDTA, and 20% glycerol (v/v). Crude homogenates from $n = 4$ separate preparations were loaded onto 4 individual hydroxylapatite columns. Brain PKC activity eluted in three separate peaks using a 20 mM - 2 M NaP_i salt gradient. PKC activity was monitored using the assay described in chapter 7. Data are means $\pm$ S.E.M. for $n = 4$ separate enzyme preparations.

DISCUSSION

It has been shown that diacylglycerol is only transiently produced in the membrane of cells due to phosphoinositide breakdown (Nishizuka, 1984). PKC is a unique protein kinase in that it apparently must be bound to the membrane of the cell in the presence of DAG to be physiologically active. Without the correct complement of membrane phospholipids and calcium the enzyme is inactive. Hence, changes in the distribution of PKC between membrane bound and free conditions should indicate the relative activity state of PKC under different physiological conditions (Table 26).

It was hypothesized that if the PKC signal transduction cascade played a part in the phosphorylation of glycolytic enzymes and the subsequent activation of liver glycogenolysis that liver PKC should show a translocation to the membrane in order to become fully activated, as is the case with mammalian forms of the enzyme. Initial studies showed that a 24 h freeze stimulated a large increase in the concentration of liver IP₃ (Table 26). The concentration of IP₃ increased 10-fold in frozen liver and began to return to control levels in thawed liver tissue, a result that supports the above hypothesis that *R. sylvatica* liver PKC is activated by a freezing stress. However, the fractionation procedure indicated that PKC was not translocated to the membrane during freezing, although predictably it should have been, if the enzyme was in an active state. Levels of membrane bound PKC actually dropped dramatically in 24 h frozen and 24 h thawed *R. sylvatica* liver (Table 26). Furthermore, the percentage of membrane bound enzyme in frog tissues, even under control conditions, was very low compared with a mean value of 31% bound in rat liver hepatocytes (Diaz-Guerra et al., 1991). It appears, then, from the data gathered in this study that the stress of freezing does not cause a translocation of PKC to the membrane in liver tissue. Perhaps, in *R. sylvatica* liver, the translocation of PKC is not needed for its activation. However, this evidence does support the proposal that PKAc is the primary protein kinase enzyme

responsible for the initial activation of GP, and therefore, glucose biosynthesis, during the early stages of the freezing process in the wood frog. Further research is required to determine whether PKC is involved in the long term activation of liver glycogenolysis and glucose production during the later stages of a freezing stress (e.g. at times greater than 1 d) in *R. sylvatica*.

IP₃, once released into the cytoplasm, is responsible for the release of Ca²⁺ from intracellular calcium stores (Berridge and Irvine, 1984). DAG and calcium function to activate PKC on the membrane surface. A large increase in liver IP₃ concentration would seem to indicate a probable concomitant rise in both liver DAG and calcium levels and hence an activation of PKC. Measurements of DAG and calcium are needed to confirm this hypothesis. However, as stated above, initial bound versus free studies of *R. sylvatica* liver PKC indicated that PKC translocation does not occur in this organ during a freezing episode.

An altogether different result was found in *R. sylvatica* brain tissue sampled at the same time points (Table 26). IP₃ levels and the percentage of PKC remained relatively constant in brain over the entire freezing event. The only similarity to wood frog liver PKC distribution was a decrease in % bound PKC in the thawed state. Furthermore, the percentage of membrane bound enzyme in frog brain, including control tissue, was very low compared with mean values of 55% bound in control rat brain (Anderson et al., 1993). Thus, it seems that freezing stress does not activate brain PKC. Instead what may be occurring is that the levels of IP₃ are kept high at all times in this tissue which ensures that brain PKC may always be in an active state despite the lack of translocation to the membrane. IP₃ concentrations in the brains of *R. sylvatica* exposed to dehydration and anoxia also remained constant (chapter 4 and 5). PKC and IP₃ concentrations are highest in vertebrate brains, whereas they are relatively low in other tissues such as liver, kidney, and skeletal muscle (Huang and Huang, 1986). Perhaps, since the levels of this second messenger do not change in the brain during

these various stresses, this data is a good indication that, at least in *R. sylvatica*, brain homeostasis is preserved under these three environmental extremes.

The data for liver are novel findings for a vertebrate form of PKC and challenge the traditional notion that an increase in cellular IP₃ and calcium concentrations cause a translocation of PKC to the membrane, initiating the activation of this kinase (Nishizuka, 1984). However, other corroborating evidence is beginning to appear that agree with the results of this study. Using a hypoglycemic rat brain model, Anderson et al. (1993) found no evidence of PKC translocation even after 40 min of exposure to severe hypoglycemia. Further studies are required to monitor the calcium flux (release) in brain cells during a freezing stress in *R. sylvatica*. Constant cellular calcium levels in brain tissues throughout this stress would demonstrate that PKC translocation is unlikely during freezing.

A possible interpretation of the fact that no PKC translocation was observed in either tissue is that any such effect may have occurred quickly within the early minutes of the freezing stress. In this experiment PKC translocation was monitored in tissues that were harvested from 24 h frozen *R. sylvatica* (Table 26). Evidence that the translocation of PKC to the membrane occurs rapidly both in liver and brain has been mounting (Diaz-Guerra et al., 1991; Louis et al., 1988). In both studies PKC redistribution from the cytosol to the membrane occurred in less than 5 min. Further studies should include the use of a freezing time course to determine whether the translocation of PKC to the plasma membrane transpires upon the initiation of a freezing event in *R. sylvatica*.

A glucose-induced translocation and activation of PKC has been shown to occur in many different mammalian cell cultures including pancreatic islets and aortic smooth muscle cells (Ganesan et al., 1990; Lee et al., 1989). Exposure of bovine vascular (retinal) cell cultures to the hyperglycemic conditions observed in diabetes mellitus, 20 mM glucose, stimulated the translocation of PKC to the membranous fraction of these

cells (Lee et al., 1989). However, this did not occur in 2 *R. sylvatica* tissues despite the large hyperglycemic conditions that exist both in liver and brain during freezing.

Perhaps in the wood frog, as glycolytic flux is restricted below PFK in order to promote glucose export, the increased production of glucose in liver suppresses PKC activation once the freezing process has been initiated. Kinetic experiments designed to test this hypothesis should include inhibition studies of PKC by glucose over the course of a freezing event.

Of the 9 known isoforms of PKC (Bell and Burns, 1991), only one isozyme was found in *R. sylvatica* liver as determined by hydroxylapatite chromatography (Fig. 9). A modified immunoblotting procedure demonstrated that liver PKC was the β -isoform and not the α -isoform (Fig. 10a and 10b). The fact that only one isozyme was found in wood frog liver is unusual. Most mammalian livers, for example rat, and other vertebrate classes like Reptilia (turtles) have more than a single PKC isozyme (Diaz-Guerra et al., 1991). Whether the β -isoform of liver PKC is the β I or the β II isozyme, or if both β -isozymes exist awaits further study. It has been hypothesized that oleate and other related fatty acids are able to activate hepatic PKC through the direct activation of the β -isozyme of PKC (Diaz-Guerra et al., 1991). The possibility that this biochemical mechanism may activate PKC during a freezing event in *R. sylvatica* is intriguing and needs further study. This mechanism may allow PKC to covalently modify GP via phosphorylation, activating the enzyme and initiating the breakdown of glycogen to glucose as a cryoprotectant.

Hydroxylapatite chromatography using *R. sylvatica* brain preparations revealed the same pattern of PKC isozyme distribution that has been detected in all vertebrate brain preparations (Huang et al., 1988; Qu et al., 1991). Three isozymes labelled type I, II, and III, or β , and α were found in *R. sylvatica* brain homogenates. The β -isoform was the most abundant in anuran brain, which was the case in mammalian brain as well (Qu et al., 1988). All *R. sylvatica* PKC brain isoforms (as well as the

liver isoform) fall under the category of Ca^{2+} and phospholipid-dependent protein kinase C enzymes. Kinetic studies on each isoform should be completed to distinguish each brain isozyme by their specific catalytic properties, providing additional supporting evidence that these 3 isozymes exist and have the same kinetic parameters as other brain PKC isozymes.

Ischemic stress produced an increase in the translocation of PKC to the membrane in rat brain preparations, but more interestingly, there was an immediate down-regulation (proteolytic degradation) of PKC activity following ischemic exposure (Wieloch et al., 1991). Perhaps the cellular changes that occur during a freezing episode in *R. sylvatica*, such as dehydration due to extracellular ice formation and increased solute concentrations due to cryoprotectant production in the form of glucose, change the membrane binding site for PKC to such an extent that PKC is unable to translocate to the membrane to become activated. The rearrangement of membrane phospholipids that presumably occurs upon freezing may have deleterious effects on the activity of this kinase while the animal remains in a frozen state.

It appears that a freezing stress in *R. sylvatica* does not initiate a redistribution of liver or brain PKC from the cytosol to the membrane, which has been shown to activate this kinase in other vertebrate systems. Indeed, only small significant decreases in % bound PKC in liver and brain were seen despite large increases in IP_3 in liver and no change in brain. The isoforms found were all calcium and phospholipid-dependent. Evidence has been presented which demonstrates that the PKC signal transduction cascade may have only a limited role in the activation of glycogenolysis during a freezing episode. In addition, this evidence may support the notion that it is the cyclic AMP-dependent protein kinase signal transduction pathway, not that of PKC, that initiates the production of glucose in response to a freezing episode.

In almost all instances an increase in cellular IP_3 results in a translocation of PKC to the membrane where it becomes activated, phosphorylating many different

cellular target proteins (Nishizuka, 1984). The fact that freezing increased IP_3 concentrations supports the contention that PKC is active during a freezing episode. However, further studies showed that there was no translocation of PKC to the membrane in two *R. sylvatica* tissues. Thus, accumulated data for IP_3 levels challenges the accepted view of PKC activation.

Perhaps this kinase is activated directly by other mechanisms, for example, by fatty acids. It may turn out that the PKA and PKC signal transduction pathways act synergistically to activate glycogenolysis at the phosphorylase locus to produce glucose, while inhibiting other regulatory enzymes in the glycolytic pathway in order to depress the animal's metabolism while in the frozen state. The activation of PKC may sustain glucose production in the later stages of a freezing stress in a number of ways including: (1) phosphorylating GP directly, (2) phosphorylating and inhibiting PFK, (3) inhibiting the enzyme responsible for the resynthesis of glycogen from glucose, glycogen synthase (GS), or (4) inhibiting other metabolic pathways so that carbon in the form of glycogen goes directly into the biosynthesis of glucose in the wood frog. Further kinetic studies must be undertaken in order to determine the exact mechanisms involved in these processes.

CHAPTER #8
GENERAL DISCUSSION

The synthesis of cryoprotectants from carbohydrate reserves, liver glycogen in freeze tolerant frogs and fat body glycogen in insects, is a well known biochemical process in many cold-hardy vertebrates and invertebrates (Storey and Storey, 1988; Zachariassen, 1985). All freeze tolerant animals have adapted a similar metabolic strategy which involves directing carbon flow from glycogen stores into the production of cryoprotectant sugars, such as glucose in *R. sylvatica* and glycerol in *Epiblema scudderiana* (Storey and Storey, 1988; Churchill and Storey, 1989).

Previous studies on freeze tolerant vertebrates provide overwhelming evidence that: (1) total GP activity is higher in livers of freeze tolerant (*R. sylvatica*) versus freeze intolerant (*R. pipiens*) frogs, (2) GP is phosphorylated into its active α form, once a freezing episode has started in the wood frog, (3) β -adrenergic hormone signals may initiate the activation of glycogenolysis in response to freezing, and 4) the apparent total activity increase of GP during freezing is due to the α to β conversion of the enzyme rather than any actual new synthesis of GP (Mommensen and Storey, 1992; Storey and Storey, 1992; Crerar et al., 1988). This evidence suggests that protein kinase enzymes may ultimately control the activation of liver glycogenolysis, leading to glucose biosynthesis, observed in wood frogs exposed to a freezing stress.

Initial investigations in this thesis were carried out to determine whether the relationships between protein kinase enzymes and their respective second messengers in response to a freezing stress were similar or different in a freeze tolerant vertebrate known to accumulate large amounts of cryoprotectant, *R. sylvatica*, and those that do not *T.s. elegans*, *C.p. marginata*, and *C.p. bellii*. The increases in the activities of PKAc and the levels of second messengers cAMP and IP₃ in liver suggest that the activation and release of these molecules is directly responsible for the coordination of cyroprotectant production in *R. sylvatica*.

Unlike freeze tolerant anurans, the percentage of free catalytic PKA decreased dramatically in the liver of *T.s. elegans*, *C.p. marginata*, and *C.p. bellii*. No large

accumulation of glucose was observed in any of the turtle organs when compared to *R. sylvatica*, which is consistent with previous results (Storey et al., 1988). Thus, at least for freeze tolerant reptiles there is no response by PKA to freezing. However, the increased concentrations of IP₃ in all organs, including liver, of these 3 freeze tolerant reptiles suggest that PKC may be the prime protein kinase enzyme system that controls the limited production of glucose in these species. Additionally, PKC may regulate other metabolic adaptations for freezing survival that are common to both turtles and frogs, such as those needed to regulate the decreases in both cell volume and whole organ water content that happen during freezing. However, without the activation of glycogenolysis via the PKA signal transduction cascade it is impossible to produce large quantities of glucose for the sole purpose of cryoprotection.

An activation of glycogenolysis at the GP enzyme locus is initiated as a response to ice crystallization on the body surface during freezing episodes in all freeze tolerant vertebrates, including frogs and turtles (Storey et al., 1988; Storey and Storey, 1992b; Storey et al., 1993). However, an inhibition of glycolytic flux at phosphofructokinase, a condition that promotes glucose synthesis in frogs, is not seen in freeze tolerant turtles (Storey et al., 1993). Although glucose levels do increase in turtle liver, the concentration of this sugar is not large enough to serve as a cryoprotectant. Instead, carbon generated by glycogenolysis during freezing in turtle liver may be largely routed through the normal glycolytic pathway to two fates: (1) lactate production as an end product under the ischemic stress of freezing, or (2) synthesis of amino acids, derived from TCA cycle intermediates, which may function as possible cryoprotectants. Further investigations, such as studies of the regulation of glycolytic enzymes by phosphorylation/dephosphorylation, in turtles are needed to provide answers to these hypotheses. For example, by incubating tissue homogenates with radioactive ³²P-ATP under conditions that promote the actions of endogenous protein kinases, specifically PKA and PKC, and then analyzing the resulting protein

phosphorylation patterns by SDS-polyacrylamide gel electrophoresis it would be possible to determine: (1) which protein kinase enzymes are responsible for phosphorylating target enzymes during a metabolic stress, (2) which target enzymes are phosphorylated, and (3) if the state of their phosphorylation changes at any point throughout a particular stress. Armed with the knowledge that the glycolytic enzymes GP and PFK are known control points that regulate glucose biosynthesis in *R. sylvatica*, initial experiments should investigate whether the phosphorylation of these two enzymes is necessary for the activation of cryoprotectant production in this and other freeze tolerant vertebrates.

Other environmental stresses that are experienced by anurans in nature are at least partially reflected during bouts of freezing in *R. sylvatica*. Extracellular freezing effectively results in cellular dehydration and also imposes oxygen limitation (hypoxia, then anoxia) on cells. Metabolic adaptations that allow the cells of freeze tolerant animals to deal with cell volume change or with anoxia may, therefore, be key to freezing survival. Comparisons of the protein kinase and second messenger responses involved in freezing with those elicited by dehydration (chapter 4) or anoxia (chapter 5) revealed that the responses to freezing and dehydration share many similarities whereas the responses to anoxia are distinctly different. The results show that the PKA signal transduction cascade response to dehydration in *R. sylvatica*, a freeze tolerant frog, is identical to that seen during freezing in this species. However, the protein kinase response to dehydration and anoxia in *R. pipiens*, a non-freeze tolerant frog, is very different than that for *R. sylvatica* subjected to the same two stresses. It is apparent that for the parameters assessed in this thesis, signal transduction at the level of second messengers and protein kinases and their effects on liver glycogenolysis, that dehydration and freezing elicit similar responses. Hence, these responses characterized in freezing are likely initiated by cellular volume regulatory mechanisms and may have evolved out of pre-existing mechanisms developed by anurans to deal with dehydration.

It takes an extended period of time at a low metabolic rate and subzero temperature to build up a substantial anoxia stress in frozen *R. sylvatica*. Therefore, the protein kinase and second messenger responses that are characterized during freezing in *R. sylvatica* do not seem to be ones that are elicited by anoxia stress in this animal. Only environmental stresses that cause a volume regulated response in *R. sylvatica* (freezing and dehydration) elicit the activation of the protein kinase enzymes that are responsible for controlling cryoprotectant production and/or eventual metabolic depression under these conditions. Thus, it must have been the natural desiccation tolerance of these animals that facilitated the development of the freeze tolerance strategy to survive over the winter months in what essentially amounts to a state of suspended animation. Although dehydration and freezing are obviously not entirely identical stresses, both stresses elicit extremely similar metabolic responses in *R. sylvatica* that are indicative of the development of these responses along similar evolutionary lines.

Kinetic data accumulated for the free catalytic subunit of *R. sylvatica* liver PKA revealed: (1) that PKAc activity at both high and low temperatures is neither inhibited by the ions from salts such as KCl and NaCl, nor by glucose, and (2) that an increase in the affinity of the enzyme for its substrates occurs at low temperatures. These data indicate that the enzyme is capable of supporting cryoprotectant synthesis in the freeze tolerant wood frog at low temperatures. *R. sylvatica* liver PKAc appears to function quite well in a cellular environment that contains high metabolite (glucose) and salt concentrations, both of which are known to occur during a freezing episode. These positive effects on PKAc activity at low temperatures may be one factor that contributes to an enhanced glycogenolysis leading to cryoprotectant synthesis in *R. sylvatica* during the early stages of a freezing episode. Additionally, the free catalytic subunit of PKA is known to phosphorylate numerous regulatory enzymes in a variety of different signal transduction pathways (Edelman et al. 1987). PKAc then, must play a critical role in

the regulation of glucose biosynthesis via glycogen phosphorylase phosphorylation while possibly inhibiting other metabolic pathways during freezing in the wood frog.

It appears that a freezing stress in *R. sylvatica* does not initiate a redistribution of liver and brain PKC from the cytosol to the membrane, which has been shown to activate this kinase in other vertebrate systems. Evidence has been presented which demonstrates that the PKC signal transduction cascade may have a limited role in the phosphorylation and activation of wood frog liver GP. In addition, this evidence further supports the notion that it is the cyclic AMP-dependent protein kinase signal transduction pathway, not that of PKC, that coordinates the initial production of glucose in response to a freezing episode in this animal. However, the activation of the PKC signal transduction cascade may have a dual function during a freezing episode in *R. sylvatica*. The first may be to sustain the long term biosynthesis of glucose, up to 48 h after freezing exposure. PKC could achieve this by phosphorylating the remaining liver GP that has not been previously activated by PKAc, converting the entire GP pool to its active α form. A second function for PKC may be to ensure that the accumulated liver and tissue glucose cryoprotectant pool is not converted back into glycogen over the winter months. PKC has been shown to be one of the several kinase enzymes that phosphorylates and inactivates liver glycogen synthase (Roach, 1990). Thus, by maintaining the phosphorylation state of GP and GS through the tight regulation of the PKC signal transduction cascade, *R. sylvatica* can keep the biosynthesis of glucose elevated during the freezing process and prevent its reconversion back to glycogen, until such a time that temperature conditions warrant this reconversion. Future studies should include the purification of PKAc and PKC from other freeze tolerant vertebrates, including both turtles and frogs. This would provide interesting comparisons with the principles suggested by these studies on the wood frog.

The data presented here challenges the accepted idea that translocation is required for the activation of PKC. An increase in cellular concentrations of IP₃ is

usually indicative of an activation of PKC. It is thought that this kinase can only be activated after it has translocated to the plasma membrane of a cell (Nishizuka, 1984). However, in wood frog liver IP_3 levels increase dramatically upon freezing whereas the percent bound PKC (percent translocated) decreases in the liver of frozen animals. Perhaps PKC in frog liver is activated directly by other mechanisms, for example, by free fatty acids, thus eliminating the need for PKC to translocate to the plasma membrane in order to become activated.

The present findings indicate that the PKA and PKC signal transduction pathways may act synergistically to activate liver glycogenolysis at the GP enzyme locus to produce glucose in freeze tolerant vertebrates. In *R. sylvatica* these kinases must inhibit other regulatory enzymes in the glycolytic pathway, such as PFK, to aid in the diversion of carbon into glucose accumulation and prevent the catabolism of the cryoprotectant. In other freeze tolerant vertebrates, such as the turtles *C.p. marginata* and *T.s. elegans*, PKA and PKC must also facilitate the shunting of carbon from glycogen to glucose. Glucose can then be utilized as an anaerobic fuel as these animals begin to depress their metabolic pathways upon entering a frozen state. Further kinetic studies must be undertaken in order to determine the exact mechanisms involved in these processes.

The responses of protein kinases examined in this thesis have attempted to elaborate the underlying principles of freezing-induced glycogenolytic activation via protein phosphorylation in a unique group of vertebrates categorized as freeze tolerant. The phosphorylation of glycolytic enzymes via protein kinases, specifically PKA and PKC, is responsible for the control of glycogenolysis and glucose cryoprotectant production during a freezing episode in *R. sylvatica*. The actions of these protein kinases during the early stages of the freezing process initiate an activation of glycogenolysis, an increase in the production of cryoprotectant, and support an extension of glucose biosynthesis even at late stages of freezing in the wood frog. In

addition, protein kinases may be responsible for the subsequent reduction in overall metabolic rate that occurs in *R. sylvatica* once it is in a frozen state.

A truly freeze tolerant vertebrate must have the metabolic machinery in place so that a rapid build up and distribution of cryoprotectant occurs just prior to, and during a freezing episode. It is clear from the data presented in this thesis that *R. sylvatica* relies on the actions of its signal transduction cascades, primarily those involving PKA and PKC, to regulate and control cyroprotectant production from liver glycogen. This is critical for coordinated control of ice formation and cryoprotectant synthesis and allows freeze tolerant organisms to survive in a frozen state for prolonged periods of time.

REFERENCES

- Anderson, L.G., D. Zhao, K.R. Dell, D.L. Severson, and R.N. Auer. 1993. Brain protein kinase C assay using MARCKS substrate reveals no translocation due to profound insulin-induced hypoglycemia. Brain Research. 606: 187-194.
- Bechtel, P.J., J.A. Beavo, and E.G. Krebs. 1977. Purification and characterization of the catalytic subunit of skeletal muscle adenosine 3':5'-monophosphate-dependent protein kinase. J. Biol. Chem. 252: 2691-2697.
- Belkin, D.A. 1963. Anoxia:tolerance in reptiles. Science. 139: 492-493.
- Bell, R.M. and D.J. Burns. 1991. Lipid activation of protein kinase C. J. Biol. Chem. 266: 4661-4664.
- Berman, D.I., A.N. Leirikh, and E.I. Mikhailova. 1984. Winter hibernation of the Siberian salamander *Hynobius keyserlingi*. J. Evol. Biochem. Physiol. 1984: 323-327.
- Berridge, M.J. 1987. Inositol triphosphate and diacylglycerol: two interacting second messengers. Ann. Rev. Biochem. 56: 159-193.
- Berridge, M.J. and R.F. Irvine. 1984. Inositol triphosphate, a novel second messenger in cellular signal transduction. Nature. 312: 315-321
- Bishoff, S.T., W.L. Combest, and L. I. Gilbert. 1990. Characterization of cAMP-dependent protein kinase from the pupal wing epidermis of *Manduca sexta*. Insect Biochem. 20: 397-404.
- Bourinet, E., F. Fournier, J. Nargeot, and P. Charnet. 1992. Endogenous *Xenopus*-oocyte Ca-channels are regulated by protein kinases A and C. FEBS Letters. 229: 5-9.
- Bradford, M.M. 1976. A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. Analyt. Biochem. 12: 248-253.
- Brooks, S.P.J. 1992. A simple computer program with statistical tests for the analysis of enzyme kinetics. Bio-Techniques. 13: 906-911.
- Brooks, S.P.J. and K.B. Storey. 1988. Anoxic brain function: molecular mechanisms of metabolic depression. Febs Lett. 232: 214-216.
- Brooks, S.P.J. and K.B. Storey. 1991. The role of protein kinases in anoxia tolerance in facultative anaerobes: purification and characterization of a protein kinase that phosphorylates pyruvate kinase. Biochem. et Biophys. Acta. 1073: 253-259.
- Brooks, S.P.J. and K.B. Storey. 1992. Bound and determined: a computer program for making buffers of defined ion concentrations. Analyt. Biochem. 201: 119-126.
- Brooks, S.P.J. and K.B. Storey. 1993. Purification and characterization of cyclic AMP dependent protein kinase from an estivating land snail. Submitted.

- Carpenter, J. and C. Crowe. 1988. The mechanism of cryoprotection of proteins by solutes. Cryobiol. 25: 244-255.
- Chakravarthy, B.R., A. Bussey, J.F. Whitfield, M. Sikorska, R.E. Williams, J.P. Durkin. 1991. The direct measurement of protein kinase C (PKC) activity in isolated membranes using a selective peptide substrate. Anal. Biochem. 196: 144-150.
- Chijiwa, T., A. Mishima, M. Hagiwara, M. Sano, K. Hayashi, T. Inoue, K. Naito, T. Tashioaka, and H. Hidaka. 1990. Inhibition of forskolin-induced neurite outgrowth and protein phosphorylation by a newly synthesized selective inhibitor of cyclic AMP-dependent protein kinase, N-[2-(p-bromocinnamylamino)ethyl]-5-isoquinolinesulfonamide (H-89), of PC12D pheochromocytoma cells. J. Biol. Chem. 265: 5267-5272.
- Chivia, J.C., M.D. Uhler, and G.S. McKnight. 1988. Characterization of genomic clones coding for the C α and C β subunits of mouse cAMP-dependent protein kinase. J. Biol. Chem. 263: 5739-5744.
- Christiansen, J. and D. Penney. 1973. Anaerobic glycolysis and lactic acid accumulation in cold submerged *Rana pipiens*. J. Comp. Physiol. 87: 237-245.
- Churchill, T.A. and K.B. Storey. 1989. Regulation of glycerol biosynthesis in a freeze avoiding insect. J. Comp. Physiol. B. 159: 461-472.
- Churchill, T.A. and K.B. Storey. 1991. Metabolic responses to freezing by organs of hatchling painted turtles *Chrysemys picta marginata* and *C.p. bellii*. Can. J. Zool. 69: 2978-2984.
- Churchill, T.A. and K.B. Storey. 1992a. Responses to freezing exposure of hatchling turtles *Trachemys scripta elegans*: factors influencing the development of freeze tolerance by reptiles. J. exp. Biol. 167: 221-233.
- Churchill, T.A. and K.B. Storey. 1992b. Freezing survival of the garter snake *Thamnophis sirtalis parietalis*. Can. J. Zool. 70: 99-105.
- Churchill, T.A., and K.B. Storey, 1993. Dehydration tolerance in wood frogs: a new perspective on the development of amphibian freeze tolerance. Am. J. Physiol. 265: R1324-R1332.
- Churchill, T.A., and K.B. Storey, 1994. Metabolic effects of dehydration on an aquatic frog *Rana pipiens*. J. Exp. Biol. in press.
- Costanzo, J.P. and D.L. Claussen. 1990. Natural freeze tolerance in the terrestrial turtle, *Terrapene carolina*. J. Exp. Zool. 254: 228-232.
- Costanzo, J.P., D.L. Claussen, and R.E. Lee. 1988. Natural freeze tolerance in a reptile. Cryo. Lett. 9: 380-385.
- Costanzo, J.P., R.E. Lee, and P.H. Lortz. 1993. Glucose concentration regulates freeze tolerance in the wood frog *Rana sylvatica*. J. Exp. Biol. 181: 245-255.

- Costanzo, J.P., M.F. Wright, and R.E. Lee. 1990. Freeze tolerance as an overwintering adaptation in Cope's gray tree frog (*Hyla chrysoscelis*). Cryobiol. 27: 678 (abstract).
- Crerar, M.M., E.S. David, and K.B. Storey. 1988. Electrophoretic analysis of liver glycogen phosphorylase activation in the freeze tolerant wood frog. Biochim. Biophys. Acta. 971: 72-84.
- Diaz-Guerra, M.J.M., M. Junco, and L. Bosca. 1991. Oleic Acid promotes changes in the subcellular distribution of protein kinase C in isolated hepatocytes. J. Biol. Chem. 266: 23568-23576.
- Duman, J.G. 1982. Insect antifreezes and ice-nucleating agents. Cryobiol. 19: 613-627.
- Edelman, A.M., D.K. Blumenthal, and E.G. Krebs. 1987. Protein serine/threonine kinases. Ann. Rev. Biochem. 56: 567-613.
- Epand, R.M. and D.S. Lester. 1990. The role of membrane biophysical properties in the regulation of protein kinase C activity. Trends Pharmacol. Sci. 11:317-320.
- Foster, J.L. and L.M. Hall. 1988. Purification of *Drosophila* cAMP-dependent protein kinase. Meth. in Enzymol. 159: 215-224.
- Ganesan, S., R. Calle, K. Zawulich, J.I. Smallwood, W.A. Zawulich, and H. Rasmussen. 1990. Glucose-induced translocation of protein kinase C in rat pancreatic islets. Proc. Natl. Acad. Sci. 87: 9893-9897.
- Grinstein, S., W. Furuya, and L. Bianchini. 1992. Protein kinases and phosphatases, and the control of cell volume. NIPS. 7: 232-236.
- Haracska, L. and A. Udvardy. 1992. Purification and characterization of the catalytic subunit of cAMP dependent protein kinase from *Drosophila melanogaster*. Insect Biochem. Molec. Biol. 22: 851-858.
- Hashimoto, E. and H. Yamamura. 1991. Protease-activated protein kinase C in rat liver. Int. J. Biochem. 23: 507-512.
- Hers, H.G. 1976. The control of glycogen in the liver. Ann. Rev. Biochem. 45: 167-189.
- Hillman, S. 1980. Physiological correlates of differential dehydration tolerance in anuran species. Copeia. 1: 125-129.
- Hillman, S. 1987. Dehydration effects on cardiovascular and metabolic capacity in two amphibians. Physiol. Zool. 60: 608-613.
- Hillman, S. 1988. Dehydration effects on brain and cerebrospinal fluid electrolytes in two amphibians. Physiol. Zool. 61: 254-259.
- Hofmann, F., J.A. Beavo, P.J. Bechtel, and E.G. Krebs. 1975. Comparisons of adenosine 3':5'-monophosphate-dependent protein kinases from rabbit skeletal and bovine heart muscle. J. Biol. Chem. 250: 7795-7801.

- Hotz, A., N. Konig, J. Kretschmer, G. Maier, H. Ponstingl, and V. Kinzel. 1989. A sequence variant in the N-terminal region of the catalytic subunit of the cAMP-dependent protein kinase. FEBS Lett. **256**: 115-117.
- Huang, K.-P. and F.L. Huang. 1986. Immunochemical characterization of rat brain protein kinase C. J. Biol. Chem. **261**: 14781-14787.
- Hunter, T. 1991. Protein kinase classification. Meth. Enzymol. **200**: 3-37.
- Hutchinson, V.H. and M.J. Dady. 1964. The viability of *Rana pipiens* and *Bufo terrestris* submerged at different temperatures. Herpt. **20**: 149-162.
- Jiang, H. and J.D. Corbin. 1991. Effect of epinephrine or cAMP on cAMP-bound protein kinase holoenzymes in rat heart. Am. J. Physiol. **260**: H722-H729.
- Katz, U. 1989. Strategies of adaptation to osmotic stress in anuran amphibia under salt and burrowing conditions. Comp. Biochem. Physiol. **93**: 499-503.
- Keiver, K.M. and P.W. Hochachka. 1991. Catecholamine stimulation of hepatic glycogenolysis during anoxia in the turtle *Chrysemys picta*. Am. J. Physiol. **261**: R1341-R1345.
- Kelly, D.A. and K.B. Storey. 1988. Organ-specific control of glycolysis in anoxic turtles. Am. J. Physiol. **255**: R774-R779.
- Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. Nature. **227**: 680-685.
- Lee, R.E., J.P. Costanzo, E.C. Davidson, and J.R. Layne. 1992. Dynamics of body water during freezing and thawing in a freeze-tolerant frog (*Rana sylvatica*). J. Therm. Biol. **17**: 263-266.
- Lee, T.-S., K.A. Saltsman, H. Ohashi, and G.L. King. 1989. Activation of protein kinase C by elevation of glucose concentration: Proposal for a mechanism in the development of diabetic vascular complications. Proc. Natl. Acad. Sci. **86**: 5141-5145.
- Livesey, S.A. and T.J. Martin. 1988. Selective activation of the cAMP-dependent protein kinase isozymes. Meth. Enzymol. **159**: 105-118.
- Louis, J.-C. E. Magal, E. Yavin. 1988. Protein kinase C alterations in the fetal brain after global ischemia. J. Biol. Chem. **263**: 19282-19285.
- Lutz, P.L., P. McMahon, M. Rosenthal, and T.J. Sick. 1984. Relationships between aerobic and anaerobic energy production in turtle brain in situ. Am. J. Physiol. **247**: R740-R744.
- McGann, L., H. Yang, and M. Walterson. 1988. Manifestations of cell damage after freezing and thawing. Cryobiol. **25**: 178-185.
- Mehrani, H. and K.B. Storey. 1994. cAMP-dependent protein kinase and anoxia survival in turtles: purification and properties of liver PKA. Submitted to the J. Biol. Chem.

- Mommsen, T.P. and K.B. Storey. 1992. Hormonal effects on glycogen metabolism in isolated hepatocytes of a freeze-tolerant frog. Gen. and Comp. Endocrin. 87: 44-53.
- Nielson, B.L. and L.R. Brown. 1984. The basis for colored silver-protein complex formation in stained polyacrylamide gels. Anal. Biochem. 141: 311-315.
- Nishizuka, Y. 1984. The role of protein kinase C in cell surface signal transduction and tumor promotion. Nature. 308: 693-697.
- Nishizuka, Y. 1986. Studies and perspectives of protein kinase C. Science. 233: 305-312.
- Okuno, S. and H. Fujisawa. 1990. Stabilization, purification, and crystallization of catalytic subunit of cAMP-dependent protein kinase from bovine heart. Biochem. Biophys. Acta. 1038: 204-208.
- Plaxton, W.C. and K.B. Storey. 1984. In vivo phosphorylation of red muscle pyruvate kinase from the channeled whelk, *Busycotypus canaliculatum*, in response to anoxic stress. Eur. J. Biochem. 143: 257-265.
- Poczopko, P. 1959. Respiratory exchange in *Rana esculenta* L. in different respiratory media. Zool. Pol. 10: 45-55.
- Qu, Y., J. Torchia, T.D. Phan, and A.K. Sen. 1991. Purification and characterization of protein kinase C isozymes from rat heart. Mol. Cell. Biochem. 103: 171-180.
- Rasmussen, H. 1989. The cycling of calcium as an intracellular messenger. Scientific American (October). pp. 66-73.
- Roach, P.J. 1990. Control of glycogen synthase by hierarchical protein phosphorylation. FASEB. 4: 2961-2968.
- Reimann, E.M. and R.A. Beham. 1983. Catalytic subunit of cAMP-dependent protein kinase. Meth. Enzymol. 99: 51-55.
- Roskoski Jr., R. 1983. Assays of protein kinase. Meth. Enzymol. 99: 3-6.
- Schmid, W.D. 1982. Survival of frogs at low temperature. Science. 215: 697-698.
- Shenolikar, S. 1988. Protein phosphorylation: hormones, drugs, and bioregulation. FASEB. J. 2: 2753-2764.
- Shoemaker, V.H. 1992. Exchange of water, ion, and respiratory gases in terrestrial amphibians. In: *Environmental Physiology of the Amphibians*. ed. Feder, M.E. and W.W. Burggren. Univ. of Chicago Press. Chicago, IL. pp. 125-159.
- Storey, J.M. and K.B. Storey. 1985a. Freeze tolerance in the grey tree frog *Hyla versicolor*. Can. J. Zool. 63: 49-54.
- Storey, J.M. and K.B. Storey. 1985b. Triggering of cryoprotectant synthesis by the initiation of ice nucleation in the freeze tolerant frog, *Rana sylvatica*. J. Comp. Physiol. B. 156: 191-195.

- Storey, K.B. 1984. Freeze tolerance in the frog *Rana sylvatica*. Experientia. 40: 1261-1262.
- Storey, K.B. 1987a. Glycolysis and the regulation of cryoprotective synthesis in liver of the freeze-tolerant wood frog. J. Comp. Physiol. B. 157: 373-380.
- Storey, K.B. 1987b. Organ specific metabolism during freezing and thawing in a freeze-tolerant frog. Am. J. Physiol. 253: R292-R297.
- Storey, K.B. 1988. Mechanisms of glycolytic control during facultative anaerobiosis in a marine mollusc: tissue specific analysis of glycogen phosphorylase and fructose-2,6-bisphosphate. Can. J. Zool. 66: 1767-1771.
- Storey, K.B. 1990. Life in a frozen state: adaptive strategies for natural freeze tolerance in amphibians and reptiles. Am. J. Physiol. 258 (Regulatory Integrative Comp. Physiol. 27: R559-R568.
- Storey, K.B. 1993. Molecular mechanisms of metabolic arrest in mollusks. In: Surviving hypoxia: Mechanisms of control and adaptation. edited by P.W. Hochachka, P.L. Lutz, T. Sick, M. Rosenthal, and G. van den Thillart. CRC Press, Boca Raton FA. pp. 253-269.
- Storey, K.B. and J.M. Storey. 1984. Biochemical adaptation for freezing tolerance in the wood frog *Rana sylvatica*. J. Comp. Physiol. B. 155: 29-36.
- Storey, K.B. and J.M. Storey. 1986. Freeze-tolerant frogs: Cryoprotectants and tissue metabolism during freeze/thaw cycles. Can. J. Zool. 64: 49-56.
- Storey, K.B. and J.M. Storey. 1988. Freeze tolerance in animals. Physiol. Rev. 68: 27-84.
- Storey, K.B. and J.M. Storey. 1990. Metabolic rate depression and biochemical adaptation in anaerobiosis, hibernation, and estivation. The Quarterly Review of Biology. 65: 145-174.
- Storey, K.B. and J.M. Storey. 1992a. Biochemical adaptations for winter survival in insects. In: Advances in Low-Temperature Biology. JAI Press Ltd. Vol 1, pp. 101-140.
- Storey, K.B. and J.M. Storey. 1992b. Natural freeze tolerance in ectothermic vertebrates. Annu. Rev. Physiol. 54: 619-637.
- Storey, K.B., and J.M. Storey, 1993. Cellular adaptations for freezing survival by amphibians and reptiles. In: Advances in Low-Temperature Biology. ed. P. Steponkus. JAI Press, London, pp. 101-129.
- Storey, K.B., D.G. McDonald, J.G. Duman, and J.M. Storey. 1991. Blood chemistry and ice nucleating activity in hatchling painted turtles. Cryo-Lett. 12: 351-358.
- Storey, K.B., J.M. Storey, S.P.J. Brooks, T.A. Churchill, and R.J. Brooks. 1988. Hatchling turtles survive freezing during winter hibernation. Proc. Nat. Acad. Sci. 85: 8350-8354.

- Storey, K.B., J.R. Layne Jr., M.M. Cutwa, T.A. Churchill, and J.M. Storey. 1993. Freezing survival and metabolism of box turtles *Terrapene carolina*. Copeia. 3: 628-634.
- Stryer, L. 1981. Biochemistry. 2nd Edition. W.H. Freeman and Co. New York.
- Suelter, C.H. 1985. A practical guide to enzymology. John Wiley & Sons, Toronto. pp. 174-177.
- Sugden, P.H., L.A. Holladay, E.R. Reimann, and J.D. Corbin. 1976. Purification and characterization of the catalytic subunit of adenosine 3':5'-cyclic monophosphate-dependent protein kinase from bovine liver. Biochem. J. 159: 409-422.
- Takhar, S. and M.R. Munday. 1992. Evidence for different isozymic forms of catalytic subunit of cyclic AMP-dependent protein kinase in heart and lactating mammary gland of the rat. Biochem. Soc. Trans. 20: 307S.
- Taylor, S.S., J.A. Beuchler, and D.R. Knighton. 1990. Cyclic AMP-dependent protein kinase: Mechanism for ATP:protein phosphotransfer. In: Peptides and protein phosphorylation. ed. B. Kemp. CRC Press, Boca Raton, FA. pp. 1-41.
- Thorson, T.B. 1955. The relationship of water economy to terrestrial amphibians. Ecology. 36: 100-116.
- Tremblay, J., R. Gerzer, and P. Hamet. 1988. Cyclic GMP in cell function. Adv. Sec. Mess. Phos. Res. 22: 319-350.
- Walsh, D.A., C.D. Ashby, C. Gonzalez, D. Calkins, E.H. Fischer, and E.G. Krebs. 1971. Purification and characterization of a protein inhibitor of adenosine 3',5'-monophosphate-dependent protein kinases. J. Biol. Chem. 246: 1977.
- Watson, P.A. 1991. Function follows form: generation of intracellular signals by cell deformation. FASEB. 5: 2013-2019.
- Wieloch, T., M. Cardell, H. Bingren, J. Zivin, and T. Saitoh. 1991. Changes in the activity of protein kinase C and the differential subcellular redistribution of its isozymes in the rat striatum during and following transient forebrain ischemia. J. Neurochem. 56: 1227-1235.
- Wolanczyk, J.R., J.G. Baust, and K.B. Storey. 1990. Seasonal ice nucleating activity in the freeze tolerant frog *Rana sylvatica*. Cryo. Lett. 11: 143-150.
- Zachariassen, K.E. 1985. Physiology of cold tolerance in insects. Physiol. Rev. 65: 799-832.
- Zoller, M.J., A.R. Kerlavage, and S.S. Taylor. 1979. Structural comparisons of cAMP-dependent protein kinase I and II from porcine skeletal muscle. J. Biol. Chem. 254: 2408-2412.

PUBLICATION LIST

- Holden, C.P. and Storey, K.B. 1994. 6-Phosphogluconate dehydrogenase from the freeze-tolerant gall fly larvae *Eurosta solidaginis* and its control of the Pentose Phosphate Pathway. Insect Biochem. and Molec. Biol. 24: 167-173.
- Holden, C.P. and Storey, K.B. 1994. Purification and characterization of aldolase in the freeze-avoiding gall moth larvae *Epiblema scudderiana* and its controlling factors during glycerol biosynthesis. Insect Biochem. and Molec. Biol. 24: 265-270.
- Brooks, S.P.J., Mancebo, M.J., Holden, C.P. and Storey, K.B. 1993. A batch elution procedure for assaying adenylate cyclase. Analyt. Biochem. 210: 419-421.
- Holden, C.P. and Storey, K.B. 1993. Purification and characterization of glycogen phosphorylase a and b from the freeze-avoiding gall moth larvae *Epiblema scudderiana*. J. Comp. Physiol. 163: 499-507.
- Holden, C.P. and Storey, K.B. 1993. The kinetic parameters of purified phosphofructokinase (PFK) during glycerol biosynthesis in the freeze-avoiding gall moth larvae *Epiblema scudderiana*. Can. J. Zool. 71: 1895-1899.
- Holden, C.P. and Storey, K.B. 1995. Control of glycerol catabolism in the freeze-avoiding gall moth larvae *Epiblema scudderiana* by the gluconeogenic enzyme fructose-1,6-bisphosphatase (FBPase). Accepted to Archives of Insect Biochemistry and Physiology.

PM-1 3½"x4" PHOTOGRAPHIC MICROCOPY TARGET
NBS 1010a ANSI/ISO #2 EQUIVALENT

PRECISIONSM RESOLUTION TARGETS

END

29-05-95

FIN