LIST OF FIGURES

Figure	Title	page
1	Structure and Numbering Schemes for the Major Nucleic Acid Components	5
2	Structural Components of Cytidylyl-(3'-5')-Guanosine ...	7
3	¹ H NMR Chemical Shift Range	22
4	³¹ P NMR Chemical Shift Range	23
5	¹⁹⁹ Hg NMR Chemical Shift Range	25
6	¹³ C NMR Chemical Shift Range	27
7	¹⁵ N NMR Chemical Shift Range	30
8	¹⁵ N NMR Results. Effects of Divalent Metal Ions on the Chemical Shifts of Guanosine	49
9	¹⁵ N NMR Results. Effects of Divalent Metal Ions on the Chemical Shifts of Inosine	50 :)
10	¹⁵ N NMR Spectrum of Inosine	51
11	¹⁵ N NMR Results. Effect of Protonation on the Chemical Shifts of Guanosine	53
12	¹⁵ N NMR Results. Effect of Protonation of the Chemical Shifts of Inosine	54
13	¹³ C NMR Spectrum of Inosine, Fully Coupled	56
14	¹³ C NMR Spectrum of Inosine, ¹ H Decoupled	57
15	¹³ C NMR Results. Effects of Divalent Metal Ions on the Chemical Shifts of Inosine	58
16	¹³ C NMR Results. Effects of Divalent Metal Ions on the Chemical Shifts of Guanosine	59
17	¹³ C NMR Results. Effect of Protonation on the Chemical Shifts of Guanosine	63
18	¹³ C NMR Results. Effect of Protonation on the Chemical Shifts of Inosine	64
19	¹³ C NMR Results. Effect of N7-Methylation on the Chemical Shifts of Guanosine	67

National Library
of Canada

Bibliothèque nationale
du Canada

Canadian Theses Service

Services des thèses canadiennes

Ottawa, Canada
K1A 0N4

CANADIAN THESES

THÈSES CANADIENNES

NOTICE

The quality of this microfiche is heavily dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to ensure the highest quality of reproduction possible.

If pages are missing, contact the university which granted the degree.

Some pages may have indistinct print especially if the original pages were typed with a poor typewriter ribbon or if the university sent us an inferior photocopy.

Previously copyrighted materials (journal articles, published tests, etc.) are not filmed.

Reproduction in full or in part of this film is governed by the Canadian Copyright Act, R.S.C. 1970, c. C-30. Please read the authorization forms which accompany this thesis.

AVIS

La qualité de cette microfiche dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

S'il manque des pages, veuillez communiquer avec l'université qui a conféré le grade.

La qualité d'impression de certaines pages peut laisser à désirer, surtout si les pages originales ont été dactylographiées à l'aide d'un ruban usé ou si l'université nous a fait parvenir une photocopie de qualité inférieure.

Les documents qui font déjà l'objet d'un droit d'auteur (articles de revue, examens publiés, etc.) ne sont pas microfilmés.

La reproduction, même partielle, de ce microfilm est soumise à la Loi canadienne sur le droit d'auteur, SRC 1970, c. C-30. Veuillez prendre connaissance des formules d'autorisation qui accompagnent cette thèse.

THIS DISSERTATION
HAS BEEN MICROFILMED
EXACTLY AS RECEIVED

LA THÈSE A ÉTÉ
MICROFILMÉE TELLE QUE
NOUS L'AVONS REÇUE

National Library
of Canada

Bibliothèque nationale
du Canada

0-315-22240-9

Canadian Theses Division Division des thèses canadiennes

Ottawa, Canada
K1A 0N4

PERMISSION TO MICROFILM — AUTORISATION DE MICROFILMER

• Please print or type — Écrire en lettres moulées ou dactylographier

Full Name of Author — Nom complet de l'auteur

MARY-JANE CONNOR BELL

Date of Birth — Date de naissance

20-04-1957

Country of Birth — Lieu de naissance

CANADA

Permanent Address — Résidence fixe

424 MCCOWIE GREEN
NEPEAN, ONT
K2G 4H4

Title of Thesis — Titre de la thèse

A MULTINUCLEAR NMR STUDY OF THE INTERACTION BETWEEN
METAL IONS AND NUCLEIC ACID DERIVATIVES IN SOLUTION

University — Université

CARLETON

Degree for which thesis was presented — Grade pour lequel cette thèse fut présentée

Ph.D.

Year this degree conferred — Année d'obtention de ce grade

1985

Name of Supervisor — Nom du directeur de thèse

DR G.W. BUCHANAN

Permission is hereby granted to the NATIONAL LIBRARY OF CANADA to microfilm this thesis and to lend or sell copies of the film

The author reserves other publication rights, and neither the thesis nor extensive extracts from it may be printed or otherwise reproduced without the author's written permission.

L'autorisation est, par la présente, accordée à la BIBLIOTHÈQUE NATIONALE DU CANADA de microfilmer cette thèse et de prêter ou de vendre des exemplaires du film

L'auteur se réserve les autres droits de publication; ni la thèse ni de longs extraits de celle-ci ne doivent être imprimés ou autrement reproduits sans l'autorisation écrite de l'auteur

Date

04-09-85

Signature

M. J. Bell

A Multinuclear NMR Study
on the Interactions
Between Metal Ions and
Nucleic Acid Derivatives in Solution.

by

© Mary-Jane Bell, B.Sc., M.Sc.

A thesis submitted to the
Faculty of Graduate Studies and Research
in partial fulfilment of
the requirements for the degree of

Doctor of Philosophy
Department of Chemistry.

Carleton University
Ottawa, Ontario
June 1985

© copyright
1985 Mary-Jane Bell

The undersigned hereby recommend to
The Faculty of Graduate Studies and Research
acceptance of the thesis, submitted by

Mary-Jane Bell, B.Sc., M.Sc.

in partial fulfilment of the requirements
for the degree of Doctor of Philosophy.

.....
Chairman, Department of
Chemistry

.....
Thesis Supervisor

.....
External Examiner

Carleton University
August 1985

ABSTRACT

A multinuclear NMR approach has been taken to investigate the metal ion complexation sites for a number of nucleic acid constituents. The regioselectivity of complexation can be obtained from analysis of the ^{13}C , ^{31}P , ^{15}N and ^{199}Hg NMR spectral data.

The complexations of the soft metal ions Hg(II) and $\text{CH}_3\text{Hg(II)}$, the borderline species Zn(II) , and the hard metal ions Ba(II) and Mg(II) were compared. The ligands involved in this study included the 6-oxopurine nucleosides and nucleotides, the adenine nucleosides and nucleotides, and the dinucleoside monophosphates; cytidylyl-(3'→5')-guanosine and 2'-deoxyribosecytidylyl-(3'→5')-2'-deoxyriboseguanosine.

In general the binding site occupied is dependent upon the polarizability of the metal ion, the solvent used; whether protic or aprotic, and the pH of the solution.

In all cases, metal ion binding to the nucleic acid derivatives was deemed to be under kinetic control, since mercury-nitrogen and mercury-oxygen bonds are known to be kinetically labile.

In the 6-oxopurine nucleosides the formation of a carbonyl complex with the hard metal ions was detected. The softer metal ions interacted with the N7 nitrogen binding site.

In aqueous solutions, at pH 8, the 6-oxopurine nucleotides exhibited an N1 binding site for the soft metal ions, involving deprotonation followed by metal ion binding. The hard metal ions were involved in outer-sphere binding at the N7 position.

The adenosine species appear to have two occupied nitrogen binding sites, with the site selectivity being governed by the nature of the metal ions.

Phosphate complexation is not observed in the monophosphates. It does, however occur with the triphosphate derivative of adenosine, where a number of pH and metal ion dependent sites have been detected using ^{31}P NMR.

In addition the metal ion effects on the conformation of two dinucleoside monophosphates were determined. The results of these model compound investigations can be extended to the RNA and DNA level.

ACKNOWLEDGEMENTS

I would like to thank Dr. G.W. Buchanan for the support and guidance provided over the duration of this work. I would also like to thank Dr. Buchanan, in association with Dr. J.B. Stothers at the University of Western Ontario, London, for the initial ^{15}N NMR spectra of the purine nucleoside-metal ion complexes.

I would like to thank Keith Bourque for the assistance provided in obtaining NMR spectra at Carleton University and Raj Capoor at Ottawa University for the ^{15}N spectra at 30.42 MHz.

The 500 MHz, ^1H NMR spectra of the dinucleosides were provided by Dr. D. Bundle, Division of Biological Sciences, National Research Council, Ottawa.

I would also like to thank Eugene Florio for the samples of 7-methylguanosine prepared in this laboratory.

TABLE OF CONTENTS

	page
ACCEPTANCE SHEET	ii
ABSTRACT	iii
ACKNOWLEDGEMENTS	v
TABLE OF CONTENTS	vi
LIST OF TABLES	ix
LIST OF FIGURES	x
 CHAPTER 1: Introduction	 1
CHAPTER 2: Nucleic Acids and Metal Ions	
2.1 Physical and Chemical Properties of Nucleic Acid Derivatives	4
2.2 Interaction of Metal Ions with Nucleic Acid Derivatives	8
2.3 Properties of Metal Ions in Solution	10
2.4 Potential Metal Ion Binding Sites on Nucleic Acid Derivatives	11
 CHAPTER 3: Multinuclear NMR	
3.1 Basic Theory of NMR	14
3.2 Specific Nuclei of Interest	20
3.21 ^1H NMR	20
3.22 ^{31}P NMR	21
3.23 ^{199}Hg NMR	24
3.24 ^{13}C NMR	26
3.25 ^{15}N NMR	28
3.3 Theory of Chemical Shifts	32
3.31 Pople's Formula	34
3.32 Chemical Shifts and Binding	40
 CHAPTER 4: 6-Oxopurine Nucleosides	 45
4.1 ^{15}N NMR Results	
4.11 Divalent Metal Ion Complexes	48
4.12 Protonation	52
4.2 ^{13}C NMR Results	
4.21 Divalent Metal Ion Complexes	55
4.22 Protonation	62
4.3 7-Methylguanosine	66
4.4 Conclusion	72
 CHAPTER 5: 6-Oxopurine Nucleotides	 74
5.1 ^{13}C NMR Results	77
5.2 ^{15}N NMR Results	80
5.3 ^{31}P NMR Results	80
5.4 Conclusion	83

	page
CHAPTER 6: Adenosine	85
6.1 ^{15}N NMR Results	87
6.2 ^{13}C NMR Results	90
6.3 Conclusion	93
CHAPTER 7: Adenosine Nucleotides	
7.1 Adenosine-5'-monophosphate	95
7.11 ^{15}N NMR Results	96
7.12 ^{13}C NMR Results	98
7.13 ^{31}P NMR Results	98
7.14 Conclusion	101
7.2 Metal Ion Complexes of Adenosine-5'-Triphosphate ...	102
7.21 ^{15}N NMR Results	103
7.22 ^{13}C NMR Results	105
7.23 ^{31}P NMR Results	108
7.24 Conclusion	113
7.3 Dephosphorylation of Adenosine-5'-Triphosphate	115
CHAPTER 8: ^{199}Hg NMR Results	119
CHAPTER 9: Dinucleoside Monophosphates	124
9.1 Homonuclear Chemical Shift Correlation of CpG	127
9.2 ^{13}C Chemical Shift Assignments of CpG	131
9.3 Conformation of CpG	140
9.31 ϵ_1 Bond Angle	141
9.32 Sugar Pucker	143
9.33 ϕ_2 Bond Angle	147
9.34 Conclusion	149
9.4 ^1H NMR dCpG	150
9.5 ^{13}C Chemical Shift Assignments of dCpG	152
9.6 Conformation of dCpG	157
9.61 Sugar Pucker	157
9.62 ϵ_1 Bond Angle	158
9.63 ϕ_2 Bond Angle	159
9.64 Conclusion	159
CHAPTER 10: Metal Ion Binding to Dinucleoside Monophosphates ..	161
10.1 Mononucleotide Model Compounds	
10.11 ^{15}N NMR Results	164
10.12 ^{13}C NMR Results	168
10.13 ^{31}P Conformational Information	172
10.2 Dinucleoside Monophosphates	
10.21 ^{13}C NMR Results	173
10.22 ^{31}P Conformational Information	177
10.3 Conclusion	180
CHAPTER 11: Conclusion	182

	page
CHAPTER 12: Experimental	
12.1 General FT-NMR Methods	188
12.2 Spectral Acquisition Parameters	191
12.3 INEPT Pulse Sequence	194
12.4 Two-Dimensional NMR	198
12.41 Homonuclear Chemical Shift Correlation Spectroscopy	199
12.5 Materials and Sample Preparation	202
12.6 Synthesis of Guanosine-N1-methylmercuric-5'- monophosphate	203
REFERENCES	204

LIST OF TABLES

Table	Title	page
1	pKa's of Nucleotide and Nucleoside Constituents	6
2	Magnetic Properties of Observed Nuclei	20
3	¹³ C Shift Data and ¹ H Coupling Information for the Base Carbons of CpG	135
4	¹³ C- ³¹ P Coupling Constants of CpG	137
5	¹³ C Chemical Shifts for CpG and the Mononucleotide Models	138
6	¹³ C- ³¹ P Coupling Constants of 3' Cytidylyl Fragments	142
7	³ E Sugar Conformer Populations for the Ribose Derivatives	144
8	P _t Populations of the Ribose Mononucleotides and CpG	148
9	¹³ C Chemical Shifts for dCpG and the Mononucleotide Models	154
10	¹³ C- ³¹ P Coupling Constants dCpG	155
11	³ E Sugar Conformer Population for the Deoxyribose Derivatives	157
12	¹³ C- ³¹ P Coupling Constants of the 3' dCytidylyl Fragments	158
13	Effect of Methylmercury(II) hydroxide on the ¹³ C- ³¹ P Coupling Constants of the 3' Cytidylyl Residue of CpG	178

LIST OF FIGURES

Figure	Title	page
1	Structure and Numbering Schemes for the Major Nucleic Acid Components	5
2	Structural Components of Cytidylyl-(3'-5')-Guanosine ...	7
3	¹ H NMR Chemical Shift Range	22
4	³¹ P NMR Chemical Shift Range	23
5	¹⁹⁹ Hg NMR Chemical Shift Range	25
6	¹³ C NMR Chemical Shift Range	27
7	¹⁵ N NMR Chemical Shift Range	30
8	¹⁵ N NMR Results. Effects of Divalent Metal Ions on the Chemical Shifts of Guanosine	49
9	¹⁵ N NMR Results. Effects of Divalent Metal Ions on the Chemical Shifts of Inosine	50 :)
10	¹⁵ N NMR Spectrum of Inosine	51
11	¹⁵ N NMR Results. Effect of Protonation on the Chemical Shifts of Guanosine	53
12	¹⁵ N NMR Results. Effect of Protonation of the Chemical Shifts of Inosine	54
13	¹³ C NMR Spectrum of Inosine, Fully Coupled	56
14	¹³ C NMR Spectrum of Inosine, ¹ H Decoupled	57
15	¹³ C NMR Results. Effects of Divalent Metal Ions on the Chemical Shifts of Inosine	58
16	¹³ C NMR Results. Effects of Divalent Metal Ions on the Chemical Shifts of Guanosine	59
17	¹³ C NMR Results.—Effect of Protonation on the Chemical Shifts of Guanosine	63
18	¹³ C NMR Results. Effect of Protonation on the Chemical Shifts of Inosine	64
19	¹³ C NMR Results. Effect of N7-Methylation on the Chemical Shifts of Guanosine	67

Figure	Title	page
20	¹³ C NMR Results. Effects of Divalent Metal Ions on the Chemical Shifts of 7-Methylguanosine	69
21	¹³ C NMR Results. Effects of Protonation and Methyl-mercuration on the Chemical Shifts of 7-Methylguanosine.	70
22	¹³ C NMR Results. Effects of Divalent Metal Ions on the Chemical Shifts of Inosine-5'-monophosphate	78
23	¹⁵ N NMR Results. Effect of Mercury(II) on the Chemical Shifts of Inosine-5'-monophosphate	81
24	³¹ P NMR Results. Effects of Divalent Metal Ions on the Chemical Shift of Inosine-5'-monophosphate	82
25	¹⁵ N NMR Results. Effects of Divalent Metal Ions on the Chemical Shifts of Adenosine	88
26	¹³ C NMR Results. Effects of Divalent Metal Ions on the Chemical Shifts of Adenosine	91
27	¹⁵ N NMR Results. Effects of Divalent Metal Ions on the Chemical Shifts of Adenosine-5'-monophosphate	97
28	¹³ C NMR Results. Effects of Divalent Metal Ions on the Chemical Shifts of Adenosine-5'-monophosphate, pH 10 ..	99
29	³¹ P NMR Results. Effect of Divalent Metal Ions on the Chemical Shifts of Adenosine-5'-monophosphate, pH 10 ..	100
30	¹⁵ N NMR Results. Effects of Divalent Metal Ions on the Chemical Shifts of Adenosine-5'-triphosphate, pH 7 and 10	104
31	¹³ C NMR Results. Effects of Divalent Metal Ions on the Chemical Shifts of Adenosine-5'-triphosphate, pH 7 and 10	106
32	¹³ C NMR Results. Effect of Zinc(II) on the Chemical Shifts of Adenosine-5'-triphosphate, pH 4	107
33	³¹ P NMR Results. Effects of Divalent Metal Ions on the Chemical Shifts of Adenosine-5'-triphosphate, pH 7 and 10	109
34	³¹ P NMR Results. Effects of Divalent Metal Ions on the Chemical Shifts of Adenosine-5'-triphosphate, pH 4	110
35	³¹ P NMR Spectrum of the Zinc(II)-Adenosine-5'-triphosphate Complex, pH 10	112

Figure	Title	page
36	^{199}Hg NMR Results. Effects of Nucleosides on the Chemical Shifts of HgCl_2	120
37	The Concentration Dependence of the ^{199}Hg Chemical Shift of HgCl_2	121
38	Base Paired Structure of CpG in Solution	125
39	^1H NMR Spectrum of CpG, Solvent Suppressed	128
40	A Two-dimensional Homonuclear- ^1H - ^1H Chemical Shift Correlation NMR Spectrum of the Ribose Region of CpG ..	129
41	Expansion of 4.0 to 4.6 ppm Region of Figure 40	130
42	The Structure, Numbering Scheme and Torsional Angles of CpG	132
43	^{13}C NMR Spectrum of CpG	133
44	Chemical Shift Correlation Between the Mononucleotides 3'CMP and 5'GMP and the Dinucleoside Monophosphate CpG ..	134
45	Pseudorotation of the Ribose Sugar Ring, Envelope and Twist Conformations	145
46	Major Ribose Sugar Conformations	146
47	^1H NMR Spectrum of DeoxyCpG	151
48	The Structure, Numbering Scheme and Torsional Angles of Deoxy CpG	153
49	Chemical Shift Correlation Between the Mononucleotides 3'dCMP and 5'dGMP and Deoxy CpG	156
50	^{15}N NMR Results. Effect of Methylmercury on the Chemical Shifts of 5'GMP and 3'CMP. The Synthetic $\text{N1-CH}_3\text{Hg(II)-5'GMP}$ Complex	165
51	^{13}C NMR Results. Effects of Methylmercury on the Chemical Shifts of 5'GMP and 3'CMP	169
52	^{13}C NMR Results. Effects of Methylmercury on the Chemical Shifts of 5'dGMP and 3'dCMP	170
53	^{13}C NMR Results. Synthetic N1 Complexes of 5'GMP and 5'dGMP with Methylmercury	171
54	^{13}C NMR Results. Effects of Methylmercury on the Chemical Shifts of CpG	174

Figure	Title	page
55	¹³ C NMR Results. Effect of Methylmercury on the Chemical Shifts of Deoxy CpG	175

Nucleic acids and nucleic acid derivatives are naturally occurring multisite ligands. The interaction of these species with metal ions have been the subject of many studies (1,2,3).

In the past this complexation has been studied largely by ^1H NMR, however protons are generally three or more bonds removed from the actual binding sites. Thus results obtained using this method are approximate.

In this present work, a multinuclear NMR approach will be taken in order to study the interaction of diamagnetic metal ions with nucleic acid derivatives. This binding process will be discussed in terms of the following specific goals: 1) the determination of site selectivity, 2) the role of metal ion polarizability, 3) the effects of solvent and pH, 4) kinetic versus thermodynamic product formation, 5) the effects of physically blocking a binding site, and 6) the nature of the complex formed; inner-sphere binding or outer-sphere, the stoichiometry of binding and the stability constant of the product.

Of major importance is the determination of site selectivity and the factors affecting the complexation process. The dependence of binding on the polarizability of the metal ion, and the solvent involved will be studied.

Binding sites for the soft metal ions Hg(II) and $\text{CH}_3\text{Hg(II)}$, will be compared to those observed for Zn(II) ; a borderline species and for Ba(II) and Mg(II) which are classified as hard metal ions.

The pH dependence and role of solvent will be investigated by comparing the results observed for the nucleotides in aqueous solution to those obtained for the nucleosides in dimethylsulfoxide (DMSO).

The observed binding site may be either the kinetically or thermodynamically controlled site of complexation depending upon the rate of equilibrium establishment between the two species. This will have important consequences when considering the role metal ions play in cellular conditions.

Secondary binding sites will be investigated by using nucleosides where the primary metal ion complexation site has been blocked by methylation. These secondary sites take on increasing importance at very high metal ion to ligand concentration ratios.

A metal ion may bind tightly in an inner-sphere complex, or outer-sphere binding through intervening solvent molecules may occur. In addition the complexation process between metal ion and nucleic acid derivative can be characterized by its stoichiometry and stability constant.

In attempting to answer these questions a clearer understanding of the metal ion-nucleic acid complex should emerge.

^{15}N NMR is expected to yield the greatest amount of information about metal ion complexation. The nitrogens of the heterocyclic bases represent primary soft metal ion binding sites (1,2,3). Large changes in chemical shift are predicted for nitrogens directly involved with a metal ion.

^{13}C and ^{31}P NMR will also be used to study metal ion complexation to a nucleic acid derivative, although these nuclei are at least one bond removed from the binding site. These results will aid in binding site determinations, in particular for the harder metal ion species

which no longer bind preferentially to the nitrogen sites.

^{199}Hg NMR will be applied to characterize the binding of Hg(II) to the nucleosides.

The nucleic acid derivatives involved in this study included the 6-oxopurine nucleosides and nucleotides. The effects of phosphorylation on metal ion binding were studied using the adenine series; adenosine, adenosine-5'-monophosphate and adenosine-5'-triphosphate. Finally, in order to provide a model for RNA and DNA interactions, the metal ion complexation properties of two self-complementary dinucleoside monophosphates were studied.

The complexation of a metal ion with a nucleic acid derivative will be detected by monitoring the chemical shifts of the ligand nuclei. An expression for the $\sigma_p(\text{loc})$ term of each shielding constant will be derived, and the effects of metal ion complexation predicted.

CHAPTER 2: NUCLEIC ACIDS AND METAL IONS

2.1 Physical and Chemical Properties of Nucleic Acid Constituents

Nucleic acids are polymers composed of repeating mononucleotide fragments. The nucleotide units themselves are composed of either a purine or pyrimidine heterocyclic base, a pentose sugar, and a phosphate or polyphosphate group (4).

There are five commonly occurring bases; two are purine and three are pyrimidine heterocycles. A third purine base occurs regularly in transfer RNA species, namely hypoxanthine. Of the pyrimidine bases, uracil occurs only with ribose sugars in RNA, while thymine associates only with 2'-deoxyribose sugars in DNA. The structures of these bases and their numbering schemes are shown in Figure 1 (i).

One of two sugars is found in the nucleic acid components. Both are pentose sugars which are in the furanose form. Either a β -D-ribose or a β -D-2'-deoxyribose sugar can be bound to the N9 position of the purines or to the N1 position of the pyrimidines to form the nucleosides. The glycosidic bond forms to the anomeric sugar carbon, as shown in Figure 1 (ii).

Nucleotides are formed when one of the free sugar hydroxyl groups is involved, through a phosphoester linkage, with a phosphate or polyphosphate group. The phosphate chain is labelled α, β, γ , etc. starting at the nucleoside end.

i. Common purine and pyrimidine bases.

ii. Ribofuranoside derivatives.

FIGURE 1: Structures and numbering schemes for the major nucleic acid components.

This phosphate group can serve to link several nucleosides together. Labelling of such a chain proceeds from the free 5' end to the 3' terminus and the location of phosphate linkages are indicated in brackets. For example, in Figure 2, the structural formula, abbreviations and numbering scheme are shown for cytidyl-(3'→5')-guanosine, or CpG.

The heterocyclic bases are generally planar, with the exocyclic amino groups lying in the plane of the ring and the lone pair of electrons delocalized into the π -system of the ring. It is generally found that the imino tautomer of the bases is not important in solution (1,4).

The sites of protonation and the pK_a values for the nucleosides and nucleotides are given in Table 1 (1). These sites are also primary metal ion interaction sites, depending upon experimental conditions.

Table 1 pK_a 's of Nucleoside and Nucleotide Constituents

	Protonation	Deprotonation
Adenosine	3.5 N1	
Guanosine	2.1 N7	9.2 N1
Inosine	2.0 N7	8.8 N1
Cytidine	4.2 N3	
sugar		12.4 (OH)
phosphate		< 1.5 pK_a 1
		~ 6 pK_a 2

FIGURE 2: Structural components of cytidylyl-(3'→5')-guanosine.

In concentrated solutions, ($>0.01M$), there are a great deal of inter-molecular interactions occurring. These include base stacking and hydrogen bonding interactions in aqueous solutions. Purine derivatives show greater base stacking than do the pyrimidines (1).

These intermolecular interactions are important to studies using dinucleoside monophosphates such as CpG. It has been found that these species will associate in aqueous solutions to form mini-helices (5). The stabilizing forces for such an interaction are hydrogen bonding and base stacking. Thus the dimer CpG can be used as a model for RNA and DNA studies.

2.2 Interaction of Metal Ions with Nucleic Acid Derivatives

Interest in the title interactions is stimulated by their potential cellular consequences. Metal ions are known to disrupt translation and transcription.¹ The results can be either mutagenic or carcinogenic. Metal ion effects are also of prime environmental concern due to their presence as industrial waste.

However metal ions are also known to act in a therapeutic fashion. For example, it was discovered that Pt(II) complexes of nucleic acids are useful antineoplastic agents (6). Copper, zinc, gold and lithium

¹Transcription involves the opening of a DNA double-helix and the building of a messenger RNA strand on the resulting template. Translation is the reading of this messenger RNA by binding specific transfer RNA's with their attached amino acids. The result is the creation of a specific protein product. Any species which affects the base structure of either DNA or RNA can cause a "mis-reading" of these templates.

are also important in disease treatment and control (3, vol. 14).

Metal ion-nucleic acid studies have thus been centered on determining binding sites on the monomer constituents of DNA. These binding sites may be studied as they occur naturally in vivo, or inert metal ion complexes can be prepared in vitro.

Applications currently being pursued in the literature include:

- 1) inert paramagnetic:nucleotide complexes as probes for enzyme active sites,
- 2) paramagnetic induced relaxation enhancements,
- 3) use of heavy metals in the electron microscopy sequencing of DNA,
- 4) heavy metal containing intercalating agents,
- 5) metal ion catalyzed degradation of DNA,
- 6) lanthanide shift reagents to aid in NMR assignments of polynucleosides,
- 7) metal ion fluorescence probes,
- 8) selective precipitation and purification of nucleic acids using metal ions, and
- 9) metal ion-heterocyclic base complexes in the synthesis of oligonucleosides (2).

Several reviews exist on the early studies, prior to 1971, of metal ion-nucleic acid complexes (3 vol. 1,7,8). However these studies dealt previously with the heterocyclic bases, and the results can not be applied to the more biologically relevant nucleosides and nucleotides.

Newer techniques, including FT-NMR have recently been applied to the problem of determining the sites of metal ion interactions. The detailed features of the metal ion binding sites on the nucleic acid derivatives, however have still not been fully determined.

2.3 Properties of Metal Ions in Solutions

The properties of the metal ion itself can determine its preferred binding site, or the ultimate effect it will have on the nucleic acids it modifies. In order to classify metal ions it is necessary to develop the concept of hard and soft acids. Several factors will affect the stability of complexes formed by metal ions. The most important of which is their acceptor behaviour toward various types of ligands. For halide ions as ligands, the order of complex stability is fluoride > chloride > bromide > iodide for the lighter s and p block cations, metals at the beginning of the transition series, the lanthanides and the actinides. These metal ions are class (a) or hard acid acceptors (9). These ions prefer to bind with the harder bases; such as oxygen, phosphate, and sulfate.

For tellurium, polonium, the platinum metals, copper, silver, gold, cadmium, mercury and thallium the order of halide stability is reversed, i.e. $I > Br > Cl > F$. These metal ions are the soft acids. They prefer soft donor ligands such as cyanides, phosphines, and mercaptans.

There are also borderline metal species such as iron, cobalt, nickel, copper and zinc, which will bind to both ligand groups.

The stability of a complex will also be increased if a stable five- or six-membered chelate ring can be formed (10).

In addition to the polarizing nature of the metal ion ligand interaction, each metal ion has different requirements in terms of coordination number and stereochemistry.

The harder metals require a greater number of coordinated ligands to form a stable complex than do the softer metal ions. This larger coordination sphere may prevent these hard metals from binding to sterically restricted binding sites.

It is of interest to note that these properties also parallel the ability of a metal ion to promote errors in the in vitro translation of a DNA sequence (3 vol. 10). This implies an in vivo mechanism of mutagenicity or carcinogenicity.

Metal ions have been divided into five groups based on their reported ability to cause cancers in human and animal studies (3 vol. 10). The most active metal ions, including chromium, cadmium, and nickel are definite carcinogens in humans. For mercury the evidence of carcinogenicity is regarded as uncertain. For zinc and magnesium no evidence of cancer promotion has been reported (3 vol. 10).

Therefore small differences in the properties of the metal ions or the sites to which they bind on nucleic acids or proteins can dramatically alter their biological effects.

2.4 Potential Metal Ion Binding Sites on Nucleic Acid Derivatives

Nucleic acid derivatives including nucleosides, nucleotides, and dinucleoside monophosphates provide a variety of potential metal ion binding sites (1,2,3).

On the heterocyclic bases, N7 and N1 in adenosine, N7 in guanosine and inosine, and N3 in cytidine are favoured binding sites at neutral pH. The protonated nitrogens, N3 in uridine and thymidine and N1 in guanosine and inosine will bind metal ions at higher pH. The amino group of guanosine, adenosine or cytidine is

not considered a potential binding site except at very high pH. The N3 position of the purine nucleosides and nucleotides does not bind metal ions due to steric hindrance by the sugar group. In the free bases N3 is a favoured binding site.

The carbonyl oxygens of guanosine, inosine, cytidine, uridine, and thymidine are potential binding sites for very hard metal ion species. These carbonyls may also form chelate complexes involving an adjacent heterocyclic nitrogen.

Metal ion binding at C5 in cytidine and uridine and at C8 in the purines can be forced using Hg(II) and Tl(III). In addition the C5-C6 double bond in the pyrimidines can be attacked by MO_4 , where $M = Mn, Ru, Os$. However these are very specialized reactions and are not important under biological conditions.

Binding to the sugar oxygens is also not a favoured interaction. However the phosphate oxygens exhibit a number of multidentate binding reactions with the hard metal ions (2).

To predict which binding sites will be favoured it is necessary to estimate the chemical reactivity of the system. One approach involves the use of generalized perturbation theory (10).

The energy difference between the highest occupied molecular orbital of the donor atom and the lowest unoccupied molecular orbital of the acceptor atom characterizes the interaction. A large amount of charge transfer is characteristic of a soft-soft interaction, whereas very little charge transfer would indicate a hard-hard complex is being formed.

The hard or soft nature of the metal ions is determined by their orbital energies. Soft metals have negative orbital energies,

whereas hard metals have positive energies. The softer the metal the greater the amount of electron transfer that will occur in the transition state.

In the same fashion it is possible to calculate the perturbation energy of a metal ion nucleoside complex. The greatest perturbation will be observed at the softest binding sites (10).

For adenosine N7 is predicted to be a softer binding site than N1. Similar results are observed for inosine and guanosine; N7 is the softest binding site, with N1 being preferred only by very hard metals.

These soft nitrogens are generally the favoured metal ion binding sites, however it is easy to shift the equilibria and favour the other binding sites by small changes in pH, ionic strength, or hydrophobic interactions. Changes to the nucleosides themselves by the addition of a phosphate group to the sugar, or the blocking of a binding site will also affect the preferred binding site.

In the present work, attempts will be made to determine the occupied binding sites of a variety of nucleotide components via the NMR chemical shift changes induced by a series of diamagnetic metal ions. The concomitant stereochemical changes will also be monitored employing various NMR spectral techniques.

CHAPTER 3: MULTINUCLEAR NMR

3.1 Basic Theory of NMR

The phenomenon of nuclear magnetic resonance (NMR) depends on the fact that some atomic nuclei behave as if they are spinning. As a result they possess both a magnetic moment (μ) and angular momentum ($\vec{p}$).

$$\vec{p} = I\hbar = \frac{I\hbar}{2\pi} \quad \dots\dots 1$$

A nucleus with a non-zero nuclear spin quantum number, I , will have a magnetic moment μ proportional to its angular momentum.

$$\vec{\mu} = \gamma \vec{p} \quad \dots\dots 2$$

The proportionality constant γ is the magnetogyric ratio and reflects nuclear properties not accounted for in the classical spinning charge model.

In this classical description the magnetic moment vector interacts with the external magnetic field $\vec{B}_0$. The magnetic moment experiences a torque $\vec{\tau}$.

$$\vec{\tau} = \frac{d\vec{p}}{dt} = \vec{\mu} \times \vec{B}_0 \quad \dots\dots 3$$

causing it to precess about the external magnetic field.

The angular velocity and direction of such a motion is given by $\vec{\omega}_0$:

$$\vec{\omega}_0 \times \vec{\mu} = \gamma \vec{\mu} \times \vec{B}_0 \quad \dots\dots 4$$

$$\vec{\omega}_0 = -\gamma \vec{B}_0 \quad \dots\dots 5$$

and the frequency of precession ν_0 ,

$$\nu_0 = \frac{|\omega_0|}{2\pi} = \frac{\gamma \bar{B}_0}{2\pi} \quad 6$$

This represents the Larmor frequency of a particular nucleus, and it is dependent only upon the strength of the applied magnetic field.

The spin angular momentum of a nucleus is quantized, with quantum number I . The angular momentum may take values of mh , where m is the discrete spin variable running from $-I$ to $+I$ in steps of one. In the presence of an external magnetic field the $2I + 1$ spin levels are separated into different energy levels,

$$E = -\vec{\mu} \times \vec{B}_0 = \gamma \hbar m \bar{B}_0 \quad 7$$

A selection rule, $\Delta m = \pm 1$ applies to transition between spin levels so the energy of such a transition is given as:

$$\Delta E = \gamma \hbar \bar{B}_0 = \frac{\gamma \hbar B_0}{2\pi} \quad 8$$

A transition is induced by the absorption of energy from an applied electromagnetic field $\vec{B}_1$ generated perpendicular to the external magnetic field.

In an NMR spectrum the mean behaviour of an ensemble of nuclear spins is important. At thermal equilibrium the spins will be distributed according to Boltzmann's Law.

$$\frac{n_l}{n_u} = \exp \frac{2\vec{\mu} \cdot \vec{B}_0}{kT} \approx 1 + \frac{2\vec{\mu} \cdot \vec{B}_0}{kT} \quad 9$$

There is only a very small population difference between the

upper state, n_0 and the lower one, n_1 . This excess in n_1 is large enough to establish a small macroscopic magnetization, $\vec{M}$, directed along $\vec{B}_0$.

$$\vec{M} = \frac{N\mu^2\vec{B}_0}{kT} \quad \dots\dots 10$$

where N is the number of nuclei per unit volume. A bulk magnetic susceptibility χ can be defined and represents the inherent sensitivity of the NMR experiment.

$$\chi = \frac{\vec{B}_0}{\vec{M}} = \frac{N\gamma^2\hbar^2 I(I+1)}{3kT} \quad \dots\dots 11$$

The nuclei in a sample will not all resonate at their Larmor frequency. Magnetically different nuclei in a sample will resonate at a characteristic frequency slightly shifted away from their Larmor frequency. These shifts are generally small, 10^{-3} to 10^{-5} MHz for most of the common nuclei.

This chemical shift difference, or screening effect observed from a free nucleus arises from the magnetic effect of induced electronic currents in the atom or molecule. These currents induced by the external field, modify the magnetic field observed at the nucleus.

$$\vec{B}_{obs} = \vec{B}_0(1 - \sigma_i) \quad \dots\dots 12$$

This in turn effects the resonance frequency of the nucleus in question,

$$\nu_i = \frac{\gamma_i \vec{B}_{obs}}{2\pi} = \frac{\gamma_i}{2\pi} (1 - \sigma_i) \vec{B}_0 \quad \dots\dots 13$$

The constant σ_i is a shielding tensor and its magnitude depends

on the electronic environment of the nucleus in question. Thus chemical shift can be used to monitor the environment of a nucleus in an NMR experiment.

A second general feature of an NMR spectrum is nuclear spin-spin coupling. A transition between nuclear spin levels will occur at a characteristic frequency, its chemical shift, as discussed previously. However, this resonance may have a multiplet structure, arising from the splitting of nuclear spin levels caused by the interaction of other nearby magnetic nuclei.

There are two general mechanisms of spin-spin interaction; through space and through chemical bonds. The through space interaction of two magnetic dipoles is dependent upon both the distance and angle between the dipoles. Thus, the rapid and random motion of small molecules in solution will average this interaction to zero.

As nuclear spin-spin coupling is generally observed in liquid samples it must arise from a mechanism independent of molecular motion. A second indirect dipole-dipole interaction occurs between bonded nuclei, using the bonding electrons and orbital momentum to convey the information.

Classical approximations involving electron point dipoles yield couplings much smaller than those observed experimentally. Quantum mechanics introduces the Fermi contact term, which allows a finite probability for finding a bonding electron at the nucleus of an atom. This principle forms the basis for a very simple model of nuclear spin-spin coupling.

Consider two nuclei, both with positive magnetogyric ratios

joined by a single chemical bond. Nucleus A has its spin orientated parallel to the magnetic field B_0 . An electron near to this nucleus will tend to orientate its spin in the opposite direction. As this electron is part of a bonding pair, the Pauli Principle requires the second electron in the bonding pair to have an opposite spin. This electron is near enough to nucleus B to influence it to line up opposite again. Thus information about the spin orientation of A is transmitted to nucleus B via the bonding electrons.

The lower energy configuration is shown above. At a slightly higher energy is the second state where the spins of A and B are parallel. However due to the small interaction energies the populations of these spin states are essentially equal.

This represents a positive coupling constant, $^1J_{AB}$, where the low energy state has spins A and B antiparallel. A negative coupling constant occurs when the low energy configuration has spins A and B parallel. For one bond couplings this is observed if the nuclei in question have opposite magnetogyric ratios.

Thus when nucleus A undergoes an NMR transition and flips its spin state with respect to the external magnetic field, the energy of this interaction will depend on the initial orientation of nucleus B relative to A. Two resonance lines will occur, their

separation a measure of the interaction energy between nuclei A and B; the nuclear spin-spin coupling.

The nuclear spin-spin coupling is independent of the external magnetic field, unlike chemical shifts which are induced by and proportional to the magnetic field applied.

Spin-spin couplings are not restricted to directly bonded nuclei. In addition the number of spin configurations is not restricted to two. For nuclei with $I > \frac{1}{2}$ more resonance lines appear. There are $2I + 1$ possible orientations of a nuclear spin relative to an external field. Thus a spin-spin coupled multiplet will be composed of $2nI + 1$ lines, where n is the number of coupled spins. Theory and practice often differ however for nuclei with $I > \frac{1}{2}$. The presence of a quadrupole moment in these nuclei can cause rapid relaxation and effectively isolate the coupled spins, so no multiplet structure is observed. In addition for nuclei with large values of I , sensitivity and resolution requirements will limit observation of complete multiplet structure.

The usefulness of coupling constants arises in the fact they can be related to a number of physical properties. For example hybridization, dihedral bond angles, and substituent electronegativity will all predictably affect the magnitude of spin-spin coupling constants.

3.2 Specific Nuclei of Interest

In this research a number of different nuclei were employed to characterize the features of binding.

The magnetic properties of these nuclei are presented in Table 2 (11).

Table 2 Magnetic Properties of Observed Nuclei

Nucleus	Spin I	Abundance %	Q.M. ^a 10^{-28}m^2	Receptivity ^b	Frequency ^c MHz
^1H	$\frac{1}{2}$	99.99	-	1.0	200.0
^{13}C	$\frac{1}{2}$	1.1	-	1.59×10^{-2}	50.28
^{15}N	$\frac{1}{2}$	0.37	-	3.40×10^{-4}	20.28
^{31}P	$\frac{1}{2}$	100	-	6.63×10^{-2}	80.96
^{199}Hg	$\frac{1}{2}$	16.8	-	8.60×10^{-2}	35.65

a) quadrupole moment

b) compared to ^1H

c) at 4.7T magnetic field

3.21 ^1H -NMR

In 1946, Purcell at Harvard first observed ^1H resonance in wax, while at Stanford Bloch found the resonance in water (12,13). The early observation of ^1H is due to its large magnetic moment and high natural abundance, making it the most easily detected NMR nucleus.

The spin lattice relaxation times of ^1H nuclei are normally between 1 and 10 secs. This relaxation proceeds mainly through

dipole-dipole interactions with other nearby protons.

^1H -NMR chemical shifts are usually confined to a 20 ppm range if paramagnetic species are omitted, eg. see Figure 3 (14).

In an ^1H -NMR spectrum the resonances are generally split by homonuclear spin-spin couplings. These couplings can be used to identify specific resonances, and to investigate stereochemistry. Karplus has defined equations for the dihedral bond angles in ethane-type fragments based on vicinal ^1H - ^1H couplings (15).

3.22 ^{31}P -NMR

There were three elements to which early high-resolution NMR was particularly applicable: ^{31}P is the second of these species for which an isotope of near 100% natural abundance has a spin of $\frac{1}{2}$. ^1H was the first and ^{19}F was the third.

^{31}P also has a strong magnetic moment making it easily observable even on the original low-field, continuous wave NMR spectrometers (16,17).

As a result of the easily observed chemical shifts and the strong couplings exhibited by ^{31}P , data on this nucleus played a key role in the development of general NMR theory.

Typical spin lattice relaxation times are .01 to 10 seconds, so generally spin saturation is not a problem.

^{31}P chemical shifts occur over a 700 ppm range, but are highly dependent upon the phosphorus oxidation state. Thus all P(III) compounds occur over the entire 700 ppm range, but the P(V) species are found within a smaller 150 ppm range, as shown in Figure 4 (18).

FIGURE 3: ^1H chemical shifts in various chemical environments.

FIGURE 4 : ^{31}P Chemical shifts in various functional groups.
Reference P_4O_6 .

The phosphorus chemical shifts are useful in structure determination and general additivity rules have been developed (19). In effect three variables are important, the bond angle, the electronegativity of the substituents, and the degree of π bonding involved.

Phosphorus couplings to other nuclei are generally quite strong, and can be used in structural determinations (19,20). Modified Karplus relationships can be used in nucleic acid chemistry to define the backbone conformation of oligonucleosides (21).

3.33 ^{199}Hg -NMR

Mercury has two stable isotopes with non-zero nuclear spins. The ^{201}Hg nucleus has a spin of $3/2$ and possesses a substantial quadrupole moment, making it very hard to observe. The more receptive spin $1/2$ ^{199}Hg isotope can be detected with little difficulty.

Mercury complexes, with the exception of organomercurials, are kinetically labile. Thus only time averaged signals are observed in the NMR spectrum. As a result these signals exhibit a marked concentration dependence.

Mercury tends to form complexes of low symmetry causing a large magnetic anisotropy to be present at the nucleus. This contributes significantly to the observed chemical shift range of 3600 ppm, as shown in Figure 5 (22).

FIGURE 5: ^{199}Hg chemical shift ranges for some different chemical environments.

The ^{199}Hg nucleus spin couples very strongly to other nuclei, $^2J_{\text{H,Hg}} \sim 100\text{-}300\text{ Hz}$ and $^3J_{\text{H,Hg}} \sim 250\text{-}350\text{ Hz}$ (23,24). ^{199}Hg -NMR has been used to follow the binding of Hg(II) to the human carbonic anhydrase B enzyme at the Zn(II) active site (25). ^{199}Hg -NMR has also been used to characterize the mercuric halide salts in solution (26) and the pH dependence of CH_3HgOH in aqueous solutions (27).

The $^2J_{\text{H,Hg}}$ values in methylmercury have also been used to determine the structure of pyridine complexes (28,29) and the effects of ligand basicity on binding.

3.24 ^{13}C -NMR

The first ^{13}C -NMR studies were reported in 1957, using rapid passage, dispersion mode techniques (30,31). With the advent of FT-NMR techniques in the early 1970's, the field of ^{13}C -NMR mushroomed (32,33, and references therein).

The popularity of ^{13}C -NMR arises from the key role carbon plays in the skeleton of organic compounds. The low natural abundance of the ^{13}C atom eliminates, to a first approximation, any complications due to ^{13}C - ^{13}C scalar interactions.

The chemical shifts of most neutral organic species are found within a 250 ppm range, as shown in Figure 6 (32).

A very large body of ^{13}C -NMR chemical shift data has been accumulated over time, largely dealing with substituent effects (32,33). The chemical shift of a carbon nucleus can be predicted through the additivity nature of these substituent effects (34).

FIGURE 6: ^{13}C chemical shift ranges in organic compounds.

Spin-spin couplings between carbon and other magnetic nuclei are well documented (32,33). Couplings to protons are the most commonly observed, and the multiplicity can be used in structure determination. The one bond couplings are the largest; 95-320 Hz, depending upon the degree of s-character the carbons possess. The geminal couplings are smaller, 2-10 Hz, with exceptions observed through a heteroatom. Similar values are found for the vicinal couplings. Long range couplings are generally only observed through an intervening π system.

The vicinal couplings between carbon and other nuclei such as phosphorus and protons can be used to yield conformational information. A number of Karplus-type relationships have been developed (35).

The ^{13}C -NMR spectra of the nucleosides and nucleotides were recorded early in the development of the technique (36,37). Corrections, however, were later made to the ribose carbon resonance assignments (38).

Recently the additive nature of ^{13}C chemical shifts has been used to investigate the interaction between nucleosides and metal ions (39,40). However care must be taken to determine the origin of these effects in the chemical shift tensor.

3.25 ^{15}N -NMR

Nitrogen also has two isotopes with non-zero nuclear spins. The more abundant ^{14}N nucleus has a spin of 1, and a relatively large quadrupole moment. Thus it is not suitable for routine high-resolution NMR studies.

^{15}N nuclei have a spin of $\frac{1}{2}$ and as such are better for structural determinations. However the very low natural abundance and negative magnetogyric ratio of this nucleus, have largely restricted its study to neat, low molecular weight liquids or enriched compounds (41).

It wasn't until the advent of FT-NMR and wide-bore superconducting magnets that natural abundance ^{15}N -NMR was found to be feasible (42,43). To obtain a spectrum several factors which influence the ^{15}N signal strength must be carefully considered. These include exchange processes, relaxation times and the nuclear Overhauser effect.

The chemical shift range of ^{15}N is slightly larger than that of ^{13}C in neutral organics, at 500 ppm. Similar to carbon, additive substituent parameters can be used to predict chemical shifts. However ^{15}N shifts are generally more sensitive to solvent changes and other intermolecular interactions. The range of chemical shifts exhibited by a variety of nitrogen compounds is shown in Figure 7 (44).

A large number of ^{15}N -NMR studies have now been done on biologically important molecules. In terms of nucleic acid studies, the ^{15}N -NMR spectra of the nucleosides and nucleotides have been recorded (45,46). ^{15}N -NMR has also been used to study the hydrogen-bonding interactions between the nucleosides (47,48).

Recently the ^{15}N -NMR spectrum of an enriched DNA sample of about 300 base pairs was obtained (49).

Applications of ^{15}N -NMR have been made to study the binding of the anti-cancer drug cis-diamminedichloroplatinum(II) to

FIGURE 7: ^{15}N Chemical shiftings in organic compounds.

nucleosides (50). Also ^{15}N NMR has been applied to the study of metal ion interactions with the nucleic acid derivatives. The interaction between the paramagnetic metal ion Mn(II) and 5'AMP was studied using the ^{15}N relaxation rates (51). A very early study using Mg(II) and Zn(II) complexed to 5'ATP pointed out the potential for using ^{15}N chemical shifts to determine binding sites for diamagnetic metal ions (52). Recent work from this laboratory has elaborated on the metal ion binding sites of the nucleosides using ^{15}N chemical shifts (53,54).

3.3 Theory of Chemical Shifts

Chemical shifts arise due to the interaction between an electron and nucleus and the effect an external magnetic field has on the electron.

In the presence of an external field the electrons will precess about the nucleus. This produces a current which will induce a secondary magnetic field (B') at the nucleus in question. This secondary field is opposite in direction to the external field (B_0).

$$B' = -\sigma B_0 \quad \dots\dots\dots 14$$

Thus to achieve resonance, a greater B_1 field must be applied to this screened nucleus. The resonance appears shifted upfield, to lower frequencies.

This is a diamagnetic screening constant as described by Lamb for individual atoms (55).

$$\sigma_d = \frac{4\pi e^2}{3mc^2} \int_0^\infty r \rho(r) dr \quad \dots\dots\dots 15$$

where $\rho(r)$ is the average electron density as a function of radial distance r from the nucleus:

However in molecules the electrons of each nucleus are hindered in their motion due to the electric fields of nearby nuclei. Thus a correction term must be applied, and the screening constant is represented as the sum of two factors.

$$\sigma = \sigma_d + \sigma_p \quad \dots\dots\dots 16$$

The paramagnetic term σ_p was developed by Ramsey using second order perturbation theory (56). This σ_p term is more complicated than that of σ_d , and depends on the electronic excited states of the nucleus and their orbital angular momentum.

$$\sigma_p = \frac{1}{3} (\sigma_p(xx) + \sigma_p(yy) + \sigma_p(zz)) \quad \dots\dots 17$$

$$\sigma_p(zz) = \frac{e^2}{2mc^2} \left\langle 0 \left| \frac{x^2 + y^2}{r^3} \right| 0 \right\rangle - \left[\frac{e\hbar}{2mc^2} \right]^2 \times$$

$$\sum \frac{\frac{\langle 0 | L_z | n \rangle \langle n | \frac{2L_z}{r^3} | 0 \rangle}{E_n - E_0} + \frac{\langle 0 | \frac{2L_z}{r^3} | n \rangle \langle n | L_z | 0 \rangle}{E_n - E_0}}{\dots\dots 18}$$

The local fields produced by the paramagnetic electron currents add to the external field $\vec{B}_0$. Thus paramagnetic shifts result in a nucleus resonating at a lower applied field $\vec{B}$, or at a higher frequency. A downfield shift is observed.

One major problem associated with this paramagnetic term is that very little is known about the energy continuum of most molecules. However limited calculation on small species suggest that the screening contributions from the continuum are at least as important as those from the discrete excited states (57,58).

This problem can be simplified in two ways, one is by applying the closure relation which holds if one is working with a completely orthogonal basis set.

$$\sum_n |n\rangle \langle n| = 1 \quad \dots\dots 19$$

The second involves choosing an average excitation energy, AEE, for all excited states. This is the most often applied approximation,

and σ_p reduces down to:

$$\sigma_p = \frac{-\mu_0 e^2}{4\pi m^2} \frac{1}{r^3} \frac{1}{AEE} \langle 0 | Q | 0 \rangle \quad \dots\dots 20$$

where Q represents the bond order of the electrons. This equation involves only the ground state function and is thus more accessible, however suitable values for AEE are often not available.

3.31 Pople's Formula

Ramsey's model of chemical shifts, and its simplified versions, have two important drawbacks; 1) the origin dependence of the screening constant and 2) the fact that σ represents the small difference between two large terms of opposite magnitude (56). In a more applicable model of nuclear shielding proposed by Pople, σ is the sum of a number of localized shielding contributions (59):

$$\sigma = \sigma_d(\text{loc}) + \sigma_p(\text{loc}) + \sigma_d(\text{nonloc}) + \sigma_p(\text{nonloc}) + \sigma_d(\text{inter}) + \sigma_p(\text{inter}) \quad \dots\dots 21$$

The various diamagnetic and paramagnetic terms do not directly compare to their counter-parts in Ramsey's theory.

This approach developed by Pople is based upon an independent electron model in which all two-electron terms vanish (60,61,62).

The local terms $\sigma_d(\text{loc})$ and $\sigma_p(\text{loc})$ arise from electronic currents localized on the atom of interest. These represent the major contributions to the screening constants of the heavier nuclei, i.e. those with p and d electrons. The non-local contributions arise from currents on neighbouring atoms. These are usually estimated by assuming the induced magnetic moments on the neighbouring atoms can

be replaced by a point dipole (60). The effect of this dipole on the screening constant of the atom in question is generally only a few ppm. This is negligible for most heavier atoms due to their very large chemical shift ranges.

Finally $\sigma_d(\text{inter})$ and $\sigma_p(\text{inter})$ arise due to screening currents not localized on any particular atom in the system, such as ring currents. This screening contribution is generally quite small for the heavier atoms, though it is significant for the lighter ^1H -nucleus.

Within the confines of Pople's independent electron model, various approaches to the theoretical solution of the shielding expression can be taken. In all cases though, a linear combination of gauge dependent atom orbitals is taken, so as to produce gauge independent results.

The problem of gauge arises due to the fundamental physical requirement that nuclear shielding be independent of gauge. It does not depend on the means of measurement involved in an experiment. Gauge-dependent contributions arise when an incomplete basis set is used, due to the supposition of a common origin, which may be a poor one for some orbitals. To avoid these problems a complete basis set must be used. Ramsey's formula however contains a distance r , which is referred to an arbitrary origin, thus it remains gauge dependent even when a complete basis set is used.

Within the confines of the independent electron model, various approaches can be made to evaluate the screening terms. Ab initio calculations are practical only for very small molecules containing light atoms. Generally one of several semi-empirical methods is

chosen to calculate the chemical shift contributions of larger molecules with heavier atoms.

For the first row nuclei, including ^{13}C and ^{15}N , it is found that $\sigma_d(\text{loc})$ remains almost constant in a range of molecular environments. The observed screening differences arise from differences in $\sigma_p(\text{loc})$ (63).

To estimate semi-empirically the non-local contribution to the diamagnetic screening constant an atom-dipole model is used (64).

$$\sigma_d(\text{nonloc}) = \frac{\mu_0 e^2}{12\pi m} \sum_{B \neq A} \frac{Z_A}{r_{AB}} \quad \dots 22$$

This correction is limited to atoms B, directly bonded to nucleus A. It is apparent that the $\sigma_d(A)$ will be increased as the number of bonded atoms increases and as the number of electrons in the molecules increases (m-magnetic component quantum number).

The local paramagnetic term can be semi-empirically calculated following the method of Jameson and Gutowsky (65). The electronic excitation energies are replaced with an average excitation energy, AEE. $\sigma_p(\text{loc})$ is expressed in terms of localized bond properties and the calculations restricted to the valence electrons. The core electrons are assumed to be unchanged going from one molecule to another. This assumption leads to the property of spherical symmetry and therefore the core electrons do not contribute to the paramagnetic term.

$\sigma_p(\text{loc})$ is expressed in terms of orbital populations in the p and d valence shells, and the effect of the angular momentum operators $\hat{I}_x$, $\hat{I}_y$, and $\hat{I}_z$ on these populations.

$$\sigma_p(\text{loc}) = \frac{-\mu_0 e^2 \hbar^2}{6\pi m^2 \text{AEE}} (\langle r^{-3} \rangle_p P_u + \langle r^{-3} \rangle_d D_u) \quad \dots 23$$

where P_u and D_u represent the "imbalance" present in the atom's p and d orbitals. The expression for P_u is given below.

$$P_u = (P_{xx} + P_{yy} + P_{zz}) - \frac{1}{2}(P_{xx}P_{zz} + P_{yy}P_{xx} + P_{yy}P_{zz}) + \frac{1}{2}(P_{xy}P_{yx} + P_{xz}P_{zx} + P_{zy}P_{yz}) \quad \dots \quad 24$$

a similar expression exist for D_u (65).

The numerical values of P_u and D_u depend on the atom's coordination number, hybridization of bonding orbitals, and ionicity of its bonds. In a spherical closed shell system, both P_u and D_u have their minimum value of zero. A maximum unbalance in the p orbitals occurs when two are filled and one empty, or one filled and two empty; P_u then has its maximum value of 2. For the d orbitals, D_u is maximum at 12 when the three t_{2g} orbitals are filled and the two e_g orbitals are empty or vice versa. The quantity $\langle 1/r^3 \rangle$ in the above expression is an average value over all the p or d valence electrons,

$$\langle r^{-3} \rangle_{np} = \frac{1}{3} \left(\frac{Z_{np}}{na_0} \right)^3 \quad \dots \quad 25$$

where a_0 is the Bohr radius and n the principal quantum number. This value can also change in going from one molecule to another, however to a lesser extent than the other terms P_u and D_u and AEE.

The third variable in the expression for $\sigma_p(\text{loc})$ is the AEE. A large number of semi-empirical methods exist for the calculation of this value, and gross chemical shift trends can be predicted with some success in a series of closely related compounds. However no a priori method is available for choosing the AEE. Atoms with

electron lone pairs have their AEE dominated by the low energy $n \rightarrow \pi^*$ transitions. For other species, significant contributions to AEE are confined to the $\sigma \rightarrow \pi^*$, $\pi \rightarrow \sigma^*$ and $\sigma \rightarrow \sigma^*$ transitions, no contribution is made by $\pi \rightarrow \pi^*$ transition (66).

The contributions from non-local σ_d and σ_p and the interatomic σ_p and σ_d are generally negligible except in a few special cases. The wider chemical shift ranges of the heavier nuclei tend to overwhelm the few ppm corrections arising from these terms.

The largest of these effects is due to nearby atoms having large and anisotropic magnetic susceptibilities. These local fields will not average to zero with molecular motion leading to large chemical shift contributions.

To find the contribution of the induced magnetic dipole at an atom, a distance r away at an angle θ ,

$$\sigma_d = (X_{||} - X_{\perp})(1 - 3\cos^2\theta) / 12\pi r^3 \quad 26$$

where $X_{\perp}$ and $X_{||}$ are the two components of the neighbouring group's magnetic susceptibility.

Anisotropy effects on chemical shifts can be very large for protons, up to 5 ppm, because the local diamagnetic and paramagnetic terms are quite small. In the heavier nuclei which have larger paramagnetic shifts these anisotropic effects are not as significant.

The chemical shift of a nucleus can also be affected if a strongly polar group is present nearby. This is particularly important for the binding of metal ions or protons to a neutral species.

The polarization effect can be either transferred through bonds, an inductive effect, or through space, an electric field effect.

The resulting shielding effect is given,

$$\sigma_e = \frac{b_{||}}{e/r} E_{||} \dots\dots 27$$

where r is the separation, e the electronic charge, $b_{||}$ the bond polarization along the axis, and $E_{||}$ the electric field generated (32).

Of an even more empirical nature are solvent induced chemical shifts. These are dependent upon the nature of solvent and compound. Solvents with a high degree of magnetic anisotropy will induce downfield shifts whereas a very polar solvent will have a preferred orientation about the solute molecule. This will enhance any diamagnetic screening effects already arising from an electric field effect.

Specific interactions between solute and solvent, such as hydrogen bonding or chemical exchange will also affect the observed chemical shifts (66).

3.32 Chemical Shifts and Binding

These semi-empirical equations for the chemical shifts of nuclei can be used to explain the spectral changes which will occur when an electrophile such as H^+ or $M(II)$ binds to a ligand.

For ^{13}C , chemical shifts are dominated by the electronegativity of the substituent. Generally it is found that $\sigma_d(\text{loc})$ remains constant, and changes in the chemical shift of a nucleus arise from changes in $\sigma_p(\text{loc})$. These changes are due to a change in the $\langle r^{-3} \rangle$ value, which is present in the $\sigma_p(\text{loc})$ term (32,33,66).

An electrophilic substituent at a nitrogen which is π bonded to a carbon atom, will affect the chemical shift of the carbon nucleus

This is generally due to the inductive effect of the nitrogen on the carbon. The average radius of the 2p orbital is thus reduced, causing an increase in the local paramagnetic term and a downfield shift of the carbon nucleus in question.

$$\sigma_p(\text{loc}) = \frac{-\mu_0 g^2 \hbar^2}{6 \pi m^2 A E E} \langle r^{-3} \rangle_p P_u \quad \dots\dots 23$$

The effects of binding at a carbonyl oxygen are discussed in terms of changes in the "center of gravity" of the π electronic charge present between the oxygen and carbon atoms (32).

Generally if the two atoms in question are directly bonded together, the change in $\sigma_p(\text{loc})$ is discussed in terms of electronegativity. If the carbon is further removed the effects are described by an electric field calculation (33).

The binding of an electropositive group such as Ba(II) or Mg(II), at the carbonyl oxygen will tend to increase the population of 2 and a downfield shift of the carbon nucleus will occur. Again, this is due generally to a change in $\langle r^{-3} \rangle$ in the $\sigma_p(\text{loc})$ contribution.

The larger chemical shift range of ^{15}N when compared to ^{13}C is due to the presence of a lone pair of electrons on the nitrogen. This gives rise to a low level $n \rightarrow \pi^*$ transition which will influence $\sigma_p(\text{loc})$ through the average excitation energy term (AEE).

The binding of electropositive species to the nitrogen nucleus can affect the $\sigma_p(\text{loc})$ term in three distinct ways; 1) it can change AEE, 2) it can cause a change in $\langle r^{-3} \rangle$, and 3) it can change the multiplicity of the nitrogen.

The extent of lone pair contributions to $\sigma_p(\text{loc})$ is observed in the deshielding of a nitrogen nucleus with respect to its protonated form (44).

This concept is illustrated using aniline and pyridine. Aniline with its lesser degree of lone pair delocalization resonates at 55 ppm.

Upon protonation to the anilinium ion, the nitrogen becomes shielded by 8 ppm. Pyridine represents a highly aromatic nitrogen and its chemical shift of 317 ppm reflects this property. When protonated to the pyridinium ion, the ^{15}N resonance is shifted upfield to 205 ppm (44).

Thus complexation of an aromatic nitrogen with an electropositive species is expected to produce a dramatic upfield shift by increasing $\langle r \rangle$ and AEE while decreasing P_u , as denoted in equation 23.

If the aromatic nitrogen being complexed is conjugated to a second nitrogen, then it will undergo a downfield shift. This is due to a contributing resonance structure where the second nitrogen can donate its electron pair via resonance to the metallated nitrogen. A reduction in $\langle r \rangle$ and an increase in Q , produce a downfield shift in $\sigma_p(\text{loc})$.

For ^{31}P chemical shifts the same equation for $\sigma_p(\text{loc})$ will apply. As the phosphate groups in nucleic acid derivatives are pentavalent, no lone pair contributions are expected. The binding of electrophiles at the phosphate oxygens is expected to cause a downfield shift of the phosphorous resonance, by decreasing the $\langle r \rangle$ value of a phosphorous electron in the P-O bond.

In principle the factors that determine relative chemical shifts in the lighter nuclei are also found in the heavier nuclei.

However some of these factors can assume more importance for the heavier nuclei.

Transition metals are by definition those elements for which d electrons play a significant role in chemical bonding. This role is manifest in those physical properties sensitive to variations in orbital angular momentum, such as nuclear shielding. It is d- and p-orbital participation, with their associated large orbital angular momenta which cause the large chemical shift ranges observed for the transition metals. For example the nuclear shielding range for Cobalt is 18,000 ppm compared to 250 for carbon (67).

The nuclear shielding range of mercury is smaller, (3600 ppm), due to the presence of a filled set of d orbitals, with its spherical symmetry and zero orbital angular momentum. The valence shell in Hg(II) complexes is mainly of s and p orbital origin.

Pople's theory as presented previously has been applied to chemical shift calculations for light nuclei in small molecules. For heavier atoms the theory begins to break down because the relativistic effects and the effects of L-S or J-J couplings are not considered. Orbital angular momentum can couple with the electron spin angular momentum and yield substantial contributions to the nuclear shielding constant (66).

Similar to the lighter nuclei, it is found that $\sigma_d(\text{loc})$ is almost constant, and $\sigma_p(\text{non-loc})$ and $\sigma_d(\text{non-loc})$ are not significant. Therefore the observed chemical shift values are due to variations in $\sigma_p(\text{loc})$.

The ability of a ligand to cause d orbital splittings in a metal ion will play a significant role in determining $\sigma_p(\text{loc})$.

Ligands are generally defined by their position in the spectrochemical series, (9)

This represents the ability of a ligand to affect the d orbital splitting of a metal. This in turn increases the AEE term in $\sigma_p(\text{loc})$, and causes an upfield shift of the metal nucleus in question. Thus from observing the chemical shifts of a transition metal, information about the nature of its ligands can be gleaned.

CHAPTER 4: OXOPURINE NUCLEOSIDES

The discovery that some metal ion complexes can function as anticancer agents, has promoted the study of metal ion nucleic acid complexes (6). In particular it is known that the antitumour agent cis-Diamminedichloroplatinum (II) interacts preferentially with the guanine base in DNA (68).

This interaction is believed to occur at N7 and involves the carbonyl oxygen, promoting deprotonation of N1 (69). However binding at O6 has never been observed in solution, and only occurs in the solid state in a few specialized cases (70). One problem in these studies is the lack of an effective spectroscopic probe to the N1 or O6 site (71).

In order to study the role of the O6 carbonyl in metal ion complexation, the two 6-oxopurine nucleosides; guanosine and inosine were used. A series of metal ions were investigated ranging from the soft Hg(II) ion, to the borderline Zn(II) species and finally, to the hard metal ion Ba(II).

The ^{15}N and ^{13}C NMR spectra of the complexes were recorded and binding sites characterized as previously outlined. Of primary importance was the characterization of the N1 binding site, if occupied, and the investigation of any carbonyl interactions.

A model for cation binding to nucleosides is obtained from ab initio calculations on the interaction of Na^+ with the purine and pyrimidine bases (72). Binding sites were calculated by minimizing the three components of the interaction energy: Coulombic, exchange and delocalization.

Guanosine

Inosine

The preferred binding sites for guanine are shown below (72).

In general there are two types of sites possible; a thermodynamically favoured site and a kinetically favoured site. It is important to note that the type of binding detected will depend upon the experimental conditions employed.

The kinetic site is occupied first, however it may not represent the most stable binding configuration. The thermodynamic site is

similar to the favoured binding sites calculated by Pullman (72) and represents the lowest energy configuration.

In solution studies, where binding is determined by metal ion titration, generally in the case of mercury-nitrogen bonds, there is a rapidly established equilibrium situation leading to the observation of thermodynamically controlled products.

In the 6-oxopurines N7 is considered the kinetically favoured site and N1 the thermodynamic site (39).

4.1 ^{15}N NMR Results

4.11 Divalent Metal Ion Complexes

The ^{15}N NMR results for the metal ion complexes of inosine and guanosine are easily interpreted due to the large chemical shift changes observed. Results for guanosine complexed with Hg(II), Zn(II) and Ba(II) were studied. These results are presented in Figures 8 and 9. The ^{15}N spectrum of inosine is shown in Figure 10. It is typical of the ^{15}N spectra obtained in this work.

Large upfield shifts of ca. 20.0 ppm are observed for N7 when guanosine and inosine are complexed with Hg(II) and Zn(II). A smaller downfield shift of 5.0 ppm is seen for N9 in both cases. The remaining resonances are shifted to a small degree, due to perturbations in the electronic structure of the purine ring.

The interaction of a metal ion with the N7 of guanosine or inosine is expected to produce an upfield shift of the resonance by decreasing P_u and increasing $\langle r \rangle$ and AEE. The conjugated N9 nitrogen resonance is expected to show a downfield shift due to a change in the hybridization of the atom.

Thus the ^{15}N results indicate a dominant N7 binding site for both Hg(II) and Zn(II) on guanosine and inosine in DMSO.

The ^{15}N results for guanosine and inosine in the presence of Ba(II) show chemical shift changes of less than 0.8 ppm. Thus the heterocyclic nitrogens are not primary binding sites for the hard metal ion.

Binding may be outer-sphere, i.e. through intervening solvent molecules, or to an oxygen site. The primary oxygen binding site is

FIGURE 8: ^{15}N Results, Effects of divalent metal ions on the chemical shifts of guanosine. Positive values are downfield.

FIGURE 9: ^{15}N Results. Effect of divalent metal ions on the chemical shifts of inosine. Positive values are downfield.

at the C6 carbonyl, though ribose interactions ~~are~~ possible. Thus, the cis-diol 2', 3'-hydroxy groups in a ribose sugar are potential binding sites for the harder metal ions, in non-aqueous solutions such as DMSO (73, 74).

The binding site of Ba(II) on guanosine and inosine will be further studied using ^{13}C NMR.

4.12 Protonation

^{15}N results for the protonation of guanosine (45) and inosine are identical to the metal ion binding results. The pKa of N7 is ca. 2 for both ~~guanosine~~ guanosine and inosine, and the N1-proton does not dissociate in the aprotic solvent used (4).

The ^{15}N spectra of guanosine and inosine in the presence of trifluoroacetic acid (TFA) show large upfield shifts of 50 to 80 ppm for N7 as well as smaller downfield shifts of ca. 4 ppm for N9. The remaining resonances undergo only very small shifts, as shown in Figures 11 and 12.

The ^{15}N chemical shifts indicate N7 is also the protonation site for guanosine and inosine in DMSO.

FIGURE 11: ^{15}N Results. Effect of protonation on the chemical shifts of guanosine. Positive values are downfield.

FIGURE 12: ^{15}N Results. Effect of protonation on the chemical shifts of inosine. Positive values are downfield.

4.2 ^{13}C NMR Results

The metal ion complexes and protonated species of inosine and guanosine can also be characterized by their ^{13}C chemical shifts. This is particularly important for the hard metal ion Ba(II) which does not bind at a nitrogen site. The formation of carbonyl complexes will be investigated by following the chemical shift of C6.

For the nucleoside inosine no literature about its interaction with Zn(II) is available. In a recent ^{13}C study on the binding of $\text{CH}_3\text{Hg(II)}$ and Hg(II) to inosine, the base carbon assignments were incorrect (75). The resonances assigned to carbons 2 and 4 need to be reversed based on present results from a fully coupled spectrum. See Figure 13.

The ^1H decoupled ^{13}C spectrum of inosine is shown in Figure 14, it is typical of the standard ^{13}C spectra obtained in this study.

4.2.1 Divalent Metal Ion Complexes

The ^{13}C results for the complexation of inosine and guanosine with the divalent metal ions Hg(II) , Zn(II) and Ba(II) are presented in Figures 15 and 16.

The ^{13}C NMR results herein are generally in accord with the few existing data in the literature (39,40,75,76,77). It is generally recognized that downfield shifts of C8 and C1' along with an upfield shift of C5 define an N7 metal ion binding site.

These titration shifts are present in the spectra recorded for

FIGURE 14 ^{13}C spectrum of inosine in DMSO, ^1H decoupled.

FIGURE 15 ^{13}C Results Effect of divalent metal ions on the chemical shifts of inosine. Positive values are downfield.

FIGURE 16 ^{13}C Results Effect of divalent metal ions on the chemical shifts of guanosine
Positive values are downfield

inosine and guanosine when complexed with the metal ions. For both guanosine and inosine the largest changes in chemical shift occur in the presence of Zn(II), followed by Hg(II), and finally only small changes are observed for the Ba(II) complexes. The studies using Ba(II) salts are limited by their low solubilities in DMSO.

The ^{13}C results indicate Hg(II) and Zn(II) are complexed at the N7 site of both guanosine and inosine, in accord with observed ^{15}N data. The interaction of Ba(II) with N7 is possible, however the small changes in chemical shift observed make firm predications impossible.

To answer the question about the role O6 is playing in the metal ion complexation process, the chemical shifts of C6 and C2 are monitored.

As discussed previously in subsection 3.32, the chemical shift of the carbonyl carbon is sensitive to the distribution of charge within the carbonyl bond. An increase in charge density at the carbon will cause an upfield shift, whereas a decrease in charge density produces a downfield shift of the ^{13}C resonance. This effect originates in the r^{-3} term of the $\sigma_p(\text{loc})$ expression.

$$\sigma_p(\text{loc}) = \frac{-\mu_e e^2 h^2}{6 \pi m^2 A E E} \langle r^{-3} \rangle_p P_u \quad 23$$

This property will be useful in defining the degree of chelation, or interaction with the C6 carbonyl that a metal ion will exhibit. As shown earlier, Pullman (72) calculated that the most stable site of Na^+ interaction on guanine involved an N7-O6 chelate structure.

This is due primarily to the favourable Coulombic term for this site.

Chelation with the carbonyl is postulated to be more important for the harder metal species $Ba(II)$ and Na^+ than for the softer metals $Zn(II)$ and $Hg(II)$. $Ba(II)$, an alkaline earth metal, also has a need for more coordinated ligands than either $Zn(II)$ or $Hg(II)$. These two d^{10} species can exhibit reasonable stability with a coordination number of two or four.

The binding of an alkaline earth metal ion to a carbonyl site in a nucleoside has been previously reported for cytidine (77). In general the binding is characterized by the downfield shift of the carbonyl carbon. This is in contrast to the upfield shift observed when a metal ion binds to a nitrogen site adjacent to the carbonyl.

The ^{13}C chemical shifts observed for the C6 resonance in guanosine and inosine are dependent upon the metal ion involved. See Figures 15 and 16. For the 6-oxopurines interaction with $Hg(II)$ causes C6 to shift upfield. In the presence of $Zn(II)$ the resonance remains unchanged, whereas it is shifted downfield in the $Ba(II)$ complexes.

Discussed in terms of carbonyl interaction the $Hg(II)$ data indicate that no oxygen complexation is occurring. $Zn(II)$ a

borderline metal ion species has some interaction with the carbonyl, and Ba(II) the hard metal ion interacts directly with the C6 carbonyl group.

Thus the C6 carbonyl may be the primary binding site in the 6-oxopurines for very hard metal ion species. It is less sterically hindered and can accommodate the larger coordination sphere of these metals.

This result is supported by the ^{15}N results for inosine complexed with Ba(II) as shown in Figure 9. Small, approximately equal upfield shifts of N7 and N1 are predicted for a primary O6 interaction of the metal ion.

4.22 Protonation

The effects of protonation on the ^{13}C chemical shifts of inosine and guanosine are presented in Figures 17 and 18.

Despite the N7 protonation site indicated by theory (4) and the ^{15}N results, small upfield shifts of 0.30 ppm are observed for C8 in both nucleosides. The upfield shifts of C5 and downfield shifts of C1' are still characteristic of N7 binding.

At C8 there is no longer an inductive electron withdrawal, which produces the downfield shift. The positive charge remains localized on the protonated nitrogen and C8 is observed to shift slightly upfield due to increased electron density.

FIGURE 17 ^{13}C Results Effect of protonation on the chemical shifts of guanosine. Positive values are downfield.

FIGURE 18 ^{13}C Results Effect of protonation on the chemical shifts of inosine
Positive results are downfield

$R_1 = \text{H}$ inosine

$R_2 = \text{NH}_2$ guanosine

The C6 resonance in the guanosine mono-cation is shifted upfield by 1.87 ppm, whereas in the inosine cation it is shifted downfield by 2.84 ppm. Thus the charge distribution of the carbonyl bond is opposite in each of these species.

In the protonated cation of inosine, the downfield shift of C6 reflects decreased electron density and an increase in P_u (bond multiplicity). An enol tautomer of inosine may be formed.

$R = \text{Ribofuranoside}$

The fact that this species is not formed for the guanosine complex may be due to the increased basicity of the N1 position (4).

4.3 N7 Blocked 6-Oxopurine Nucleosides

The metal ion binding properties of 7-methylguanosine were studied using ^{13}C NMR, to determine the effects of blocking the primary binding site.

7-Methylguanosine exists as a zwitterion in DMSO solutions (78).

This will bring about changes in the ^{13}C chemical shifts when compared to guanosine, as shown in Figure 19.

The largest changes occur for the C6 and C2 resonances. The downfield shift of C6 is in response to the decrease in electron density at C6, which in turn causes a reduction in ρ , and an increase in $\rho_p(\text{loc})$ (33). A strictly steric interaction between the methyl substituent at N7 and C6 would have resulted in an upfield shift of the resonance (33).

The C2 resonance also undergoes a dramatic downfield shift of 3.0 ppm. This is due to the increased delocalization of the carbon atom electrons, and a reduction in ρ (33).

FIGURE 19 ^{13}C Results. Effect of N7-methylation on the chemical shifts of guanosine. Positive values are downfield.

It will be of interest to note the effects metal ions such as Hg(II), Zn(II), Mg(II) and $\text{CH}_3\text{Hg(II)}$ will have on this nucleoside. Protonation effects on the zwitterion will also be studied.

7-Methylguanosine allows one to investigate the presence of secondary metal ion complexation sites, because the primary N7 binding site is physically blocked.

The ^{13}C results for the metal ion complexation and protonation of 7-methylguanosine are presented in Figures 20 and 21.

It is interesting to note that the divalent metal ions have a maximum binding ratio of 1:2 (metal:ligand) before precipitation occurs, whereas the univalent species will bind up to a 1:1 ratio.

This suggests the formation of a 2:1 complex with Hg(II), Zn(II) and Mg(II). Complexation does not depend on the polarizability of the metal ion. Identical results are observed with the very soft metal ion $\text{CH}_3\text{Hg(II)}$ when compared with the Mg(II) complex.

The greatest changes observed involve the chemical shifts of C6 and C2, both of which were shifted upfield. An N1 site of complexation is predicted, which will cause the C6 carbonyl bond to reform.

$\text{R} = \text{Ribofuranoside}$

FIGURE 20: ^{13}C Results. Effect of divalent metal ions on the chemical shifts of 7-methylguanosine. Positive values are downfield.

FIGURE 21: ^{13}C Results Effect of protonation and methylmercury on the chemical shifts of 7-methylguanosine. Positive results are downfield.

4. H_2O and CO_2 dentate long are formed at high temperatures and
exposure to H_2O and CO_2 .

4.4 Conclusion

An initial study on the metal ion complexes of the 6-oxopurine nucleosides has served to answer a number of questions about metal ion nucleic acid interactions in general.

The determination of binding sites on inosine and guanosine for protonation and soft metal ion complexation, is made facile by ^{15}N NMR. Very large upfield shifts (ca. 10-80 ppm) are observed for the nitrogen atom involved in complexation.

The soft metal ions Hg(II) and Zn(II) , and the proton bind exclusively to the N7 position of inosine and guanosine in DMSO.

^{13}C NMR chemical shift changes, while not as large or unambiguous are in agreement with the ^{15}N NMR results for the divalent metal ions. However an important drawback to the use of ^{13}C chemical shifts for determining binding sites is demonstrated by the protonation results. The ^{13}C nuclei are generally one to three bonds removed from the actual nitrogen binding sites of the soft metal ions. Thus small changes occurring in the molecule far removed from the actual binding site will affect the ^{13}C chemical shift changes observed.

^{13}C NMR is however very important for determining the complexation site of the hard metal Ba(II) . Binding to a carbonyl group can be observed easily using ^{13}C NMR.

The binding site selected is dependent upon the polarizability of the metal ion involved. This is illustrated by comparing the degree of interaction a metal ion has with the carbonyl group of inosine or guanosine. With the soft metal ion Hg(II) no carbonyl

binding is detected, the primary binding site is N7. For the borderline (Zn(II) and Cu(II) some carbonyl interaction is seen, but the primary binding site remains N7. The hard metal ion Ba(II) interacts preferentially with the carbonyl oxygen. Very small effects on N1 and N3 are observed.

$R_1 =$ H inosine

$=$ NH_2 guanosine

$R_2 =$ ribofuranoside

If the favoured N7 binding site is blocked, as in the case of 7-methylguanosine, then binding at a secondary N1 site is observed.

The metal ion complexation properties of the 6-oxopurine nucleotides were also investigated. Once again ^{15}N and ^{13}C NMR chemical shifts were used to determine binding site selectivity. In addition, ^{31}P NMR was employed.

Several additional questions can be probed in this study:

1) the role of solvent, 2) the pH dependence of binding and 3) the effect of a single phosphate group on the complexation process.

The sodium salts of inosine-5'-monophosphate (5'IMP) and guanosine-5'-monophosphate (5'GMP) are no longer soluble in DMSO, so water becomes the solvent of choice. Particular attention must be paid to the ionization state of any potential metal ion binding site. The pH of the solutions was initially adjusted so that the base had the same electronic properties that it possessed in the DMSO-nucleoside solutions. The nucleotide samples were initially run at a pH between 7 and 8, where N1 is protonated and N7 deprotonated. This will mean however, that the phosphate group will be ionized and carry a charge of -2.

At pH 8 results obtained with the nucleotides in water will be compared to previous results using the nucleosides in DMSO. This will allow an estimation of solvent effects on the binding process.

The pH dependence of binding will be investigated by comparing the complexation sites observed at pH 8 and pH 2.

The dependence of complexation on the polarizability of the metal involved was investigated using the soft metal ion Hg(II) .

R = H inosine - 5' - monophosphate

= NH₂ guanosine - 5' - monophosphate

and the hard Mg(II) metal ion. Mg(II) replaces Ba(II) in the aqueous complexation studies due to its greater solubility.

Studies using Zn(II) can no longer be carried out in solution for the purine mononucleotides as insoluble complexes result. Zn(II), because of its borderline properties of hardness and softness occupies a unique position in nucleic acid chemistry. The balanced affinity of Zn(II) for a nitrogen or oxygen donor ligand means that it is most effective at promoting base-stacking (79). This unique binding interaction also serves to facilitate total DNA rewinding after denaturation (80).

In addition to these structural effects Zn(II) also promotes degradation of nucleic acid components (81). This will be further discussed in Section 7.3, where Zn(II) is observed to promote dephosphorylation of 5'ATP.

A recent solid-state X-ray structural characterization of the 5'IMP and 5'GMP complexes of Zn(II) have shown a base-stacked, N7

metallated structure. The phosphate group is involved in intramolecular hydrogen bonds to some of the coordinated water ligands (82).

It is known that very hard metal ions such as Ba(II) and Mg(II) show a preference to bind with the phosphate groups in DNA and the nucleoside triphosphates (83,84). However softer metal ions such as Hg(II), Zn(II), and Pt(II) will still interact primarily with the heterocyclic nitrogens (1,2,3).

The N1 nitrogen becomes a potential binding site in aqueous solutions, particularly at higher pH (40). Both N1 and N7 binding by Cu(II) to 5'IMP has been observed using paramagnetic relaxation techniques (85). FT-IR studies on 5'GMP have found that Cu(II) binds simultaneously to the phosphate group and N7, whereas Cd(II), Pt(II), and Mg(II) will bind only at N7 (86,87).

A few ^{15}N NMR studies have been done on the metal ion complexes of adenosine nucleotides. The 5'AMP complex with $\text{Pt}(\text{NH}_3)_2^{+2}$ has been characterized, and binding is found to occur at N7 (50). The N7 complex of Mn(II) and 5'AMP has also been studied using ^{15}N relaxation data (51). A pioneering study in ^{15}N NMR involved the complexation of 5'ATP with Zn(II) and Mg(II) (52). However ^{15}N studies on the metal ion complexes of the 6-oxopurine nucleotides have not yet been carried out.

^{31}P NMR has also been used previously to follow metal ion binding to adenosine-5'-triphosphate (87,88,89). In general a downfield shift of the phosphorous resonance indicates a metal ion interaction. This trend also holds for metal ion binding to the phosphorous groups in a DNA backbone (90).

5.1 ^{13}C NMR Results

It was immediately apparent that solvent and a phosphate group had a pronounced effect on the metal ion binding properties of 5'GMP. The nucleotide itself was insoluble in solutions below a pH of 5, and at pH 8 in the presence of divalent metal ions (i.e. Hg(II), Cd(II), Ba(II) and Zn(II)) an insoluble polymeric species was formed. This precipitate was soluble at higher pH, (approximately 12) however ^{13}C NMR indicated a breakdown of the nucleotide structure under these conditions.

Due to the limited solubility of 5'IMP in aqueous solution, ca. 0.5 M at pH 8, ^{13}C titration studies were initially undertaken. ^{15}N spectra were obtained at 30.41 MHz to identify the site of metal ion complexation.

The effects of Hg(II) and Mg(II) on the ^{13}C resonances of 5'IMP at pH 8 and 2 are presented in Figure 22.

In the case of Mg(II) complexed with 5'IMP at pH 8, downfield shifts of 0.3 and 0.1 ppm are observed for C1' and C8 respectively, and an upfield shift of 0.1 ppm is observed for C5. These shifts, while characteristic of an N7 metal ion binding site, are too small to unequivocally assign an N7 binding site for Mg(II) on 5'IMP at pH 8. The interaction of Mg(II) at N7 may be outer-sphere, or solvent mediated.

The upfield shifts of C2 and C6 indicate that no N1 or carbonyl interactions are occurring.

If the ^{13}C results obtained for Hg(II) interacting with 5'IMP at pH 8 are compared to the results obtained for the nucleoside, as

FIGURE 22: ^{13}C Results. Effect of divalent metal ions on the chemical shifts of inosine-5'-monophosphate. Positive results are downfield.

9.

downfield shift of 16 occur upon the addition of 1.0 equivalent of AgCl.

The downfield shifts observed arise due to changes in ρ_{loc} ; E and r decrease while an increase in ρ_{loc} occurs.

$$\sigma_p(\text{loc}) = \frac{-\mu_0 e^2 \hbar^2}{6\pi m^2 AEE} \langle r^{-3} \rangle_p P_u \quad 23$$

These changes are due to the deprotonation of N1 and the binding of a metal ion at this position. The deprotonation at N1 leads to an increase in conjugation at C6, and increased delocalization of charge at both C2 and C6. The binding of a metal ion will moderate these effects and residual downfield shifts are expected.

This unexpected and previously unstudied binding site will be further characterized by ^{15}N NMR.

The interaction between Hg(II) and 5'IMP at pH 2 produces small (less than 0.25 ppm) upfield shifts of all the base carbon resonances. These shifts indicate that Hg(II) does not interact with the purine base, or the ribose sugar. The N1 binding site is blocked by lowering the pH of the solution.

5.2 ^{15}N NMR Results

In order to confirm the presence of N1 bound Hg(II) in 5'IMP solutions at pH 8, ^{15}N spectra were recorded at 30.41 MHz. Results are presented in Figure 23.

At 0.5 equivalents of Hg(II), a 2.2 ppm upfield shift of N7 is observed. Downfield shifts of 26.7, 1.7 and 0.6 ppm are observed for N1, N3 and N9 respectively. Clearly N1 is the resonance most affected by the addition of Hg(II). As will be discussed more fully in Chapter 10, the results indicate a deprotonation of N1 followed by metal ion complexation.

The ^{15}N shifts are in agreement with the ^{13}C results for an N1 Hg(II) binding site at pH 8. A small amount of N7 complexation may also be occurring given the shifts of N7 and N9.

5.3 ^{31}P NMR Results

The ^{31}P spectra of 5'IMP complexed with Mg(II) and Hg(II) at pH 8 show only upfield shifts, 0.7 ppm with Mg(II) and 2.6 ppm with Hg(II). See Figure 24.

Metal ion binding at the phosphate group is expected to cause a downfield shift of the ^{31}P resonance, due to a decrease in $\langle r^{-2} \rangle$; (refer to sub-section 3.32).

Therefore no phosphate binding is apparent between Hg(II) or Mg(II) and 5'IMP at pH 8.

FIGURE 23: ^{15}N Results. Effect of $\text{Hg}(\text{II})$ on the chemical shifts of inosine-5'-monophosphate. pH 8.
Positive values are downfield.

FIGURE 24: ^{31}P Results. Effect of divalent metal ions on the chemical shift of inosine-5'-monophosphate. pH 8. Negative values are upfield.

2

5.4 Conclusion

The effects of solvent on the metal ion complexes of the hypoxanthine base can be studied by comparing the binding properties of inosine in DMSO to those observed for 5'IMP in aqueous solutions.

In contrast to the N7 binding observed for the soft metal ions on the purine nucleosides in DMSO, an N1 complex of Hg(II) and 5'IMP is detected by ^{15}N and ^{13}C NMR at pH 8.

Deprotonation of N1, followed by metal ion complexation is facilitated in the slightly basic aqueous solution. Previous N1 binding of Hg(II) had been suggested for inosine in the presence of a nitrogen base, however complications arise due to interaction between metal ion and base (40). No such problems occur in the aqueous samples of 5'IMP.

N1 has been identified as the thermodynamically favoured metal ion binding site (40).

The hard metal ion Mg(II) no longer shows a direct interaction with the heterocyclic base, or the phosphate. Indirect, solvent-mediated, or outer-sphere binding at N7 is possible however. No carbonyl binding is observed in aqueous solution. The formation of metal ion hydrates (e.g. $\text{Mg}(\text{H}_2\text{O})_6^{+2}$) and solvent-ligand hydrogen bonds have reduced the complexing ability of the hard metal ion in aqueous solutions.

Binding of Hg(II) in aqueous solutions has a pH dependence related to the ionization states of the potential nitrogen donor sites.

If the pH of a solution is substantially lower than the pK_a of a potential binding site, no interaction is observed.

The presence of a single phosphate group in the mononucleotide does not influence the complexation process. No metal ion-phosphate interactions are detected. Even the $Mg(II)$ -5'IMP mixture, which shows no binding site on the nucleoside fragment, does not exhibit any phosphate interactions.

CHAPTER 6:

ADENOSINE

A third purine nucleoside, adenosine was complexed with the three divalent metal ions Hg(II) , Zn(II) and Mg(II) . The binding sites of these metal ions were characterized by their ^{15}N and ^{13}C NMR spectra.

Adenosine differs from the 6-oxapurines in that the heterocyclic base is fully aromatic. Thus there are two potential nitrogen binding sites, N1 and N7.

The theoretical calculations of Pullman (72) on the preferred binding sites of Na^+ on the adenine base indicate N1 is the most stable binding site, followed by N3 and then N7, as shown below.

N3 is not a potential binding site in the nucleosides or nucleotides due to steric hinderance by the ribose ring. N7 and N1 exhibit similar reactivities in the adenosine species due to the lower pKa of N1 when compared to the 6-oxopurines. See Table 1 (1). Competition between N1 and N7 for a metal ion is expected (3).

Adenosine-metal ion interactions have been extensively studied in the past (2, 39, 91, 92). Many of these studies occurred prior to 1972 and involved techniques such as pH titration, UV spectroscopy, thermometric titrations and temperature jump methods. These methods however, cannot identify specific metal ion binding sites, only association constants. Improvements are seen with the introduction of X-ray diffraction and ^1H NMR techniques (91). Adenosine has two non-exchangeable proton resonances on the purine ring. Thus binding to the 6-membered or 5-membered ring could be distinguished (2,91).

Further refinements in determining the site of metal ion interaction are observed with ^{13}C NMR (39, 76). The ^{13}C NMR parameters for an N7 binding site have been clearly established (39, 76). However the extent that an N1 binding reaction may be occurring has not been fully answered (39, 76).

6.1 ^{15}N NMR Results

The ^{15}N NMR results for the complexation of adenosine with Hg(II) and Zn(II) have been previously published by workers from this laboratory (53). Additional results with the hard metal Mg(II) were recorded for this study. Large chemical shift changes are observed in the presence of Zn(II) and Hg(II) , whereas only small shifts are observed upon the addition of Mg(II) . See Figure 25.

In the presence of the soft metal ion Hg(II) , a large upfield shift of 9.3 ppm is observed for N1, with a smaller 1.0 ppm upfield shift of N7. The two conjugated nitrogens N3 and N9 are shifted downfield, 1.0 and 3.0 ppm respectively. The amino resonance also undergoes a 2.4 ppm downfield shift upon the addition of Hg(II) (53).

In the Zn(II) -adenosine complex, N7 is shifted upfield by 7.5 ppm, while N9 moves downfield. Very small upfield shifts are observed for the N1 and N3 resonances, while the NH_2 group is shifted downfield 1.0 ppm (53).

In the presence of Mg(II) , the N7 and N1 resonances of adenosine are shifted upfield 1.5 ppm. The N9 and N3 resonances also shift upfield, though less than 0.5 ppm, while the amino resonance is shifted downfield 0.3 ppm.

In the presence of the softer metal ions Hg(II) and Zn(II) two distinct binding sites are observed. As discussed previously the binding of an electropositive species at an sp^2 nitrogen, is expected to increase $\langle r \rangle$ and AEE while decreasing P_u . This will render the $\sigma_p(\text{loc})$ term less negative, thereby shifting the resonance upfield.

$$\sigma_p(\text{loc}) = \frac{-\mu_0 e^2 h^2}{6 \pi m^2 AEE} \langle r^{-3} \rangle_p P_u \quad 23$$

FIGURE 25: ^{15}N Results. Effect of divalent metal ions on the chemical shifts of adenosine. Positive values are downfield.

82
A nitrogen conjugated to this binding site is expected to shift downfield due to a reduction in $\langle r \rangle$.

Thus in adenosine two binding sites are observed: Mg(II) binds largely to N1, with a small degree of N7 complex being formed. Zn(II), on the other hand binds exclusively to the N7 site.

The ^{15}N shifts of the adenosine Mg(II) complex are too small to conclude that a direct nitrogen-metal ion interaction is occurring. A secondary, or outer-sphere binding mode may be preferred by this hard metal ion. An outer sphere complex has been suggested for the Ba(II)-5'AMP complex on the basis of ^{15}N spin lattice relaxation measurements (51). Binding may also be occurring at the harder oxygen donor atoms of the ribose sugar. Further studies on the Mg(II) complex of adenosine will be made using ^{13}C NMR.

6.2 ^{13}C NMR Results

As discussed earlier, in Section 4.2 an N7 metal ion binding site is characterized by a downfield shift of C8 and C1' accompanied by an upfield shift of C5.

There is a problem however in defining an N1 binding site using ^{13}C chemical shifts (39,76). Using the clear example of N1 complexation afforded by the Hg(II)-adenosine complex, this question will be investigated.

The ^{13}C NMR results for adenosine complexed with Hg(II), Zn(II) and Mg(II) are presented in Figure 26.

One can use theoretical ^{13}C chemical shift substituent effects to accurately predict the effects of metal ion complexation at N1 in adenosine.

Protonation and metallation will produce similar changes in the carbon resonances adjacent to the nitrogen binding site. Shielding of these resonances upon protonation is attributed to a decrease in the N-C bond order and an increase in the AEE term in $\sigma_p(\text{loc})$ (32).

This is indeed what is observed for the Hg(II) complex of adenosine, as shown in Figure 26. The upfield shifts of C2 and C6 are now greater than the shift of C5, in contrast to the previous results for an N7 complex.

FIGURE 2.6: ^{13}C Results. Effect of divalent metal ions on the chemical shifts of adenosine. Positive values are downfield.

Thus the N1 binding site of Hg(II) on adenosine indicated by ^{15}N NMR can be confirmed using ^{13}C chemical shifts. The existence of some N7 complexed Hg(II) is also shown by the ^{13}C NMR results.

For the Zn(II) complex with adenosine, C2 no longer moves upfield, therefore no N1 binding is indicated and the results confirm the ^{15}N NMR data. The downfield shifts of C8 and C1' and the upfield shift of C5 are consistent with the existence of an N7 binding site.

The ^{13}C results for Hg(II) and adenosine indicate neither a direct N1 nor N7 binding site is occupied. An indirect, solvent mediated interaction with N7, is possible however. Complexation with a ribose oxygen is ruled out, based on the very small chemical shift changes observed, less than 0.10 ppm.

6.3 Conclusion

Adenosine, with its aromatic purine base has two potential metal ion binding sites N1 and N7. By observing the ^{15}N chemical shifts of an adenosine-metal ion complex, binding to each of these sites was distinguished. This allowed also, the development of a ^{13}C NMR probe to the N1 binding site, not previously available (39,75).

It was observed that Hg(II) will bind preferentially to the N1 of adenosine, with some N7 product being detected, whereas, Zn(II) complexes exclusively at the N7 binding site.

The N7 site of adenosine remains theoretically softer than the N1 site, by virtue of its lower pK_a (1,10). The N1 site however, produces the most stable metal ion complex, based on the *ab initio* calculations of Pullman (72).

Thus N7 can be treated as the kinetically favoured binding site and N1 the thermodynamically favoured site.

The polarizability of the metal ions; Hg(II) and Zn(II) are similar enough, as is the relative softness of N7 versus N1, that it is not a controlling factor in the binding of these metal ions to adenosine.

Site selectivity for Hg(II) and Zn(II) is controlled by complex stability and steric hindrance. The larger metal ion Hg(II) prefers the less hindered N1 binding site.

Metal ion polarizability is however important when discussing the hard metal ion Mg(II) , and its interaction with adenosine. The lack of a hard binding site on adenosine prevents the formation of a direct Mg(II) complex.

Previously, N1 complexes of adenosine and the soft metal Pd(II) have been observed (93). In addition, dimetallated species of adenosine and Pt(II) have been reported at greater than 1.0 equivalent of metal ion (94).

The effects of a phosphate group on the metal ion binding properties of adenosine were determined using ^{15}N , ^{13}C , and ^{31}P NMR. Two adenosine nucleotides were employed in this study; adenosine-5'-monophosphate and adenosine-5'-triphosphate.

7.1 Adenosine-5'-monophosphate

As observed previously for the 6-oxopurine nucleosides, the addition of a phosphate group can change the metal ion binding properties of the base. See Chapter 5.

The sodium salt of adenosine-5'-monophosphate (5'AMP) was used in this study. Its limited solubility in DMSO and acidic solutions, required that NMR samples be prepared at pH 10. Though this leaves the phosphate group with a -2 charge, the base is isoelectric with its counterpart in DMSO.

The binding of both a hard metal ion $Mg(II)$ and a soft metal ion $Hg(II)$ will be examined.

The 5'AMP complex with $Pt(NH_3)_2^{2+}$ has recently been characterized using ^{15}N NMR, with an N7 binding site identified (50). The N7 site is also preferred by $Mn(II)$ (51).

^{13}C and 1H NMR have also been used to characterize metal ion binding sites of 5'AMP. $Pd(II)$ and $Pt(II)$ are found to prefer an N7 binding site (95).

7.11 ^{15}N NMR Results

The ^{15}N NMR results for the interaction of divalent metal ions with 5'AMP are shown in Figure 27.

The interaction of $Zn(II)$ and 5'AMP produces an immediate precipitate. This has been observed previously and is specific for $Zn(II)$ and $Pb(II)$ (79). This precipitate is due to polymer formation, caused by base stacking interactions and simultaneous base and phosphate binding by the metal ion (96).

In the $Hg(II)$ -5'AMP complex, the 27.0 ppm upfield shift of N1 and the 4.0 ppm downfield shift of N3, indicate that the metal ion binds preferentially at the N1 site. The ca. 2 ppm downfield shifts of both N7 and N9 are characteristic of no N7 complex formation.

With the hard metal ion $Mg(II)$, the limited solubility of the $Mg(II)$ salt restricted studies to 0.30 equivalents of metal ion to nucleotide. A 2.5 ppm upfield shift of N1, and a 0.2 ppm upfield shift of N7 are observed. In addition the N3, N9, and amino resonances are shifted downfield ca. 0.3 ppm.

FIGURE 27: ¹⁵N Results. Effect of divalent metal ions on the chemical shifts of adenosine-5'-monophosphate. pH 10. Positive values are downfield

These results would seem to indicate a degree of N1-Mg(II) complex is being formed, with a very minor component of N7 complexed metal ion observed.

7.12 ^{13}C NMR Results

The ^{13}C results for the Mg(II) and Hg(II) complexes of 5'AMP are shown in Figure 28. These results confirm the binding sites identified by ^{15}N NMR.

In the case of Hg(II), the 5'AMP complex shows 0.2 ppm upfield shifts at C2 and C6, characteristics of an N1 binding site. The 0.6 ppm downfield shift of C5 indicates that no N7 Hg(II) bound 5'AMP is present.

The chemical shift changes observed for the Mg(II)-5'AMP complex are limited by the solubility of the salt. The 0.2 ppm upfield shifts of C2 and C6 are characteristic of an N1 complex. Binding of the hard metal ion to a ribose oxygen is not considered likely due to the very small shifts observed, less than 0.02 ppm.

7.13 ^{31}P NMR Results

The ^{31}P NMR results for the Hg(II) and Mg(II) complexes of 5'AMP, are shown in Figure 29.

As discussed previously, the interaction between a metal ion and a phosphate oxygen is expected to cause a downfield shift of the ^{31}P resonance, due to a decrease in $\langle r^{-2} \rangle$.

Hg(II) causes the phosphorus resonance of 5'AMP to shift upfield 0.7 ppm at 1.0 equivalent of metal ion. In the presence of

FIGURE 28 : ^{13}C Results. Effect of divalent metal ions on the chemical shifts of adenosine-5'-monophosphate. pH 10. Positive values are downfield.

FIGURE 29: ^{31}P Results. Effect of divalent metal ions on the chemical shift of adenosine-5'-monophosphate. pH 10. Negative values are upfield.

0.30 equivalents of Mg(II) the phosphorus signal is shifted upfield 0.25 ppm.

No phosphate binding is observed when Hg(II) or Mg(II) are complexed with 5'AMP.

7.14 Conclusion

Both ^{15}N and ^{13}C NMR indicate that N1 is the preferred site of Mg(II) and Hg(II) complexation on 5'AMP.

The N1 site combined with a dominant anti conformation of the nucleotides prohibits any phosphate binding from occurring.

7.2 Metal Ion Complexes of Adenosine-5'-Triphosphate

Metal ions play an important role in the hydrolysis of adenosine-5'-triphosphate (5'ATP) and its release of energy for cellular reactions (97). The origin of 5'ATP as a source of energy has been proposed to lie in the relative orientations of the base and triphosphate chain (97). The presence of a metal ion serves to promote and stabilize this particular conformation, as shown below (98). Thus the metal ion

complexes of 5'ATP have been extensively studied in the past (2,87,88, 89,90,98,99,100,101).

A series of metal ions were studied; Hg(II), Zn(II), and Mg(II) to determine the effects of metal ion polarizability on the metal ion binding site. The role of the phosphate group as a multi-site metal ion ligand was also investigated. The effect of pH on the binding process can also be studied, as 5'ATP is sufficiently soluble for NMR

studies over a wide range of pH.

Similar to adenosine and 5'AMP, the N7 site is the kinetically favoured binding site, and N1 the thermodynamically favoured binding site. However the formation of a chelated metal ion site between the phosphate chain and purine base will also be energetically favoured over single ligand metal ion binding sites, such as N1 or N7 (9).

7.21 ^{15}N NMR Results

To allow comparison with the results obtained for 5'AMP, the ^{15}N NMR spectra of 5'ATP complexed with Hg(II), Zn(II) and Mg(II) were recorded at pH 10. Results are presented in Figure 30.

When in the presence of Hg(II), all the ^{15}N resonances of 5'ATP were shifted downfield, thus no interaction with the purine nitrogens at pH 10 is indicated.

At pH 7, Raman difference studies have indicated an N1 Hg(II) complex of 5'ATP is formed (100), although N7 binding was not ruled out.

At pH 7, 5'ATP is not sufficiently soluble to obtain an ^{15}N NMR spectrum, even at 30.42 MHz. Accordingly, studies on the Hg(II) complex of 5'ATP will be done using ^{13}C and ^{31}P NMR only.

The 5'ATP complex formed in the presence of 0.30 equivalent of Zn(II) at pH 10, produced a 4.0 ppm upfield shift of N7. A 0.4 ppm downfield shift of N9 was also observed. Upfield shifts of less than 0.2 ppm were observed for N1 and N3, while the amino group is shifted downfield 1.1 ppm.

FIGURE 30: ¹⁵N Results. Effect of divalent metal ions on the chemical shifts of adenosine-5'-triphosphate, pH 7 and 10. Positive values are downfield.

The ^{15}N data indicate Zn(II) is complexed at the N7 position of 5'ATP at pH 10, with no N1 binding detected. This N7 complex between Zn(II) and 5'ATP has been previously observed by ^{15}N NMR (52).

In the presence of 1.0 equivalent of Mg(II) at pH 10, the N1 and N7 resonances of 5'ATP were shifted upfield 0.9 ppm. The amino and N9 resonances are shifted downfield 0.2 ppm, and N3 undergoes a slight, 0.1 ppm upfield shift. From these shifts it is possible to conclude that Mg(II) does not bind directly to either N1 or N7.

To determine if Mg(II) will complex with 5'ATP at pH 10, further studies using ^{13}C and ^{31}P NMR are required.

7.22 ^{13}C NMR Results

The ^{13}C results obtained for 5'ATP complexed with Zn(II) and Mg(II) at pH 10 are presented in Figure 31. As the Hg(II) complex of 5'ATP is not stable at high pH, it was observed at pH 7. See Figure 31.

The ^{31}P results will indicate a pH dependence for the interaction of Zn(II) with 5'ATP exists. Therefore, the ^{13}C results for Hg(II) and Zn(II) were also obtained at pH 4. See Figure 32.

The Hg(II) complex of 5'ATP at pH 7 shows ca. 0.15 ppm upfield shifts of C2 and C6, and downfield shifts of 0.44, 0.56 and 0.79 ppm for C1', C8 and C5 respectively. These results are similar to those observed for the Hg(II) 5'AMP complex as shown in Figure 28.

As previously discussed, these shifts are characteristic of an N1 bound metal ion, with no minor N7 component.

FIGURE 31: ^{13}C Results. Effect of divalent metal ions on the chemical shifts of adenosine-5'-triphosphate, pH 7 and pH 10. Positive values are downfield.

FIGURE 32: ^{13}C Results. Effect of $Zn(II)$ on the chemical shifts of adenosine-5'-triphosphate, pH 4. Positive values are downfield.

At pH 4, no binding between Hg(II) and 5'ATP is observed. All the adenine base carbons are shifted upfield. This may be attributed to competition between the metal ion and protons for the N1 binding site. The pK_a of N1 is 3.5 and significant protonation is expected to occur at pH 4.

The Zn(II) complex of 5'ATP shows the ^{13}C NMR characteristics of an N7 binding site at both pH 4 and pH 10. The C8 and C1' resonances are shifted downfield 0.7 to 1.0 ppm, while C5 is shifted upfield. There is no indication of any N1 binding, in agreement with observed ^{15}N NMR shifts.

For the hard metal ion Mg(II) the ^{13}C shift differences were too small, (less than 0.08 ppm) to conclude that binding to the purine base or ribose sugar was occurring. The interaction between Mg(II) and 5'ATP therefore appears to occur at the phosphate group (vide infra).

7.23 ^{31}P NMR Results

The ^{31}P spectra of the complexes between 5'ATP and Zn(II) and Mg(II) were recorded at pH 10 and pH 4. The results are presented in Figures 33 and 34. The ^{31}P spectra were recorded at pH 7 for the Hg(II)-5'ATP species, as shown in Figure 33.

The upfield shifts of all three 5'ATP phosphorous resonances in the presence of Hg(II) at pH 7, confirms the lack of a phosphate binding site for the soft metal ion species. This is appropriate for an N1 bound metal species.

FIGURE 33: ^{31}P Results. Effect of divalent metal ions on the chemical shifts of adenosine-5'-triphosphate, pH 7 and pH 10. Positive values are downfield.

FIGURE 34 : ^{31}P Results. Effect of divalent metal ions on the chemical shifts of adenosine-5'-triphosphate, pH 4. Positive values are downfield.

The ^{31}P NMR spectrum of 5'ATP in the presence of Zn(II) at pH 10 is shown in Figure 35. In addition to phosphorous peaks arising from 5'ATP, additional peaks are present due to the unstable nature of the Zn(II)-5'ATP complex.

As discussed previously, the phosphorous resonance will be shifted downfield when complexed with a metal ion, due to a decrease in $\langle r^2 \rangle$.

Previous ^{31}P NMR studies on the interaction of metal ions with 5'ATP have resulted in different interpretations for the site of complexation on the triphosphate chain (89,102). When the changing pK_a of the phosphate group is taken into consideration, Mg(II) is found to bind only at the β phosphate residue (89).

The results presented in Figures 33 and 34 confirm this. At pH 4 and 10, only the β phosphorous resonance is shifted downfield, ca. 0.5 and 1.0 ppm respectively.

The ^{31}P results for the Zn(II) complex of 5'ATP indicate a pH dependence for the binding process. At pH 10, the γ phosphate residue is shifted upfield 2.0 ppm, while the β resonance shifts downfield 1.0 ppm and the α resonance is shifted 0.5 ppm downfield. This would indicate Zn(II) is involved with both the α and β phosphate positions of 5'ATP at pH 10.

At pH 4, a different Zn(II) complex of 5'ATP is formed, according to the ^{31}P results. A 2.0 ppm downfield shift of the β resonance and a smaller downfield shift of the γ phosphate group, indicate complexation is occurring at these sites. The α phosphate group now shifts 0.2 ppm upfield, a sign that it is no longer involved

FIGURE 35: ^{32}P spectrum of adenosine-5'-triphosphate with 0.50 eq. Zn(II) present.
pH 10.

with the Zn(II) metal ion.

7.24 Conclusion

In contrast to the results obtained for the monophosphate species; 5'IMP and 5'AMP, the phosphate group is now a potential metal ion binding site. Binding to the phosphate chain, which is a ~~hard~~ ligand is controlled by the polarizability, or softness, of the metal ion.

The very soft metal ion Hg(II) binds exclusively to the N1 position of 5'ATP at pH 7. This is the same binding site as was observed for the Hg(II) complexes of adenosine and 5'AMP. The phosphate group does not affect the binding of a soft metal ion to a soft nitrogen ligand.

The borderline metal ion species Zn(II) exhibits some very interesting binding phenomena. At pH 10 and 4, the ^{15}N and ^{13}C NMR data indicate that Zn(II) occupies an N7 binding site on the purine base. However, because of its borderline properties between hardness and softness, the Zn(II) metal ion can interact simultaneously with the triphosphate chain (52,102). This interaction is confirmed by the ^{31}P NMR results for the complex. The nature of the complex formed is pH dependent. At pH 10, the Zn(II) ion is bound at N7 and the α and β phosphate groups. At pH 4, the β and γ phosphate groups and N7 are involved in the Zn(II)-5'ATP complex. These macrochelate complexes of Zn(II) may play an important role in cellular reactions, as will be discussed in the next section.

The hard metal ion $Mg(II)$, does not have a complexation site on the purine base or ribose ring, as indicated by ^{15}N and ^{13}C NMR. It does however, bind to the α phosphate group of the triphosphate chain.

Binding is not dependent upon pH, as long as the pH is greater than the pK_1 of the phosphate (1.2) (1).

7.3 Dephosphorylation of 5'ATP

In studying the process of metal ion complexation with 5'ATP, using ^{31}P NMR, time dependent spectra were obtained. The complex formed between Zn(II) and 5'ATP at pH 10 was found to be unstable. This is illustrated in the ^{31}P spectrum of the complex, as shown in Figure 35. A resonance at 3 ppm, corresponding to free HPO_4^{-2} increases in intensity with increasing time. In addition phosphate resonances corresponding to an adenosine-5'-diphosphate-Zn(II) species appear.

The hard metal ion Mg(II) did not promote this dephosphorylation. In addition both metal ions formed stable 5'ATP complexes at pH 4.

The dephosphorylation of 5' trinucleotides in the presence of metal ions has been observed previously (101,103). The increased concentration of inorganic phosphate was measured as a function of metal ion presence. No structural details about the reaction intermediate were collected, however in these studies (101,103).

In the present ^{31}P NMR study on the metal ion promoted dephosphorylation of 5'ATP, a reaction intermediate is proposed. The role of metal ion polarizability is also discussed.

Phosphoryl transfer in nucleotides can be described as a loosely coupled bimolecular $\text{S}_{\text{N}}2$ reaction (101). A trigonal bipyramidal geometry is required and a metal ion can play several roles (3 vol. 5).

In general terms, the reaction pathway can be represented as below

where for example X, Y and Z are all oxygen atoms and R a phosphorous.

The metal ion may act purely in a charge neutralization capacity, allowing the approach of a nucleophile to the anionic phosphate chain (104). In many systems the coordination of a single, or second metal ion to the leaving group itself enhances its rate of leaving by increasing its acidity (101).

The metal ion may also serve to bring the reacting species together, as a template does. This is particularly true for hydrolysis reactions, where the nucleophile is itself a metal ion ligand. In some cases, the nucleophile is activated in this process, dependent upon the polarizability of the metal ion involved.

The hard metal ions, those with very low polarizing power are located further to the left and top of the periodic table (9). The softer metal ions, to the right and lower in the periodic table are much more polarizable.

Thus Zn(II) is a much more polarizing metal ion than Mg(II), both are at the extremes of the transition metal group (d^{10} and d^0 respectively). Zn(II) is a good example of a strong Lewis acid catalyst, it will fully accept electron pairs. Mg(II), while still a good catalyst, is best for substrates involving weaker base centers.

In an aqueous solution, the polarizing ability of a metal ion is very important. Metal ions will always exist in a solvated form, but their polarizing power dictates the nature of the ligands.

For metals with low polarizability, the hard metal ions such as Mg(II), a simple hexahydrate is observed (9). As the polarizing power of the metal ion increases it will fully gain an electron pair from the ligands. In the case of water, a proton is lost, and the resulting ligand is the hydroxide ion. Thus Zn(II), but not Mg(II) can serve to activate water as a nucleophile.

The next step, in which a second electron pair is lost to the metal ion and an oxide ligand produced is not observed with the higher d metals (d^8 , d^9 , d^{10}). This is due to the instability of the higher metal ion oxidation states necessary to achieve a high enough effective charge on the metal (9).

Another metal ion property which is important in catalysis, is the rate at which ligand substitution occurs. The rate of hydrolysis is much greater for Zn(II) compared with Mg(II). Also required of metal ion catalysts is a degree of coordination flexibility, again Zn(II) possesses more bonding freedom than Mg(II).

These properties are reflected in the degree to which a metal ion will promote the dephosphorylation of 5'ATP. The rate of metal ion catalysis of dephosphorylation is found to be: (98)

The most important factor however, is that each metal ion promoting dephosphorylation interacts with both N7 on the base and

the phosphate chain.

The formation of this mixed ligand complex is a property of the softer, more polarizable metal ions to the right of the periodic table, excluding Mg(II).

This interaction tends to stabilize the complex formed, bringing a metal ion with its activated water ligand close enough to cause γ -dephosphorylation.

The proposed reaction intermediate is supported by ^{31}P , ^{15}N and ^{13}C NMR studies, as discussed previously. Thus with a multinuclear NMR approach the interaction between metal ions and nucleic acid derivatives can be probed not only from a structural aspect, but also in terms of function.

Another method of studying the interaction of the soft Hg(II) species with the nucleosides is to examine ¹⁹⁹Hg NMR spectral behaviour. Shifts of this nucleus are expected to be sensitive to ligand substitutions, similar to the other heavy nuclei, ¹³C and ³¹P. The equation for $\sigma_p(\text{loc})$ developed by Pople (59) can be applied to ¹⁹⁹Hg chemical shifts also.

The ¹⁹⁹Hg NMR spectra were recorded for the purine nucleosides adenosine, guanosine, and inosine at a number of metal:nucleoside ratios. The results are presented in Figure 36.

As ¹⁹⁹Hg chemical shifts are known to have a large dependence upon concentration (22,23,27) and this effect was studied as well. See Figure 37.

A reference value for 1.0 M HgCl₂ in DMSO of -1501.6 ppm relative to external, neat dimethylmercury was used to calibrate the ¹⁹⁹Hg chemical shifts (23). All results reported are converted to this reference scale, although experimentally an external reference of .55 M HgCl₂ in DMSO was used.

An infinite dilution shift of -1464.0 ppm and a titration shift equal to 18.7 ppm upfield per mmole of Hg(II) were observed for HgCl₂ in DMSO.

Differences observed in these parameters in the presence of the nucleosides are taken to be indicative of a complexation reaction. These differences in the ¹⁹⁹Hg chemical shifts in the presence of a nucleoside are presented in Figure 36. An expected shift based only on the concentration of HgCl₂ is calculated using the expression:

FIGURE 36: ^{199}Hg Results. Effect of nucleosides on the chemical shift of HgCl_2 .

FIGURE 37: The concentration dependence of the ^{199}Hg chemical shifts in HgCl_2 .

$$\delta_{ex} = -14644 + -18.7 \times \text{mmoles Hg(II)} \quad \dots, 28$$

The difference between this value and the observed shift will provide information about the complexation process. A positive value represents a downfield shift of the ^{199}Hg resonance.

In the ^{199}Hg spectra of these complexes only a single resonance line is observed, due to a fast exchange rate on the NMR time scale, between solvated and ligand bound mercury. At no time are ^{199}Hg - ^{15}N couplings observed.

Each of the nucleosides used affected the ^{199}Hg chemical shift in a linear fashion, as shown in Figure 36. It is of interest to note the change in slope occurring in the inosine samples when more than 1.0 eq of metal is added. This is interpreted to mean that the first equivalent of Hg(II) is bound specifically to the N7 site as indicated by ^{15}N and ^{13}C NMR (54). At greater than 1.0 eq of metal other sites may be complexed to a varying degree.

Although its observation is limited by solubility, the same phenomenon is observed in the presence of guanosine. A change in slope is found at 1.0 eq metal ion. ^{15}N NMR results again indicate a dominant N7 binding site for the first equivalent of metal ion (53).

Only adenosine does not show any change in slope at greater than 1.0 eq of metal ion. Hg(II) is known to bind at the N1 position in adenosine, as determined by ^{15}N and ^{13}C NMR (53).

The N7 and N1 binding sites in adenosine are expected to have similar reactivities, based on their pK_a values. Thus no change in slope will be observed with the binding of a second equivalent of metal ion.

Previous ^{13}C results have suggested that both N1 and N7 binding is possible for inosine. These studies were however, carried out in aqueous solution in the presence of a base (40). ^{199}Hg NMR has thus provided the first hints that secondary, weaker binding sites may be important at very high metal ion concentrations.

CHAPTER 9:

DINUCLEOSIDE MONOPHOSPHATES

The aim of most studies involving metal ions and the nucleic acid derivatives is to determine the binding sites of these species in a living cell (in vivo). Thus the logical progression from nucleoside and nucleotide studies is to the dinucleoside monophosphate or oligonucleoside level.

Several investigations have been carried out on the metal binding properties of ApU and dApdT (105). Most studies involve the soft metal species such as Hg(II) or cis-Diammine-dichloroplatinum (II), where binding is likely to occur at the nitrogen sites of the bases.

The dinucleoside monophosphate cytidylyl-(3'→5')-guanosine (CpG) was chosen for this study because of its ability to form a base-paired mini helix in concentrated aqueous solutions, as depicted in Figure 38 (105,106).

This complex will provide a model system for RNA and DNA metal ion interactions. Longer segments of oligonucleosides would provide a more stable duplex model, but the complexity of their NMR spectra rules out their use as probes of metal ion binding sites, using presently available spectrometers.

The precipitation reaction observed between 5'GMP and the divalent metal ions such Hg(II), Zn(II) and Mg(II) is also observed at the dinucleoside level. Thus, a unifunctional soft metal ion species was desired for these studies.

The methylmercury ion was chosen and because of its very soft

FIGURE 38 Base paired structure of CpG.

nature (9), it is expected to interact exclusively with the nitrogen binding sites on the bases.

Methylmercury hydroxide, CH_3HgOH , is known to cause the denaturation of both animal and bacterial DNAs (106). The most probable binding sites are the imino NH groups of thymine or guanine. Organomercurials can cause chromosome abnormalities leading to mutagenic or teratogenic effects (106).

Through the use of ^{13}C NMR it is hoped to define the binding site of methylmercury on the dinucleoside CpG. It is also possible to determine if this binding is affecting the conformation of the dinucleoside by using vicinal carbon-phosphorus, and proton-proton coupling constants. The role of the sugar moiety will also be explored when the binding of methylmercury to 2'-deoxycytidylyl-(3'-5')-2'-deoxyguanosine (dCpG) is studied.

9.1 Two-Dimensional ^1H - ^1H Chemical Shift Correlation Spectroscopy

The literature contains a number of ^1H NMR studies on the dinucleosides CpG and dCpG (107,108,109). Extensive overlap occurs in the ^1H - ^1H coupled ribose region of the spectrum at 200 MHz. See Figure 39. For this reason a two dimensional ^1H - ^1H homonuclear chemical shift correlated NMR spectrum (2-D ^1H -COSY) was recorded for CpG.

In a 2-D ^1H -COSY spectrum, a typical one dimensional ^1H spectrum occurs along the diagonal axis. The presence of off-diagonal peaks, however indicates that a scalar J coupling exists between the two resonances. Thus a highly overlapped region of an NMR spectrum can be analyzed by running a single experiment rather than several selectively decoupled ones.

The 2-D ^1H COSY spectrum of CpG at 500 MHz is presented in Figures 40 and 41. This spectrum allows the identification of the ribose protons for later use in ^{13}C heteronuclear selective decoupling experiments.

FIGURE 40: A two dimensional homonuclear ^1H - ^1H chemical shift correlation NMR spectrum of the ribose region in cytidyl-(3'→5')-guanosine dinucleoside monophosphate.

FIGURE 41: Expansion of 4.0 to 4.6 ppm region of Figure 40.

9.2 ^{13}C Chemical Shift Assignments for CpG

As no literature results were available concerning the ^{13}C NMR spectrum of the dinucleoside monophosphate CpG, initial assignments had to be made.

The structure and numbering scheme for CpG is depicted in Figure 42. Since 13 of the 19 carbons of this molecule are protonated, selective ^1H decoupling in the ^{13}C spectrum can be used to identify directly bound carbon atoms. The ^1H NMR chemical shifts were verified via an ^1H - ^1H chemical shift correlation experiment at 500 MHz. ^{13}C assignments were also made by comparison with the original nucleoside and mononucleotide spectra (36,37,38). The influence of an attached phosphorus group is also considered. Carbons directly attached to the phosphate undergo a downfield shift of ca. 2.5 ppm relative to their resonance position in the nucleosides. The other ribose carbons are not greatly affected.

The presence of carbon-phosphorus couplings provide an important clue in the identification of the ribose carbons. The two-bond coupling constant, $^2J_{\text{C,P}}$ is independent of conformation and takes on values of 5.0 ± 1.0 Hz. The three bond vicinal carbon-phosphorus couplings are, however conformation dependent and will vary from 0 to 10.0 Hz.

The ^{13}C NMR spectrum of CpG is presented in Figure 43. In addition the ^{13}C chemical shift correlation between the mononucleotides 3'CMP and 5'GMP, and the dinucleoside monophosphate CpG is shown in Figure 44.

FIGURE 42: The structure, numbering scheme, and torsional angles of CpG.

FIGURE 44 : ^{13}C Chemical shift correlation between the mononucleotides, 3'CMP and 5'GMP, and the dinucleoside monophosphate, CpG. In D_2O , pH 6.8.

In the fully coupled, nOe enhanced ^{13}C spectrum of CpG several long range ^1H couplings are observed. The chemical shifts and coupling behaviour of the base resonances of CpG are given in Table 3.

Table 3: ^{13}C shift data and ^1H coupling information for the base carbons of CpG

Chemical shift ^a , ppm	Coupling behaviour $^1\text{H}-^{13}\text{C}$
161.50	doublet 9.4 Hz
158.37	
153.63	
151.52	unresolved multiplet
151.16	unresolved multiplet
142.29	doublet 190 Hz
136.98	deuterium broadening
115.82	
95.06	doublet 190 Hz

a. reference to external TMS

The resonance at 136.98 ppm can be assigned as Gua 8, due to the observed deuterium broadening in a decoupled ^{13}C spectrum. The CpG sample was lyophilized from D_2O several times prior to running the ^{13}C spectrum. The C8-H bond of the purine ring is reasonably labile and significant deuterium exchange will occur (137).

The protonated carbon assignments for CpG followed directly from selective decoupling experiments at the cytidine H5 resonance frequency.

The downfield resonance at 161.50 ppm exhibits a long-range coupling of 9.4 Hz which is also removed upon irradiation of cytidine H5. Thus this resonance is assigned as Cyt 4.

The two resonances located at ca. 151 ppm show several long range interactions. The downfield peak however can be effectively decoupled by applying the decoupler field at the frequency of cytidine H1'. The upfield peak is sharpened when the decoupler frequency is equal to the resonance frequency of guanosine H1'. Thus the peak at 151.16 ppm is assigned to Gua 4 and the peak at 151.52 ppm corresponds to Cyt 2.

The resonance at 115.82 ppm is separated by more than 20 ppm from any other in CpG. It can therefore, by analogy with the results for 5'GMP, be confidently assigned to the 5 position of the guanosine residue.

The remaining two base resonances of CpG appear at 153.63 and 158.37 ppm respectively and they show no apparent splittings in the ^1H coupled ^{13}C spectrum in D_2O . This is consistent with expectation for the 2 and 6 positions of guanosine, the only two base carbons unassigned. Specific designation of these two resonances was made using 5'GMP as a model and the agreement is excellent.

The ribose carbons were assigned using carbon-phosphorus coupling data. See Table 4.

Table 4: ^{13}C - ^{31}P Coupling Constants CpG
(Hz $\pm$.1)

Compound	Pathway	2J	3J
CpG	G4'-5'-O-P		8.6
CpG	G5'-O-P	2.8	
CpG	C4'-3'-O-P		5.7
CpG	C2'-3'-O-P		3.1
	C3'-O-P	4.6	
5'GMP	G4'-5'-O-P		9.2
3'GMP	G5'-O-P	5.5	
3'CMP	G4'-3'-O-P		6.0
3'CMP	C2'-3'-O-P		2.4
3'CMP	C3'-O-P	3.0	

The results for the model compounds 3'CMP and 5'GMP are in good agreement with published findings (37,110).

To identify the two anomeric carbons the effects of selective irradiation at the two H1' resonance frequencies were considered.

In Table 5 are presented the complete ^{13}C chemical shift data along with those of the related mononucleotides. For 3'CMP there has been a recent report of the ^{13}C - ^{31}P couplings, but no ^{13}C shift data were included (110). Our results for 5'GMP are in good agreement with the literature (37), with the appropriate reversal of the original 2' and 3' carbon resonance assignments (38).

Table 5: ^{13}C Chemical Shifts for CpG and the Mononucleotide Models
(ppm from TMS : .01)

Site	CpG .10M	3' CMP .50M	5' GMP .50M
Cyt 2	151.52	156.93	
Cyt 4	161.50	165.60	
Cyt 5	95.06	96.18	
Cyt 6	142.29	141.65	
Cyt 1'	90.28	89.81	
Cyt 2'	73.06	73.53	
Cyt 3'	72.01	72.06	
Cyt 4'	82.68	83.17	
Cyt 5'	59.55	60.56	
Gua 2	153.63		153.49
Gua 4	151.16		150.95
Gua 5	115.82		115.46
Gua 6	158.37		158.29
Gua 8	136.98		137.05
Gua 1'	87.56		86.80
Gua 2'	73.49		74.02
Gua 3'	69.47		70.22
Gua 4'	82.71		83.88
Gua 5'	64.46		63.35

General overlap is observed between the resonances of 5'GMP and the guanyl resonances of CpG. The cytidylyl resonances however are shifted upfield from their corresponding frequencies in 3'CMP. This is due to the increase ring current effects on the cytidylyl peaks brought about by the adjacent purine ring.

Ring currents contribute to $\sigma_p(\text{nonloc})$ in special circumstances. If an aromatic bond system is orientated perpendicular to the applied magnetic field, an inter-atomic circulation of the π electrons will result. This will set up a current around the outside of the ring system, which gives rise to a magnetic field opposing the applied field within the ring and adding to the external field outside the ring.

The purine bases have a greater aromatic character than the pyrimidine bases, and thus have a larger ring current. The base-stacking which occurs in a self-complementary dinucleoside monophosphate, such as CpG, causes the pyrimidine base carbons to be over the purine ring. See Figure 38. These carbon resonances accordingly show an upfield shift from their frequencies in the nucleotides.

2.3 Conformational Properties CnG

Base sequence dependent changes in nucleic acid conformation may be important in the determination of binding sites for a large number of drugs and metal ion species. The conformation of a base stacked, self-complementary ribodinucleoside monophosphate such as CpG will provide a good model of RNA structures (111,112).

Despite the great deal of freedom theoretically possessed by the singly bonded sugar-phosphate backbone of a polynucleotide, as shown in Figure 42 it will adopt one of only a few preferred conformations when confined to a helix structure in solution.

Recent NMR studies using ^1H - ^1H vicinal coupling constants have elucidated these conformational details for the monomers, dimers and several oligonucleotides (21). ^1H NMR has provided the largest body of conformational data through the use of Karplus relationships, chemical shift temperature dependence, spin lattice relaxation times (107,108), two dimensional NMR (113) and lanthanide shift reagents (114).

Recently ^{13}C NMR has begun to be used to study dinucleoside monophosphates, though to date only on derivatives involving adenosine and uracil (38,115,116,117). Conformational analysis using the three-bond phosphorous carbon couplings has been applied to the mononucleotides (110,118,119) and the adenosine and uracil dimers (118,120):

Using the established Karplus relationships (21) for the carbon-phosphorous couplings, two dihedral angles in the dinucleoside

monophosphates can be determined from three observed $^3J_{C,P}$ values.

9.31 ϵ_1 ; P-O3'-C3'-C4'

ϵ_1 , the torsional angle between C3' and O3' on the 3' phosphate cytidyl residue, can be examined via couplings between C2' and C4' and the phosphorus atom.

Theoretically three staggered conformations are possible about the O3'-C3' bond, as shown below. However, 1H NMR results indicate that this 3 model system is not appropriate to describe the observed couplings (115). To solve the problem the trans, $\epsilon_1 = 60^\circ$ conformer is discarded, due to two gauche interactions it does not have an equal weighting when compared to the other two species.

Recently a new set of Karplus parameters was derived which provides better agreement between dinucleoside conformations obtained

by NMR and those observed by x-ray diffraction (121).

$$^3J_{C,P} = 6.9 \cos^2 \theta - 3.4 \cos \theta + 0.7 \quad 29$$

This leads to a gauche coupling constant of 0.73 Hz, and a coupling of 11.0 Hz between trans carbon and phosphorus atoms.

If all three populations were equally weighted, then three simultaneous equations would describe the θ_1 torsional angle.

$$P_1 + P_2 + P_3 = 1 \quad 30$$

$$^3J_{P,C2'}(\text{obs}) = 0.73 P_1 + 0.73 P_2 + 11.0 P_3 \quad 31$$

$$^3J_{P,C4'}(\text{obs}) = 0.73 P_1 + 11.0 P_2 + 0.73 P_3 \quad 32$$

However because P_1 is not a likely conformation, only the dominant conformation can be determined, either P_2 or P_3 . If $^3J_{P,C4'} = ^3J_{P,C2'}$, then $\theta_1 = 300^\circ$ is dominant, however if $^3J_{P,C2'} = ^3J_{P,C4'}$, then the opposite is true and $\theta_1 = 180^\circ$ is the preferred conformer.

The results for the cytidyl residue in the mononucleotide and the dinucleoside monophosphate are given in Table 6.

Table 6: ^{13}C - ^{31}P Coupling Constants (Hz)

	$^3J_{P,C2'}$	$^3J_{P,C4'}$	Ref.
3' CMP	5.7 ± .6	3.1 ± .6	this work
3' CMP	6.0 ± .2	3.0 ± .2	110
3' CMP	3.9 ± .3	5.2 ± .3	118
CpG	2.4 ± .2	6.0 ± .2	this work

Thus one finds the ϵ_{tg} population dominates in the ribose mononucleotides, but in the dimer the preferred conformation is ϵ_{gt} . This has been noticed previously in other cytidine polynucleosides, unlike those involving adenosine or uracil, conformation is not conserved when going from the monomer to the polymer (122).

A relationship exists between this ϵ_1 torsional angle and the ribose sugar pucker (118). If ϵ_1 is found in the ϵ_{gt} range then the sugar is 3' endo, and if ϵ_1 is in the ϵ_{tg} region the sugar is 2' endo. See Figure 46.

9.32 Ribose Sugar Pucker

In the ribose sugar ring there are twenty different pseudorotation conformations, one every 18° (21). The twist conformations (T) occur at even multiples of 18° , whereas envelope (E) species occur at odd multiples. To simplify the discussion usually only the two minimum energy conformers are discussed, 2E and 3E . A standard conformation, $P = 0^\circ$ is defined when the torsional angle $C1'-C2'-C3'-C4'$ has its maximum positive value (21). See Figure 45.

From this definition 3E , or the 3' endo conformation occurs at 18° . It is an N type conformation with the 3' carbon above the plane of the other carbons, i.e. on the side of the base. The 2E , or 2' endo occurs at 162° , and belongs to the S class of sugar conformers where the 2' carbon is above the plane of the remaining sugar carbons. These conformations are shown in Figure 45.

A large number of methods exist for determining the sugar conformer populations from vicinal proton-proton couplings (107,108,21).

The simplest and most approximate involves a Karplus relationship using only $^3J_{H1',H2'}$. This is the easiest method to apply because of extensive overlap in the 1H NMR spectrum of a dinucleoside monophosphate, even at 500 MHz. The solution of two simultaneous equations will give the relative populations of each conformation,

$$^3E + ^2E = 1 \quad \dots\dots 33$$

$$10.1^2E + 10.3^3E = ^3J_{H1',H2'}(\text{obs}) \quad \dots\dots 34$$

Data and results are given in Table 7, included are reference values found using a more complete analysis of the ribose coupling constants (107).

Table 7: 3E Sugar Conformer Populations for the Ribose Derivatives

	$^3J_{H1',H2'} \text{ (Hz)}$	$^3E(\%)$	$^3E(\%)^{107}$
5'GMP	5.5 $\pm$.5	46 $\pm$ 5	37
3'CMP	3.8 $\pm$.5	63 $\pm$ 5	60
CpG			
Cp	2.8 $\pm$.5	73 $\pm$ 5	66
pG	4.2 $\pm$.5	59 $\pm$ 5	52

For ribose mononucleotides, such as 3'CMP and 5'GMP a firm relationship between ϵ_1 and the ribose pucker does not exist. Unlike the deoxyribose mononucleotides, ϵ_1 is dependent upon the nature of the base in the ribose series not the sugar pucker.

The relationship between P , the phase angle of pseudorotation and the twist (T) and envelope (E) sugar conformations.

Nomenclature		P°	
C(1')-endo-C(2')-exo	$1T$	324	N type
C(2')-exo	$1E$	342	
C(3')-endo-C(2')-exo	$2T$	0	
C(3')-endo	$2E$	18	
C(3')-endo-C(4')-exo	$3T$	36	
C(4')-exo	$3E$	54	
O(4')-endo-C(4')-exo	$4T$	72	
O(4')-endo	$4E$	90	
O(4')-endo-C(1')-exo	$5T$	108	
C(1')-exo	$5E$	126	
C(1')-exo-C(2')-endo	$6T$	144	S type
C(2')-endo	$6E$	162	
C(2')-endo-C(3')-exo	$7T$	180	
C(3')-exo	$7E$	198	
C(4')-endo-C(3')-exo	$8T$	216	
C(4')-endo	$8E$	234	
C(4')-endo-O(4')-exo	$9T$	252	
O(4')-exo	$9E$	270	
C(1')-endo-O(4')-exo	$10T$	288	
C(1')-endo	$10E$	306	

FIGURE 45: Pseudorotation of the ribose sugar ring. All envelope and twist conformations.

FIGURE 46: Major ribose conformations.

9.33 β_2 ; P-05'-C5'-C4'

The second backbone torsional angle which can be studied using carbon-phosphorous vicinal couplings is β_2 . β_2 is the torsional angle between P-05'-C5'-C4' and describes the orientation of the two monomer residues relative to each other in the dimer.

Again there are three possible conformations allowed as shown below, looking down the 05-C5' bond.

$$\beta_2 = 60^\circ$$

$$P_1 \beta_{g+}$$

$$\beta_2 = 180^\circ$$

$$P_2 \beta_1$$

$$\beta_2 = 300^\circ$$

$$P_3 \beta_{g-}$$

The same Karplus relationship used for ϵ_1 is applied in the case of β_2 . However with only one three-bond carbon-phosphorus coupling, $^3J_{C4',P}$ the three simultaneous equations cannot be solved. The system is reduced to two conformations by combining the two gauche species.

$$P_1 + P_3 = P_g \quad 35$$

$$P_2 = P_t \quad 36$$

Only two equations must now be solved using the one coupling constant.

$$P_g + P_t = 1 \quad 37$$

$$^3J_{C4',P}(\text{obs}) = .73 P_g + 11.0 P_t \quad 38$$

Data and results for the determination of β_2 are presented in

Table 8.

Table 8: P_t populations of ribose mononucleotides, and CpG

	$^3J_{C4',P}(\text{Hz})$	$P_t(^{\circ})$
5'GMP	$8.6 \pm .6$	77
CpG	$9.2 \pm .2$	82

The torsional angle β_2 is not greatly dependent on the monomer to dimer shift, other than to observe a further preference for the trans conformation.

9.34 Conclusion

The ^{13}C NMR spectrum of CpG yields additional conformational information from the observed vicinal carbon-phosphorus coupling constants. The two backbone angles α_1 and α_2 determined from this data agree with results from ^1H - ^{31}P couplings.

The ribose pucker is 3'endo for 3'CMP and the dinucleoside CpG, but is 2'endo for 5'GMP. This also agrees with literature values. The sugar pucker is independent of α_1 but dependent on the nature of the base for the ribose monomers and polymers.

It is possible now to determine the effects of metal binding on the conformation of the mononucleotides and dinucleoside monophosphate using ^{13}C NMR.

9.4 ^1H -NMR Spectrum Deoxyribodinucleoside monophosphate

The ^1H NMR spectrum of 2'-deoxyribosecytidyl-(3'→5')-2'-deoxy-ribo-guanosine (dCpG) is better dispersed than that of the corresponding ribose dinucleoside monophosphate. The sugar protons are readily assigned at 200 MHz, as shown in Figure 47.

These proton resonances are used for selective decoupling in the ^{13}C spectrum of dCpG. The ^1H - ^1H couplings in the deoxyribose sugar will also provide conformational information.

9.5 ^{13}C Chemical Shift Assignments dCpG

The ^{13}C NMR spectrum of dCpG does not appear in the literature, so an initial assignment of the resonances is again necessary. The structure and numbering scheme of dCpG is shown in Figure 48.

Similar to the strategy used for assigning the ^{13}C resonances of rCpG, selective ^1H decoupling of the ^{13}C spectrum was employed to identify the directly bonded carbons of the bases.

The ribose carbons were assigned based upon their observed carbon-phosphorous couplings and comparison with the monomer model compounds 5'dGMP and 3'dCMP. The anomeric carbons were assigned using selective ^1H irradiation at guanosine H1.

The remaining 6 quaternary carbons were assigned using several methods. The resonance at 115.63 ppm is well removed from any other quaternary peak and is therefore assigned as Gua 5, by analogy to the results for 5'dGMP.

In the totally coupled ^{13}C spectrum of dCpG several $^2\text{J}_{\text{C,H}}$ and $^3\text{J}_{\text{C,H}}$ interactions are observed. These interactions can be removed using selective ^1H decoupling, and lead to the assignment of 3 resonances.

Specifically Cyt 4 at 165.28 ppm ($^2\text{J}_{\text{C,H5}} = 8.3 \text{ Hz}$), Cyt 2 at 156.29 ppm ($^3\text{J}_{\text{C,H1}}$, $^3\text{J}_{\text{C,H6}} = 6.4 \text{ Hz}$) and Gua 4 at 151.28 ppm ($^3\text{J}_{\text{C,H1}} = 5.1 \text{ Hz}$) were assigned in this manner.

The remaining two resonances at 158.54 and 153.63 ppm show no long range couplings in D_2O solution. This is consistent with expectations for Gua 2 and Gua 6, the only two remaining unassigned carbons. Specific designation of these two resonances was made using 5'dGMP as a model and the agreement is quite good.

FIGURE 48: The structure, numbering scheme, and torsional angles of dCpG.

The ^{13}C shift data for dCpG along with those for the related mononucleotide models 3'dCMP and 5'dGMP are given in Table 9. For 3'dCMP there has been a recent report of the ^{13}C - ^{31}P couplings, but no ^{13}C shift data were included (110). Our results for 5'dGMP are in agreement with the literature (37,38).

Table 9: ^{13}C Shift Data (C ppm from TMS, ± 0.01) for dCpG, and Mononucleotide Models.

Site	dCpG	3'dCMP	5'dGMP
Cyt 2	156.29	157.27	
Cyt 4	165.28	166.09	
Cyt 5	96.12	96.29	
Cyt 6	140.77	141.70	
Cyt 1'	85.29	86.08	
Cyt 2'	37.55	38.71	
Cyt 3'	75.37	73.22	
Cyt 4'	85.51	86.17	
Cyt 5'	60.91	61.42	
Gua 2	153.63		153.67
Gua 4	151.28		150.72
Gua 5	115.63		115.65
Gua 6	158.54		158.48
Gua 8	137.08		137.08
Gua 1'	82.76		83.05
Gua 2'	38.06		38.83
Gua 3'	70.43		71.28
Gua 4'	85.12		86.08
Gua 5'	64.76		63.82

Similar to CpG, the cytidyl base resonances are shifted upfield when compared to the mononucleotide, due to the increased ring current effect of the adjacent guanyl residue. The guanyl peaks generally overlap those of the mononucleotide model 5'dGMP. See Figure 49.

The geminal and vicinal ^{13}C - ^{31}P couplings observed for specific sites in dCpG and the two model mononucleotides are presented in Table 10. Our results for 3'dCMP and 5'dGMP are in good agreement with published findings [37,110].

Table 10: ^{13}C - ^{31}P Coupling Constants Hz $\pm$ 0.1

Compound	Pathway	2	3
dCpG	(C4')-3'-O-P		7.4
dCpG	(C2')-3'-O-P		1.9
dCpG	(C3')-O-P	5.9	
dCpG	(G4')-5'-O-P		8.9
dCpG	(G5')-O-P	3.6	
3'dCMP	(C4')-3'-O-P		not resolved
3'dCMP	(C2')-3'-O-P		2.9
3'dCMP	(C3')-O-P	4.3	
5'dGMP	(G4')-5'-O-P		7.8
5'dGMP	(G5')-O-P	3.8

FIGURE 49: ^{13}C Chemical shift correlation between the deoxymononucleotides, 3'dCMP and 5'dGMP, and the deoxydinucleoside monophosphate, dCpG. In D_2O , pH 6.8.

9.6 Conformation of dCpG

As was the case with CpG, the ^{13}C NMR spectrum of dCpG, once assigned, will yield conformational information about the duplex.

Similar to CpG, dCpG will form a base-stacked, hydrogen bonded duplex model for DNA (123). However the lack of a 2'-OH function on the ribose has far reaching conformational consequences.

9.61 Sugar Pucker

The primary difference between ribose and deoxyribose polynucleotides occurs in the sugar torsional angles. As described previously there are two major populations of sugar conformers, ^2E and ^3E , as shown in Figure 46. These populations are estimated from $^3\text{J}_{\text{H}1',\text{H}2'}$ in the ^1H spectrum of the dinucleoside in question. Results are given in Table 11 and compared to results obtained using $^3\text{J}_{\text{H},\text{P}}$ and $^3\text{J}_{\text{H},\text{H}}$ (109).

Table 11: ^3E Sugar Populations for the Deoxyribose Derivatives

	$^3\text{J}_{\text{H}1',\text{H}2'}$ $\pm 2 \text{ Hz}$	$^3\text{E}(\%)$ ± 5	$^3\text{E}(\%)^{109}$
5'dGMP	7.2	29	
3'dCMP	6.7	34	
dCpG			
Cp	7.2	29	33
pG	7.2	29	37

In contrast to the ribodinucleoside, the preferred sugar conformation of dCpG is 2E .

This change in sugar pucker also affects other torsional angles in the molecule. The glycosidic bond, χ for example is increased, although still within the anti range (123).

In deoxyribose monomers and polymers the ${}^2E(2'endo)$ conformation is strongly preferred, independent of base nature and sequence and intermolecular forces such as base-stacking. This is in sharp contrast to the ribonucleotides where the sugar pucker is determined by a number of intra- and inter-molecular forces.

For both ribo- and deoxyribo-nucleotides the τ_1 torsional angle is dependent upon the sugar pucker (120). However, for deoxyribonucleotides this dependence is base sequence independent.

9.62 τ_1 ; P-O3'-C3'-C4'

As previously described the backbone torsional angle τ_1 can be investigated from the carbon-phosphorus couplings in a ${}^{13}C$ spectrum of the dinucleoside monophosphate. The data are presented in Table 12.

Table 12: ${}^{13}C$ - ${}^{31}P$ Coupling Constants (Hz).

	${}^3J_{P,C2'}$	${}^3J_{P,C4'}$	ref.
3'dCMP	2.9 $\pm$.6	-	this work
3'dCMP	3.2 $\pm$.2	6.1 $\pm$.2	110
dCpG _n	1.9 $\pm$.2	7.4 $\pm$.2	this work

ϵ_1 is found in the ϵ_{tg} range for both the monomer and dimer.

This is in contrast to the ribose series, where ϵ_{gt} is preferred in the dinucleotide. The change observed for the deoxyribose dimer is in response to the sugar conformation and an inter-dependence of these two conformation angles.

ϵ_{gt} is generally associated with an 3E sugar pucker, whereas ϵ_{tg} occurs when an 2E sugar pucker dominates.

9.63 β_2 ; P-O5'-C5'-C4'

The torsional angle β_2 , between O5' and C5' in the 5'guanyl residue of dCpG can also be determined from the carbon-phosphorus coupling data.

In the monomer, 5'dGMP ${}^3J_{C4',P}$ is $7.8 \pm .6$ Hz, and in the dimer dCpG a value of $8.3 \pm .2$ Hz is obtained. This indicates a dominant trans population for both species; 69% in the case of the monomer and 76% for the dimer.

As expected, the β_2 angle being farther removed from the sugar and base region, no dependence on sugar population or type is observed. Polymer formation serves only to remove some freedom from the rotational motion in both deoxyribose and ribose dinucleoside monophosphates.

9.64 Conclusion

From ${}^{13}C$ - ${}^{31}P$ couplings it has been shown that in dCpG, unlike CpG, the angle ϵ_1 is no longer dependent upon base type. The ϵ_{tg} conformer dominates in both the monomer and dimer species. This is due

to the dominant 2E sugar pucker associated with 2'-deoxyribose sugar derivatives.

For dCpG, the β_2 torsional angle remains in the trans domain, similar to CpG.

CHAPTER 10: METAL ION BINDING TO DINUCLEOSIDE MONOPHOSPHATES

With complete ^{13}C chemical shift assignments of CpG and dCpG in hand, a study on the metal ion binding properties of these species was subsequently undertaken.

In the present study the soft methylmercury (II) cation was used to probe the binding sites on the two dinucleoside monophosphates. The unifunctional nature of $\text{CH}_3\text{Hg(II)}$ makes it an ideal probe and in addition, its pH dependence makes it possible to select for and synthesize specific model compounds (75,124,125).

The sites of metal ion binding on the mononucleotide model compounds 3'CMP, 5'GMP, 3'dCMP and 5'dGMP were also observed using ^{13}C and ^{15}N NMR titration shifts. Previously, Raman difference spectroscopy had been used to determine the binding sites of $\text{CH}_3\text{Hg(II)}$ on the ribose nucleotides (126,127,128).

Specific N1 metal bound complexes of the guanosine mononucleotides were synthesized, in a manner similar to the preparation of the nucleoside complexes (75). These complexes were studied using ^{13}C and ^{15}N NMR.

Finally the binding reactions between $\text{CH}_3\text{Hg(II)}$ and the dinucleoside monophosphates CpG and dCpG, were studied by ^{13}C NMR. Any conformational changes occurring upon binding were determined from the vicinal carbon-phosphorus coupling constants.

Methylmercury is known to complex preferentially to the N3 site of uridine or thymidine in polynucleotides and cause mutagenic effects (129), a secondary site is the N1 position of guanosine. At very low concentrations, $\text{CH}_3\text{Hg(II)}$ binding to DNA need not cause denaturation,

or bring about conformational changes (130).

The binding of CH_3HgOH to the monomer or dimer at N1 occurs by proton exchange and is thus very pH dependent (126,127,128). The species distribution of $\text{CH}_3\text{Hg(II)}$ itself is a function of pH in aqueous solution and is given below.

At pH 8 and higher mercuriation at N7 of 5'GMP is blocked by the hydroxide coordination of $\text{CH}_3\text{Hg(II)}$. At lower pH, the mercuriation of N1 is blocked by ligand protonation. At pH 8.5, 92% N1 complex has been found to form in solution, as determined by Raman difference spectroscopy (127).

For 3'CMP, complex formation at N3 is maximized at pH 4 (127). At very high pH, greater than 10, a small amount of binding will occur at the 4- NH_2 group through proton replacement (128).

Pullman has calculated the primary metal ion binding sites for the hard metal ion Na^+ and the cytosine base (72). As shown below, the preferred site is a chelated position between the carbonyl oxygen and N3. Of secondary importance is the single oxygen site, followed by N3.

Significant changes in site preference are expected for the softer $\text{CH}_3\text{Hg(II)}$ species, which does not ordinarily interact with oxygen ligands.

As illustrated previously for guanine, the chelated site between O6 and N7 is preferred by Na^+ , followed by O6 and N7 alone (20). For the nucleoside, N1 is the thermodynamically favoured binding site and N7 the kinetic site (39).

10.1 Mononucleotide Model Compounds

10.11 ^{15}N NMR Results

The ^{15}N NMR results for the binding of CH_3Hg^+ (I) to the ribose mononucleotides are presented in Figure 5. The results for the synthetic diphosphate-Ni-ethylurea (1:1) complex (Ni-CH₃Hg) (II) (5) are also included.

For (I) (MF), a 410 ppm upfield shift of N3 is observed, while the N1 and N9 resonances are shifted slightly downfield, 1.5 and 1.2 ppm respectively.

These results indicate a binding site for CH_3Hg^+ (I) which preferentially involves the N1 site of (I) (MF), as shown below. The nearly equal downfield shifts of the N9 and N1 resonances are consistent with these sites being conjugated with the bound nitrogen site.

R = ribose - 3' - monophosphate

FIGURE 50: ^{15}N Results. Effect of $\text{CH}_3\text{Hg(II)}$ on the chemical shifts of the ribose mononucleotides, 5' GMP and 3' CMP. Synthetic $\text{NI-CH}_3\text{Hg(II)}$ complex of 5' GMP. Positive values are downfield.

Similar ^{15}N results were obtained when $\text{Hg}(\text{II})$ was bound at the N3 position of cytosine in DMSO (53). The same upfield shift of a metallated sp^2 nitrogen resonance is observed for the purine nucleosides.

These upfield shifts are due to an increase in the P_u term and a decrease in the r and AEE terms of $\sigma_p(\text{loc})$.

$$\sigma_p(\text{loc}) = \frac{-\mu \cdot e^2 h^2}{6 \pi m^2 \text{AEE}} (\langle r^{-3} \rangle_p P_u) \quad 23$$

The interaction between $\text{CH}_3\text{Hg}(\text{II})$ and 5'GMP was studied in two ways: one a binding titration was carried out, and two, a synthetic N1 complexed species was prepared and characterized by NMR.

The binding titration was limited to the addition of 0.16 equivalents of metal due to solubility constraints at pH 8. A downfield shift of 21.3 ppm is observed for N1, in addition a small 0.7 ppm downfield shift occurs at N3. Small upfield shifts, less than 2 ppm are observed for the N7, N9 and NH_2 resonances.

The ^{15}N spectrum of the N1- $\text{CH}_3\text{Hg}(\text{II})$ -5'GMP complex exhibits a 49.9 ppm downfield shift for N1 compared to the parent mononucleotide 5'GMP. Smaller downfield shifts are also observed for the other nitrogen resonances.

The ^{15}N results certainly indicate that N1 is the most affected nitrogen of 5'GMP when $\text{CH}_3\text{Hg}(\text{II})$ is added at pH 8. The observed downfield shift is however opposite to shifts observed for all other coordinated nitrogens.

There are several factors which contribute to this observation, the most important of which is that N1 must deprotonate before it binds $\text{CH}_3\text{Hg}(\text{II})$. It has been shown that deprotonation of 5'GMP does begin to occur at pH 8. Full deprotonation at N1 will lead to a 60 ppm downfield

shift of the ^{15}N resonance 46.

The presence of a methyl group on the metal ion can also lead to a downfield shift of N1. The downfield effect of a methyl group is well documented for ^{13}C 33. The magnitude of deshielding is generally larger for a nitrogen nucleus than the corresponding carbon nucleus 44.

For example consider the downfield nitrogen shift observed in N-methylacetamide when a methyl group is introduced 33.

δ_{N} 105.8 ppm

δ_{N} 124.8 ppm

The observed ca. 50 ppm downfield shift of N1 probably arises from these two effects combined with the normal upfield shift observed when a metal ion binds at an sp^2 nitrogen center.

The observed shifts of N9 and N7 in both $\text{CH}_3\text{Hg(II)}$ species of 5'GMP confirm that N7 is not a primary binding site at pH 8.

The pH dependence of $\text{CH}_3\text{Hg(II)}$ binding to 5'GMP cannot be studied by natural abundance ^{15}N NMR due to the low solubilities of CH_3HgOH and 5'GMP in aqueous solutions below pH 8.

The ^{15}N results do indicate however, that at pH 6 the primary site of $\text{CH}_3\text{Hg(II)}$ complexation on 3'CMP is N3. While at pH 8 the $\text{CH}_3\text{Hg(II)}$ species will bind at N1 in 5'GMP.

¹³C NMR Results

The ¹³C NMR results for the binding of CH₃Hg(II) to the monomers 3'GMP, 5'GMP, 3'dGMP and 5'dGMP are presented in Figures 51 and 52. Results for the synthetic complexes N1-CH₃Hg(II)-5'GMP and N1-CH₃Hg(II)-5'dGMP are presented in Figure 53.

It is apparent from these results that the nature of the sugar moiety in the mononucleotide does not affect metal ion complexation. Accordingly, only the results for the ribose mononucleotide will be discussed in detail.

The ¹³C NMR characteristics of an N3 binding site in cytosine complexed with Hg(II) have been previously determined (39). A downfield shift of C5 and upfield shifts of C4 and C2 are observed when N3 complexation occurs. These shifts are also observed when CH₃Hg(II) is added to 3'GMP at pH 7, as shown in Figure 51.

The predicted N3 complexation site is in agreement with previous ¹⁵N results. The upfield shift of C2 indicates that chelation of N3 bound CH₃Hg(II) to the carbonyl oxygen is not an important contributing mechanism in the binding process, despite the predicted stability of such a complex involving a harder metal ion (72).

For the interaction of CH₃Hg(II) with 5'GMP, the ¹³C titration shifts are identical to the shifts observed for the synthesized N1 complexes. See Figures 51, 52 and 53.

Downfield shifts of 2 to 3 ppm are observed for the C2 and C6 resonances, while a smaller downfield shift of C5 occurs. The C8 resonance undergoes very small shifts either upfield or downfield,

5' GMP pH 8

3' CMP pH 7

FIGURE 51: ^{13}C Results. Effect of $\text{CH}_3\text{Hg(II)}$ on the chemical shifts of the ribose mononucleotides, 5' GMP and 3' CMP.

Positive values are downfield

FIGURE 52: ^{13}C Results Effect of $\text{CH}_3\text{Hg(II)}$ on the chemical shifts of the deoxymononucleotides, 5'dGMP and 3'dCMP. Positive values are downfield.

FIGURE 53: ^{13}C Results. Synthetic NI complexes of 5'GMP and 5'dGMP with $\text{CH}_3\text{Hg}(\text{I.I})$. Positive values are downfield.

depending on the ligand involved, and C4 is shifted slightly upfield in all cases.

The large downfield shifts of C2 and C6 were previously observed when a metal ion was bound to the N1 position of 7-methylguanosine. This binding mechanism also involves the loss of the N1-proton followed by metal ion substitution. Thus both ^{15}N and ^{13}C NMR support a concomitant deprotonation mechanism for $\text{CH}_3\text{Hg(II)}$ binding to 5'GMP at pH 8.

The ^{13}C chemical shifts of the C5 and C8 resonances indicate that no N7 complex is being formed under these conditions.

10.13 Carbon Phosphorus Couplings - Conformational Results

The ϕ and ψ torsional angles in the mononucleotides were monitored via the magnitude of the vicinal carbon-phosphorus couplings. The effect of $\text{CH}_3\text{Hg(II)}$ complexation on these angles was determined from the titration data.

For 5'GMP and 5'dGMP no changes in $^3J_{\text{C4}',\text{P}}$ were observed. A coupling of $8.8 \pm .6$ Hz is indicative of a dominant trans population of the B bond angle.

Similarly for 3'CMP, $^3J_{\text{C4}',\text{P}}$ and $^3J_{\text{C2}',\text{P}}$ remain constant at $4.6 \pm .6$ Hz and $5.8 \pm .6$ Hz respectively. For the 3'dCMP titration with $\text{CH}_3\text{Hg(II)}$, $^3J_{\text{C2}',\text{P}}$ remains constant at $3.0 \pm .6$ Hz and $^3J_{\text{C4}',\text{P}}$ cannot be resolved due to peak overlap.

For the ribose 3'CMP, ϵ remains in the ϵ_{gt} range when $\text{CH}_3\text{Hg(II)}$ is added. It is probable that the deoxy derivative 3'dCMP also remains in its preferred conformation range ϵ_{tg} .

10.2 Dinucleoside Monophosphates

10.21 ^{13}C Chemical Shifts - Metal Ion Binding Sites

The ^{13}C NMR results obtained for the addition of $\text{CH}_3\text{Hg(II)}$ to CpG and dCpG are presented in Figures 54 and 55. Once again the nature of the ribose sugar does not influence the binding of $\text{CH}_3\text{Hg(II)}$ to the nucleoside bases, and only the ribose CpG results will be discussed in detail.

Beginning with the 3'-cytidyl residue, the chemical shifts observed in response to $\text{CH}_3\text{Hg(II)}$ addition differ from what is observed for 3'-CMP under similar conditions.

Downfield shifts of 2 to 4 ppm are observed for the C2 and C4 resonances, smaller downfield shifts, less than 1.0 ppm occur for C5 and C1'. An upfield shift of ca. 1.0 ppm is observed at C6.

These shifts do not indicate the presence of N3 complexed metal ion on the 3'-cytidyl residue of CpG. The relatively large downfield shifts of C2 and C4 may be due to decreased base-stacking with the guanine base of CpG (132), as $\text{CH}_3\text{Hg(II)}$ is a known denaturation agent of DNA (133).

Base binding by a metal ion disrupts the two stabilizing forces of a double helix; hydrogen bonding and base-stacking. The effect is similar to the temperature dependent changes in conformation, shown below, which remove base-stacking interactions (1,123).

FIGURE 54: ^{13}C Results. Effect of $\text{CH}_3\text{Hg(II)}$ on the chemical shifts of cytidyl-(3'→5')-guanosine. pH6. Positive values are downfield.

FIGURE 55: ^{13}C Results. Effect of $\text{CH}_3\text{Hg(II)}$ on the chemical shifts of 2'-deoxycytidyl - (3'→5') - 2'-deoxyguanosine. pH 6. Positive values are downfield.

Heat denaturation of CpG

The ca. 3 ppm downfield shifts observed for C4 and C2 in the 3' cytidylyl residue are equal and opposite to the changes in shift observed going from 3'CMP to CpG, as shown in Table 5.

Thus no $\text{CH}_3\text{Hg(II)}$ binding site is found on the 3' cytidylyl residue of CpG. However, binding at some other location is affecting the base-stacking properties of the dinucleoside monophosphate.

The ^{13}C chemical shifts of the 5'-guanyl residue of CpG in the presence of $\text{CH}_3\text{Hg(II)}$ are shown in Figure 54. At pH 8, ca. 1 ppm downfield shifts of C5, C6 and C6' are observed. 5 ppm upfield shifts are recorded for the C4 and C4' resonances while C1 remains essentially unchanged. As discussed previously, these results are indicative of both N1 and N7 binding taking place.

N7 binding in the 5'-guanyl residue is observed at pH 6, in contrast to the exclusive N1 complex formed between the mononucleotide 5'-GMP and $\text{CH}_3\text{Hg(II)}$ at pH 8. The N7 complex occurs because in the pH dependence profile of $\text{CH}_3\text{Hg(II)}$ species, the charged $[\text{CH}_3\text{Hg}(\text{OH})]^{2+}$ occurs in equilibrium with CH_3HgOH at pH 6. The cationic species is able to coordinate at N7, while the methylmercuric (II) hydroxide binds at N1 in a two step process involving deprotonation.

The binding of $\text{CH}_3\text{Hg(II)}$ to the 5'-guanyl residue of CpG will interrupt the base-stacking interactions of the dinucleoside monophosphate. This will indirectly affect the ^{13}C chemical shifts of the 3'-cytidyl residue, as observed.

Thus, we conclude, that in the absence of a uridine or thymidine residue, guanosine becomes the preferred site of $\text{CH}_3\text{Hg(II)}$ interaction in polynucleosides at pH 6.

10.22 Carbon-Phosphorus Couplings - Conformational Changes

For CpG complexed with $\text{CH}_3\text{Hg(II)}$ no change is observed in the $^3\text{J}_{\text{G4',P}}$ coupling in the 5'-guanyl residue. It remains constant at 9.2 ± 0.2 Hz.

In the deoxyribose derivative, the coupling of the α proton of the deoxyribose is constant at 11.2 Hz. In the β conformational angle is not affected by the addition of $\text{CH}_3\text{Hg(II)}$. It remains in the preferred tg conformational range.

The first ribonucleoside coupling of the α deoxyribose residue of dCpG is in the tg conformational range, while those of $\text{dCpG} \cdot 0.35\text{eqCH}_3\text{Hg(II)}$ are in the gt conformational range (Table I).

Table I. Effect of $\text{CH}_3\text{Hg(II)}$ on the coupling of the α deoxyribose residue of dCpG and dCp

	$\text{CH}_3\text{Hg(II)}$	$\text{CH}_3\text{Hg(II)}$
dCp	6.1 Hz	2.4 Hz
$\text{dCp} \cdot 0.35\text{eqCH}_3\text{Hg(II)}$	4.6 Hz	4.2 Hz
dCpG	8.9 Hz	1.9 Hz
$\text{dCpG} \cdot 0.35\text{eqCH}_3\text{Hg(II)}$	9.3 Hz	2.5 Hz

In the ribose derivative the near equal coupling constants in the complexed dinucleoside monophosphate, indicate that the two conformations tg and gt occur to almost the same extent. Thus the dimer is becoming more like the monomer, in terms of its α_1 conformation. Base-stacking is decreased, in agreement with ^{13}C chemical shift data presented previously.

The complexation of dCpG with $\text{CH}_3\text{Hg(II)}$ produces no change in the α_1 populations, the tg conformer remains dominant. This is due to the interdependence of α_1 and the sugar pucker 2E in the deoxyribose.

polynucleotides. Unlike the ribose derivatives, ρ_1 is now independent of base sequence and base-stacking effects.

Zn(II) still occupies a borderline position binding to both N7 and the phosphate chain. However, with the trinucleotide these are now intra-molecular chelates and no insoluble polymers are formed. The Zn(II)-adenosine-5'-triphosphate complexes exhibit a dramatic pH dependence. A small change in binding sites, dependent upon pH, leads to significant changes in complex stability.

The hard metal ion species Mg(II), now has a hard donor site at the β phosphate position and a stable complex is observed, independent of pH.

Independent of the factors which govern site-selectivity; such as metal ion polarizability, solvent and pH, is the classification of a binding site as kinetically or thermodynamically favoured. The thermodynamic site can be calculated from the energies of interaction between a metal ion and potential binding site (10,72). Generally the N1 site in the purines is identified as the thermodynamic site and N7 the kinetic site, since it is readily accessible under neutral conditions (40,95).

In the binding titrations used in this study no activation energy is provided above that available at 20°C.

These experimental conditions, coupled with the known kinetic lability of mercury-nitrogen and mercury-oxygen bonds, leads to the rapid establishment of equilibrium and hence the likely observation of thermodynamically favoured products.

However, the formation of thermodynamic products was impossible to verify since the approach to an identical equilibrium state from two opposite sides could not be carried out.

10.3 Conclusion

Using ^{13}C and ^{15}N NMR it has been possible to determine the binding sites of $\text{CH}_3\text{Hg}(\text{II})$ on the dinucleoside monophosphates CpG and dCpG, and on their constituent mononucleotides 3'CMP, 3'dCMP, 5'GMP and 5'dGMP.

The primary metal ion binding site, at pH 6 on 3'CMP was determined to be N3 and no chelation with the carbonyl was observed for the soft $\text{CH}_3\text{Hg}(\text{II})$ species. This binding site was independent of the type of sugar residue present and caused no change in the torsional angle.

The binding of $\text{CH}_3\text{Hg}(\text{II})$ to 5'GMP at pH 8, was found to be a two step process as studied by ^{15}N and ^{13}C chemical shifts. Initial deprotonation at N1 is followed by metal ion binding.

R - ribose - 5' - monophosphate

The binding does not depend on the sugar type, or affect the torsional angle, but it is highly dependent upon the pH of the solution. Lower pH blocks the reaction and allows the formation of some N7 bound complex, while higher pH favours the interaction of the exocyclic amino group (127).

The binding sites of Cu^{2+} II on the mononucleotides were used to study the metal ion binding properties of the dinucleoside monophosphates.

The primary binding sites were identified as N1 and N7 on the 5'-guanylate residue of both CpG and dCpG at pH 6. Slight N^7 complexation is observed due to the lower pH of the samples. pH 6. Binding does not affect the γ torsional angle, but does reduce the amount of base-stacking experienced by the 3'-cytidylate residue. This decreased base-stacking is apparent from chemical shift changes and a change in J for the 3'-cytidylate residue in CpG.

No change in J is observed for dCpG, because of the angles dependence on sugar pucker and independence of base-stacking effects.

Thus, we have shown ^{13}C NMR to be an effective method for studying the interaction of soft metal ions and dinucleoside monophosphates. Natural abundance ^{15}N NMR studies are not yet possible for these systems.

Over the course of this work, the utility of multinuclear NMR for the study of metal-ion-nucleic acid derivative interactions has been amply demonstrated. A more detailed picture of the complexes formed can be gleaned from examining the ^{13}C , ^{15}N and where applicable

^{31}P and ^{199}Hg spectral data than was possible in earlier work, where ^1H NMR and potentiometric titrations were the primary investigational tools (2,3,7,8).

The primary goal of most studies involving metal ions and nucleic acids is the determination of a binding site and the factors affecting its selection.

Theoretical calculations, taking into account the Coulombic, exchange and delocalization energies of a system, can be used to predict the interaction sites of a metal ion on a base (10,72). These sites are generally the same as those occupied in the nucleosides and nucleotides; with the exception of N3 in the purines which is sterically blocked by the sugar group.

Experimentally, the binding site occupied by a metal ion depends upon the polarizability of the metal ion, the solvent involved and its pH, and the presence of polyphosphate esters on the sugar group.

The nature of the metal ion involved strongly influences the preferred binding site. Soft metal ions such as Hg(II) and $\text{CH}_3\text{Hg(II)}$ interact only with the heterocyclic nitrogens. The borderline species Zn(II) can bind to both the heterocyclic nitrogens and harder oxygen donor species, depending on the nucleic acid derivative involved. The

hard metal ions such as $Mg(II)$ and $Ba(II)$ prefer the harder oxygen donor species, however they will associate, in solvent-mediated complexes, to the ring nitrogens.

These effects are observed for the 6-oxopurine nucleosides, where $Hg(II)$ and $Zn(II)$ bind to the soft N7 donor site. The hard metal ion $Mg(II)$ forms a carbonyl complex at O6 of the purine base.

Solvent effects and the presence of a single 5'-phosphate group are found to alter the metal ion complexes of the 6-oxo-purine nucleotides. $Zn(II)$ now exhibits its borderline polarizability by forming insoluble polymeric species involving N7 and PO_4 binding (82). Insoluble $Zn(II)$ complexes are also formed with adenosine-5'-monophosphate.

The soft metal ion $Hg(II)$ in a slightly basic solution (pH 8) will cause N1 deprotonation and subsequent metallation. This mode of binding is pH dependent, and is effectively blocked at lower pH. The softer $CH_3Hg(II)$ species will bind at N7 at a pH lower than 6, while binding exclusively to N1 at a pH of 8.

The binding of hard metal ions such as $Mg(II)$ or $Ba(II)$ is reduced or prevented in aqueous solutions. The carbonyl site on the 6-oxopurine nucleotides is involved in hydrogen bonding and does not bind the hard metal species. The hard ions themselves are effectively solvated and only outer-sphere binding to the nitrogen sites is observed.

The effects of pH and the extent of phosphorylation on binding site selectivity are further illustrated by the adenosine derivatives: adenosine, adenosine-5'-monophosphate and adenosine-5'-triphosphate.

The two potential binding sites on adenosine, N1 and N7, are similar in their softness, as are the two metal ions $Hg(II)$ and $Zn(II)$

in DMSO. Thus steric effects play a role in site selectivity. The larger ion, Hg(II) preferring to bind at the less hindered N1 position. Zn(II) on the other hand binds exclusively to the N7 position.

The lack of a hard donor site on adenosine effectively prevents the formation of a direct Mg(II) complex.

The addition of a monophosphate group and going to aqueous solutions at pH 10, changes the site selectivity properties of the metal ions. As mentioned previously, Zn(II) with its ability to bind both phosphate and N7 positions produces insoluble complexes. Hg(II) remains complexed at N1, seemingly independent of phosphate addition and solvent changes.

The hard metal ion Mg(II) can now form indirect, solvent mediated complexes with the soft nitrogen sites of adenosine-5'-monophosphate. The N1 site is slightly preferred due to its larger coordination sphere.

At no time with the 6-oxopurine mononucleotides, or the adenosine mononucleotide was binding at the phosphate group detected. The only exception to this trend may be in the insoluble Zn(II) complexes (79,82).

For the triphosphate derivatives of adenosine, binding at the phosphate group is now observed, controlled by the polarizability of the metal ion involved.

The binding of Hg(II) still occurs at N1 for adenosine-5'-triphosphate at pH 7, however at higher or lower pH (i.e. 4 and 10) binding does not occur.

Zn(II) still occupies a borderline position binding to both N7 and the phosphate chain. However, with the trinucleotide these are now intra-molecular chelates and no insoluble polymers are formed. The Zn(II)-adenosine-5'-triphosphate complexes exhibit a dramatic pH dependence. A small change in binding sites, dependent upon pH, leads to significant changes in complex stability.

The hard metal ion species Mg(II), now has a hard donor site at the β phosphate position and a stable complex is observed, independent of pH.

Independent of the factors which govern site-selectivity; such as metal ion polarizability, solvent and pH, is the classification of a binding site as kinetically or thermodynamically favoured. The thermodynamic site can be calculated from the energies of interaction between a metal ion and potential binding site (10,72). Generally the N1 site in the purines is identified as the thermodynamic site and N7 the kinetic site, since it is readily accessible under neutral conditions (40,95).

In the binding titrations used in this study no activation energy is provided above that available at 20°C.

These experimental conditions, coupled with the known kinetic lability of mercury-nitrogen and mercury-oxygen bonds, leads to the rapid establishment of equilibrium and hence the likely observation of thermodynamically favoured products.

However, the formation of thermodynamic products was impossible to verify since the approach to an identical equilibrium state from two opposite sides could not be carried out.

Secondary binding sites were observed by employing nucleosides, such as 7-methylguanosine, where the primary metal ion binding site is blocked. Binding at a secondary N1 site is observed, independent of metal ion polarizability.

^{15}N NMR was found to facilitate the identification of metal ion binding sites on the nucleic acid derivatives, in particular when ^{13}C NMR gave ambiguous results.

^{15}N NMR allowed the observation of an N1 binding site in the 6-oxopurine nucleotides at pH 8. Because the ^{15}N shifts were downfield, as opposed to the normal upfield shift of a complexed nitrogen, the mechanism was determined to involve concomitant deprotonation.

The combination of ^{15}N and ^{31}P NMR allowed one to determine the mechanism and pH dependence of the metal ion catalyzed dephosphorylation of adenosine-5'-triphosphate.

The parameters of N1 binding in the adenosine derivatives were also determined. Previously only N7 binding was well characterized by NMR methods.

^{199}Hg NMR chemical shift titration curves were found to yield significant information about the complexation of $\text{Hg}(\text{II})$ with the nucleosides.

The ^{13}C and ^{15}N results on the monomer species, were then successfully applied to the study of dinucleoside monophosphates, previously unreported in the literature.

The binding site of methylmercury(II) on cytidyl-(3'→5')-guanosine and 2'-deoxycytidyl-(3'→5')-2'-deoxyguanosine was determined. In addition the denaturing ability of the metal ion was observed.

The application of ^{15}N NMR to this final problem was not possible with the available NMR spectrometers or without ^{15}N enriched materials. Based on our previous findings, however, ^{15}N NMR would be expected to provide facile identification of metal ion complexation sites in dinucleosides and higher oligomers.

12.1 General FT NMR Methods

All NMR spectra were obtained using pulsed Fourier Transform (FT) methods (134). The applied rf pulse can excite all possible resonances at one-time due to its rectangular shape.

Quadrature detection is used to enhance the observed S/N. The frequency of the rf pulse carrier is selected to be one-half the spectral width under investigation.

The pulse width, or duration the rf pulse is applied to the system is directly related to the signal strength generated in the receiver coil. The pulse angle θ is defined by the equation,

$$\begin{aligned}\theta &= \gamma B_1 t_p \quad (\text{radians}) \\ &= \frac{360}{2\pi} \gamma B_1 t_p \quad (\text{degrees})\end{aligned}\quad 39$$

where t_p is the pulse width in microseconds.

The acquisition time (AT), during which sampling of the FID occurs controls the ultimate resolution of the spectrum.

$$\text{Resolution} = 1/AT \quad 40$$

This acquisition time is limited by the spectrometer itself and the amount of computer memory available. The highest frequency observable, without fold-back, is equal to the sweep width of the spectrum. The Nyquist theorem requires two sampling points per cycle, thus

$$AT \times 2 \times SW = NP \quad 41$$

where NP is the number of points which must be stored in the computer.

Prior to the application of the next rf pulse a waiting period D1 is required to allow the magnetization to relax back to its

equilibrium value. This process is controlled by the spin-lattice relaxation time T_1 . Generally $5T_1$'s must be allowed, to ensure complete relaxation following a $\pi/2$ pulse. This is prohibitively long in most cases and results in decreased S/N.

A balance between the delay period $AT + D1$ and the pulse width PW must be found to ensure maximum S/N (135). The optimum pulse angle θ_M is defined by the equation where $T_{t1} = AT + D1$ (136).

$$\theta_M = \arccos \left[\exp(-T_{t1}/T_1) \right] \quad \dots 42$$

In all heteronuclear NMR spectra, ^{13}C , ^{15}N , ^{31}P and ^{199}Hg , ^1H noise decoupling was routinely applied. This leads to the decoupling of ^1H spins from the heteronucleus in question, and simplified spectral analyses.

If the ^1H spin in question is coupled by a dipole-dipole relaxation mechanism to the heteronucleus A, an additional effect is observed: The nuclear Overhauser effect arises from a polarization transfer in the spin levels of the heteronucleus and enhances the observed signal, according to the equations shown below.

$$\text{NOEF} = 1 + \frac{\gamma_H}{2\gamma_A} \times \frac{R_1^d}{R_1} \quad \dots 43$$

$$R_1' = R_1^d + R_1^0 \quad \dots 44$$

The nuclear Overhauser enhancement factor (NOEF) is dependent upon the magnetogyric ratio of the nucleus in question, γ_A , and the extent of dipole-dipole relaxation occurring R_1^d with the other mechanisms R_1^0 .

For ^{13}C -NMR spectra, the NOEF is often a maximum of 3 and thus provides an additional mechanism for S/N enhancement.

For ^{31}P and ^{199}Hg spectra, the NOEF is not much greater than the original factor of one due to the lack of dipole-dipole relaxation with the ^1H spins. Thus no significant S/N enhancement is obtained.

The ^{15}N nucleus however has a negative magnetogyric ratio and thus in some circumstances the NOEF can be zero or less than 1, resulting in loss of signal strength. Accordingly for ^{15}N NMR spectra, the NOE is generally suppressed using a gated decoupler sequence. The decoupler is on during the observe pulse and data acquisition period, but remains off during the longer delay periods.

12.2 Spectral Acquisition Parameters

NMR spectra were obtained on a Varian XL-200 superconducting FT NMR spectrometer, except where noted otherwise.

The 4.7 T wide-bore superconducting magnet will accept both 5 mm and 10 mm broad band tunable probes. Frequencies between 20 and 110 MHz, as well as 200 MHz are possible under computer control.

The A/D converter has 12-bits and 32 k of memory are available for data acquisition. Fourier transforms with integer mathematics are used for greatest speed.

Specialized pulse sequence programs are available in the hard disk computer storage unit.

^1H NMR spectra of the dinucleoside monophosphates were obtained using samples lyophilized five times from 99.8% D_2O . Final concentrations were 0.10 M with samples contained in 5 mm tubes.

Standard parameters included a sweep width of 2000 Hz, acquisition times of 8 seconds, and a transmitter offset equal to 500 Hz. A pulse width of 58° or 9 μsec was used and no additional delay between pulses was required. Typically 100 scans were accumulated.

Processing included a resolution enhancement factor equal to 1/3 of the FID length and an apodization factor 3 to 5 times the applied resolution enhancement. This provided increased spectral resolution, to identify overlapping resonances, at the expense of some signal to noise.

Solvent suppression pulse sequences were also applied to some samples where lyophilization was not sufficient to remove the residual H₂O peak. Solvent suppression involves the use of a repetitive two pulse sequence (136) and depends on the relative relaxation times of solvent and sample resonances.

The pulse sequence $(\pi - t - \pi/2 + AT - D1)_n$ is used and as long as the solvent T_1 is greater than 1.4 times the solute T_1 , then the solvent peak can be removed by saturation. The value of the delay t can be calculated based on the T_1 of the solvent (T_{1p}), the desired length of the experiment and the required resolution ($D' = D1 + AT$).

$$\left[2 - \exp(-D/T_{1p}) \right] \exp(-\tau/T_{1p}) = 1 \quad \dots \quad 45$$

The H₂O present in the CpG samples had a T_1 of $12.2 \pm .4$ sec, and thus a total delay D' of 7 seconds was required to suppress the solvent peak.

Samples for ^{31}P NMR analysis were prepared in 10 mm tubes and contained a 5 mm external reference of 85% H_3PO_4 .

Standard acquisition parameters were a sweep width of 4000 Hz, an acquisition time of 1 second, a pulse width of 30° and a delay of 1 second. An internal lock was provided by the D_2O solvent. Generally less than 100 scans were required to provide adequate S/N. High power ^1H noise decoupling was standard.

^{13}C samples were prepared in both 5 and 10 mm tubes. Samples in the 10 mm tubes were referenced to a 4 mm external methanol solution. Samples in the 5 mm tubes were referenced to a 3 mm external TMS reference.

Standard acquisition parameters were a sweep width of 10,000 Hz

and an acquisition time of 1.6 seconds for maximum resolution. A pulse width of 30° degrees and a delay of 1 second were used with high power ^1H noise decoupling. A small amount of line broadening 0.8 Hz was used to improve the overall S/N.

Specialized off-resonance ^{13}C spectra were also obtained using low power square wave ^1H decoupling centered at the proton resonance frequency.

Spectral sweep widths were reduced to 2000 Hz and the acquisition times increased to provide maximum resolution for the measurement of the carbon-phosphorus coupling constants.

^{199}Hg spectra were run at 35.78 MHz in 10 mm tubes. A sweep width of 30,000 Hz and an acquisition time of 0.53 seconds were used. A delay of 1 second and a 70° pulse width were used in the presence of high power ^1H noise decoupling. Samples were referenced using an external reference of 0.55 M HgCl_2 in a 5 mm tube.

Line broadening of 0.53 Hz was employed, and adequate S/N ratios were generally obtained after less than 2000 scans.

^{15}N was the least sensitive nucleus employed in this study and special care was taken in obtaining spectra.

Samples were prepared in 10 mm tubes and referenced to an enriched 2.9 M NH_4Cl in 1.0 M HCl sample run before and after each spectrum. Standard conditions employed an nOe suppressed ^1H decoupler sequence, a 90° pulse width and a 40 second delay. A sweep width of 10,000 Hz, a transmitter offset of -5000 Hz, and an acquisition time of 1.6 sec were used.

A S/N ratio of 5 was generally achieved in 16 hours using 0.1 M solutions. Large amounts of line broadening were required in some cases, up to 5 Hz.

To achieve greater S/N ratios in shorter time periods for the protonated nitrogens the INEPT pulse sequence was used.

12.3 INEPT

Specialized pulse sequences exist to perform specific functions. As an example, for a nucleus with low sensitivity, dipole-dipole coupled to a more sensitive nucleus the INEPT pulse sequence will transfer polarization from the strong nucleus to the weaker one, causing an enhancement in the observed S/N ratio.

INEPT is an acronym for insensitive nuclei enhancement by polarization transfer and was developed in 1979 (137). It is a particularly useful method for systems with unfavourable γ 's, due to the negative magnetogyric ratio of the nuclei in question. Thus the method is often applied to the study of ^{15}N -NMR resonances.

The enhancement arises by transferring spin polarization from a nucleus with a large Boltzmann population difference, usually protons, to the nucleus under investigation. The basic pulse sequence applied is illustrated below (138).

Here PP represents a ^1H - 90° pulse applied using the decoupler coils, PW is a 90° ^{15}N pulse, and D1, D2, and Δ are delay times.

The method in general relies on the generation of spin-echos and Δ is fixed at $1/(4J)$, where J is the spin-spin coupling constant between ^1H and ^{15}N .

The ^{15}N - ^1H coupled spin system can be represented as an AX system. One can observe the greater population difference for the

proton transitions and thus their greater sensitivity.

Considering only the proton spins in the ^{15}N - ^1H coupled system, an initial 90° pulse flips the spins into the x,y plane. The two components of the proton magnetization are labelled α and β . After a time D2 equal to $1/4J$, the two spin vectors are 90° out of phase. A 180° pulse at this point will interchange α and β and reverse their direction of precession. Therefore a second D2 delay would cause a spin-echo to be formed. However this is not the case because a simultaneous 180° ^{15}N pulse is applied to the AX spin system, which inverts the ^{15}N spin states, essentially reversing the ^1H spin state

labels. Thus the spin vectors continue to diverge, at a time D2

away the spins will be 180° out of phase along the x-axis. A third proton pulse of 90° is now applied, however this time phase shifted to the y-axis. This rotates the two vectors to the z-axis

at D2.

This pulse sequence has inverted the ^1H - α spin population while leaving the β population unchanged. A new population diagram for the

spin system, shows how this has enhanced the population difference for one of the ^{15}N transitions, while not changing the second.

Because the ^{15}N and ^1H transitions now share common energy levels, the population difference for the ^{15}N spins has been increased by a factor $1/2$.

A second and often more important consequence of applying this INEPT pulse sequence is that the repetition rate is now governed by the much shorter $^1\text{H}-T_1$'s.

Thus for ^{15}N resonances coupled strongly to a proton spin system the INEPT pulse sequence offers considerable advantage to the generally used ^1H suppression conditions, with their long delay times.

12.4 2-Dimensional NMR Spectroscopy

Previous NMR techniques have produced a linear response as a result of perturbing a linear system. In a classical, linear NMR experiment only amplitude, perhaps with phase information is considered as a function of frequency.

A whole new branch of NMR is just now opening up, which studies the non-linear properties of a system. These non-linear parameters include relaxation effects and the connectivity (J coupling) of the energy levels in a spin system. In addition, multiple quantum transitions can be observed. To do this, a second parameter must be observed as a function of frequency. A number of methods have been developed using rf field strength, time, or a second frequency as this extra parameter. The graphical representation of such an experiment leads to a 2-dimensional (2-D) NMR spectrum.

The most useful of these techniques involves the use of two independent time variables as a function of which a signal amplitude can be measured (139).

In any two dimensional NMR experiment there are four important time periods: 1) a preparation period, 2) an evolution period, 3) a mixing period, which may be absent, and 4) a detection period. The two time variables important in signal detection are t_1 and t_2 , the evolution and detection periods respectively. It is the correlation between a spin's behaviour in these periods which provides the added parameter in a 2-D NMR spectrum.

A two dimensional response of a spin detected during time t_2 requires that it "remember" what has happened to it during a previous

time t_1 . This corresponds to an effect on the magnetization amplitude or phase of the spin during t_1 . Thus both phase and amplitude modulation occur in two dimensional NMR. The type of modulation controls the method of presentation and the line shape involved.

For phase modulation, the phase of a magnetization component at the beginning of the detection period t_2 is linearly dependent upon the length of the evolution period t_1 . Amplitude modulation occurs if the magnetization component in the transverse plane at the beginning of t_2 is an oscillating function of the time t_1 . The phase of the detected signal is independent of t_1 . Another important form of amplitude modulation occurs from the addition or subtraction of two phase modulated signals prior to Fourier transformation.

12.41 Homonuclear Chemical Shift Correlation Spectroscopy

In all two dimensional shift correlation experiments there is a relation or correlation between two nuclei at particular chemical shifts. This mutual interaction is shown by a cross peak at (Ω_A, Ω_B) in the 2-D spectrum. Nuclei A and B may be of the same type or different; homonuclear and heteronuclear correlation. This correlation may occur through direct or long-range scalar couplings, chemical exchange, cross-relaxation or dipole-dipole couplings in non-isotropic samples.

The original two dimensional experiment proposed by Jeener(140), was a homonuclear chemical shift correlation experiment (COSY). It provides an alternative to selective decoupling, to sort through a

highly coupled homonuclear spin system. The 2-D NMR spectrum will reveal the connectivity patterns of the spin system.

The basic pulse sequence is shown below:

where T is the experimental delay between the end of data acquisition and the start of the next sequence. This ensures a thermal equilibrium exists in the spin system at this point. The application of a $\pi/2$ pulse at this point will create transverse magnetization components for all allowed transitions, mn . These components will precess during the evolution time t_1 , with their characteristic frequencies Ω_{mn} .

At the end of this time period a second pulse is applied. This pulse is generally less than 90° , so it only generates single quantum transitions, kl . This redistributes the transverse magnetization from mn to kl . The amount of coherence between the states is proportional to $\sin(\Omega_{mn} t_1 + \theta_{kl,mn})$, where θ is a phase constant dependent upon the connectivity of the two transitions. Any pair of transitions in a weakly-coupled spin- $\frac{1}{2}$ AX system are either parallel or progressively or regressively connected (141).

If the transitions kl and mn belong to the same multiplet, they are said to be parallel and the phase constant $\theta_{kl,mn}$ is 0° . The signal detected during t_2 is modulated as a function of t_1 and an off-diagonal peak will show up in the 2-D NMR spectrum at (δ_x, δ_A) .

If the two transitions are not in the same multiplet, they are either progressively or regressively connected and the phase constant

is -90° or $+90^\circ$. These transitions give rise to signals detected during t_2 which are not modulated as a function of t_1 . These peaks therefore occur on the axis of a 2-D spectrum.

Thus in a homonuclear chemical shift correlated 2-D NMR spectrum a typical 1-D NMR spectrum occurs along the diagonal axis, whereas off-diagonal peaks indicate a J-coupling between the two resonances. Thus a highly overlapped region of an NMR spectrum can be analyzed by running only a single experiment rather than several selective decoupling ones.

A COSY ^1H - ^1H 2-D NMR spectrum of CpG was run at 500 MHz on a Bruker 500. A sweep width of 625 Hz in both dimensions was centered over the ribose proton resonances between 3.6 and 6.2 ppm.

A data matrix of 256 x 512 data points was Fourier transformed to yield the final 2-D spectrum. A resolution of 4.88 Hz was obtained under these conditions.

12.5 Materials and Sample Preparation

All nucleosides and nucleotides were purchased from Sigma Chemical Company and used without further purification. The dinucleoside monophosphates were also obtained from Sigma Chemical Company.

The Silanor DMSO- d_6 , 99.9% pure and the D_2O , 99.8 minimum purity were obtained from Merck, Sharn and Dohme Isotopes.

$HgCl_2$ was obtained from Fisher Scientific and was purified by vacuum sublimation to remove paramagnetic impurities. $Zn(NO_3)_2 \cdot 6H_2O$, $Ba(NO_3)_2$, and $BaCl_2$ were also obtained from Fisher Scientific.

$Mg(NO_3)_2 \cdot 6H_2O$ and NaOH were obtained from Anachemia. The concentrated HCl was purchased from the J.T. Baker Chemical Co.

CH_3HgI and $[CH_3Hg_3O]OH$ were received from Alpha Products, while CH_3HgCl was obtained from Pfatty and Bauer Inc.

All samples were prepared on a Mettler Type H16 analytical balance with an uncertainty of ± 0.0005 g. The pH of the resulting solutions was determined using Accutint indicator papers from Anachemia. When necessary the pH of solutions was measured more precisely with a Fisher Acumet pH meter fitted with a combination glass electrode.

The pH of a solution was adjusted using 50% HCl and NaOH solutions, prepared in D_2O .

If a 1H spectrum was to be recorded for a sample in D_2O , the sample was lyophilized at least 3 times from 99.8% D_2O .

12.6 Synthesis of N1-CH₃Hg(II)-5'GMP

2.0 mmoles (0.976g) 5'GMP were dissolved in 20.0 mL water, 2.0 mmoles of CH₃HgOH (0.453 g of [CH₃Hg₃O]OH) were added and the solution was stirred for 2 hours. The pH remained constant at 8.5.

The solution was slowly evaporated and the resulting solid was recrystallized from ethanol.

The yield was 1.3 g (100%) of white, plate-like crystals, which decomposed at 195°C. The product was confirmed as N1-CH₃Hg(II)-5'GMP using MS, ¹H NMR and ¹³C NMR data.

The ¹H NMR spectrum of the product was obtained on a Varian 360 spectrometer. It contained the resonances characteristic of guanosine-5'-monophosphate, as the sample was run in D₂O the presence of an N1 proton could not be confirmed or denied. In addition a singlet CH₃ peak was observed at 1.5 ppm, characteristic of a CH₃Hg(II) substituent.

The ¹³C NMR spectrum was recorded under the standard conditions. Ten resonances due to the nucleotide are observed and the methylmercury peak occurs at -0.5 ppm relative to TMS, as expected (35).

The mass spectrum was obtained on a VG770E fast atom bombardment mass spectrometer in a glycerol carrier matrix. A parent ion at m/e = 621 is observed, consistent with C₁₁H₁₄O₈N₅HgPNa₂. Major peaks are observed at 598 and 575 corresponding to the loss of sodium. Fragment ion peaks corresponding to guanosine, guanine, ribose, and methylmercury are also observed.

This procedure was repeated to prepare the N1-methylmercury derivatives of 5'dGMP and 5'IMP.

REFERENCES

- 1) L.G. Marzilli, T.J. Kistenmacher, G.L. Eichhorn in Nucleic Acid Metal Ion Interactions, ed. T. Spiro, John Wiley & Sons, Toronto, 1980.
- 2) L.G. Marzilli. Prog. Inorg. Chem. 23 255 (1977).
- 3) Metal Ions in Biological Systems, Vol. 1-18, ed. H. Sigel, Marcel Dekker Inc., New York. 1974-1985.
- 4) P.O.P. Ts'o (ed.). Basic Principles of Nucleic Acid Chemistry, Vol. 1 (1974), Vol. 2 (1974), Academic Press, New York.
- 5) P.O.P. Ts'o, N.S. Kondo, M.P. Schweizer, D.P. Hollis. Biochem 8 997 (1969).
- 6) B. Rosenberg. Cancer Chemother. Rep. 59 589 (1975).
- 7) R.M. Izatt, J.J. Christensen, J.H. Rytting. Chem. Rev. 71 439 (1971).
- 8) G.L. Eichhorn in Inorganic Biochemistry, Vol. 2, ed. G.L. Eichhorn, Elsevier, New York, 1973.
- 9) "Inorganic Chemistry" A.G. Sharpe. Longman, New York, 1981.
- 10) T.A. Glassman, G.K. Klopman, C. Cooper. Biochem. 12 5013 (1973).
- 11) M.L. Martin, J.-J. Delpuech, G.J. Martin. "Practical NMR Spectroscopy" Heyden, 1980.
- 12) E.M. Purcell, H.C. Torrey, R.V. Pound. Phys. Rev. 69 37 (1946).
- 13) F. Bloch, W.W. Hansen, M. Packard. Phys. Rev. 69 127 (1946).
- 14) E.D. Becker. "High Resolution NMR Theory and Chemical Applications" Academic Press, 1980.
- 15) M. Karplus. J. Chem. Phys. 30 11 (1959); J. Am. Chem. Soc. 85 2870 (1963).
- 16) W.C. Dickenson. Phys. Rev. 81 717 (1951).
- 17) H.S. Gutowsky, D.W. McCall. Phys. Rev. 82 748 (1951).
- 18) B.E. Mann in "NMR and the Periodic Table" ed. R.K. Harris, B.E. Mann, Academic Press, New York, 1978.
- 19) M. Crutchfield, C.H. Dungan, J.H. Letcher, V. Mark, J. Van Wazer in "Topics in Phosphorous Chemistry-5" ed. M. Grayson, E. Griffith, Interscience Publ., New York, 1967.

- 20) J.F. Nixon, A. Pidcock in Ann. Rev. NMR Spec. 2 ed. E.F. Mooney, Academic Press, New York, 1969.
- 21) D.B. Davies. Prog. NMR Spec. 12 291 (1978).
- 22) R.G. Kidd in Ann. Rep. in NMR Spec. 1QA ed. G.A. Webb, Academic Press, 1980.
- 23) M.A. Sens, N.K. Wilson, P.D. Ellis, J.D. Odom. J. Mag. Res. 19 323 (1975).
- 24) M.F. Roberts, D.A. Vidusek, G. Bodenhausen. FEBS Lett. 117 311 (1980).
- 25) J.L. Sudmeier, T.G. Perkins. J. Am. Chem. Soc. 99 7732 (1977).
- 26) P. Peringer. Inorg. Chim. Acta 39 67 (1980).
- 27) J.L. Sudmeier, R.R. Birge, T.G. Perkins. J. Mag. Res. 30 491 (1978).
- 28) A.J. Canty. J. Organometallic Chem. 144 371 (1978).
- 29) A.J. Canty. J. Organometallic Chem. 179 447 (1979).
- 30) R.C. Lauterbur. J. Chem. Phys. 26 217 (1957).
- 31) C.H. Holm. J. Chem. Phys. 26 707 (1957).
- 32) J.B. Stothers. "¹³C NMR Spectroscopy", Academic Press, New York, 1972.
- 33) "Topics in ¹³C NMR Spectroscopy" ed. G.C. Levy, Wiley-Interscience, New York; Vol. 1 (1974), Vol. 2 (1976).
- 34) J.W. Copper. "Spectroscopic Techniques for Organic Chemists.", John-Wiley & Sons, New York, 1980.
- 35) P.L. Laszlo, P. Stang. "Organic Spectroscopy: Principles and Applications.", Harper & Row Publ., New York, 1980.
- 36) A.J. Jones, M.W. Winkley, D.M. Grant, R.K. Robins. Proc. Nat. Acad. Sci. USA. 65 27 (1970).
- 37) D.E. Dorman, J.D. Roberts. Proc. Nat. Acad. Sci. USA. 65 19 (1970).
- 38) H.H. Mantsch, I.C.P. Smith. Biochem. Biophys. Res. Comm. 46 808 (1972).
- 39) L.G. Marzilli, B. de Castro, J.P. Caradonna, R.C. Stewart, C.P. van Vuuren. J. Am. Chem. Soc. 102 916 (1980).
- 40) L.G. Marzilli, B. de Castro, C. Solrzano. J. Am. Chem. Soc. 104 461 (1982).
- 41) J.B. Lambert, G. Binsch, J.D. Roberts. Proc. Nat. Acad. Sci. USA. 61 735 (1964).

- 42) R.L. Lichter, J.D. Roberts. J. Am. Chem. Soc. 94 2495 (1972).
- 43) J.M. Briggs, L.F. Farnell, E.W. Randall. J. Chem. Soc., Chem. Comm. 680 (1971).
- 44) G.C. Levy, R.L. Lichter. "Nitrogen-15 Nuclear Magnetic Resonance Spectroscopy." Wiley-Interscience Publ., New York, 1979.
- 45) V. Markowski, G.R. Sullivan, J.D. Roberts. J. Am. Chem. Soc. 99 714 (1977).
- 46) P. Buchner, W. Maurer, H. Ruterjans. J. Mag. Res. 29 45 (1978).
- 47) C. Dyllick-Brenzinger, G.R. Sullivan, P.P. Pang, J.D. Roberts. Proc. Nat. Acad. Sci. USA. 77 5580 (1980).
- 48) M. Watanabe, H. Sugeta, H. Swahashi, Y. Kyogoku, M. Kainosho. Eur. J. Biochem. 117 553 (1981).
- 49) T.L. James, J.L. James, A. Lapidot. J. Am. Chem. Soc. 103 6748 (1981).
- 50) I.M. Ismail, P.J. Sadler in Metal Chemotherapeutic Agents, Am. Chem. Soc. Publ., 1983.
- 51) G.C. Levy, J.L. Dechter. J. Am. Chem. Soc. 102 6191 (1980).
- 52) J.A. Happe, M. Morales. J. Am. Chem. Soc. 88 2077 (1966).
- 53) G.W. Buchanan, J.B. Stothers. Can. J. Chem. 59 291 (1981).
- 54) G.W. Buchanan, M. Bell. Can. J. Chem. 61 2445 (1983).
- 55) W.E. Lamb, Jr. Phys. Rev. 60 817 (1941).
- 56) N.F. Ramsey. Phys. Rev. 78 699 (1950); 77 567 (1950); 83 540 (1951).
- 57) H.F. Hameka. J. Chem. Phys. 40 3127 (1964).
- 58) L.C. Snyder, R.G. Parr. J. Chem. Phys. 34 827 (1961).
- 59) J.A. Pople. Disc. Faraday Soc. 34 7 (1962).
- 60) J.A. Pople. J. Chem. Phys. 37 53 (1962).
- 61) J.A. Pople. J. Chem. Phys. 37 60 (1962).
- 62) J.A. Pople. Mol. Phys. 7 301 (1964).
- 63) K.A.K. Ebraheem, G.A. Webb, M. Witanowski. Org. Mag. Res. 8 317 (1976).
- 64) W.H. Flygare, J. Goodisman. J. Chem. Phys. 49 3122 (1968).
- 65) C.J. Jameson, H.S. Gutowsky. J. Chem. Phys. 40 1714 (1964).
- 66) G.A. Webb in NMR and the Periodic Table ed. R.K. Harris, B.E. Mann. Academic Press, New York, 1978.

- 67) G.A. Webb in NMR of Newly Accessible Nuclei, Vol. 1 Chemical and Biochemical Applications. ed. P. D. Barto, Academic Press, New York, 1983.
- 68) J.P. Janowski, J.P. Macquet, J.L. Butour. Biochimie **60** 901 (1978).
- 69) L.G. Marzilli, K. Wilkowsky, C. Chiang, T. Kistenmacher. J. Am. Chem. Soc. **101** 7504 (1979).
- 70) R. Gillert, B. Fisher, R. Bau. J. Am. Chem. Soc. **102** 7812 (1980).
- 71) L.G. Marzilli. Adv. Inorg. Biochem. **3** 47 (1981).
- 72) D. Perahia, A. Pullman, B. Pullman. Theoret. Chim. Acta **43** 207 (1977).
- 73) S.T. Rao, M. Sundaralingam. J. Am. Chem. Soc. **91** 1210 (1969).
- 74) E. Shifter, K. Trueblood. Acta Crystallogr. **18** 1067 (1965).
- 75) E. Buncel, A. Norris, W. Racz, S. Taylor. Inorg. Chem. **20** 98 (1981).
- 76) K.W. Jennette, S.J. Lippard, D.A. Ucko. Biochem. Biophys. Acta **402** 403 (1975).
- 77) L. Marzilli, R. Stewart, C. Van Vuuren, B. de Castro, J. Caradonna. J. Am. Chem. Soc. **100** 3967 (1978).
- 78) C.B. Reese, R. Saffhill. J.C.S. Perkin I 2937 (1972).
- 79) J. Rifkind, G. Eichhorn. J. Am. Chem. Soc. **94** 6526 (1972).
- 80) G. Eichhorn, Y. Shin. J. Am. Chem. Soc. **90** 7323 (1968).
- 81) J. Butzow, G. Eichhorn. Biopolymers **3** 95 (1965).
- 82) S. Miller, D. VanDervier, L.G. Marzilli. J. Am. Chem. Soc. **107** 1048 (1985).
- 83) G. Hammes, D. Miller. J. Chem. Phys. **46** 1533 (1967).
- 84) J.A. Happe, R.L. Ward. J. Phys. Chem. **83** 3457 (1979).
- 85) K. Mashos. Acta Biochim. Polonica **25** 113 (1978); **28** 183, 317 (1981).
- 86) H.A. Tajmir-Riahi, T. Theophanides. Can. J. Chem. **61** 1813 (1983).
- 87) J. Bock. J. Inorg. Biochem. **12** 119 (1980).
- 88) J. Bock, D. Ash. J. Inorg. Biochem. **12** 119 (1980).
- 89) T.-D. Son, M. Roux, M. Ellenberger. Nuc. Acid Res. **2** 1101 (1975).
- 90) D. Wilson, B. Heyl, R. Reddy, L. Marzilli. Inorg. Chem. **21**, 2527 (1982).

- 91) R. Phillips. Chem. Rev. 66, 501 (1966).
- 92) S. Shimokawa, H. Fukui, K. Hotta. J. Am. Chem. Soc. 95 1777 (1973).
- 93) K. Schiller, V. Schiller-Krattiger, R. Martin. J. Am. Chem. Soc. 103 6833 (1981).
- 94) N. Farrell. J.C.S. Chem. Comm. 1014 (1980).
- 95) P.V. Vestues, R.B. Martin. J. Am. Chem. Soc. 103 806 (1981).
- 96) "Biochemistry", A. Lehninger, Worth Publ. Inc., 1975.
- 97) A. Szent Gyorgyi. Bioenergetics, Chapter 10, Academic Press, New York, 1957.
- 98) T. Glassman, C. Copper, L. Harrison, T. Swift. Biochem. 10 843 (1971).
- 99) Y.H. Mariam, R.B. Martin. Inorg. Chim. Acta 35 23 (1979).
- 100) A. Lanir, N.-T. Yu. J. Biol. Chem. 254 5882 (1979).
- 101) H. Sigel, P.E. Amsler. J. Am. Chem. Soc. 98 7390 (1976).
- 102) M. Cohn, T. Hughes. J. Biol. Chem. 237 176 (1962).
- 103) H. Sigel, F. Hofstittér, R.B. Martin, R. Milburn, V. Scheller-Krattiger, K. Sheller. J. Am. Chem. Soc. 106 7935 (1984).
- 104) T.G. Spiro in Inorg. Biochem. Vol. 1 ed. G.L. Eichhorn, Elsevier, New York, 1973.
- 105) P. Yong, U. Shankar Nandi, N. Kallenbach. Biochem. 21 62 (1982).
- 106) B.L. Vallee, D.D. Ulmer. Annual Rev. Biochem. 41 91 (1972).
- 107) J.M. Neumann, W. Guschlbauer, S. Tran-Dinh. Eur. J. Biochem. 100 141 (1979).
- 108) S. Tran-Dinh, J.M. Neumann, J. Borrel. Biochem. Biophys. Acta 655 167 (1981).
- 109) D.M. Cheng, R.H. Sarma. J. Am. Chem. Soc. 99 7333 (1977).
- 110) W.J.P. Blonski, F.E. Hruska, K.L. Sadana, P.C. Lowen. Biopolymers 22 605 (1983).
- 111) P.N. Borer, L.S. Kan, P.O.P. Ts'O. Biochem. 14 4847 (1975).
- 112) L.S. Kan, P.N. Borer, P.O.P. Ts'O. Biochem. 14 4864 (1975).
- 113) D.R. Kearns, P.A. Meraw, N. Assa-Munt, R.W. Behling in Nucleic Acids: The Vectors of Life, p. 113 1983.
- 114) C.D. Barry, J.A. Glasel, A.C.T. North, R.J.P. Williams, A.V. Xavier. Biochem. Biophys. Acta 262 101 (1972).

- 115) J.L. Alderfer, P.O.P. Ts'0. *Biochem.* 16 2410 (1977).
- 116) T. Schleich, B.P. Cross, I.C.P. Smith. *Nuc. Acid Res.* 3 355 (1976).
- 117) I.C.P. Smith, H.J. Jennings, R. Deslauriers. *Accts. Chem. Res.* 8 306 (1975).
- 118) D.B. Davies, H. Sadihot. *Biopolymers* 22 1843 (1983).
- 119) R.D. Lapper, H.H. Mantsch, I.C.P. Smith. *J. Am. Chem. Soc.* 94 6243 (1972).
- 120) D.B. Davies in *Nucleic Acids: The Vectors of Life*. p. 141 (1983).
- 121) P. Lankhorst, C.A.G. Haasnoot, C. Erkelens, C. Altona. *J. Biomolecular Structure Dynamics* 1 1387 (1984).
- 122) *Nature (London)* 263 567 (1976).
- 123) *Nucleic Acid Geometry and Dynamics*. Ed. Sarma, Pergamon Press, New York, 1980.
- 124) R.B. Simpson. *J. Am. Chem. Soc.* 86 2059 (1964).
- 125) A. Norris, R. Kumar, E. Buncel, A. Beauchamp. *J. Inorg. Biochem.* 21 277 (1984).
- 126) S. Mansy, T. Wood, J. Sprowles, R. Tobias. *J. Am. Chem. Soc.* 96 1762 (1974).
- 127) S. Mansy, R. Tobias. *J. Am. Chem. Soc.* 96 6874 (1974).
- 128) S. Mansy, R. Tobias. *J. Frick. Biochim. Biophys. Acta* 378 319 (1975).
- 129) J.J. Mulvihill. *Science* 176 132 (1972).
- 130) R. Anderson, A. Maki, C. Ott. *Biochem.* 19 4412 (1980).
- 131) G. Fronza, R. Mondelli, F.W. Randall, G.P. Gardini. *J.C.S. Perkin II* 1746 (1977).
- 132) H. Swahashi, Y. Kyogoku. *J. Am. Chem. Soc.* 99 7761 (1977).
- 133) A. Jack, J.E. Ladner, A. Klug. *J. Mol. Biol.* 108 619 (1976).
- 134) D. Shaw. *Fourier Transform NMR Spectroscopy*, Elsevier, Amsterdam, 1976.
- 135) R.R. Ernst, W.A. Anderson. *Rev. Sci. Instrum.* 37 93 (1966).
- 136) "Practical NMR Spectroscopy", M.L. Martin, J.J. Delpuech, G.J. Martin. Heyden, London 1980.
- 137) G.A. Morris, R. Freeman. *J. Am. Chem. Soc.* 101 760 (1979).

138) Varian XL200 Operator Manual.

139) A. Bax. 2-D NMR in Liquids, Delft University Press, Holland, 1982.

140) J. Jeener. Ampere Int. Summer School, Basko Polji, Yugoslavia, 1971.

141) W.P. Aue, E. Bartholdi, R.R. Ernst. J. Chem. Phys. 64 2229 (1976).

END

0 9 | • 0 4 | • 8 6 |

FIN