1

PM-1 3 1/4" x 4" PHOTOGRAPHIC MICROCOPY TARGET
NBS 1010a ANSI/ISO #2 EQUIVALENT

National Library
of Canada

Acquisitions and
Bibliographic Services Branch

385 Wellington Street
Ottawa, Ontario
K1A 0N4

Bibliothèque nationale
du Canada

Direction des acquisitions et
des services bibliographiques

385, rue Wellington
Ottawa (Ontario)
K1A 0N4

Your file Votre référence

Our file Notre référence

NOTICE

The quality of this microform is heavily dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to ensure the highest quality of reproduction possible.

If pages are missing, contact the university which granted the degree.

Some pages may have indistinct print especially if the original pages were typed with a poor typewriter ribbon or if the university sent us an inferior photocopy.

Reproduction in full or in part of this microform is governed by the Canadian Copyright Act, R.S.C. 1970, c. C-30, and subsequent amendments.

AVIS

La qualité de cette microforme dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

S'il manque des pages, veuillez communiquer avec l'université qui a conféré le grade.

La qualité d'impression de certaines pages peut laisser à désirer, surtout si les pages originales ont été dactylographiées à l'aide d'un ruban usé ou si l'université nous a fait parvenir une photocopie de qualité inférieure.

La reproduction, même partielle, de cette microforme est soumise à la Loi canadienne sur le droit d'auteur, SRC 1970, c. C-30, et ses amendements subséquents.

Statistical Modeling and Analysis of the *Salmonella*/Microsome Assay

by
Qun Wang

A thesis submitted to
the Faculty of Graduate Studies and Research
in partial fulfilment of the requirements for the degree of

Doctor of Philosophy

Department of Mathematics and Statistics

**Carleton University
Ottawa, Canada
December, 1994**

**©Copyright 1994
Qun Wang**

National Library
of Canada

Acquisitions and
Bibliographic Services Branch

395 Wellington Street
Ottawa, Ontario
K1A 0N4

Bibliothèque nationale
du Canada

Direction des acquisitions et
des services bibliographiques

395, rue Wellington
Ottawa (Ontario)
K1A 0N4

Your file Votre référence

Our file Notre référence

**THE AUTHOR HAS GRANTED AN
IRREVOCABLE NON-EXCLUSIVE
LICENCE ALLOWING THE NATIONAL
LIBRARY OF CANADA TO
REPRODUCE, LOAN, DISTRIBUTE OR
SELL COPIES OF HIS/HER THESIS BY
ANY MEANS AND IN ANY FORM OR
FORMAT, MAKING THIS THESIS
AVAILABLE TO INTERESTED
PERSONS.**

**L'AUTEUR A ACCORDE UNE LICENCE
IRREVOCABLE ET NON EXCLUSIVE
PERMETTANT A LA BIBLIOTHEQUE
NATIONALE DU CANADA DE
REPRODUIRE, PRETER, DISTRIBUER
OU VENDRE DES COPIES DE SA
THESE DE QUELQUE MANIERE ET
SOUS QUELQUE FORME QUE CE SOIT
POUR METTRE DES EXEMPLAIRES DE
CETTE THESE A LA DISPOSITION DES
PERSONNE INTERESSEES.**

**THE AUTHOR RETAINS OWNERSHIP
OF THE COPYRIGHT IN HIS/HER
THESIS. NEITHER THE THESIS NOR
SUBSTANTIAL EXTRACTS FROM IT
MAY BE PRINTED OR OTHERWISE
REPRODUCED WITHOUT HIS/HER
PERMISSION.**

**L'AUTEUR CONSERVE LA PROPRIETE
DU DROIT D'AUTEUR QUI PROTEGE
SA THESE. NI LA THESE NI DES
EXTRAITS SUBSTANTIELS DE CELLE-
CI NE DOIVENT ETRE IMPRIMES OU
AUTREMENT REPRODUITS SANS SON
AUTORISATION.**

ISBN 0-612-02965-4

Canada

Name Qian Wang

Dissertation Abstracts International is arranged by broad, general subject categories. Please select the one subject which most nearly describes the content of your dissertation. Enter the corresponding four-digit code in the spaces provided.

Public Health

SUBJECT TERM

0573

SUBJECT CODE

U·M·I

Subject Categories

THE HUMANITIES AND SOCIAL SCIENCES

COMMUNICATIONS AND THE ARTS

Architecture 0729
Art History 0377
Cinema 0900
Dance 0378
Fine Arts 0357
Information Science 0773
Journalism 0391
Library Science 0399
Mass Communications 0708
Music 0413
Speech Communication 0459
Theater 0465

EDUCATION

General 0515
Administration 0514
Adult and Continuing 0516
Agricultural 0517
Art 0273
Bilingual and Multicultural 0282
Business 0688
Community College 0275
Curriculum and Instruction 0727
Early Childhood 0518
Elementary 0524
Finance 0277
Guidance and Counseling 0519
Health 0680
Higher 0745
History of 0520
Home Economics 0278
Industrial 0521
Language and Literature 0279
Mathematics 0280
Music 0522
Philosophy of 0998
Physical 0523

Psychology 0525
Reading 0535
Religious 0527
Sciences 0714
Secondary 0533
Social Sciences 0534
Sociology of 0340
Special 0529
Teacher Training 0530
Technology 0710
Tests and Measurements 0288
Vocational 0747

LANGUAGE, LITERATURE AND LINGUISTICS

Language 0679
General 0289
Ancient 0290
Linguistics 0291
Modern 0291
Literature 0401
General 0294
Classical 0295
Comparative 0297
Medieval 0298
Modern 0316
African 0591
American 0305
Asian 0352
Canadian (English) 0355
Canadian (French) 0593
English 0311
Germanic 0312
Latin American 0315
Middle Eastern 0313
Romance 0314
Slavic and East European 0314

PHILOSOPHY, RELIGION AND THEOLOGY

Philosophy 0422
Religion 0318
General 0321
Biblical Studies 0319
Clergy 0320
History of 0322
Philosophy of 0469
Theology 0323

SOCIAL SCIENCES

American Studies 0323
Anthropology 0324
Archaeology 0326
Cultural 0327
Physical 0310
Business Administration 0272
General 0770
Accounting 0454
Banking 0338
Management 0385
Marketing 0501
Canadian Studies 0503
Economics 0505
General 0508
Agricultural 0509
Commerce Business 0510
Finance 0511
History 0358
Labor 0366
Theory 0351
Folklore 0578
Geography 0578
Gerontology 0578
History 0578
General 0578

Ancient 0579
Medieval 0581
Modern 0582
Black 0328
African 0331
Asia, Australia and Oceania 0332
Canadian 0334
European 0335
Latin American 0336
Middle Eastern 0333
United States 0337
History of Science 0585
Law 0398
Political Science 0615
General 0616
International Law and 0617
Relations 0616
Public Administration 0814
Recreation 0452
Social Work 0626
Sociology 0627
General 0938
Criminology and Penology 0631
Demography 0628
Ethnic and Racial Studies 0629
Individual and Family 0630
Studies 0700
Industrial and Labor 0344
Relations 0709
Public and Social Welfare 0999
Social Structure and 0453
Development 0629
Theory and Methods 0709
Transportation 0999
Urban and Regional Planning 0453
Women's Studies 0453

THE SCIENCES AND ENGINEERING

BIOLOGICAL SCIENCES

Agriculture 0473
General 0285
Agronomy 0475
Animal Culture and 0476
Nutrition 0359
Animal Pathology 0478
Food Science and 0479
Technology 0480
Forestry and Wildlife 0817
Plant Culture 0777
Plant Pathology 0746
Plant Physiology 0306
Range Management 0287
Wood Technology 0308
Biology 0309
General 0379
Anatomy 0329
Biostatistics 0353
Botany 0369
Cell 0793
Ecology 0410
Entomology 0307
Genetics 0317
Limnology 0433
Microbiology 0821
Molecular 0778
Neuroscience 0472
Oceanography 0786
Physiology 0760
Radiation 0425
Veterinary Science 0996
Zoology 0996
Biophysics 0786
General 0760
Medical 0425
Earth Sciences 0996

Geodesy 0370
Geology 0372
Geophysics 0373
Hydrology 0388
Mineralogy 0411
Paleobotany 0345
Paleoecology 0426
Paleontology 0418
Paleozoology 0985
Palynology 0427
Physical Geography 0368
Physical Oceanography 0415

HEALTH AND ENVIRONMENTAL SCIENCES

Environmental Sciences 0768
Health Sciences 0566
General 0300
Audiology 0992
Chemotherapy 0567
Dentistry 0350
Education 0769
Hospital Management 0758
Human Development 0982
Immunology 0564
Medicine and Surgery 0347
Mental Health 0569
Nursing 0570
Nutrition 0380
Obstetrics and Gynecology 0354
Occupational Health and 0381
Therapy 0571
Ophthalmology 0419
Pathology 0572
Pharmacology 0382
Pharmacy 0573
Physical Therapy 0574
Public Health 0575
Radiology 0575
Recreation 0575

Speech Pathology 0460
Toxicology 0383
Home Economics 0386

PHYSICAL SCIENCES

Pure Sciences 0485
Chemistry 0749
General 0486
Agricultural 0487
Analytical 0488
Biochemistry 0738
Inorganic 0490
Nuclear 0491
Organic 0494
Pharmaceutical 0495
Physical 0754
Polymer 0405
Radiation 0605
Mathematics 0986
Physics 0606
General 0608
Acoustics 0748
Astronomy and 0507
Astrophysics 0798
Atmospheric Science 0759
Atomic 0609
Electronics and Electricity 0610
Elementary Particles and 0752
High Energy 0756
Fluid and Plasma 0611
Molecular 0463
Nuclear 0798
Optics 0759
Radiation 0609
Solid State 0610
Statistics 0346
Applied Sciences 0984
Applied Mechanics 0984
Computer Science 0984

Engineering 0537
General 0538
Aerospace 0539
Agricultural 0540
Automotive 0541
Biomedical 0542
Chemical 0543
Civil 0544
Electronics and Electrical 0348
Heat and Thermodynamics 0545
Hydraulic 0546
Industrial 0547
Marine 0794
Materials Science 0548
Mechanical 0743
Metallurgy 0551
Mining 0552
Nuclear 0549
Packaging 0765
Petroleum 0554
Sanitary and Municipal 0790
System Science 0428
Geotechnology 0796
Operations Research 0795
Plastics Technology 0994
Textile Technology 0994

PSYCHOLOGY

General 0621
Behavioral 0384
Clinical 0622
Developmental 0620
Experimental 0623
Industrial 0624
Personality 0625
Physiological 0989
Psychobiology 0349
Psychometrics 0632
Social 0451

The Faculty of Graduate Studies and Research
acceptance of the thesis.

Statistical Modeling and Analysis of the *Salmonella*/Microsome Assay

submitted by

Qun Wang

in partial fulfilment of the requirements
for the degree of Doctor of Philosophy

Chairman, Department of Mathematics and Statistics

Thesis Supervisor

External Examiner

Carleton University

December, 1994

Abstract

This thesis considers a series of problems relating to the statistical design and analysis of the Ames *Salmonella*/microsome assay. First, a biologically based dose-response model is developed, allowing for histidine diffusion in the Ames assay. The proposed model reduces to the two previous models proposed in literature. An exact solution for the model is presented; conditions under which previously published approximations hold are established using both analytical and numerical methods.

Second, a generalized score test for trend is proposed allowing for extra-Poisson variation both between and within laboratories. The asymptotic properties of the proposed statistic are derived. Limited simulation studies demonstrate that the use of historical control data enhances the power of the test for trend.

Third, optimal experimental designs for the Ames assay are developed. Emphasis is placed on C, D, and A-optimal designs under a linear-exponential dose-response model. Suboptimal designs that may be used in practice are recommended following the analysis of 1120 datasets from a recent collaborative trial conducted by the International Programme on Chemical Safety. Bayesian and minimax optimal designs are also considered briefly.

Finally, the Rao-Scott transformation is used to study tests for trend, parameter estimation, and goodness-of-fit. With the Rao-Scott transformation, no assumptions about the nature of the mean-variance relationship of the data are necessary. Limited simulation studies showed that the goodness-of-fit test using the transformed data has better operating characteristics than that based on the original data.

Acknowledgements

First of all, my sincere thanks go to my thesis supervisor, Dr. D. Krewski. Without his insight into the topics included in my research, this thesis may not have come into existence. He is also chiefly responsible for my interest in biostatistics, especially in the area of environmental health. I deeply grateful for his direction, encouragement, time and efforts. I have thoroughly enjoyed working under his supervision.

I would also like to express my thanks to Dr. E. Saleh, my thesis co-supervisor for his support that helped to make it possible for me to come to Carleton University. I benefitted a lot from his consistent encouragement throughout my study.

I would like to acknowledge Dr. B. Margolin, Dr. J.N.K. Rao and Dr. M. Csörgö for the time they put in for my thesis examination. As well, I would like to acknowledge Dr. B. Leroux for his helpful comments and suggestions on certain aspects of the thesis.

I appreciate the Department of Mathematics and Statistics for the rigorous academic training.

Last, but not least, I owe my achievement to my wife Shaorun Li, my parents, and my grandmother for their love and support.

Contents

Acceptance Sheet	i
Abstract	iii
Acknowledgements	iv
Contents	v
List of Tables	vii
1 Introduction	1
1.1 Motivation and Objectives	1
1.2 Literature Review	2
1.2.1 Modeling the Ames Assay	2
1.2.2 Extra-Poisson Variation and Test for Trend	4
1.2.3 Optimal Experimental Designs	6
1.2.4 Rao-Scott Transformation	7
1.2.5 Background and Outline	8
2 Biologically Based Dose Response Models	16
2.1 Introduction	16
2.1.1 The Ames <i>Salmonella</i> /Microsome Assay	17

2.2	A Stochastic Model for the Ames Assay	18
2.3	A Histidine Diffusion Model	20
2.3.1	Model Formulation	20
2.3.2	An Exact Solution for the Histidine Diffusion Model	21
2.3.3	Simulation Results	25
2.4	Summary	27
3	Tests for Trend with Historical Controls	28
3.1	Introduction	28
3.2	Mean-Variance Function	29
3.3	Generalized Score Tests	32
3.4	Asymptotic Distribution	35
3.5	Calculation of Score Statistic	54
3.6	Some Simple Properties	55
3.7	Summary	57
4	Optimal Experimental Designs	59
4.1	Introduction	59
4.2	C, A, and D-Optimal Designs	60
4.2.1	C-Optimal Designs	61
4.2.2	A-Optimal Designs	66
4.2.3	D-Optimal Designs	67
4.3	Suboptimal Designs	71
4.3.1	Efficiency of Suboptimal Designs	72
4.3.2	Designs in the IPCS Study	74
4.4	Bayesian Optimal Designs	94
4.4.1	Bayesian D-Optimal Designs	94
4.4.2	Bayesian C-Optimal Designs	96
4.4.3	Bayesian A-Optimal Designs	97

4.5	Optimal Design under Minimax Criteria	98
4.6	Summary	101
5	Rao-Scott Transformation	103
5.1	Test for Trend	104
5.1.1	Test Statistics	104
5.1.2	Simulation Results	107
5.2	Parameter Estimation	107
5.2.1	Parameter Estimation without Using the Rao-Scott Transfor- mation	108
5.2.2	Parameter Estimation Using the Rao-Scott Transformation . .	109
5.2.3	Simulation Results	110
5.3	Goodness-of-Fit Test	111
5.3.1	Goodness-of-Fit Tests without Using the Rao-Scott Transfor- mation	112
5.3.2	Goodness-of-Fit Test Using the Rao-Scott Transformation . .	115
5.3.3	Simulation Results	117
5.3.4	The Asymptotic Distribution of the Proposed Goodness-of-Fit Test	121
5.4	Summary and Conclusions	123
6	Summary and Further Research	125

List of Tables

Table	Title	Page
Table 2.1	Ratio Performance of R_0 , R_1 and R_2	27
Table 3.1	Type I and Type II Error of Test for Trend with and without Historical Controls	58
Table 4.1	Two Examples of C-optimal Designs	66
Table 4.2	Two Examples of A-optimal Designs	67
Table 4.3	Two Examples of D-optimal Designs	71
Table 4.4	C-optimal Designs for the Ames Assay	78
Table 4.5	D-optimal Designs for the Ames Assay	79
Table 4.6	A-optimal Designs for the Ames Assay	80
Table 4.7	Efficiency of Original 6-Dose Design and Recommended 3-Dose Design Under C-optimality Relative to C-optimal 3-Dose Design	81
Table 4.8	Efficiency of Original 6-Dose Design and Recommended 3-Dose Design Under D-optimality Relative to D-optimal 3-Dose Design	82
Table 4.9	Efficiency of Original 6-Dose Design and Recommended 3-Dose Design Under A-optimality Relative to A-optimal 3-Dose Design	83
Table 4.10	Efficiency of Fixed 3-6 Dose Designs with Equal Allocation (C-optimality)	84
Table 4.11	Optimal Allocation for Fixed 3-6 Dose Designs (C-optimality)	85
Table 4.12	Efficiency of Fixed 3-6 Dose Designs with Unequal Allocation (C-optimality)	86
Table 4.13	Efficiency of Fixed 3-6 Dose Designs with Equal Allocation (D-optimality)	87
Table 4.14	Optimal Allocation for Fixed 3-6 Dose Designs (D-optimality)	88
Table 4.15	Efficiency of Fixed 3-6 Dose Designs with Unequal Allocation (D-optimality)	89
Table 4.16	Efficiency of Fixed 3-6 Dose Designs with Equal Allocation (A-Optimality)	90
Table 4.17	Optimal Allocation for Fixed 3-6 Dose Designs (A-optimality)	91
Table 4.18	Efficiency of Fixed 3-6 Dose Designs with Unequal Allocation (A-optimality)	92

Table	Title	Page
Table 4.19	Efficiency of Fixed 3-6 Dose Designs	93
Table 4.20	An Example of a Bayesian D-optimal Design	96
Table 4.21	An Example of a Bayesian C-optimal Design	97
Table 4.22	An Example of a Bayesian A-optimal Design	98
Table 4.23	An Example of a Minimax Optimal Design	101
Table 5.1	Type I and II Errors of Test for Trend at a Significance level of 5%	108
Table 5.2	Estimates of the Parameter in the Linear-exponential Model Before and After Applying the Rao-Scott Transformation	110
Table 5.3	Comparison of Parameter Estimates Before and After Applying the Rao-Scott Transformation	112
Table 5.4	Relative Bias, Relative Standard Deviation and Relative Mean Squared Errors of Regression Parameters Estimation with and without the Rao-Scott Transformation	113
Table 5.5	Type I and Type II Error Rates of Goodness-of-Fit Test Before (B) and After (A) Applying the Rao-Scott Transformation	122

Chapter 1

Introduction

1.1 Motivation and Objectives

Genetic toxicology became established as a recognized discipline in the late 1960's and early 1970's. Toxicological tests are now available for genetic alterations in bacteria, fungi, plants, insects, cultured mammalian cells or small mammals. The tests are relatively short-term, and can identify both gene and chromosomal mutations.

A mutagen is a chemical or radiological agent that is capable of causing heritable changes in genetic material. Since mutation is a necessary component of neoplastic transformation, tests for mutagenicity have also been used to identify potential carcinogens. Traditionally, testing chemicals for carcinogenic potential has been done using long-term animal experiments. These experiments are time consuming and expensive (Krewski & Goddard, 1990). Consequently, there is considerable interest in short-term tests which have been developed for genetic alternations (Kilbey et al., 1984).

The rapid adoption by many laboratories of *Salmonella* and other assays led to an explosion of work in environmental mutagenesis. The Ames *Salmonella*/microsome test (Ames et al., 1975) is the foremost example of a microbial mutagenicity test system (Margolin et al., 1989). Based on validation studies, Ames et al. (1975)

concluded that the Ames assay is the only established test for the prediction of carcinogenicity.

In the Ames test, auxotrophic and prototrophic strains of *Salmonella typhimurium* are used. The latter can synthesize histidine, an amino acid needed for bacterial growth, whereas the former cannot grow without histidine. The auxotrophic strain can, however, revert to histidine independence either spontaneously or by the induction of a mutagen. Therefore, if a reversion occurs on a plate containing *Salmonella*, the test agent and histidine, the resulting revertant and its descendants that arise by cell division can grow and divide and eventually form a visible colony. Those auxotrophs not becoming revertant may continue to divide until all of the histidine in the plate is exhausted. When no histidine is left, no more visible colonies form because auxotrophic cell division and growth terminates. In the Ames test, the mutagenic response is quantified by counting the number of visible colonies on plates containing approximately 10^8 microbes (Broekhoven & Nestmann, 1991).

This thesis addresses several issues in the statistical design and analysis of the *Salmonella*/microsome assay. First, modeling of the Ames assay based on fundamental biological principles is considered. Second, tests for trend in mutation rates utilizing historical controls are developed. Third, optimal experimental designs for the Ames assay are constructed. And fourth, the use of Rao-Scott transformation to simplify the statistical analysis of Ames assay data is explored.

1.2 Literature Review

1.2.1 Modeling the Ames Assay

A number of different dose-response models have been developed for analyzing Ames assay data. We will briefly review six of these models which were proposed by Margolin et al. (1981), Bernstein et al. (1982), Stead et al. (1981), Snee & Irr (1984), and Krewski et al. (1993).

The response to mutagenic chemical substance may be an increase in the number of revertant colonies as the dose of the test agent is increased. However, if the

chemical is toxic, then the number of revertant colonies might decline at high doses due to cytotoxicity. Thus, the dose response curve might have an initial rise followed by a decline.

Using single hit kinetics, Bernstein et al. (1982) argued that the dose-response curve for mutagenicity should be approximately linear, at least in the initial portion of the dose response curve. Under this model, the probability that a microbe mutates at dose d is given by

$$P(d, \beta) = 1 - \exp(-\beta_0 - \beta_1 d) \approx \beta_0 + \beta_1 d,$$

where $\beta = (\beta_0, \beta_1)$.

Stead et al. (1982) modeled the *Salmonella* assay using an empirical function of sufficient complexity to cope with the variety of data which might result from the test. The model assumes that the number of revertant colonies at any dose follows a Poisson distribution, while the mean number of revertants per plate is a nonlinear function of up to four parameters. Snee and Irr (1984) also used an empirical approach to modeling based on the Box-Cox transformation.

Myers et al. (1981) suggested that the mean number μ of revertants per plate can be influenced by mutagenicity, toxicity, and saturation. The most general form of their model is given by

$$\mu(d) = C[1 - e^{-f(d)}]e^{-g(d)} \approx C f(d)e^{-g(d)}.$$

Here, $f(d)$ and $g(d)$ are nonnegative functions of d , measuring the mutagenic and toxic effects at dose d , respectively, both of which may incorporate saturation effects. The special case of linear mutagenicity, $f(d) = \beta_0 + \beta_1 d$, attenuated by exponentially decreasing toxicity, $g(d) = \beta_2 d$, can be handled with this model.

Margolin et al. (1981) developed a mechanistic model to describe and model the Ames assay. A mechanistic model is one which is developed from fundamental biological principles which are believed to underlie the process which is being studied. This model will be reviewed in chapter 2.

Krewski et al. (1993) proposed another biologically based model by allowing for histidine diffusion within the plate agar. It is assumed that the slope of the initial

(linear) component of the dose response curve provides a measure of mutagenic potency. In order to allow for toxic effects which may be nonlinear at low doses, it is suggested that correcting for toxicity can be accomplished by using the factor $\exp(-\beta_2 d^\theta)$ with $\theta > 1$. This model is consistent with the generally accepted view that most toxic effects are not likely to occur below a certain dose threshold. Krewski et al. (1992) illustrate the application of their model using data from a collaborative trial using the *Salmonella* assay coordinated by the International Programme on Chemical Safety (IPCS). A value of $\theta = 2$ is used, reflecting a sublinear dose-response relationship for toxic effects in the low dose region.

Broekhoven and Nestmann (1991) conducted a detailed comparison of these models and concluded that for statistical purposes, the choice between models can be left to the preference of the analyst. However, they expressed preference for biologically based models of the type proposed by Margolin et al. (1981) and Krewski et al. (1993). Most (1993) conducted a limited empirical comparison of the above statistical models, and calculated the predicted mutagenic potency based on each model. In most cases, all models generated comparable estimates of mutagenic potency, and tend to fit equally well.

1.2.2 Extra-Poisson Variation and Test for Trend

Margolin et al. (1981) observed that because the variance among replicate plates exceeded the mean for these plates, the distribution of the number of revertants per plate did not appear to follow a Poisson distribution. They conjectured that laboratories are unable to maintain a high degree of homogeneity among replicate plates because of variability in pipetting the microbes, the plate agar, the S9 liver homogenate and the test compound.

The Poisson mean can be considered as a random variable that varies from plate to plate. Based on the above conjecture, a negative binomial model (a gamma mixing of the Poisson distribution) was adopted. As a result, the relationship between the mean μ and the variance $Var(Y)$ of the observed number of counts Y can be written as $Var(Y) = \mu + \mu^2/c$ for some constant $c > 0$. Another method of modeling extra-

Poisson variation in the data is based on the generalized Poisson family described by Consul (1989) and employed by Bernstein et al. (1982). This model implies that the variance σ^2 is proportional to a power p of the mean: $Var(Y) = \sigma^2 \mu^p$. An alternative approach is to use only the mean and variance structure implied by the mixed Poisson model, estimating the regression coefficients by quasi-likelihood, and the variance parameter by the method of moments (Breslow, 1984; Williams, 1982). Quasi-likelihood leads quite generally to consistent estimates of the regression coefficients, even if the variance function is misspecified (Moore and Tsiatis, 1991). Three tests are available under likelihood and quasi-likelihood theory: the Wald test based on the comparison of estimated coefficients with their standard errors; a quasi-likelihood ratio test based on comparison of deviances under full and reduced models (McCullagh and Nelder, 1983, section 8.4); and the quasi-likelihood score test (McCullagh and Nelder, 1983, sec.11.4) that uses estimating equations for inference. Unlike the score test method, the Wald and deviance tests require the fitting of the full model. In addition, the deviance is not always explicitly calculable, and the Wald test is not invariant under model reparameterization (Breslow, 1984).

Tests for trend in bioassay data have been considered by many authors. Following the earliest work of Cochran (1954) and Armitage (1955), Tarone (1982) considered the use of historical control counts in the test for trend. Kikuchi & Yanagawa (1988) proposed a score statistic based on the conditional likelihood. Krewski et al. (1991) also considered conditional and unconditional tests with historical controls. In the Ames assay study, Leroux et al. (1994) proposed several methods to test for trend with historical controls using both ordinary likelihood theory and an estimating equation approach to inference. Prentice et al. (1991) developed tests for trend for quantal response data assuming certain relations between the current and historical data. Recently, Fung et al. (1994) studied tests for trend in developmental toxicity using the Rao-Scott transformation (1992). Simpson & Margolin (1986, 1990) considered nonparametric tests for trend designed to handle the dose-response curves that exhibit a downturn at high doses. Such downturns are attributable to toxicity at high doses, and are quite common in the Ames assay.

1.2.3 Optimal Experimental Designs

The information obtained from any bioassay can be enhanced through a judicious choice of the experimental design. The experimenter has considerable latitude in the selection of the number of dose groups, the actual dose levels, and the allocation of the number of observations to each dose group.

The general theory of optimal experimental design has undergone vigorous development since 1950, with early contributions by Elfving (1952, 1955, 1959), Chernoff (1953), de la Garza (1954, 1956), Guest (1958) and Kiefer (1958, 1959, 1960). Further developments were reported by Fedorov & Malynov (1972), Whittle (1973), and Kiefer (1974). Optimal designs for linear models have been extensively studied. In the case of generalized linear models, Wu (1988), Ford et al. (1992) considered both C and D-optimal designs. Sitter and Wu (1993a,b) used an F-optimal criterion. Sitter & Torsney (1994) considered experiments with two design variables.

Optimal experimental design in carcinogenicity experiments has received considerable attention. For a recent overview, see Portier (1991) and Pukelsheim (1993). Krewski and Kovar (1982) studied optimal designs for the one-hit model and the probit model with known slope parameter. They also considered the efficiency of sub-optimal designs with doses near, but not at, the optimal dose. The results for the one-hit model were later extended to include the two-parameter logit, probit and extreme-value models with zero background (Krewski et al., 1986). Hoel and Jennrich (1979) used the same approach to find optimal bioassay designs for a model similar to the multistage model of Armitage (1954). Krewski et al. (1984) studied the probit model, logistic model, gamma multi-hit model and the Weibull model, allowing for non-zero background. Some related problems concerning experimental designs for extrapolation were also studied by Lawless (1987).

In the nonlinear situation, optimal designs depend on the unknown model parameters. However, these parameters are generally unknown. Chaloner (1984) and Sitter (1992) introduced Bayesian and minimax optimality criteria, respectively, to handle uncertainty in the unknown model parameters.

1.2.4 Rao-Scott Transformation

Overdispersion in bioassay data is quite common (Margolin et al. 1981). The effect of ignoring overdispersion in the satisfied analysis has been studied by Cox (1983) and others. In particular, if overdispersion is not properly considered, the standard errors of the parameter estimates may be underestimated. As noted earlier, Margolin et al. (1981) used the negative binomial regression model for the analysis of the Ames assay data. Under quasi-likelihood theory, only the mean and variance structure are specified, and the corresponding estimating equations are solved simultaneously with the moment equation for the overdispersion parameters.

Krewski & Zhu (1994) studied developmental toxicity using the generalized estimating equations. Breslow (1984, 1990), Dean et al. (1989a,b), Moore & Tsiatis (1991), and Krewski et al. (1993) employed quasi-likelihood methods for overdispersed Poisson data. Breslow (1990) proposed both model-based and empirical score tests, as well as an empirical covariance matrix, similar to that used by Liang and Zeger (1986). These empirical methods are robust to misspecification of the variance function, unlike the model-based approach which exactly specifies the variance structure. Random effects methods that model the inter-individual variability directly have also been developed to handle the overdispersion problem (Lawless, 1987a, 1987b; Breslow, 1984).

In contrast to the above methods, Rao and Scott (1992) developed a simple method for clustered binary data, based on the concept of design effect used in sample surveys. They proposed the use of a simple data transformation for binary data before conducting the statistical analysis. Since the transformed data is not subject to overdispersion, standard statistical methods can then be applied. Fung et al. (1994) performed a simulation study comparing the Rao-Scott method with quasi-likelihood and other methods. Their simulation suggested that the Rao-Scott method performs well in terms of size and power of test for trend. Krewski & Zhu (1995) also used the Rao-Scott transformation to estimate benchmark doses in developmental toxicity experiments. They compared the estimates based on generalized estimating equations applied to the original data with estimates obtained after using

the Rao-Scott transformation, and found good agreement between the two methods. Recently, Scott & Rao (1994) extended the Rao-Scott method for clustered binary data to the case of overdispersed Poisson count data.

1.2.5 Background and Outline

In chapter 2, we review previous models of Margolin et al. (1981) and Krewski et al. (1993), and propose a biologically-based model for the Ames assay, that takes into account histidine diffusion within the plate agar. Although the proposed model is not easily used in practice due to its complicated form, it reduces to the same model given by Margolin et al. (1981) and Krewski et al. (1993) when identical model assumptions are made. We examine the difference between the proposed model and that developed by Krewski et al. (1993) both analytically and by computer simulation. Our results indicate that the latter model is a good approximation to that developed here. Since the model given by Krewski et al. (1993) has a simple form, it is easier to use in practice.

In chapter 3, we examine tests for trend in response with increasing dose. In the Ames *Salmonella*/microsome assay (Broekhoven and Nestmann, 1991), the number of revertant colonies of bacteria observed in replicate culture plates are counted at a series of increasing dose levels $d_0 < d_1 \cdots < d_K$, including the unexposed control at $d_0 = 0$. Let Y_{1j} denote the number of revertant colonies observed in plate $j = 1, \dots, n_1$, at dose $i = 0, 1, \dots, K$ and let Y_{1i} denote the total number of revertant colonies at dose d_i . If each observation Y_{1j} follows a Poisson distribution with mean $\mu_i = \beta_0 + \beta_1 d_i$, then Y_{1i} follows the Poisson distribution with mean equal to $n_1 \mu_i$. For the problem of testing for a linear trend in count data, Armitage (1955) proposed the statistic

$$T_0 = \sum_{i=0}^K Y_{1i} d_i - \bar{Y} \sum_{i=0}^K n_{1i} d_i$$

with variance estimated by

$$\hat{V}(T_0) = \bar{Y} \sum_{i=0}^K n_{1i} d_i^2 - n_1^{-1} \left(\sum_{i=0}^K n_{1i} d_i \right)^2,$$

where $\bar{Y} = \sum_{i=0}^K Y_{1i}/n_1$, and $n_1 = \sum_{i=0}^K n_{1i}$. Under the null hypothesis of no trend, $\beta_1 = 0$, the standardized statistic $Z_{CA} = T_0/\sqrt{\hat{V}(T_0)}$ is asymptotically normally distributed (Leroux et al., 1994).

Suppose now that information on the response rate in the control groups from other experiments is available. These historical controls provide some information on the rate of spontaneous mutation. Tarone (1982) proposed a statistical test for trend which makes use of the available historical control information. Let Y_j denote the total number of revertants across the n_j plates in historical control group $j = 2, \dots, s$, and let θ_j denote the expected response rate in group j . Assuming that the θ_j are independent gamma random variates with mean μ and variance $\mu\phi$, Y_j follows a negative-binomial distribution with mean $n_j\mu$ and variance $n_j\mu(1 + \mu\phi)$, where the parameter $\phi > 0$ represents overdispersion with respect to the Poisson distribution ($\phi = 0$ for the Poisson distribution). If the parameters μ and ϕ are known, Tarone (1982) showed that the score test for trend based on the negative-binomial likelihood of the data is

$$T = \sum_{i=0}^K Y_{1i} - \bar{Y} \sum_{i=0}^K n_{1i} d_i,$$

where $\bar{Y} = (Y_1 + \mu\phi^{-1})/n$, with $n = n_1 + \phi$. An estimate of the variance of T is given by

$$\hat{V}(T) = \bar{Y} \left[\sum_{i=0}^K n_{1i} d_i^2 - n^{-1} \left(\sum_{i=0}^K n_{1i} d_i \right)^2 \right].$$

When μ and ϕ are replaced by their maximum likelihood estimates, $\hat{\mu}$ and $\hat{\phi}$, Tarone's statistic $Z_T = T/\sqrt{\hat{V}(T)}$ is asymptotically normally distributed.

Following Prentice et al. (1991), Leroux et al. (1994) extended Tarone's method by applying an estimating equation approach to inference. With this approach, it is only necessary to specify the first two moments of the data, instead of the complete distribution of the data. Leroux et al. (1994) assume that the mean μ of the historical control response rates is equal to the current control rate and thus the data from current experiment follow Poisson distribution; only the historical controls were considered to be subject to extra Poisson variation of between studies.

Margolin et al. (1981) observed that because the variance among replicate plates exceeds the mean for these plate within the same experiment, the distribution of the number of revertant colonies per plate did not appear to follow a Poisson distribution. A possible explanation for this finding is that the expected number of revertant colonies per plate varies among replicate plates, resulting in extra-Poisson variation. Therefore, we consider two types of overdispersion: we use ϕ_1 to denote the overdispersion parameter within each experiment, and ϕ_2 to denote the overdispersion parameter between experiments. Analysis of the Ames assay data generated by the International Programme on Chemical Safety (IPCS) (Krewski et al. 1992) showed that between laboratory variation is at least three times larger on average than within laboratory variation on a logarithmic scale. This empirical finding indicates that within and between laboratory variation needs to be handled separately.

Breslow (1990) gave a comprehensive review of the theory of hypotheses testing in the overdispersed Poisson case. In chapter 3, we propose a test for trend which considers both between and within laboratory overdispersion. The proposed test for trend extends the results given by Leroux et al. (1994). The asymptotic distribution of the test statistic is shown to be normally distributed, using arguments similar to those employed by Moore (1986). A simulation study is also conducted to examine type I error and power of the proposed test for trend, with and without using the historical control data.

The information obtained from the Ames *Salmonella*/microsome assay can be enhanced through a judicious choice of the experimental design. The experimenter has considerable latitude in the selection of the number of dose groups, the dose levels and the allocation of the available plates to each dose group. In chapter 4, the topic of optimal design of the Ames assay is studied.

Suppose that the observation Y follow a certain probability distribution, $p(y|d, \theta, \tau)$ with nuisance parameter τ , at dose level d , and we wish to estimate θ . If we assume sufficient regularity of the family $\{p(y|d, \theta, \tau); u, \tau\}$, then for a single observation on y , the partitioned Fisher information matrix for (θ, τ) is

$$I(d; \theta, \tau) = \begin{pmatrix} I_{\theta\theta}(d) & I_{\theta\tau}(d) \\ I_{\tau\theta}(d) & I_{\tau\tau}(d) \end{pmatrix},$$

where $I_{\theta\theta}(d)$ is the $k \times k$ matrix whose (i, j) th element is $E\{-\partial^2 \log p(y|d, \theta, \tau) / \partial \theta_i \partial \theta_j\}$. Similar definitions hold for $I_{\theta\tau}(d)$ and $I_{\tau\tau}(d)$. The information for N independent observations taken respectively at $d_1, d_2, \dots, d_N$ is

$$I(d; \theta, \tau) = \sum_{i=1}^N I(d_i; \theta, \tau),$$

where $d = (d_1, \dots, d_N)$.

Assume that d is such that $I(d; \theta, \tau)$ is non-singular. Then $I^{-1}(d; \theta, \tau)$ provides a lower bound for the variance matrix of the unbiased estimator of (θ, τ) . Its leading $k \times k$ sub-matrix

$$V(d; \theta, \tau) = [I_{\theta\theta}(d) - I_{\theta\tau}(d) L_{\tau\tau}^{-1}(d) I_{\tau\theta}^T(d)]^{-1}$$

is a lower bound for the variance matrix of an unbiased estimator of θ . Note the dependence of this lower bound on the design d . Note also that if N is large, the variance covariance matrix of the maximum-likelihood estimation of θ is approximately $V(d; \theta, \tau)$. The basic idea is that we should choose d to make $V(d; \theta, \tau)$ as small as possible in some sense. By doing so we are choosing d to get as much information as possible about θ , in a Fisherian sense.

Two questions arise in the above discussion. One is that how to define the term "as small as possible". We will be discussing different criteria, including C-, A- and D- optimality, in chapter 4. The other question is whether or not we can find d , independent of θ and τ , which makes $V(d; \theta, \tau)$ small in the desired sense for all θ and τ . The answer to this question is yes for a certain class of problems, namely linear regression problems. The main results concerning linear optimal design are summarized by Silvey (1980). However, for nonlinear problems the criteria generally depend on the parameters θ or τ .

In the non-linear situation, the optimal design depends on the unknown model parameters. To obtain an optimal design, we thus must know the value of the

parameters. One way to circumvent this difficulty is to assume that good initial estimates are available, either from previous experiments or expert opinion. These designs are termed “locally” optimum. Chaloner & Larntz (1989) proposed designs which formally account for prior uncertainty in the parameter values. They derived a general theory for concave design criteria for non-linear models and then applied the theory to logistic regression. They introduced prior distributions for the uncertain model parameters; the new criteria were considered to be Bayesian, although no posterior distributions are involved. They used a prior distribution for the unknown parameters rather than a single guess as to their most likely values. The theory of optimal Bayesian design for non-linear problems parallels the theory for linear problems as presented in Silvey (1980) and Fedorov (1972, 1981). One result obtained by Chaloner & Larntz (1989) is that as uncertainty in the prior distribution increases, so does the number of support points in the optimal design. Previous work in this area by Tsutakawa (1972) and Abdelbasit and Plackett (1983) is restricted to designs with observations at equally spaced points with equal number of observations at each point. Sitter (1992) proposed a minimax procedure for choosing designs that are robust against poor initial estimates of the parameters. As in the case of Bayesian criterion, the minimax method shows that the more uncertain the experimenter is about the initial parameter estimates, the more spread out is the proposed design.

In chapter 4, optimal design are studied for count data obtained in the Ames assay. The discussion is limited to the theory and application of designs optimal for parameter estimation. Three criteria, namely C-, A- and D-optimality, are studied in detail. Suppose β_0 , β_1 and β_2 are the parameters of interest in the linear-exponential model given by Krewski et al. (1993), with

$$\mu(d_i) = EY_{i,j} = (\beta_0 + \beta_1 d_i) \exp(-\beta_2 d_i).$$

The objective of C-optimality is to minimize $Var(\hat{\beta}_1)$, that of A-optimality is to minimize $\sum_{i=0}^2 Var(\hat{\beta}_i)$, and that of D-optimality is to minimize the determinant of the variance-covariance matrix of $(\hat{\beta}_0, \hat{\beta}_1, \hat{\beta}_2)^T$. Under these three criteria, the number of design points, the spacing of the design points, and the allocations of the

observations to each design point are investigated and calculated using the general theory of locally optimal designs. Numerical methods are used whenever an analytic solution is not available.

Next, we develop optimal designs for dose-response models fitted to the IPC's database. Twenty laboratories worldwide participated in this study and provided the bioassay data. The U.S. National Institute of Standards and Technology provided homogeneous reference samples of urban air and diesel particle and a coal tar solution (test agents) to each participating laboratory, along with samples of benzo(a)pyrene and 1-nitropyrene which served as positive controls. Krewski et al. (1993) fit the linear-exponential model to these data, using quasi-likelihood methods. Using the estimated values of the model parameters, we find the locally optimal design for each experiment. All of these optimal designs are three points design (i.e. designs with exactly three dose levels, including the control). In practice, a three point design is seldom used for two reasons. First, construction of an optimal design requires knowledge of the parameters of the underlying stochastic model from which the experimental data arise, which is generally unavailable in advance of the study. Second, since optimal design involve a relatively small number of critically placed doses, poor dose selection or the loss of an entire treatment group may compromise the integrity of the original design. Using the three point designs as benchmarks, we investigate the efficiencies of a number of suboptimal designs with up to six doses.

Bayesian optimal (Chaloner & Larntz, 1989) and minimax optimal (Sitter, 1992) criteria designs are also considered briefly. Prior distributions of the uncertain model parameters are introduced. The simplex method is used to find the optimal number of dose levels, dose spacing, and number of observations allocated to each dose level. In the non-Bayesian case, there are bounds for the number of dose levels in an optimal design (Chernoff, 1953). However, for Bayesian or minimax criterion, no such bound exists. Therefore we restrict the number of dose levels under study to be less than six in both the Bayesian and minimax cases. Results from several examples show that the optimal designs are still three point designs regardless of the dispersion in the prior distribution. This phenomenon is different from the binary data situation (Chaloner & Larntz, 1989, Sitter, 1992) in which the number of design

points becomes larger as the prior distribution becomes more distributed.

The existence of extra-Poisson or extra-Binomial data in the toxicological experiments requires the introduction of an overdispersion parameter and the correct mean and variance relationship when modeling the data or conducting regression analysis. Two types of models dealing with count data are proposed in the literature. In the first type, the Poisson mean of individual observed counts is assumed to follow a gamma distribution. As a result, we obtain the negative-binomial model for the count data with larger variance than assumed by the Poisson model (Lawless, 1987a). Margolin et al. (1981) modeled the Ames *Salmonella*/microsome data using the negative-binomial distribution. Lawless (1987a) obtained maximum likelihood estimators of the regression parameters as well as likelihood ratio and Wald tests for detecting overdispersion. The generalized Poisson family has also been proposed to handle the extra-Poisson data (cf. Krewski et al., 1993). Since the distribution of the data is fully specified, maximum likelihood methods may be used for inference.

The second type of model is based on quasi-likelihood or generalized estimating equation methods. In this case, only the mean and variance structure of the data needs to be specified, rather than the entire probability distribution of the data. The regression parameters are then estimated by quasi-likelihood methods, while the overdispersion parameters are estimated by moment methods. Since this method only concerns the first two moments, it is robust to misspecification of the mixing distribution in the fully parametric formulation. Numerous authors, including Breslow (1984, 1990), Dean et al. (1989a,b), Moore & Tsiatis (1991), and Krewski et al. (1993) studied this method. In terms of applications, Krewski et al. (1993) applied this methodology to the analysis of the Ames assay data. Krewski et al. (1992) conducted a detailed analysis of the Ames assay data collected from 1121 experiments in the IPCS database.

Although methods exist for handling overdispersion, simpler or more efficient methods of analysis are desirable. Rao and Scott (1992) proposed a simple method for clustered binary data, based on sample survey techniques. The idea is to transform the original data, and then apply standard binomial-based methods. In a large scale simulation study, Fung et al. (1994) demonstrated that the Rao-Scott method

performs well in terms of size and power in testing for trend. Recently, Scott and Rao (1994) extended their method to the case of overdispersed Poisson count data.

In chapter 5, the Rao-Scott transformation is applied to the Ames assay. A straightforward application of the Rao-Scott method is presented to test for trend and to estimate the regression parameters in the linear-exponential model. Comparisons are also made between the results from Rao-Scott method and those based on a direct application of generalized estimating equation (GEE) methods applied to the original data. A small simulation study shows that the results of the two different methods are comparable in both testing for trend and parameter estimation. The issue of goodness-of-fit is also studied. It is of interest to develop a method to examine whether the mean of the data is really in a certain curvature form (e.g. linear-exponential).

Suggestions for goodness-of-fit test have been given by Leroux et al. (1993), although no detailed formula was developed. This test, based on the deviance function, is developed in chapter 5. We also use the Rao-Scott transformation to develop a goodness-of-fit test based on the transformed data. Simulations are conducted to examine the type I and type II errors of the goodness-of-fit tests with and without using the Rao-Scott transformation. We found that both tests had satisfactory type I and type II errors when the sample size is moderately large. The goodness-of-fit test based on the Rao-Scott transformation appears to perform better than the test based on the original data, when the assumed mean-variance relationship is not satisfied.

Chapter 2

Biologically Based Dose Response Models

2.1 Introduction

In this chapter, we discuss biologically based dose-response models for the Ames *Salmonella*/microsome assay. In particular, we review the stochastic model developed by Margolin et al. (1981), and the extension of the model proposed by Krewski et al. (1993). An exact solution for the latter model is derived, and the accuracy of the approximate solution given by the latter authors confirmed.

We begin with a detailed description of the biological basis of the Ames assay (section 2.2). In section 2.3, a biologically based model for the Ames assay developed by Margolin et al. (1981) is reviewed. This stochastic model is based on a series of fundamental biological assumptions about the way in which microbes evolve. An extension to this model given by Krewski et al. (1993) allowing for diffusion of histidine about the plate agar is discussed in section 2.4, and an exact solution derived. The validation of the approximate solution given by Krewski et al. (1993) is confirmed using computer simulation. A brief summary of the results of this chapter is given in section 2.5.

2.1.1 The Ames *Salmonella*/Microsome Assay

The objective of the Ames *Salmonella*/microsome assay is to detect reverse mutations from auxotrophic cells requiring histidine for growth (His-) to histidine-independent prototrophic cells (His+). According to the description in the paper by Maron and Ames (1983), 0.1ml of a bacterial culture grown overnight in nutrient broth is added to each plate as the initial inoculum. The actual number N_0 of cells added is between 10^7 and 10^8 . Variation within these limits is not considered to seriously affect the number of colonies produced (Maron and Ames, 1983). A trace amount of histidine is incorporated into each soft agar overlay to allow residual growth of the auxotrophic bacteria during mutagenic treatment. Growing cells are generally more sensitive to mutagen. With an initial inoculum of 10^8 cells, it has been estimated that the number of population doubling is between 4 and 5, corresponding to $3.1 - 6.3 \times 10^9$ cell generations (A cell generation is the period of time between cell divisions. For example, if a single cell divides four times, there will be $1 + 2 + 4 + 8 + 16 = 31$ cell generations). These His- cells form a background lawn that is opaque to the eye, and the small individual microcolonies that comprise the lawn are distinguishable with a dissecting microscope (McGregor et al., 1984). The occurrence of a reverse mutation to histidine prototrophy results in a His+ colony which overgrows the background lawn, and is clearly visible even to the naked eye.

In addition to inducing mutations, the test chemical may also cause mutations. Irrespective of its mutagenicity, it can also kill parts or all of the target cells. Cell lethality may result in a reduction in the density of the His- background lawn or decreasing the number of revertant colonies due to death of His+ cells. Recently two methods to quantitative cell killing in the *Salmonella* test have been introduced (Waleh et al., 1982; McGregor et al., 1984).

The quantitative aspects of the bacterial test depend upon the chemical nature of the test compound and its active ability to induce a mutagenic and/or lethal response. Certain chemical substances may be quite stable and remain at its original concentration in the plate agar and exert their effects through the entire experiment. Others may be highly volatile and have an effect only in the initial stages of the

study. Test chemicals may also require metabolic activation by bacterial enzymes or exogenous mammalian enzymes added to the aqueous milieu in which the cells are grown. Metabolic activation is commonly achieved using a 900g homogenate (S9) of rat liver cells (Krewski et al., 1991).

2.2 A Stochastic Model for the Ames Assay

Margolin et al. (1981) developed a mechanistic dose response model for the Ames assay using fundamental principles which are believed to underlie the mutagenic and toxic processes governing colony formation. The specific assumptions involved in the development of this stochastic model are summarized as follows:

- 1) The lifespan of a microbe is defined as the interval between the time of its formation by the division of its parent cell and the time the microbe itself divides. The lifespan is assumed to be constant, and corresponds to a single generation of cells.
- 2) Sufficient histidine is provided for k generations of cells, with k typically in the range 1-6.
- 3) The test chemical is mutagenic for $m \leq k$ generations.
- 4) During the first m generations, a cell has a constant probability $p_m > 0$ of mutation. Mutations are assumed not to occur in subsequent generation during which the test agent is no longer effective. The values of p_m are usually small, on the order of 10^{-6} .
- 5) The chemical is toxic for $t \geq k$ generations. The probability that the chemical kills an individual cell is p_t in the first t generations and zero thereafter. The values of p_t are not necessarily small. The value of p_t is the same for both normal and mutant cells.
- 6) The values of p_m and p_t depend on the dose of the chemical being tested though t and m do not. For any particular system it is assumed that the values of t and m are known in advance.
- 7) Mutation and toxicity are independent of each other.

The above assumptions indicate that sufficient histidine is provided for k gener-

ations, with a constant period of time between cell division. The test chemical is mutagenic for m generations and cytotoxic for t generations. It is assumed that the values of m and t do not depend on the dose of the chemical being tested, and that mutagenic and toxic effects are independent.

Given the values $m = k$ and t , we let $P(m, t)$ denote the probability that a plated microbe will give rise to a visible colony of mutant microbes. Then the expected number of visible colonies is

$$\mu = N_0 P(m, t),$$

where N_0 is the number of initial cells being plated. The function $P(m, t)$ is in general expressed in terms of a complex difference equation in m and t . Its derivation is described below.

Consider a single plated microbe. The microbe or its descendants can mutate with probability p_m during the first m generations. The microbe or any of its descendants may die with probability p_t per generation during the first t generations. Let $Q(s)$ denote the probability that all the descendants of a particular microbe become extinct in s generations due to toxicity. To calculate $P(m, t)$, consider the following two pathways for the development of a viable revertant colony. First, the microbe itself can mutate and survive to divide, and not all its descendants become extinct. Second, the microbe may not mutate, but it does survive to divide with at least one of its two daughters giving rise to a revertant colony. For the daughter cells, mutation and toxicity persist for the next $m - 1$ and $t - 1$ generations, respectively. In the first event, the probability of mutation is p_m , the probability of survival is $1 - p_t$, and the probability that not all descendants of the two daughter cell become extinct in the next $t - 1$ generations is $1 - Q^2(t)$. Since these are independent, the probability of the event is the product of these terms. In the second event, the parent cell does not mutate with probability $1 - p_m$, and survives with probability $1 - p_t$; the probability that at least one of its offspring gives rise to a revertant colony is $1 - \{1 - P(m - 1, t - 1)\}^2$. These are also independent, so the probability of the second event is the product of three terms. This leads to the difference equation

$$P(m, t) = p_m(1-p_t)(1-Q^2(t-1)) + (1-p_m)(1-p_t)\{1-(1-P(m-1, t-1))^2\}. \quad (2.2.1)$$

The function $Q(s)$ satisfies the difference equation

$$Q(s) = p_t + (1-p_t)Q^2(s-1), \quad (2.2.2)$$

where $Q(0) = 0$, and $Q(\infty) = \min\{1, p_t/(1-p_t)\}$. Note that since $P(r, v)$ is very small for all $r \leq m$ and $v \leq t$, (2.2.1) may be approximated by

$$\begin{aligned} P(m, t) \approx & p_m(1-p_t)[1-Q^2(t-1)] \\ & + 2(1-p_t)(1-p_m)P(m-1, t-1). \end{aligned} \quad (2.2.3)$$

If $m = k$, then (2.2.3) admits the following solution:

$$\begin{aligned} P(m, t) = & p_m(1-p_t)\left\{\sum_{j=0}^{m-1} [2(1-p_m) \right. \\ & \left. \times (1-p_t)]^j \times [1-Q^2(t-1-j)]\right\}. \end{aligned} \quad (2.2.4)$$

2.3 A Histidine Diffusion Model

2.3.1 Model Formulation

In formula (2.2.1), it is assumed that sufficient histidine is provided to support up to k generations of cell division. This result is based on the assumption that histidine does not diffuse through the plate agar in which the bacteria grow, so that each cell has only a local supply of histidine which cannot be used by other cells. It may be more reasonable to assume that diffusion occurs, and that cell division will continue until all the histidine is exhausted, or until all cells die. Krewski et al. (1991) proposed a model which incorporates histidine diffusion. In their model, N_h denotes the maximum number of cell divisions that can be supported by the amount h of

histidine plated. To a good approximation, the expected number of mutant colonies is approximately

$$\begin{aligned}\mu &= N_h p_m (1 - p_t) (1 - Q^2(\infty)) \\ &= N_h p_m (1 - 2p_t)_+ / (1 - p_t),\end{aligned}\tag{2.3.1}$$

where $(1 - 2p_t)_+ = \max\{(1 - 2p_t), 0\}$, when the toxic and mutagenic effects persist throughout the experiment ($m = t = \infty$). This approximation ignores the event that all normal cells become extinct prior to depletion of the supply of histidine which is extremely unlikely. Note that (2.3.1) corresponds to (2.2.4) with long-term toxicity and sufficient histidine for only one generation ($m = k = 1, t = \infty$) (Krewski et al., 1993).

Suppose now that the test chemical exerts its toxic effects only for a short period of time, say during the first generation of cell divisions, and that there is sufficient histidine to support only one generation i.e. $N_h = N_1$. Regardless of the time period during which mutations induced by the test chemical occur, the expected number of revertant colonies is

$$\mu = N_1 p_m (1 - p_t).\tag{2.3.2}$$

Model (2.3.2) represents a slight generalization of the model (2.2.4) for $m = k = t = 1$ in that the constraint on the duration of mutagenic activity is avoided.

2.3.2 An Exact Solution for the Histidine Diffusion Model

In this section we develop an exact solution for the histidine diffusion model. The solution is developed for the general case in which m and k are positive intergers. Once plated, *Salmonella* undergo a birth-death-mutation process until the end of the experimental period. In the absence of death due to toxicity, each cell survives for a fixed period of time corresponding to the lifespan of a single generation. Mutation may occur at the time of auxotrophic cell division, leading to the production of one histidine dependent and one histidine independent daughter cell.

Suppose that there are $N_{(i)}$ cell divisions between the $(i-1)th$ and the ith generation. A microbe can mutate and survive to further divide, and not all of its descendants may become extinct. This occurs with probability $p_m(1-p_t)[1-Q^2(t-i)]$, so that the expected number of revertant colonies generated as a result of cell divisions occurring between the $(i-1)th$ and ith generations is

$$\mu_i = E[N_{(i)} - M_{(i)}]p_m(1-p_t)(1-Q^2(t-i)), \quad (2.3.3)$$

where $M_{(i)}$ denotes the number of mutant cells in the ith generation. Since p_m is small, we have

$$\mu_i \approx E[N_{(i)}]p_m(1-p_t)[1-Q^2(t-i)].$$

The expected number of revertant colonies generated by divisions occurring all generations is then

$$\begin{aligned} \mu &= \sum_{i=1}^{N_h} E[N_{(i)} - M_{(i)}]p_m(1-p_t)[1-Q^2(t-i)] \\ &\approx \sum_{i=1}^{N_h} EN_{(i)}p_m(1-p_t)[1-Q^2(t-i)], \end{aligned} \quad (2.3.4)$$

where $Q(s) = p_t + (1-p_t)Q^2(s-1)$, and $Q(0) = 0$. Notice that the term $Q(s)$ remains regardless of whether or not the supply of histidine is exhausted, since mutant cells do not need histidine for continued growth. We further note that in general, $N_h > E(\sum_{i=1}^{N_h} N_{(i)})$. When none of the cells die before the supply of histidine is exhausted, $N_h \equiv \sum_{i=1}^{N_h} N_{(i)}$. Provided p_t is not large, we expect $E(\sum_{i=1}^{N_h} N_{(i)}) \approx N_h$. (This approximation is investigated by computer simulation in section 2.5.).

A number of special cases can be developed from formula (2.3.4) depending on the value of m and t in relation to N_h . Specifically, we consider the following six cases.

(i) When $t \geq N_h$,

$$\mu_{(i)} = \begin{cases} [EN_{(i)} - M_{(i)}]p_m(1-p_t)(1-Q^2(t-i)) & \text{if } i \leq m \\ 0 & \text{otherwise.} \end{cases}$$

Therefore,

$$\mu = \sum_{i=1}^{\min(m, N_h)} E[N_{(i)} - M_{(i)}] p_m (1 - p_t) (1 - Q^2(t - i)).$$

(ii) When $m \leq t < N_h$, $p_t = p_m = 0$ and $Q(t - i) = 0$ if $i > t$. Therefore,

$$\mu = \sum_{i=1}^m E[N_{(i)} - M_{(i)}] p_m (1 - p_t) (1 - Q^2(t - i)).$$

(iii) When $t < m \leq N_h$,

$$\mu = \sum_{i=1}^t E[N_{(i)} - M_{(i)}] p_m (1 - p_t) (1 - Q^2(t - i)) + \sum_{i=t+1}^m (E N_{(i)}) p_m.$$

Note that $N_{t+1} = 2N_t$, $N_{t+2} = 2N_{t+1}, \dots$ since there are no toxic effects when $t + 1 \leq i \leq m$.

(iv) We claim that (2.3.4) reduces to the result (2.3.1) given by Krewski et al. (1993) when both $t = m = \infty$. Since $Q^2(t - i) = Q^2(\infty)$ for $t = \infty$,

$$\begin{aligned} \mu &= \sum_{i=1}^{\min(\infty, N_h)} E[N_{(i)} - M_{(i)}] p_m (1 - p_t) (1 - Q^2(t - i)) \\ &= \sum_{i=1}^{\min(\infty, N_h)} E[N_{(i)} - M_{(i)}] p_m (1 - p_t) (1 - Q^2(\infty)) \\ &\approx E\left(\sum_{i=1}^{N_h} N_{(i)}\right) p_m (1 - p_t) (1 - Q^2(\infty)) \\ &= N_h p_m (1 - p_t) (1 - Q^2(\infty)) \end{aligned}$$

for moderate p_t , corresponding to (2.3.1).

(v) We now show the relationship between (2.3.4) and (2.2.4) as established by Margolin et al. (1981). If there is sufficient histidine to maintain k generations of cell

deviations, as assumed in (2.2.4), then it is easy to see that $EN_{(i)} = N_0[2(1 - p_t)]^{i-1}$. Substituting this into formula (2.3.4) yields

$$\begin{aligned}\mu &= \sum_{i=1}^k N_0[2(1 - p_t)]^i p_m (1 - p_t) (1 - Q^2(t - i)) \\ &= N_0 p_m (1 - p_t) \sum_{i=1}^k [2(1 - p_t)]^{i-1} [1 - Q^2(t - i)].\end{aligned}$$

Since $p_m \approx 10^{-6}$ is much less than 1, the above expression is identical to (2.2.4), when the term $(1 - p_m)$ in (2.2.4) is replaced by 1 (cf. Margolin et al. (1981), equation [11]). In the remainder of this section, we justify the approximation $N_{(i)} - M_{(i)} \approx N_{(i)}$, provided p_m is small.

In the first generation, the expected number of surviving cell is $S_1 = N_0[2(1 - p_t)]$, and the expected number of mutant cells is $M_1 = S_1 p_m$. In the second generation, the expected number of surviving cells is

$$S_2 = S_1[2(1 - p_t)] = N_0[2(1 - p_t)]^2$$

and the expected number of mutant cells is

$$M_{(2)} = M_{(1)}[2(1 - p_t)] + (S_1 - M_{(1)})[2(1 - p_t)]p_m,$$

where the first term on the right hand side is the expected number of surviving mutant cells from the last generation and the second term represents the expected number of newly mutated cells from the normal cells of previous generation. Similarly, the expected number of normal cells and mutant cells in the i th generation are given by

$$S_i = S_{i-1}[2(1 - p_t)] = N_0[2(1 - p_t)]^i$$

and

$$M_{(n)} = M_{(n-1)}[2(1 - p_t)] + (S_{n-1} - M_{(n-1)})[2(1 - p_t)]p_m$$

respectively. When p_m is much less than 1,

$$\begin{aligned}M_{(n)} &= S_{n-1}[2(1 - p_t)]p_m + M_{(n-1)}[2(1 - p_t)](1 - p_m) \\ &\approx S_{n-1}[2(1 - p_t)]p_m + M_{(n-1)}[2(1 - p_t)] \\ &= S_1[2(1 - p_t)]^{n-1} p_m.\end{aligned}$$

Therefore,

$$\begin{aligned} N_{(n)} - M_{(n)} &\approx S_n - M_{(n)} \\ &= S_1[2(1 - p_t)]^{n-1}(1 - np_m). \end{aligned}$$

Since np_m is also much less than one ($p_m \approx 10^{-6}$), we obtain

$$N_{(n)} - M_{(n)} \approx S_1[2(1 - p_t)]^{n-1} = S_n = N_{(n)},$$

which is what we want to show.

2.3.3 Simulation Results

In this section, the relationship among (2.3.4), (2.3.1), and (2.3.2) is explored through Monte-Carlo simulation. The purpose of this simulation is to evaluate the accuracy of the approximation $E(\sum_{i=1}^{N_h} N_{(i)}) \approx N_h$, and compare the predicted expected number of colonies by using approximations (2.3.1), (2.3.2), and the exact solution (2.3.4).

Given the value of p_t (the probability of cell death due to toxicity), N_0 (the initial number of cells plated) and N_h (the total number of cell divisions supported by the amount h of histidine plated), we may follow the three steps below to simulate the process of cell division.

Step I: Generate N_0 pseudo-random numbers from a standard uniform distribution $U(0,1)$ to obtain the number n_1 of values falling into the interval $(0, (1 - p_t))$. We then define $N_{(1)} = 2n_1$.

Step II: Generate $N_{(1)}$ pseudo-random numbers from $U(0,1)$, and determine the number n_2 of values falling into the interval $(0, (1 - p_t))$. Then, we let $N_{(2)} = 2n_2$.

Step III: Generate $N_{(2)}$ pseudo-random number from $U(0,1)$, and repeat Step II to obtain a sequence $N_{(1)}, N_{(2)}, N_{(3)}, \dots$ stopping with $\sum_{i=1}^{N_h} N_{(i)} \geq N_h$.

To obtain the values of $EN_{(i)}$, we may repeat Steps I-III K times to obtain K sequences $\{N_{(1)}^{(k)}, N_{(2)}^{(k)}, \dots\}_{k=1}^K$. We may then estimate $E(N_{(i)})$ by $K^{-1} \sum_{k=1}^K N_{(i)}^{(k)}$.

In the present simulation, we set $N_0 = 10^8$, $p_m = 10^{-6}$ and $N_h = 60 \times N_0$. These values are typical of those encountered in Ames test (Broekhoven & Nestmann, 1991, p. 207). A range of possible values of $p_t = 0.1, 0.2, 0.4$ are also considered.

This procedure is used to estimate the ratio

$$R_0 = \sum_{i=1}^{N_h} E(N_{(i)})/N_h,$$

which measures the accuracy of the approximation $E(\sum_{i=1}^{N_h}) \approx N_h$. The accuracy of (2.3.1) as an approximation to (2.3.4) is evaluated by the ratio

$$\begin{aligned} R_1 &= \frac{\sum E(N_i)p_m(1-p_t)(1-Q^2(t-i))}{N_h p_m(1-p_t)(1-Q^2(\infty))} \\ &= \frac{\sum E N_{(i)}(1-Q^2(t-i))}{N_h(1-Q^2(\infty))}. \end{aligned}$$

We also calculate the ratio, R_2 , of (2.3.4) to (2.3.2):

$$\begin{aligned} R_2 &= \frac{\sum E(N_i)p_m(1-p_t)(1-Q^2(t-i))}{N_h p_m(1-p_t)} \\ &= \frac{\sum E N_{(i)}(1-Q^2(t-i))}{N_h}. \end{aligned}$$

Recall that (2.3.2) is obtained from (2.3.1) when $t = m = 1$, and corresponds to the linear-exponential dose-response model which will be used in subsequent chapters.

Since $Q(\infty) = 1$ for $p_t \geq 0.5$, the denominator of R_2 is zero in this case. Consequently, we restrict our attention to the case $p_t < 0.5$. Estimates of R_0 , R_1 , and R_2 based on $K = 1000$ simulation are given in Table 2.1. When $p_t < 0.4$, R_1 is close to 1; however for $0.4 \leq p_t < 0.5$, R_1 exceeds 2. Hence, (2.3.1) is a good approximation when $p_t < 0.4$. Both (2.3.4) and (2.3.1) yield close approximations to the expected number of revertant colonies. R_2 is closer to unity compared to R_1 , indicating that (2.3.2) is a better approximation to (2.3.4).

Table 2.1 Ratios of R_0 , R_1 and R_2 Based on Computer Simulation

p_t	R_0	R_1	R_2
0.1	1.00	1.00	0.988
0.2	1.00	1.05	0.984
0.4	0.99	2.34	1.30

2.4 Summary

In this chapter, the Ames assay is modeled from biological principles when histidine diffusion exists in the test plate agar. First, two models developed by Margolin et al. (1981) and Krewski et al (1993) are reviewed. Then a model considering the occurrence of histidine diffusion in the test plate agars is proposed, under more general conditions. It is also shown that the proposed model can be reduced to exactly the models developed both by Margolin et al. (1981) and by Krewski et al. (1993) when the same restrictions of their models are imposed on the proposed model. In this sense, the proposed model is a generalization of the two previous models. Finally, the simulation results indicate that the proposed model and Krewski et al. (1993) model predict almost the same expected number of revertant colonies when the toxic effect is not very strong.

Chapter 3

Tests for Trend with Historical Controls

3.1 Introduction

The need to test for trend in response arises in many areas of biomedical research. Studies of the carcinogenicity of chemicals or other substances in recent bioassays constitute an area of particular importance (Leroux et al., 1994). In this chapter, we develop score-type statistics to test for trend using an estimating equation approach (Breslow, 1990), taking into account any available information on historical control response rates. Historical control data contains useful information about the spontaneous response rate, and can be used to enhance the power of tests for trend (Krewski et al., 1991; Fung et al., 1994; Hoel & Yanagawa, 1986).

Leroux et al. (1994) have previously considered tests for trend in count data using historical control data, but did not distinguish between within-laboratory and between-laboratory sources of extra-Poisson variation. Since within-laboratory variability and between-laboratory variability may be quite different in magnitude, it is of interest to use different parameters to model overdispersion within and between studies (Krewski et al., 1992; Knuiman et al., 1987).

In section 3.2, we introduce a moment structure for the observations in the experiment at hand. Since an estimating equation approach to inference is developed, only the first two moments of the data require specification. The expected value of the observations is assumed to be a linear function of the dose levels; the variance is a quadratic function of the expected value of the observations, which contains

the overdispersion parameter. Two variations on this theme are presented, both of which allow for extra-Poisson variation within the current experiment, and an additional overdispersion parameter to allow for inter-study variation among the historical controls.

In section 3.3, an estimating equation approach is used to estimate the unknown regression parameters as well as the overdispersion parameters. Numerical methods for solving these equations are discussed. In order to test for trend, a generalized score-type statistic is obtained from one of these estimating equations.

In section 3.4, the asymptotic properties of parameter estimates and the score statistic are discussed. First, following a similar method employed by Moore (1985, 1986), we show that the estimators of the unknown regression and overdispersion parameters are consistent. The inverse function theorem plays a key role in the proof of the consistency. Then, we use Taylor expansions and the Linderbeg central limit theorem to show that the score statistic given in section 3.3 is asymptotically normally distributed. The theoretical arguments in this section are based on the ideas of Moore (1985, 1986) and Foutz (1977).

In section 3.5, a computable formula for the standardized score statistic is derived. The variance of the score statistic is also obtained. In section 3.6, a small computer simulation study is used to examine the type I and type II error rates of the score statistic used to test for trend. This limited study suggests that the use of historical data will lead to increased power. The main results of this chapter are summarized in section 3.7.

3.2 Mean-Variance Function

Let Y_{1t} denote the number of revertant colonies in plate $t = 1, \dots, n_1$, at dose d_1 , and let $Y_{1i} = \sum_{t=1}^{n_1} Y_{1t}$ denote the total number of revertant colonies across the n_1 plates at dose d_i ($i = 1, \dots, K$).

Method 1

Following Leroux et al. (1994), we assume

$$\left. \begin{aligned} EY_{1,t} &= \mu_t = \beta_0 + \beta_1 d_t \\ \text{Var}(Y_{1,t}) &= (\beta_0 + \beta_1 d_t)[1 + (\beta_0 + \beta_1 d_t)\tau_1] \end{aligned} \right\}. \quad (3.2.1)$$

with $\text{Cov}(Y_{1,t}, Y_{1,t'}) = 0$ for $t \neq t'$. The moment assumptions are based on a classical approach which treats the Poisson means associated with each observed count as latent variables that are sampled from a gamma distribution. The gamma mixing distribution leads to a negative binomial distribution for the observed data (Margolin et al., 1981). This moment structure applies to observations $\{Y_{1,t}\}$ within the experiment of current interest.

For the historical control counts $Y_{j,t}$ ($j = 2, \dots, s; t = 1, \dots, n_j$), we assume

$$\left. \begin{aligned} E(Y_{j,t}|\theta_j) &= \theta_j \\ E\{(Y_{j,t} - \theta_j)^2|\theta_j\} &= \theta_j(1 + \theta_j\tau_1) \end{aligned} \right\}, \quad (3.2.2)$$

where the θ_j are independent random variable reflecting random differences among the historical experiments $j = 2, \dots, s$. We further assume

$$\left. \begin{aligned} E\theta_j &= \beta_0 \\ \text{Var}(\theta_j) &= \beta_0^2\tau_2 \end{aligned} \right\}, \quad (3.2.3)$$

Note that the distribution of θ_j 's is centered on the expected response in the concurrent control group. It follows that

$$\left. \begin{aligned} EY_{j,t} &= \beta_0 \\ \text{Var}Y_{j,t} &= \beta_0\{1 + \beta_0(\tau_1 + \tau_2 + \tau_1\tau_2)\} \end{aligned} \right\}, \quad (3.2.4)$$

with $\text{Cov}(Y_{j,t}, Y_{j,t'}) = \beta_0^2\tau_2$, and $\text{Cov}(Y_{j,t}, Y_{j',t'}) = 0$ for $t \neq t'$. The moments of $Y_j = \sum_{t=1}^{n_j} Y_{j,t}$ are then given by

$$\left. \begin{aligned} E(Y_j) &= E(\sum_{t=1}^{n_j} Y_{j,t}) = n_j\beta_0 \\ \text{Var}(Y_j) &= \text{Var}(\sum_{t=1}^{n_j} Y_{j,t}) = n_j\beta_0\{1 + \beta_0(\tau_1 + \tau_1\tau_2 + n_j\tau_2)\} \end{aligned} \right\}. \quad (3.2.5)$$

This formulation differs from that considered by Leroux et al. (1994) in that overdispersion among the concurrent controls (reflected by the parameter τ_1 in (3.2.1)) will be less than the overdispersion among the historical controls (reflected by the term

$(1 + \tau_1 + \tau_2 + \tau_1 \tau_2)$ in (3.2.4)). This reflects the fact that variation among the historical controls includes inter-study variation, which is not present in the concurrent control group.

In the next section, we will use the mean and variance structure assumed in (3.2.1) and (3.2.4) to construct estimating equations on which the generalized score-type test for trend is based. Before proceeding, however, we discuss an alternative moment structure.

Method 2

For the observations Y_{1it} in the current experiment, we assume

$$\left. \begin{aligned} EY_{1it} &= \mu_{1i} = \beta_0 + \beta_1 d_i \\ \text{Var}(Y_{1it}) &= \mu_{1i}(1 + \tau_1) \end{aligned} \right\}. \quad (3.2.6)$$

Similarly, for the historical control observations

$$\left. \begin{aligned} E(Y_{jt}|\theta_j) &= \theta_j \\ \text{Var}(Y_{jt}|\theta_j) &= \theta_j(1 + \tau_1) \end{aligned} \right\}. \quad (3.2.7)$$

In fact, if we assume that the data come from a Poisson/gamma mixture, then the moment structures of the data will be exactly the same as that given by (3.2.6) or (3.2.7). Specifically, if the Poisson mean is a random variable having a two-parameter gamma distribution, and we assume the shape parameter is held to be constant but allow the scale parameter to vary as the mean of the gamma varies with dose, we obtain the moment structure (3.2.6). If the scale parameter is held to be constant but the shape parameter varies, then we obtain (3.2.1) (Krewski et al., 1993).

Assuming that the historical control expected mutation rate follows a distribution with moments

$$\left. \begin{aligned} E\theta_j &= \beta_0 \\ \text{Var}(\theta_j) &= \beta_0 \tau_2 \end{aligned} \right\}, \quad (3.2.8)$$

we can obtain the unconditional moments.

$$\left. \begin{aligned} EY_{jt} &= \beta_0 \\ \text{Var}(Y_{jt}) &= \beta_0(1 + \tau_1 + \tau_2) \\ \text{Cov}(Y_{jt}, Y_{jt'}) &= \beta_0^2 \tau_2 \end{aligned} \right\}. \quad (3.2.9)$$

Letting $Y_j = \sum_{i=1}^{n_j} Y_{ji}$, we have

$$\left. \begin{aligned} E(Y_j|\theta_j) &= n_j\theta_j, \\ \text{Var}(Y_j|\theta_j) &= n_j\theta_j(1 + \tau_1) \end{aligned} \right\}. \quad (3.2.10)$$

Unconditionally,

$$\left. \begin{aligned} EY_j &= n_j\beta_0 \\ \text{Var}Y_j &= n_j\beta_0(1 + \tau_1 + n_j\tau_2) \\ \text{Cov}(Y_{ji}, Y_{j'i'}) &= \beta_0^2\tau_2 \end{aligned} \right\}. \quad (3.2.11)$$

3.3 Generalized Score Tests

In this section, estimating equations are used to develop generalized score statistics to test for trend in response with increasing dose in the current experiment, taking into account the historical control information. Initially, we will assume the moment structure (3.2.1)-(3.2.4) given in section 3.2.

Let $Y_{1i} = \sum_{j=1}^{n_{1i}} Y_{1ij}$ ($i = 1, 2, \dots, k$), $Y_j = \sum_{i=1}^{n_j} Y_{ji}$, and

$$\mathbf{Y} = (Y_{111}, \dots, Y_{11n_{11}}, Y_{121}, \dots, Y_{12n_{12}}, \dots, Y_{1K1}, \dots, Y_{1Kn_{1K}}, Y_2, Y_3, \dots, Y_s)^T.$$

We then have

$$E\mathbf{Y} = \boldsymbol{\mu}(\boldsymbol{\beta})$$

where $\boldsymbol{\beta} = (\beta_0, \beta_1)^T$.

An estimate of $\boldsymbol{\beta}$ can be obtained by solving the estimating equation:

$$\mathbf{D}^T \mathbf{V}^{-1}(\mathbf{Y} - \boldsymbol{\mu}(\boldsymbol{\beta})) = 0, \quad (3.3.1)$$

where $\mathbf{D}^T = d\boldsymbol{\mu}/d\boldsymbol{\beta}$. More precisely, (3.3.1) can be written as

$$\mathbf{D}_1^T \mathbf{V}_1^{-1}(\mathbf{Y}_1 - \boldsymbol{\mu}_1) + \sum_{j=2}^s \mathbf{D}_j^T \mathbf{V}_j^{-1}(\mathbf{Y}_j - \boldsymbol{\mu}_j) = 0, \quad (3.3.2)$$

where $\mathbf{Y}_1 = (Y_{10}, Y_{11}, \dots, Y_{1k})^T$, $\mathbf{Y}_j = (Y_{j1}, \dots, Y_{jn_j})^T$, $\boldsymbol{\mu}_j = (\beta_0, \dots, \beta_0)^T$, $\boldsymbol{\mu}_1 = (n_{10}(\beta_0 + \beta_1 d_0), \dots, n_{1k}(\beta_0 + \beta_1 d_k))^T$,

$$\mathbf{D}_1^T = \begin{bmatrix} \partial \boldsymbol{\mu}_1 / \partial \beta_0 \\ \partial \boldsymbol{\mu}_1 / \partial \beta_1 \end{bmatrix} = \begin{bmatrix} n_{10} & n_{11} & \dots & n_{1k} \\ 0 & n_{11}d_1 & \dots & n_{1k}d_k \end{bmatrix},$$

and

$$\mathbf{V}_1^{-1} = \text{diag}\{n_1, (\beta_0 + \beta_1 d_1)(1 + \tau_1(\beta_0 + \beta_1 d_1))\}^{-1}.$$

To complete the specification of (3.3.2), we have

$$\mathbf{D}_j^T = \begin{bmatrix} 1, & 1, & \dots & 1 \\ 0, & 0, & \dots & 0 \end{bmatrix}_{2 \times n_j},$$

and

$$\begin{aligned} \mathbf{V}_j &= \beta_0\{1 + \beta_0(\tau_1 + \tau_1\tau_2)\}\mathbf{I}_{n, x_{n_j}} + \beta_0^2\tau_2\mathbf{J}_{n, x_{n_j}}, \\ &= a\mathbf{I}_{n, x_{n_j}} + b\mathbf{J}_{n, x_{n_j}}, \end{aligned}$$

with $a = \beta_0\{1 + \beta_0(\tau_1 + \tau_1\tau_2)\}$ and $b = \beta_0^2\tau_2$, where $\mathbf{I}_{n, x_{n_j}}$ is the identity matrix, and $\mathbf{J}_{n, x_{n_j}}$ is a matrix with all elements equal to unity. Therefore,

$$\mathbf{V}_j^{-1} = \{a^{-1}\mathbf{I}_{n, x_{n_j}} + n_j^{-1}(\frac{1}{a + n_j b} - \frac{1}{a})\mathbf{J}_{n, x_{n_j}}\}.$$

With some algebra, it can be shown that

$$\mathbf{D}_j^T \mathbf{V}_j^{-1} (\mathbf{Y}_j - \mu_j) = [\frac{1}{a + b n_j} (Y_j - n_j \beta_0), 0]^T.$$

Substituting the above results into the left hand side of (3.3.2) (under $\beta_1 = 0$), and assuming $\mathbf{D}^T \mathbf{V}^{-1} (\mathbf{Y} - \mu) = (U_1, U_2)^T$, we can write

$$U_1(\beta_0, \tau_1, \tau_2) = \frac{1}{\beta_0(1 + \beta_0\tau_1)} \sum_{i=1}^k \sum_{t=1}^{n_{1i}} (Y_{1it} - \beta_0) + \sum_{j=2}^s \frac{Y_j - n_j \beta_0}{\beta_0(1 + \beta_0(\tau_1 + n_j \tau_2 + \tau_1 \tau_2))}$$

and

$$U_2(\beta_0, \tau_1, \tau_2) = \frac{1}{\beta_0(1 + \beta_0\tau_1)} \sum_{i=1}^k \sum_{t=1}^{n_{1i}} (Y_{1it} - \beta_0) d_{1i}.$$

Under $H_0 : \beta_1 = 0$, the estimating equation

$$f_1(\theta) \equiv U_1(\beta_0, \tau_1, \tau_2) = 0 \quad (3.3.3)$$

provides the basis for estimating the regression parameter β ; $U_2(\beta_0, \tau_1, \tau_2)$ is the score statistic used to test for trend. Under the null hypothesis $H_0 : \beta_1 = 0$, we can

set $U_1(\beta_0, \tau_1, \tau_2) = 0$ when τ_1 and τ_2 are known to find the GEE estimate $\hat{\beta}_0$ of β_0 and use $U_2(\hat{\beta}_0, \tau_1, \tau_2)$ as the test for trend (Breslow, 1990). We can also use the two moment equations (3.3.4) and (3.3.5) below to estimate τ_1 and τ_2 .

Let $\theta_0 = (\beta_0, \tau_{10}, \tau_{20})^T$ denote the true values of the model parameters under H_0 . The following two moment equations can be used to estimate the overdispersion parameters τ_1 and τ_2 (Breslow, 1990):

$$f_2(\theta) = \sum_{i=1}^k \sum_{t=1}^{n_{1i}} \frac{(Y_{1it} - \beta)^2}{\beta(1 + \beta\tau_1)} - \sum_{i=1}^k n_{1i} = 0 \quad (3.3.4)$$

and

$$f_3(\theta) = \sum_{j=2}^s \frac{(Y_j - n_j\beta)^2}{n_j\beta(1 + \beta(\tau_1 + \tau_1\tau_2 + n_j\tau_2))} - (s - 1) = 0. \quad (3.3.5)$$

Solving the equations $f_1 = 0$, $f_2 = 0$ and $f_3 = 0$ simultaneously yields the estimated values $\hat{\beta}_0$, $\hat{\tau}_1$ and $\hat{\tau}_2$ of β_0 , τ_{10} and τ_{20} respectively.

The generalized score-type statistic is then given by

$$T = U_2(\hat{\beta}_0, \hat{\tau}_1, \hat{\tau}_2) = \frac{1}{\hat{\beta}_0(1 + \hat{\beta}_0\hat{\tau}_1)} \sum_{i=1}^k \sum_{t=1}^{n_{1i}} (Y_{1it} - \hat{\beta}_0)d_i. \quad (3.3.6)$$

In the next section, it will be shown that $T/\sqrt{\text{Var}(T)}$ is asymptotically normally distributed.

Under the null hypothesis the estimated values of $\beta_0, \tau_{10}, \tau_{20}$ can be obtained by solving equations (3.3.3)–(3.3.5) in an iterative fashion. Letting $\tau_2^* = \tau_1 + n_j\tau_2 + \tau_1\tau_2$, it follows from (3.3.3) that

$$(1 + \beta_{20}\tau_{20}^*)Y_1 - \beta_0(1 + \beta_0\tau_{20}^*)n_1 + (1 + \beta_0\tau_{10})(\sum_{j=2}^s Y_j) - \beta_0(1 + \beta_0\tau_{10})n_2 = 0,$$

where $n_1 = \sum_{i=1}^k n_{1i}$ and $n_2 = \sum_{j=2}^s n_j$. This can be re-expressed as the quadratic form

$$(n_1\tau_{20} + \tau_{10}n_2)\beta_0^2 + [n_1 + n_2 - Y_2\tau_{10} - Y_1\tau_{20}^*]\beta_0 - (Y_1 + Y_2) = 0,$$

where $Y_1 = \sum_{i=1}^k Y_{1i}$ and $Y_2 = \sum_{j=2}^s Y_j$. The solution to this quadratic form is

$$\beta_0 = \frac{[Y_1\tau_{20}^* + Y_2\tau_{10} - (n_1 + n_2)] + \sqrt{(Y_1\tau_{20}^* + Y_2\tau_{10} - n_1 - n_2)^2 + P}}{2[n_1\tau_{20}^* + n_2\tau_{10}]}, \quad (3.3.7)$$

where $P = 4(n_1\tau_{20}^* + n_2\tau_{10})(Y_1 + Y_2)$.

Using the moment equation (3.3.4), we have

$$(K + 1 - 2)(1 + \beta_0\tau_{10}) = \sum_{i=0}^k \frac{(Y_{1i} - n_{1i}\beta_0)^2}{n_{1i}\beta_0},$$

which yields the estimator

$$\hat{\tau}_{10} = \left[\sum_{i=0}^k \frac{(Y_{1i} - n_{1i}\hat{\beta}_0)^2}{n_{1i}\hat{\beta}_0} - (K + 1 - 2) \right] / [(K + 1 - 2)\hat{\beta}_0]. \quad (3.3.8)$$

Similarly, (3.3.5) is written as

$$(S - 1 - 2)(1 + \beta_0\tau_{20}^*) = \sum_{j=2}^s \frac{(Y_j - n_j\beta_0)^2}{n_j\beta_0}$$

which yields

$$\hat{\tau}_{20} = \left[\sum_{j=2}^s \frac{(Y_j - n_j\hat{\beta}_0)^2}{n_j\hat{\beta}_0} - (S - 2 + 1) \right] / [(S - 2 + 1)\hat{\beta}_0]. \quad (3.3.9)$$

The final estimates of β_0 , τ_{10} , and τ_{20} are then obtained by solving the three equations simultaneously in an iterative fashion. Specifically, the iterative procedure is implemented as follows: (1) choose initial values $\hat{\beta}_0$; (2) insert $\hat{\beta}_0$ into (3.3.8) to get $\hat{\tau}_{10}$; (3) insert $\hat{\beta}_0$ into (3.3.9) to get $\hat{\tau}_{20}$; (4) substitute $\hat{\tau}_{10}$ and $\hat{\tau}_{20}$ obtained from steps (2) and (3) into (3.3.7) to obtain $\hat{\beta}_0$. Steps (2)-(4) are repeated until convergence. We use $\hat{\theta}_m = (\hat{\beta}_0, \hat{\tau}_1, \hat{\tau}_2)^T$ to denote the estimated parameter values at m times of repeating steps (2)-(4). Convergence is achieved when $\|(\theta_m - \theta_{m-1})/\theta_{m-1}\| \leq \epsilon$, where the norm $\|\cdot\|$ is defined as the maximal relative errors of the three corresponding components between vector θ_{m-1} and θ_m .

3.4 Asymptotic Distribution

In this section, we will show that under the null hypothesis $\beta_1 = 0$, the estimators of the parameters β_0 , τ_{10} , and τ_{20} obtained from equations (3.4.2)-(3.4.5) are consistent as n_1 and $s \rightarrow \infty$ with $n_1/s \rightarrow c > 0$. Methods, similar to those employed by Moore (1985, 1986) are used to show the consistence of the parameter estimates. The use

of inverse function theorem plays a key role in the proof of the consistence, which was first proposed by Foutz (1977). In this section, some modifications of Moore's method are made in order to take into account of historical controls. We will also show that the asymptotic distribution of the standardized score-type statistic given in (3.4.6) is normal.

Define

$$\nu_1(\beta, \tau_1) = [\beta(1 + \beta\tau_1)]^{-1}$$

and

$$\nu_2(\beta, \tau_1, \tau_2, n_j) = [\beta(1 + \beta(\tau_1 + n_j\tau_2 + \tau_1\tau_2))]^{-1}.$$

The following assumptions (C1)-(C7) will be used in the reminder of this section.

- (C1) n_1 and $s \rightarrow \infty$ with $n_1/s \rightarrow c > 0$, where $n_1 = \sum_{i=1}^k n_{1i}$.
- (C2) $\sum_{i=1}^k n_{1i}d_i/n_1 \rightarrow \bar{d} > 0$, as $n_1 \rightarrow \infty$.
- (C3) n_j ($j = 2, \dots, s$) are uniformly bounded with $\max_{2 \leq j \leq s} \{n_j\} \leq N$, $2 \leq s \leq \infty$.
- (C4) For all $(\beta, \tau_1, \tau_2)^T$ in a neighborhood U_δ of the true parameters $(\beta_0, \tau_{10}, \tau_{20})^T$.
 $|\nu_1(\beta, \tau_1)| < M$, $|\nu_2(\beta, \tau_1, \tau_2, n_j)| < M$, $j = 2, 3, \dots, s$, $2 \leq s \leq \infty$.
- (C5) $\sup_{\theta \in U_\delta} \left| \frac{\partial \nu_l}{\partial \theta} \right| \leq M < \infty$, for all j , where $\theta = (\beta, \tau_1, \tau_2)^T$. $l = 1, 2$.
- (C6) For all i, j, t , $|E(Y_{1it} - \beta_0)^4| \leq M$, $|E(Y_j - n_j\beta_0)^4| \leq M$.
- (C7) $\lim_{K^* \rightarrow \infty} \frac{1}{K^*} \text{Cov}(f(\theta_0)) \rightarrow \Sigma_0$, where $K^* = \sum_{i=1}^k n_{1i} + (s-1)$.

C1 indicates that the number of historical controls increases at the same rate of total number of current observations. C2 requires that the average dose converges to a constant. C3 states that the number of observations within each historical control is uniformly bounded and remains so as the number of historical controls goes to infinity. C4 holds in the present case since, β , τ_1 , τ_2 and n_j are non-negative, bounded, and β is bounded away from zero. To show that C5 holds, we note that

$$\begin{aligned} \frac{\partial \nu_1(\beta, \tau_1)}{\partial \beta} &= (-1)[\beta(1 + \beta\tau_1)]^{-2}(1 + 2\beta\tau_1), \\ \frac{\partial \nu_1(\beta, \tau_1)}{\partial \tau_1} &= (-1)[\beta(1 + \beta\tau_1)]^{-2}\beta^2, \end{aligned}$$

$$\frac{\partial \nu_2(\beta, \tau_1, \tau_2, n_j)}{\partial \beta} = (-1)[\beta(1 + \beta(\tau_1 + n_j\tau_2 + \tau_1\tau_2))]^{-2}[1 + \tau_1 + n_j\tau_2 + \tau_1\tau_2],$$

$$\frac{\partial \nu_2(\beta, \tau_1, \tau_2, n_j)}{\partial \tau_1} = (-1)[\beta(1 + \beta(\tau_1 + n_j\tau_2 + \tau_1\tau_2))]^{-2}[\beta^2 + \beta^2\tau_2],$$

and

$$\frac{\partial \nu_2(\beta, \tau_1, \tau_2, n_j)}{\partial \tau_2} = (-1)[\beta(1 + \beta(\tau_1 + n_j\tau_2 + \tau_1\tau_2))]^{-2}[n_j\beta^2 + \beta^2\tau_1].$$

Since, the parameter β is positive and bounded away from zero, and $0 < \tau_i < \infty$ ($i = 1, 2$), the partial derivatives of $\nu_1(\beta, \tau_1)$ and $\nu_2(\beta, \tau_1, \tau_2, n_j)$ with respect to β , τ_1 and τ_2 are uniformly bounded in a neighbourhood U_δ of the true parameter values β_0 , τ_{10} and τ_{20} . C6 asserts the forth moment of both current and historical control observations are bounded, a reasonable assumption for observation that are subject to a natural upper bound. C7 is required in the asymptotic normality proof, and is similar to condition C4 of Krewski & Rao (1981).

Let $\mathbf{f}(\theta_0) = (f_1(\theta_0), f_2(\theta_0), f_3(\theta_0))^T$, where functions f_1 , f_2 , and f_3 are given by (3.3.3), (3.3.4), and (3.3.5) respectively.

Using the notations $\nu_1(\beta, \tau_1)$ and $\nu_2(\beta, \tau_1, \tau_2, n_j)$, we may write

$$f_1(\theta) = \sum_{i=1}^k \sum_{t=1}^{n_{1i}} (Y_{1it} - \beta) \nu_1(\beta, \tau_1) + \sum_{j=2}^s (Y_j - n_j \beta) \nu_2(\beta, \tau_1, \tau_2, n_j),$$

$$f_2(\theta) = \sum_{i=1}^k \sum_{t=1}^{n_{1i}} (Y_{1it} - \beta)^2 \nu_1(\beta, \tau_1) - \sum_{i=1}^k n_{1i},$$

and

$$f_3(\theta) = \sum_{j=2}^s (Y_j - n_j \beta)^2 n_j^{-1} \nu_2(\beta, \tau_1, \tau_2, n_j) - (s - 1),$$

where $\theta = (\beta, \tau_1, \tau_2)^T$. The partial derivatives of the functions f_r ($r = 1, 2, 3$) are obtained below by direct calculation.

$$\frac{\partial f_1}{\partial \beta} = \sum_{i=1}^k \sum_{t=1}^{n_{1i}} [(Y_{1it} - \beta) \frac{\partial \nu_1}{\partial \beta} + (-1) \nu_1(\beta, \tau_1)] + \sum_{j=2}^s [(Y_j - n_j \beta) \frac{\partial \nu_2}{\partial \beta} + (-n_j) \nu_2(\beta, \tau_1, \tau_2, n_j)].$$

$$\frac{\partial f_1}{\partial \tau_1} = \sum_{i=1}^k \sum_{t=1}^{n_{1i}} [(Y_{1it} - \beta) \frac{\partial \nu_1}{\partial \tau_1}] + \sum_{j=2}^s (Y_j - n_j \beta) \frac{\partial \nu_2}{\partial \tau_1}.$$

$$\begin{aligned}
\frac{\partial f_1}{\partial \tau_2} &= \sum_{j=2}^{\cdot} (Y_j - n_j \beta) \frac{\partial \nu_2}{\partial \tau_2}. \\
\frac{\partial f_2}{\partial \beta} &= \sum_{i=1}^k \sum_{t=1}^{n_{1i}} [2(Y_{1it} - \beta)(-1)\nu_1(\beta, \tau_1) + (Y_{1it} - \beta)^2 \frac{\partial \nu_1}{\partial \beta}]. \\
\frac{\partial f_2}{\partial \tau_1} &= \sum_{i=1}^k \sum_{t=1}^{n_{1i}} [(Y_{1it} - \beta)^2 \frac{\partial \nu_1}{\partial \tau_1}]. \\
\frac{\partial f_2}{\partial \tau_2} &= 0. \\
\frac{\partial f_3}{\partial \beta} &= \sum_{j=2}^{\cdot} [2(Y_j - n_j \beta)(-n_j)n_j^{-1}\nu_2(\beta, \tau_1, \tau_2, n_j) + (Y_j - n_j \beta)^2 n_j^{-1} \frac{\partial \nu_2}{\partial \beta}]. \\
\frac{\partial f_3}{\partial \tau_1} &= \sum_{j=2}^{\cdot} (Y_j - n_j \beta)^2 n_j^{-1} \frac{\partial \nu_2}{\partial \tau_1}. \\
\frac{\partial f_3}{\partial \tau_2} &= \sum_{j=2}^{\cdot} (Y_j - n_j \beta)^2 n_j^{-1} \frac{\partial \nu_2}{\partial \tau_2}. \tag{3.4.1}
\end{aligned}$$

Define

$$G(\theta) = \frac{1}{n_1} \mathbf{f}'(\theta) = \frac{1}{n_1} (\mathbf{f}'_1, \mathbf{f}'_2, \mathbf{f}'_3),$$

where $\mathbf{f}'_r = [\partial f_r / \partial \beta, \partial f_r / \partial \tau_1, \partial f_r / \partial \tau_2]$ for $r = 1, 2, 3$, and let $\Gamma(\theta) = EG(\theta)$.

We will need the following version of the Inverse Function Theorem given by Foutz (1977). Here the spectral norm $\|\mathbf{A}\|$ of a matrix is defined $\mathbf{A}$ to be $\sup_{|a| < 1} |\mathbf{A}a|$.

Theorem 3.1 Suppose $\mathbf{f}$ is a mapping from an open set H in E^r into E^r , the partial derivatives of $\mathbf{f}$ exist and are continuous on H , and the matrix of derivatives $\mathbf{f}'(\theta^*)$ has inverse $\mathbf{f}'(\theta^*)^{-1}$ for some $\theta^* \in H$. Write

$$\lambda = 1/(4 \|\mathbf{f}'(\theta^*)^{-1}\|).$$

Use the continuity of the elements of $\mathbf{f}'(\theta^*)$ to fix a neighborhood U_δ of θ^* of sufficiently small radius $\delta > 0$ to insure $\|\mathbf{f}'(\theta) - \mathbf{f}'(\theta^*)\| \leq 2\lambda$, whenever $\theta \in U_\delta$. Then (1) for every θ_1, θ_2 in U_δ ,

$$|\mathbf{f}(\theta_1) - \mathbf{f}(\theta_2)| \geq 2\lambda |\theta_1 - \theta_2|,$$

and (2) the image set $\mathbf{f}(U_\delta)$ contains the open neighborhood with radius $\lambda\delta$ about $\mathbf{f}(\theta^*)$.

The first conclusion insures that $\mathbf{f}$ is one-to-one on U_δ , and the second that $\mathbf{f}^{-1}$ is well defined on the image set $\mathbf{f}(U_\delta)$.

Using (3.4.1) it is obvious that

$$\Gamma(\theta_0)^T = \begin{bmatrix} n_1^{-1} \sum_{i=1}^k \sum_{t=1}^{n_{1i}} (-1) \nu_1(\beta, \tau_1) + & 0 & 0 \\ (cs)^{-1} \sum_{j=2}^s (-n_j) \nu_2(\beta, \tau_1, \tau_2, n_j) & & \\ n_1^{-1} \sum_{i=1}^k \sum_{t=1}^{n_{1i}} \nu_1^{-1} \frac{\partial \nu_1}{\partial \beta} & n_1^{-1} \sum_{i=1}^k \sum_{t=1}^{n_{1i}} \nu_1^{-1} \frac{\partial \nu_1}{\partial \tau_1} & 0 \\ (cs)^{-1} \sum_{j=2}^s \nu_2^{-1} n_j^{-1} \frac{\partial \nu_2}{\partial \beta} & (cs)^{-1} \sum_{j=2}^s \nu_2^{-1} n_j^{-1} \frac{\partial \nu_2}{\partial \tau_1} & (cs)^{-1} \sum_{j=2}^s \nu_2^{-1} n_j^{-1} \frac{\partial \nu_2}{\partial \tau_2} \end{bmatrix} \quad (3.4.2)$$

In (3.4.2), the parameters β , τ_1 and τ_2 take their true values β_0 , τ_{10} and τ_{20} respectively.

Similar to the argument of Moore (1985), we introduce

Lemma 3.1 Given conditions C3-C5, $\Gamma(\theta_0)$ is of full rank.

Proof: It is obvious that the determinant of $\Gamma(\theta_0)$ equals the product of diagonal elements in (3.4.2), which is not equal to zero. $\square$

We wish to show that, as n_1 and $s \rightarrow \infty$ with $n_1/s \rightarrow \infty$, $G(\theta) \rightarrow \Gamma(\theta)$ in probability, uniformly in a neighborhood of θ_0 (lemma 3.2).

The proof of the following lemma is similar to Lemma 5.2 of Moore (1985). Some modifications are made in order to take into account of historical controls.

Lemma 3.2: Under C3 – C6, as n_1 and $s \rightarrow \infty$ with $n_1/s \rightarrow c > 0$, $\sup_{\theta \in U_\delta} \|G(\theta) - \Gamma(\theta)\| \rightarrow 0$ in probability.

Proof: Let $\theta_0 = (\beta_0, \tau_{10}, \tau_{20})^T$ be the true value of the parameter $\theta = (\beta, \tau_1, \tau_2)^T$.

(i) We first show that, as n_1 and $s \rightarrow \infty$ with $n_1/s \rightarrow c > 0$, $\sup_{\theta \in U_\delta} |G_{11} - \Gamma_{11}| \rightarrow_p 0$.

Using (3.4.1), we have

$$\begin{aligned} |G_{11} - \Gamma_{11}| &= n_1^{-1} \left| \sum_{i=1}^k \sum_{t=1}^{n_{1i}} (Y_{1it} - \beta) \frac{\partial \nu_1}{\partial \beta} + \sum_{j=2}^s (Y_j - n_j \beta) \frac{\partial \nu_2}{\partial \beta} \right. \\ &\quad \left. - \sum_{i=1}^k \sum_{t=1}^{n_{1i}} (\beta_0 - \beta) \frac{\partial \nu_1}{\partial \beta} - \sum_{j=2}^s (n_j \beta_0 - n_j \beta) \frac{\partial \nu_2}{\partial \beta} \right| \\ &= n_1^{-1} \left| \sum_{i=1}^k \sum_{t=1}^{n_{1i}} (Y_{1it} - \beta_0) \frac{\partial \nu_1}{\partial \beta} + \sum_{j=2}^s (Y_j - n_j \beta_0) \frac{\partial \nu_2}{\partial \beta} \right| \\ &\leq n_1^{-1} \left| \sum_{i=1}^k \sum_{t=1}^{n_{1i}} (Y_{1it} - \beta_0) \frac{\partial \nu_1}{\partial \beta} \right| \end{aligned}$$

$$+(cs)^{-1} \left| \sum_{j=2}^s (Y_j - n_j \beta_0) \frac{\partial \nu_2}{\partial \beta} \right|. \quad (3.4.3)$$

We now have, as $n_1 \rightarrow \infty$,

$$\begin{aligned} \sup_{\theta \in U_s} |n_1^{-1} \sum_{i=1}^k \sum_{t=1}^{n_{1i}} \frac{\partial \nu_1}{\partial \beta} (Y_{1it} - \beta_0)| &\leq \sup_{\theta \in U_s} \left| \frac{\partial \nu_1}{\partial \beta} \right| n_1^{-1} \left| \sum_{i=1}^k \sum_{t=1}^{n_{1i}} (Y_{1it} - \beta_0) \right| \\ &\leq M n_1^{-1} \left| \sum_{i=1}^k \sum_{t=1}^{n_{1i}} (Y_{1it} - \beta_0) \right| \rightarrow_p 0, \end{aligned} \quad (3.4.4)$$

under C5, by the law of large numbers. Next, we show that, as $s \rightarrow \infty$,

$$\sup_{\theta \in U_s} \left| s^{-1} \sum_{j=2}^s \frac{\partial \nu_2}{\partial \beta} (Y_j - n_j \beta_0) \right| \rightarrow_p 0.$$

For each fixed $s \in [2, \infty)$, there are at most N distinct values of n_j ($j = 2, \dots, s$). Let s_r denote the number of cases in which $n_j = r$. We then have $s-1 = s_1 + s_2 + \dots + s_N$, where $0 \leq s_r \leq s-1$ ($r = 1, 2, \dots, N$). We can now write

$$\sum_{j=2}^s \frac{\partial \nu_2}{\partial \beta} (Y_j - n_j \beta_0) = \sum_{r=1}^N \left\{ \sum_{j=2}^s l_{[n,=r]} \frac{\partial \nu_2}{\partial \beta} (Y_j - n_j \beta_0) \right\},$$

where $l_{[n,=r]} = 1$ if $n_j = r$ and 0 otherwise. We need to show that, for each r , $\sup_{\theta \in U_s} s^{-1} \left| \sum_{j=2}^s l_{[n,=r]} \frac{\partial \nu_2}{\partial \beta} (Y_j - n_j \beta_0) \right| \rightarrow_p 0$, as $s \rightarrow \infty$. Since $\frac{\partial \nu_2}{\partial \beta}$ is constant for r fixed, we may write

$$\begin{aligned} \sup_{\theta \in U_s} s^{-1} \left| \sum_{j=2}^s l_{[n,=r]} \frac{\partial \nu_2}{\partial \beta} (Y_j - n_j \beta_0) \right| &\leq \sup_{\theta \in U_s} \left| \frac{\partial \nu_2}{\partial \beta} \right| \frac{s_r}{s} \left| \frac{1}{s_r} \sum_{j=2}^s l_{[n,=r]} (Y_j - n_j \beta_0) \right| \\ &\leq \frac{s_r}{s} M \left| \frac{1}{s_r} \sum_{j=2}^s l_{[n,=r]} (Y_j - n_j \beta_0) \right|. \end{aligned}$$

If $s_r \rightarrow \infty$ as $s \rightarrow \infty$, then $\left| \frac{1}{s_r} \sum_{j=2}^s l_{[n,=r]} (Y_j - n_j \beta_0) \right| \rightarrow_p 0$, by the law of large numbers. If s_r remains finite as $s \rightarrow \infty$, we have

$$P \left\{ \frac{s_r}{s} M \left| \frac{1}{s_r} \sum_{j=2}^s l_{[n,=r]} (Y_j - n_j \beta_0) \right| > \varepsilon \right\} \leq \left[\frac{M}{s} \right]^2 s_r \frac{\text{Var}(Y_j)}{\varepsilon^2} \rightarrow 0,$$

as $s \rightarrow \infty$ by the Chebyshev inequality, under C3 and C5. Therefore, as $s \rightarrow \infty$,

$$\sup_{\theta \in U_s} \left| s^{-1} \sum_{j=2}^s \frac{\partial \nu_2}{\partial \beta} (Y_j - n_j \beta_0) \right| \rightarrow_p 0.$$

Hence, as n_1 and $s \rightarrow \infty$ with $n_1/s \rightarrow c > 0$.

$$\sup_{\theta \in U_s} |G_{11} - \Gamma_{11}| \rightarrow_p 0.$$

(ii) We next show $\sup_{\theta \in U_s} |G_{21} - \Gamma_{21}| \rightarrow_p 0$, as n_1 and $s \rightarrow \infty$ with $n_1/s \rightarrow c > 0$.

The same argument used in (i) above can be used to show

$$|G_{21} - \Gamma_{21}| \leq n_1^{-1} \left| \sum_{i=1}^k \sum_{t=1}^{n_{1i}} (Y_{1it} - \beta_0) \frac{\partial \nu_1}{\partial \tau_1} \right| + (cs)^{-1} \left| \sum_{j=2}^s (Y_j - n_j \beta_0) \frac{\partial \nu_2}{\partial \tau_1} \right|,$$

and

$$\sup_{\theta \in U_s} |G_{21} - \Gamma_{21}| \rightarrow_p 0,$$

as n_1 and $s \rightarrow \infty$ with $n_1/s \rightarrow c > 0$.

(iii) $\sup_{\theta \in U_s} |G_{31} - \Gamma_{31}| \rightarrow_p 0$, as $s \rightarrow \infty$.

Observe that

$$|G_{31} - \Gamma_{31}| = (cs)^{-1} \left| \sum_{j=2}^s (Y_j - n_j \beta_0) \frac{\partial \nu_2}{\partial \tau_2} \right|,$$

which implies that, as $s \rightarrow \infty$, $\sup_{\theta \in U_s} |G_{31} - \Gamma_{31}| \rightarrow_p 0$, under C3 and C5, by the same argument used in (i).

(iv) $\sup_{\theta \in U_s} |G_{12} - \Gamma_{12}| \rightarrow_p 0$, as $n_1 \rightarrow \infty$.

$$\begin{aligned} |G_{12} - \Gamma_{12}| &= \left| n_1^{-1} \left[\sum_{i=1}^k \sum_{t=1}^{n_{1i}} \{ 2(Y_{1it} - \beta_0)(-1)\nu_1(\beta, \tau_1) + ((Y_{1it} - \beta)^2 - E(Y_{1it} - \beta)^2) \frac{\partial \nu_1}{\partial \beta} \} \right] \right| \\ &\leq T_{12} + TT_{12}, \end{aligned} \tag{3.4.5}$$

where

$$T_{12} = 2 \left| \sum_{i=1}^k \sum_{t=1}^{n_{1i}} n_1^{-1} \nu_1(\beta, \tau_1) (Y_{1it} - \beta_0) \right|,$$

and

$$\begin{aligned} TT_{12} &= n_1^{-1} \left| \sum_{i=1}^k \sum_{t=1}^{n_{1i}} [(Y_{1it} - \beta)^2 - E(Y_{1it} - \beta)^2] \frac{\partial \nu_1}{\partial \beta} \right| \\ &= n_1^{-1} \left| \sum_{i=1}^k \sum_{t=1}^{n_{1i}} [(Y_{1it} - \beta_0)^2 - E(Y_{1it} - \beta_0)^2 + 2(\beta_0 - \beta)(Y_{1it} - \beta_0)] \frac{\partial \nu_1}{\partial \beta} \right| \\ &\leq TTT_1 + TTT_2, \end{aligned}$$

with

$$TTT_1 = n_1^{-1} \left| \sum_{i=1}^k \sum_{t=1}^{n_{1i}} [(Y_{1it} - \beta_0) 2(\beta_0 - \beta) \frac{\partial \nu_1}{\partial \beta}] \right|,$$

and

$$TTT_2 = n_1^{-1} \left| \sum_{i=1}^k \sum_{t=1}^{n_{1i}} [(Y_{1it} - \beta_0)^2 - E(Y_{1it} - \beta_0)^2] \frac{\partial \nu_1}{\partial \beta} \right|.$$

Following the same argument leading up to (3.4.3), we can easily show that $\sup_{\theta \in U_s} |T_{12}| \rightarrow_p 0$ and $\sup_{\theta \in U_s} |TTT_1| \rightarrow_p 0$, as $n_1 \rightarrow \infty$.

We also have

$$\begin{aligned} \sup_{\theta \in U_s} |TTT_2| &= \sup_{\theta \in U_s} \left| n_1^{-1} \sum_{i=1}^k \sum_{t=1}^{n_{1i}} [(Y_{1it} - \beta_0)^2 - E(Y_{1it} - \beta_0)^2] \frac{\partial \nu_1}{\partial \beta} \right| \\ &\leq \sup_{\theta \in U_s} \left| \frac{\partial \nu_1}{\partial \beta} \right| \cdot \left| n_1^{-1} \sum_{i=1}^k \sum_{t=1}^{n_{1i}} [(Y_{1it} - \beta_0)^2 - E(Y_{1it} - \beta_0)^2] \right| \rightarrow_p 0, \end{aligned} \quad (3.4.6)$$

as $n_1 \rightarrow \infty$, under C3, C5, and C6, by the law of large numbers.

Therefore, we obtain $\sup_{\theta \in U_s} |TT_{12}| \rightarrow_p 0$, as $n_1 \rightarrow \infty$. Since we also have shown that, as $n_1 \rightarrow \infty$, $\sup_{\theta \in U_s} |T_{12}| \rightarrow_p 0$, it follows from (3.4.5) that

$$\sup_{\theta \in U_s} |G_{12} - \Gamma_{12}| \rightarrow_p 0,$$

as $n_1 \rightarrow \infty$.

(v) $\sup_{\theta \in U_s} |G_{22} - \Gamma_{22}| \rightarrow_p 0$, as $n_1 \rightarrow \infty$.

Note that

$$|G_{22} - \Gamma_{22}| = n_1^{-1} \left| \sum_{i=1}^k \sum_{t=1}^{n_{1i}} [(Y_{1it} - \beta)^2 - E(Y_{1it} - \beta)^2] \frac{\partial \nu_1}{\partial \tau_1} \right|.$$

Following the same argument used in (3.4.6), it is straightforward to show $\sup_{\theta \in U_s} |G_{22} - \Gamma_{22}| \rightarrow_p 0$, as $n_1 \rightarrow \infty$.

(vi) $\sup_{\theta \in U_s} |G_{32} - \Gamma_{32}| \rightarrow_p 0$, as n_1 and $s \rightarrow \infty$ with $n_1/s \rightarrow c > 0$.

The proof for $|G_{32} - \Gamma_{32}| \rightarrow_p 0$, as n_1 and $s \rightarrow \infty$ with $n_1/s \rightarrow c > 0$, is trivial, since it equals zero.

(vii) $\sup_{\theta \in U_s} |G_{13} - \Gamma_{13}| \rightarrow_p 0$, as $s \rightarrow \infty$.

Write

$$\begin{aligned} |G_{13} - \Gamma_{13}| &= (cs)^{-1} \left| \sum_{j=2}^s 2(Y_j - n_j \beta_0) \nu_2(\beta, \tau_1, \tau_2, n_j) \right. \\ &\quad \left. + \sum_{j=2}^s [(Y_j - n_j \beta)^2 - E(Y_j - n_j \beta)^2] n_j^{-1} \frac{\partial \nu_2}{\partial \beta} \right| \\ &\leq S_{13} + SS_{13}, \end{aligned}$$

where

$$S_{13} = |(cs)^{-1} \sum_{j=2}^s 2\nu_2(\beta, \tau_1, \tau_2, n_j)(Y_j - n_j \beta_0)|$$

and

$$SS_{13} = |(cs)^{-1} \sum_{j=2}^s [(Y_j - n_j \beta)^2 - E(Y_j - n_j \beta)^2] n_j^{-1} \frac{\partial \nu_2}{\partial \beta}|.$$

Under assumptions C3, and C4, it is easy to see that, as $s \rightarrow \infty$, $\sup_{\theta \in U_s} |S_{13}| \rightarrow_p 0$, by the same argument used in (i).

To show that, as $s \rightarrow \infty$, $\sup_{\theta \in U_s} |SS_{13}| \rightarrow_p 0$, we follow the proof in (iv) and write

$$\begin{aligned} SS_{13} &= |(cs)^{-1} \sum_{j=2}^s n_j^{-1} \frac{\partial \nu_2}{\partial \beta} [(Y_j - n_j \beta_0)^2 - E(Y_j - n_j \beta_0)^2 + 2n_j(\beta - \beta_0)(Y_j - n_j \beta_0)]| \\ &\leq SSS_1 + SSS_2, \end{aligned}$$

where

$$SSS_1 = |(cs)^{-1} \sum_{j=2}^s n_j^{-1} \frac{\partial \nu_2}{\partial \beta} [2n_j(\beta - \beta_0)]|$$

and

$$SSS_2 = |(cs)^{-1} \sum_{j=2}^s n_j^{-1} \frac{\partial \nu_2}{\partial \beta} [(Y_j - n_j \beta_0)^2 - E(Y_j - n_j \beta_0)^2]|.$$

It is straightforward to see that, as $s \rightarrow \infty$, $\sup_{\theta \in U_s} |SSS_1| \rightarrow_p 0$, under C3, and C5, by the same argument in (i). Also following the argument in (i) and under assumptions C3, C5, and C6, we have $\sup_{\theta \in U_s} |SSS_2| \rightarrow_p 0$, as $s \rightarrow \infty$. Hence, as $s \rightarrow \infty$, $\sup_{\theta \in U_s} |G_{13} - \Gamma_{13}| \rightarrow_p 0$.

(viii) $\sup_{\theta \in U_s} |G_{23} - \Gamma_{23}| \rightarrow_p 0$, as $s \rightarrow \infty$.

Note that

$$\begin{aligned} |G_{23} - \Gamma_{23}| &= (cs)^{-1} \left| \sum_{j=2}^s [(Y_j - n_j\beta_0 + n_j\beta_0 - n_j\beta)^2 - E(Y_j - n_j\beta_0 + n_j\beta_0 - n_j\beta)^2] n_j^{-1} \frac{\partial \nu_2}{\partial \tau_1} \right| \\ &\leq T_{23} + TT_{23}, \end{aligned}$$

where

$$T_{23} = (cs)^{-1} \left| \sum_{j=2}^s 2(Y_j - n_j\beta_0)(n_j(\beta_0 - \beta)) n_j^{-1} \frac{\partial \nu_2}{\partial \tau_1} \right|$$

and

$$TT_{23} = (cs)^{-1} \left| \sum_{j=2}^s [(Y_j - n_j\beta_0)^2 - E(Y_j - n_j\beta_0)^2] n_j^{-1} \frac{\partial \nu_2}{\partial \tau_2} \right|.$$

Following the argument in (vii), we have that, as $s \rightarrow \infty$, $\sup_{\theta \in U_\delta} |G_{23} - \Gamma_{23}| \rightarrow_p 0$ under C3, C5 and C6.

(ix) $\sup_{\theta \in U_\delta} |G_{33} - \Gamma_{33}| \rightarrow_p 0$, as $s \rightarrow \infty$.

Note that $|G_{33} - \Gamma_{33}| = |n_1^{-1} \sum_{j=2}^s [(Y_j - n_j\beta) - E(Y_j - n_j\beta)] n_j^{-1} \frac{\partial \nu_2}{\partial \tau_2}|$. Under C3, C5, C6, and following the same argument in (vii), the result follows.

Based on the results (i)-(ix), we conclude that, as n_1 and $s \rightarrow \infty$ with $n_1/s \rightarrow c > 0$, $\sup_{\theta \in U_\delta} \|G(\theta) - \Gamma(\theta)\| \rightarrow_p 0$. $\square$

Let $\hat{\theta}$ be the solution to the equation $\mathbf{f}(\theta) = [f_1(\theta), f_2(\theta), f_3(\theta)]^T = 0$.

Again, similarly to Theorem 5.4 of Moore (1985), we introduce the following theorem which establishes the consistency of $\hat{\theta}$.

Theorem 3.2: Let $\mathbf{F}_{n_1}(\theta) = n_1^{-1}\mathbf{f}(\theta)$, and recall that Let $\Gamma(\theta) = E\mathbf{F}'_{n_1}(\theta)$. Let $\mathbf{F}_{n_1}^{-1}(\mathbf{0})$ be the root of $\mathbf{F}_{n_1}(\theta) = \mathbf{0}$. Under C1 and C3-C6, we have the following results.

i) The root $(\hat{\beta}, \hat{\tau}_1, \hat{\tau}_2)$ of $\mathbf{F}_{n_1}(\theta) = \mathbf{0}$, $\mathbf{F}_{n_1}^{-1}(\mathbf{0})$ exists in U_δ with probability approaching to one as $n_1 \rightarrow \infty$.

ii) Since δ may be arbitrarily small, $\mathbf{F}_{n_1}^{-1}(\mathbf{0})$ converges in probability to θ_0 , as $n_1 \rightarrow \infty$.

iii) Since $\mathbf{F}_{n_1}$ is a one-to-one function on U_δ , any other sequence $\hat{\theta}^* = (\hat{\beta}^*, \hat{\tau}_1^*, \hat{\tau}_2^*)$ of the root to $\mathbf{F}_{n_1}(\theta) = \mathbf{0}$ necessarily lies outside of U_δ with probability approaching one as $n_1 \rightarrow \infty$. This implies $\hat{\theta}^*$ does not converge to θ_0 .

Proof: Since $\Gamma(\theta_0)$ is of full rank (Lemma 3.1), we may define

$$\lambda = \frac{1}{4 \|\Gamma(\theta_0)^{-1}\|}$$

and select U_δ sufficiently small so that

$$\|\Gamma(\theta) - \Gamma(\theta_0)\| < \lambda/2$$

whenever $\theta \in U_\delta$. By lemma 3.2, the derivative of $\mathbf{F}_{n_1}(\theta)$, $\mathbf{F}'_{n_1}(\theta)$, converges uniformly to $\Gamma(\theta)$, as $n_1 \rightarrow \infty$, for any θ such that $\|\theta - \theta_0\| < \delta$. Therefore, we have

$$\begin{aligned} \|\mathbf{F}'_{n_1}(\theta) - \mathbf{F}'_{n_1}(\theta_0)\| &\leq \|\mathbf{F}'_{n_1}(\theta) - \Gamma(\theta)\| \\ &\quad + \|\Gamma(\theta_0) - \mathbf{F}'_{n_1}(\theta_0)\| + \|\Gamma(\theta) - \Gamma(\theta_0)\| < \lambda. \end{aligned} \quad (3.4.7)$$

Since $\mathbf{F}'_{n_1}(\theta_0) \rightarrow \Gamma(\theta_0)$ in probability, as $n_1 \rightarrow \infty$ (lemma 3.2), and $\Gamma(\theta_0)$ is invertible (Lemma 3.1), and since matrix inversion is a continuous function, it follows that $\mathbf{F}'_{n_1}(\theta_0)^{-1}$ exists with probability approaching to one as $n_1 \rightarrow \infty$. We may thus write

$$\lambda_{n_1} = \frac{1}{4 \|\mathbf{F}'_{n_1}(\theta_0)^{-1}\|}$$

whenever the inverse function $\mathbf{F}'_{n_1}(\theta_0)$ exists. Recall the definitions of λ, δ and use the uniform convergence of $\mathbf{F}'_{n_1}(\theta)$ to $\Gamma(\theta)$ to argue $\lambda_{n_1} \rightarrow \lambda$ in probability, as $n_1 \rightarrow \infty$, and if

$$\|\theta - \theta_0\| < \delta,$$

we have, by (3.4.7),

$$\|\mathbf{F}'_{n_1}(\theta) - \mathbf{F}'_{n_1}(\theta_0)\| < \lambda < 2\lambda_{n_1}$$

Then, from the Inverse Function theorem (Theorem 3.1), we have that $\mathbf{F}_{n_1}(\theta)$ is a one-to-one function from U_δ onto $\mathbf{F}_{n_1}(U_\delta)$ and that the image set $\mathbf{F}_{n_1}(U_\delta)$ contains the open neighborhood of radius $\lambda_n \delta$ about $\mathbf{F}_{n_1}(\theta_0)$. Hence it contains, with probability one, the neighborhood of radius $\lambda \delta/2$.

We may see that $\mathbf{0} \in \mathbf{F}_{n_1}(U_\delta)$ with probability approaching one by noting that

$$E\mathbf{F}_{n_1}(\theta_0) = \mathbf{0}$$

and that

$$|F_{n_1}(\theta_0) - 0| < \lambda\delta/2$$

with probability approaching one as $n_1 \rightarrow \infty$ by the strong law of large number.

Consider the inverse function $F_{n_1}^{-1} : F_{n_1}(U_\delta) \rightarrow U_\delta$. It is well defined whenever $F_{n_1}^{-1}$ is one-to-one, i.e., with probability going to one. Because $0 \in F_{n_1}(U_\delta)$ with probability approaching one as $n_1 \rightarrow \infty$, Theorem 3.2 is established. $\square$

Conclusion (iii) implies the uniqueness of $\hat{\theta}$ satisfying $F_{n_1}(\hat{\theta}) = 0$, (ii) shows $\hat{\theta} \rightarrow \theta_0$ in probability as $n_1 \rightarrow \infty$.

Next, we will show the asymptotic normality of the test for trend given in (3.3.5). We need to show first that $n_1^{-1/2}(f_1(\theta_0), f_2(\theta_0), f_3(\theta_0))^T$ is asymptotically normal. We will employ the following version of Lindeberg central limit theorem (see e.g. Billingsley, 1986).

Theorem 3.3: Assume $\{X_i\}_{i=1}^n$ are mutually independent random variables with

$$EX_i = 0, \quad \sigma_i^2 = EX_i^2, \quad s_n^2 = \sum_{i=1}^n \sigma_i^2,$$

and $S_n = \sum_{i=1}^n X_i$. If

$$\lim_{n \rightarrow \infty} \frac{1}{s_n} \sum_{i=1}^n E[X_i^2 I_{\{|X_i| > \epsilon s_n\}}] \rightarrow 0,$$

Then (i)

$$\frac{S_n}{s_n} \rightarrow N(0, 1)$$

in distribution as $n \rightarrow \infty$. In addition, if

$$\lim_{n \rightarrow \infty} \frac{1}{n} s_n^2 = \sigma_0^2,$$

(ii) then, as $n \rightarrow \infty$, $S_n/n \rightarrow N(0, \sigma_0)$.

We will also require the following lemma.

Lemma 3.3: Let Z_r , ($r = 1, 2, \dots, n$) be independent random vectors of finite dimension. Assume

$$\Sigma_r = \text{Cov}(Z_r), \text{ and } \left\| \frac{1}{n} \sum_{r=1}^n \Sigma_r - \Sigma_0 \right\| \rightarrow 0 \text{ in probability,}$$

as $n \rightarrow \infty$, where $\|\cdot\|$ denotes the spectral norm. Then, as $n \rightarrow \infty$,

$$\frac{1}{n} \sum_{r=1}^n Z_r \rightarrow N(0, \Sigma_0)$$

in distribution.

Proof: For any real vector λ , we have, as $n \rightarrow \infty$,

$$\| \lambda' [\frac{1}{n} \sum_{r=1}^n \Sigma_r] \lambda - \lambda' \Sigma_0 \lambda \| \rightarrow 0.$$

By Theorem 3.3, we have, as $n \rightarrow \infty$,

$$Y_n = \lambda' \frac{\sum_{r=1}^n Z_r}{n} \rightarrow N(0, \lambda' \Sigma_0 \lambda).$$

Therefore the characteristic function of Y_n

$$E[\exp\{itY_n\}] \rightarrow \exp\{-\frac{1}{2}t^2 \lambda' \Sigma_0 \lambda\},$$

as $n \rightarrow \infty$. For $t = 1$, we have, as $n \rightarrow \infty$,

$$E[\exp\{i\lambda' \frac{\sum_{r=1}^n Z_r}{n}\}] \rightarrow \exp\{-\frac{1}{2}\lambda' \Sigma_0 \lambda\}$$

for all λ . Since the distribution function and the characteristic function are one-to-one, we have

$$\frac{\sum_{r=1}^n Z_r}{n} \rightarrow N(0, \Sigma_0),$$

as $n \rightarrow \infty$. $\square$

Again, we introduce a theorem which is similar to Theorem 5.2 in Moore (1985).

Theorem 3.4: Let $K^* = \sum_{i=1}^k n_{1i} + (s-1)$. Under the conditions C1 and C3-C6,

$$K^{*-1/2}(f_1(\theta_0), f_2(\theta_0), f_3(\theta_0))^T \rightarrow N_3(0, \Sigma_0)$$

as $K^* \rightarrow \infty$, where $\Sigma_0 = \lim_{K^* \rightarrow \infty} [Cov(f(\theta_0))/K^*]$.

Proof: For $t = 1, \dots, n_{1i}$, $i = 1, \dots, K$, and $j = 2, \dots, s$, let $Y_{1it}^* = (Y_{1it} - \beta_0)\nu_1(\beta_0, \tau_{10})$, $Y_{1it}^{**} = (Y_{1it} - \beta_0)^2\nu_1(\beta_0, \tau_{10}) - 1$, $Y_j^* = (Y_j - n_j\beta_0)\nu_2(\beta_0, \tau_{10}, \tau_{20}, n_j)$, and $Y_j^{**} = (Y_j - n_j\beta_0)^2n_j^{-1}\nu_2(\beta_0, \tau_{10}, \tau_{20}, n_j) - 1$.

Define $(Z_{11}, Z_{12}, \dots, Z_{1K^*})^T$ as

$$(Y_{111}^*, \dots, Y_{11n_{11}}^*, Y_{121}^*, \dots, Y_{12n_{12}}^*, \dots, Y_{1K1}^*, \dots, Y_{1Kn_{1K}}^*, Y_2^*, Y_3^*, \dots, Y_s^*)^T.$$

Define $(Z_{21}, Z_{22}, \dots, Z_{2K^*})^T$ as

$$(Y_{111}^{**}, \dots, Y_{11n_{11}}^{**}, Y_{121}^{**}, \dots, Y_{12n_{12}}^{**}, \dots, Y_{1K1}^{**}, \dots, Y_{1Kn_{1K}}^{**}, 0, 0, 0, \dots, 0, 0)^T.$$

Define $(Z_{31}, Z_{32}, \dots, Z_{3K^*})^T$ as

$$(0, \dots, 0, 0, \dots, 0, 0, \dots, 0, Y_2^*, \dots, Y_s^*)^T.$$

Further define $\mathbf{Z}_r = (Z_{1r}, Z_{2r}, Z_{3r})^T$ for $r = 1, \dots, K^*$. $\mathbf{S}_{K^*} = \sum_{r=1}^{K^*} \mathbf{Z}_r$, and $S_{K^*} = \boldsymbol{\lambda}' \mathbf{S}_{K^*}$, where $\boldsymbol{\lambda} = (\lambda_1, \lambda_2, \lambda_3)^T$. Note that

$$\begin{aligned} \text{Var}(S_{K^*}) &= \boldsymbol{\lambda}' \text{Cov}(\mathbf{S}_{K^*}) \boldsymbol{\lambda} \\ &= \boldsymbol{\lambda}' \left(\sum_{r=1}^{K^*} \Sigma_r \right) \boldsymbol{\lambda} \\ &= \sum_{r=1}^{K^*} \boldsymbol{\lambda}' \Sigma_r \boldsymbol{\lambda} \\ &= \sum_{r=1}^{K^*} \sigma_r^2 = s_{K^*}^2. \end{aligned}$$

where $\Sigma_r = \text{Cov}(\mathbf{Z}_r)$, $\boldsymbol{\lambda}' \Sigma_r \boldsymbol{\lambda} = \sigma_r^2$.

We need to verify the Lindeberg condition in Theorem 3.3:

$$\frac{1}{s_{K^*}} \sum_{r=1}^{K^*} E |\boldsymbol{\lambda}' \mathbf{Z}_r|^2 I_{\{|\boldsymbol{\lambda}' \mathbf{Z}_r| \geq \varepsilon s_{K^*}\}} \rightarrow 0$$

for any real vector $\boldsymbol{\lambda}$ as $K^* \rightarrow \infty$. This is not difficult in our case. It follows from C6 and C7 that the elements of $\Sigma_r = \text{Cov}(\mathbf{Z}_r)$ are bounded by $M^{**} < \infty$ for all $r = 1, 2, \dots, K^*$. Therefore,

$$s_{K^*}^2 / K^* = \frac{1}{K^*} \sum_{r=1}^{K^*} \sigma_r^2 = O(1),$$

for each $\boldsymbol{\lambda}$. Further note that

$$\begin{aligned} \{|\boldsymbol{\lambda}' \mathbf{Z}_r| \geq \varepsilon s_{K^*}\} &= \{|\lambda_1 Z_{1r} + \lambda_2 Z_{2r} + \lambda_3 Z_{3r}| \geq \varepsilon s_{K^*}\} \\ &\subseteq \{M_1(|\lambda_1| + |\lambda_2| + |\lambda_3|) \geq \varepsilon s_{K^*}\}, \end{aligned}$$

and

$$|\boldsymbol{\lambda}' \mathbf{Z}_r| \leq M_1 \max[\lambda_i] = M^*$$

for $i = 1, 2, 3$. Hence

$$\frac{1}{s_{K^*}} \sum_{r=1}^{K^*} E |\boldsymbol{\lambda}' \mathbf{Z}_r|^2 I_{\{|\boldsymbol{\lambda}' \mathbf{Z}_r| \geq \varepsilon s_{K^*}\}} \leq M^{*2} E I_{\{M^* \geq \varepsilon s_{K^*}\}} \rightarrow 0$$

as $K^* \rightarrow \infty$, since $s_{K^*} = O(\sqrt{K^*})$. Using C7. result (ii) of Theorem 3.3 and Lemma 3.3, asymptotic normal follows. $\square$

Since, as $n_1 \rightarrow \infty$, $\hat{\theta} \rightarrow \theta_0$ in probability, applying a Taylor expansion, we have

$$0 = F_{n_1}(\hat{\theta}) = F_{n_1}(\theta_0) + F'_{n_1}(\theta^*)(\hat{\theta} - \theta_0)$$

where θ^* lies between $\hat{\theta}$ and θ_0 . Equivalently, we have

$$-\sqrt{n_1}f(\theta_0) = \sqrt{n_1}f'(\theta^*)(\hat{\theta} - \theta_0).$$

Using Theorem 3.2 and Theorem 3.4, we now have the following result.

Corollary 3.1: $\sqrt{n_1}(\hat{\theta} - \theta_0) \rightarrow N_3(0, \hat{\psi})$ in distribution, as $n_1 \rightarrow \infty$, where $\hat{\psi} = \Gamma^{-1}(\theta_0)\Sigma_0(\Gamma^{-1}(\theta_0))^T$.

Recall that in section 3.3 we introduced U_1 and U_2 as:

$$D^T V^{-1}(Y - \mu) = (U_1(\beta, \tau_1, \tau_2), U_2(\beta, \tau_1, \tau_2))^T.$$

In the remaining part of this section, we will derive the asymptotic distribution of the score statistic $U_2(\hat{\beta}_0, \hat{\tau}_1, \hat{\tau}_2)$ given in (3.3.6).

Applying a Taylor expansion to $n_1^{-1}U_1(\hat{\theta}) = 0$ about the point θ_0 , we have

$$\begin{aligned} 0 &= n_1^{-1}U_1(\hat{\beta}, \hat{\tau}_1, \hat{\tau}_2) \\ &= n_1^{-1}\{U_1(\beta_0, \tau_{10}, \tau_{20}) + \frac{\partial U_1}{\partial \beta_0}(\hat{\beta} - \beta_0) + \frac{\partial U_1}{\partial \tau_{10}}(\hat{\tau}_1 - \tau_{10}) \\ &\quad + \frac{\partial U_1}{\partial \tau_{20}}(\hat{\tau}_2 - \tau_{20}) + \frac{\partial^2 U_1}{\partial \beta^2}(\hat{\beta} - \beta_0)^2 + \frac{\partial^2 U_1}{\partial \beta' \partial \tau_1'}(\hat{\beta} - \beta_0)(\hat{\tau}_1 - \tau_{10}) + \dots\}, \end{aligned} \quad (3.4.8)$$

where β' , τ_1' , and τ_2' lie in the intervals $(\beta_0, \hat{\beta})$, $(\tau_{10}, \hat{\tau}_1)$, and $(\tau_{20}, \hat{\tau}_2)$ respectively.

Recall that $\hat{\theta} = (\hat{\beta}, \hat{\tau}_1, \hat{\tau}_2)^T \rightarrow \hat{\theta}_0 = (\beta_0, \tau_{10}, \tau_{20})^T$ in probability, as $n_1 \rightarrow \infty$ (Theorem 3.2), and by corollary 3.1 $\sqrt{n_1}(\hat{\theta} - \hat{\theta}_0)$ is asymptotic normal, as $n_1 \rightarrow \infty$. Corollary 3.1 implies that the convergence rates for $\hat{\beta}$, $\hat{\tau}_1$, and $\hat{\tau}_2$ are same. Also note that the partial derivatives of $n_1^{-1}U_1$ are bounded. Then

$$n_1^{-1} \frac{\partial U_1}{\partial \tau_{10}} = n_1^{-1} \left\{ \sum_{i=1}^k \sum_{t=1}^{n_{1i}} (Y_{1it} - \beta_0) \frac{\partial \nu_1}{\partial \tau_1} + \sum_{j=2}^k (Y_j - n_j \beta_0) \frac{\partial \nu_2}{\partial \tau_1} \right\} \rightarrow 0$$

in probability as $n_1 \rightarrow \infty$. Also

$$n_1^{-1} \frac{\partial U_1}{\partial \tau_{20}} = n_1^{-1} \sum_{j=2}^k (Y_j - n_j \beta_0) \frac{\partial \nu_2}{\partial \tau_2} \rightarrow 0$$

in probability as $n_1 \rightarrow \infty$.

Using these two results and Theorem 3.2, we can ignore the terms involving $\partial U_1 / \partial \tau_{i0}$ ($i = 1, 2$) as $n_1 \rightarrow \infty$, as well as the higher order terms. Therefore (3.4.8) can be written as

$$0 = U_1(\hat{\theta}) = U_1(\beta_0, \tau_{10}, \tau_{20}) + \frac{\partial U_1}{\partial \beta_0}(\hat{\beta} - \beta_0) + o_p\left(\frac{\partial U_1}{\partial \beta_0}(\hat{\beta} - \beta_0)\right).$$

We now have

$$\hat{\beta} - \beta_0 = -\frac{U_1(\beta_0, \tau_{10}, \tau_{20})}{\partial U_1 / \partial \beta_0} + o_p(\hat{\beta} - \beta_0). \quad (3.4.9)$$

Similarly, applying a Taylor expansion to $U_2(\hat{\theta})$ about θ_0 yields

$$\begin{aligned} U_2(\hat{\beta}, \hat{\tau}_1, \hat{\tau}_2) &= U_2(\beta_0, \tau_{10}, \tau_{20}) + \frac{\partial U_2}{\partial \beta_0}(\hat{\beta} - \beta_0) \\ &\quad + \frac{\partial U_2}{\partial \tau_{10}}(\hat{\tau}_1 - \tau_{10}) + \frac{\partial U_2}{\partial \tau_{20}}(\hat{\tau}_2 - \tau_{20}) \\ &\quad + \frac{\partial^2 U_2}{\partial \beta^2}(\hat{\beta} - \beta_0)^2 + \dots \end{aligned} \quad (3.4.10)$$

Note that although $n_1^{-1} |\partial U_2 / \partial \tau_{i0}| \rightarrow 0$ ($i = 1, 2$) in probability as $n_1 \rightarrow \infty$, $n_1^{-1} / \partial U_2 / \partial \beta_0$ does not converge to zero. From (3.4.10), we have

$$U_2(\hat{\beta}, \hat{\tau}_1, \hat{\tau}_2) = U_2(\beta_0, \tau_{10}, \tau_{20}) + \frac{\partial U_2}{\partial \beta_0}(\hat{\beta} - \beta_0) + o_p\left(\frac{\partial U_2}{\partial \beta_0}(\hat{\beta} - \beta_0)\right). \quad (3.4.11)$$

Substituting (3.4.9) into (3.4.11) yields the approximation

$$U_2(\hat{\beta}, \hat{\tau}_1, \hat{\tau}_2) = U_2(\beta_0, \tau_{10}, \tau_{20}) + \frac{\partial U_2}{\partial \beta_0} \left[-\frac{\partial U_1}{\partial \beta_0} \right]^{-1} U_1(\beta_0, \tau_{10}, \tau_{20}) + o_p\left(\frac{\partial U_2}{\partial \beta_0}(\hat{\beta} - \beta_0)\right). \quad (3.4.12)$$

Under the null hypothesis $\beta_1 = 0$, define $\partial U_i = \partial U_i / \partial \beta_0$ ($i = 1, 2$). By direct calculation, it can be seen that

$$E\partial U_1 = -C_{11}^* n_1 - C_{21}^* \sum_{j=2}^I n_j,$$

$$E\partial U_2 = -C_1^{**} \sum_{i=1}^K n_{1i} d_i,$$

and

$$\begin{aligned} \frac{E\partial U_2}{E\partial U_1} - \frac{\partial U_2}{\partial U_1} &= \frac{E\partial U_2 \partial U_1 - \partial U_1 \partial U_2 + \partial U_1 \partial U_2 - \partial U_2 E\partial U_1}{E\partial U_1 \partial U_1} \\ &= \frac{\partial U_1 (E\partial U_2 - \partial U_2) + \partial U_2 (\partial U_1 - E\partial U_1)}{E\partial U_1 \partial U_1}. \end{aligned}$$

Under the assumptions C1-C6, it follows from Slutsky's theorem that both $(E\partial U'_1 - \partial U_1)/E\partial U_1$ and $(E\partial U'_2 - \partial U_2)/E\partial U_1$ converge to zero in probability, as $n_1 \rightarrow \infty$. Thus

$$\frac{E\partial U'_2}{E\partial U'_1} - \frac{\partial U'_2}{\partial U'_1} \rightarrow 0$$

in probability, as $n_1 \rightarrow \infty$. Replacing $\partial U'_2$, and $\partial U'_1$ by their expected value $E\partial U'_2$, and $E\partial U'_1$ respectively in (3.4.12), and using corollary 3.1 to evaluate the $o_p(\frac{\partial U'_2}{\partial \beta_0}(\hat{\beta} - \beta_0))$, we have

$$\begin{aligned} (1/\sqrt{n_1})U_2(\hat{\beta}, \hat{\tau}_1, \hat{\tau}_2) &= (1/\sqrt{n_1})U_2(\beta_0, \tau_{10}, \tau_{20}) \\ &= (1/\sqrt{n_1})\{U_2(\beta_0, \tau_{10}, \tau_{20}) - E\frac{\partial U_2}{\partial \beta_0}[E\frac{\partial U_1}{\partial \beta_0}]^{-1}U_1(\beta_0, \tau_{10}, \tau_{20})\} + o_p(1) \end{aligned} \quad (3.4.13)$$

Note that the elements of

$$U(\beta_0, \tau_{10}, \tau_{20}) = \begin{pmatrix} U_1(\beta_0, \tau_{10}, \tau_{20}) \\ U_2(\beta_0, \tau_{10}, \tau_{20}) \end{pmatrix} = \mathbf{D}^T \mathbf{V}^{-1}(\mathbf{Y} - \boldsymbol{\mu})$$

involve a sum of n_1 random variables. Each such random variable has zero mean and finite variance (see C6). Following the proof of Theorem 3.4, we have that $\frac{1}{\sqrt{n_1}}[U_1(\beta_0, \tau_{10}, \tau_{20}), U_2(\beta_0, \tau_{10}, \tau_{20})]^T$ has multivariate normal limiting distribution under C1-C7. Therefore, using (3.4.13),

$$\begin{aligned} \frac{1}{\sqrt{n_1}}U_2(\hat{\beta}, \hat{\tau}_1, \hat{\tau}_2) &= \frac{1}{\sqrt{n_1}}\{U_2(\beta_0, \tau_{10}, \tau_{20}) - (E\frac{\partial U_2}{\partial \beta_0})(E\frac{\partial U_1}{\partial \beta_0})^{-1}U_1(\beta_0, \tau_{10}, \tau_{20})\} + o_p(1) \\ &= \frac{1}{\sqrt{n_1}}[-(E\frac{\partial U_2}{\partial \beta_0})(E\frac{\partial U_1}{\partial \beta_0})^{-1}, 1][U_1(\beta_0, \tau_{10}, \tau_{20}), U_2(\beta_0, \tau_{10}, \tau_{20})]^T + o_p(1) \end{aligned}$$

has a univariate normal distribution. Let

$$\text{Cov}(U_1(\boldsymbol{\theta}_0), U_2(\boldsymbol{\theta}_0)) = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}$$

where $\theta_0 = (\beta_0, \tau_{10}, \tau_{20})^T$. Then

$$\begin{aligned}
 & \text{Cov}\{U_2(\hat{\beta}, \hat{\tau}_1, \hat{\tau}_2), U_2(\hat{\beta}, \hat{\tau}_1, \hat{\tau}_2)\} \\
 &= \left[\left(E \frac{\partial U_2}{\partial \beta_0} \right) \left(E \frac{\partial U_1}{\partial \beta_0} \right)^{-1}, \quad 1 \right] \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix} \begin{bmatrix} - \left(E \frac{\partial U_2}{\partial \beta_0} \right) \left(E \frac{\partial U_1}{\partial \beta_0} \right)^{-1} \\ 1 \end{bmatrix} \\
 &= \left[-E \frac{\partial U_2}{\partial \beta_0} \left(E \frac{\partial U_1}{\partial \beta_0} \right)^{-1} A_{11} + A_{21}, -E \frac{\partial U_2}{\partial \beta_0} \left(E \frac{\partial U_1}{\partial \beta_0} \right)^{-1} A_{12} + A_{22} \right] \begin{bmatrix} -E \frac{\partial U_2}{\partial \beta_0} \left(E \frac{\partial U_1}{\partial \beta_0} \right)^{-1} \\ 1 \end{bmatrix} \\
 &= \left(E \frac{\partial U_2}{\partial \beta_0} \right)^2 \left(E \frac{\partial U_1}{\partial \beta_0} \right)^{-2} A_{11} - E \frac{\partial U_2}{\partial \beta_0} \left(E \frac{\partial U_1}{\partial \beta_0} \right)^{-1} A_{21} - E \frac{\partial U_2}{\partial \beta_0} \left(E \frac{\partial U_1}{\partial \beta_0} \right)^{-1} A_{12} + A_{22}.
 \end{aligned}$$

It can be shown that at $\theta = \theta_0$

$$EU_1^2 = \frac{n_1}{\beta_0(1 + \beta_0\tau_{10})} + \sum_{j=2}^s \frac{n_j}{\beta_0(1 + \beta_0(\tau_{10} + n_j\tau_{20} + \tau_{10}\tau_{20}))}$$

and

$$\begin{aligned}
 E \frac{\partial U_1}{\partial \beta_0} &= -\frac{n_1}{\beta_0(1 + \beta_0\tau_{10})} - \sum_{j=2}^s \frac{n_j}{\beta_0(1 + \beta_0(\tau_{10} + n_j\tau_{20} + \tau_{10}\tau_{20}))} \\
 EU_1^2 &= -E \frac{\partial U_1}{\partial \beta_0}.
 \end{aligned}$$

We also have

$$EU_1U_2 = \sum_{i=1}^K \frac{n_{1i}d_i}{\beta_0(1 + \beta_0\tau_{10})}$$

and

$$E \frac{\partial U_2}{\partial \beta_0} = -\sum_{i=1}^K \frac{n_{1i}d_i}{\beta_0(1 + \beta_0\tau_{10})}.$$

Therefore

$$EU_1U_2 = -E \frac{\partial U_2}{\partial \beta_0}.$$

Since $EU_i = 0$ ($i = 1, 2$), we have

$$A_{11} = EU_1^2 = -E \frac{\partial U_1}{\partial \beta_0},$$

$$A_{12} = EU_1U_2 = -E\frac{\partial U_2}{\partial \beta_0},$$

$$A_{21} = EU_2U_1 = A_{12},$$

and

$$A_{22} = EU_2^2.$$

Substituting A_{11} , A_{12} , and A_{22} for $E\frac{\partial U_i}{\partial \beta_0}$ ($i = 1, 2$) in $Cov\{U_2(\hat{\beta}), \hat{\tau}_1, \hat{\tau}_2\}$ we obtain

$$\begin{aligned} Cov(U_2(\hat{\theta}))|_{\beta_1=0} &= A_{12}A_{11}^{-1}A_{21} - 2A_{12}A_{11}^{-1}A_{21} + A_{22} \\ &= A_{22} - A_{21}A_{11}^{-1}A_{12}. \end{aligned} \quad (3.4.14)$$

Therefore,

$$Cov[(1/\sqrt{n_1})U_2(\hat{\theta})] \rightarrow C_{22} - C_{21}C_{11}^{-1}C_{12},$$

as $n_1 \rightarrow \infty$, where C_{ij} are the elements of matrix

$$C = \lim_{n_1 \rightarrow \infty} n_1^{-1} \mathbf{A}_n$$

Since

$$\mathbf{A}_n = -E\frac{\partial U(\theta)}{\partial \theta} = \sum_{i=1}^k \frac{1}{V(\mu_i, \cdot)} \frac{\partial \mu_i}{\partial \theta} \frac{\partial \mu_i}{\partial \theta^T},$$

we have

$$\mathbf{A}_n = \mathbf{D}^T \mathbf{V}^{-1} \mathbf{D}.$$

Therefore C is given by

$$C = \lim_{n_1 \rightarrow \infty} n_1^{-1} \mathbf{D}^T \mathbf{V}^{-1} \mathbf{D}. \quad (3.4.15)$$

The above results can be summarized in the following theorem.

Theorem 3.5: Let C be defined by (3.4.15). Replacing θ_0 by $\hat{\theta}$ in C and denoting the estimate of C by $\hat{C}$.

Let $T = (1/\sqrt{n_1})U_2(\hat{\theta})$ and $Z_T = T/\sqrt{\hat{V}(T)}$, where $\hat{V}(T) = \hat{C}_{22} - \hat{C}_{21}\hat{C}_{11}^{-1}\hat{C}_{12}$ and $\hat{C}$ is obtained by replacing θ_0 in (3.4.15) by $\hat{\theta}$. Under C2-C7, as n_1 and $s \rightarrow \infty$ with $n_1/s \rightarrow c > 0$,

$$Z_T = T/\sqrt{\hat{V}(T)} \rightarrow_d N(0,1), \quad (3.4.16)$$

under the null hypothesis $\beta_1 = 0$.

3.5 Calculation of Score Statistic

In this section, we will derive a useful formula for calculating the standardized score-type statistic. Since the statistic T is given by the left hand-side of the second estimating equation in (3.3.2), T is given by (3.3.6). We only need to replace the parameters $\beta_0, \tau_{10}, \tau_{20}$ by their consistent estimators. This section will be focusing on calculating $Var(T)$. For simplicity we use τ_{20}^* to denote $\tau_{10} + n, \tau_{20} + \tau_{10}\tau_{20}$.

Write

$$\mathbf{A}_n = \mathbf{D}^T \mathbf{V}^{-1} \mathbf{D} = \mathbf{D}_1^T \mathbf{V}_1^{-1} \mathbf{D}_1 + \sum_{j=2}^s \mathbf{D}_j^T \mathbf{V}_j^{-1} \mathbf{D}_j,$$

$$\mathbf{D}_1^T = \begin{bmatrix} n_{10} & n_{11} & \cdots & n_{1k} \\ 0 & n_{11}d_1 & \cdots & n_{1k}d_k \end{bmatrix},$$

and

$$\mathbf{V}_1^{-1} = [\text{diag}[n_{1i}(\beta_0 + \beta_1 d_i)(1 + \tau_1(\beta_0 + \beta_1 d_i))]]^{-1}.$$

Letting $f(d_i) = [(\beta_0 + \beta_1 d_i)(1 + \tau_1(\beta_0 + \beta_1 d_i))]^{-1}$, we have

$$\mathbf{D}_1^T \mathbf{V}_1^{-1} = \begin{bmatrix} f(d_0) & f(d_1) & \cdots & f(d_k) \\ 0 & f(d_1)d_1 & \cdots & f(d_k)d_k \end{bmatrix}.$$

Furthermore, recalling that $\mu_{1i} = \beta_0 + \beta_1 d_i$, we have

$$\mathbf{D}_1^T \mathbf{V}_1^{-1} \mathbf{D}_1 = \begin{bmatrix} \sum_{i=0}^k \frac{n_{1i}}{\mu_{1i}(1+\tau_1\mu_{1i})} & \sum_{i=0}^k \frac{n_{1i}d_i}{\mu_{1i}(1+\tau_1\mu_{1i})} \\ \sum_{i=0}^k \frac{n_{1i}d_i}{\mu_{1i}(1+\tau_1\mu_{1i})} & \sum_{i=0}^k \frac{n_{1i}d_i^2}{\mu_{1i}(1+\tau_1\mu_{1i})} \end{bmatrix},$$

and

$$\begin{aligned} \sum_{j=2}^s \mathbf{D}_j^T \mathbf{V}_j^{-1} \mathbf{D}_j &= \sum_{j=2}^s (n_j, 0)^T (n_j, \beta_0(1 + \tau_{20}\beta_0))^{-1} (n_j, 0) \\ &= \sum_{j=2}^s \begin{bmatrix} n_j^2 / (n_j \beta_0 (1 + \tau_{20} \beta_0)) & 0 \\ 0 & 0 \end{bmatrix}. \end{aligned}$$

Therefore,

$$\begin{aligned} D^T V^{-1} D &= D_1^T V_1^{-1} D_1 + \sum_{j=2}^s D_j^T V_j^{-1} D_j \\ &= \begin{bmatrix} \sum_{i=0}^k \frac{n_{1i}}{\mu_{1i}(1+\tau_{10}\mu_{1i})} + \sum_{j=2}^s \frac{n_j}{\beta_0(1+\tau_{20}^s\beta_0)} & \sum_{i=0}^k \frac{n_{1i}d_i}{\mu_{1i}(1+\tau_{10}\mu_{1i})} \\ \sum_{i=0}^k \frac{n_{1i}d_i}{\mu_{1i}(1+\mu_{1i})} & \sum_{i=0}^k \frac{n_{1i}d_i^2}{\mu_{1i}(1+\tau_{10}\mu_{1i})} \end{bmatrix}, \end{aligned}$$

which is equivalent to the matrix A_{n_1} , the variance-covariance matrix of $(l'_1(\theta_0), l'_2(\theta_0))^T$.

Therefore, by (3.4.14), the variance of $U_2(\hat{\beta}_0, 0)$ is

$$\begin{aligned} \text{Var}(U_2(\hat{\beta}_0, 0)) &= A_{22} - A_{21}A_{11}^{-1}A_{12} \\ &= \sum_{i=0}^k \frac{n_{1i}d_i^2}{\mu_{1i}(1+\tau_{10}\mu_{1i})} - \frac{(\sum_{i=0}^k \frac{n_{1i}d_i}{\mu_{1i}(1+\tau_{10}\mu_{1i})})^2}{\sum_{i=0}^k \frac{n_{1i}}{\mu_{1i}(1+\tau_{10}\mu_{1i})} + \sum_{j=2}^s \frac{n_j}{\beta_0(1+\tau_{20}^s\beta_0)}} \\ &= \frac{1}{\beta_0(1+\tau_{10}\beta_0)} \sum_{i=0}^k n_{1i}d_i^2 - \frac{[\frac{1}{\beta_0(1+\tau_{10}\beta_0)}]^2 (\sum_{i=0}^k n_{1i}d_i)^2}{\frac{1}{\beta_0(1+\tau_{10}\beta_0)} \sum_{i=0}^k n_{1i} + \frac{1}{\beta_0(1+\tau_{20}^s\beta_0)} \sum_{j=2}^s n_j} \\ &= \frac{1}{(\beta_0(1+\tau_{10}\beta_0))^2} [(\beta_0(1+\tau_{10}\beta_0)) \sum_{i=0}^k n_{1i}d_i^2 - \frac{\beta_0(1+\tau_{10}\beta_0)(\sum_{i=0}^k n_{1i}d_i)^2}{n_1 + \frac{1+\beta_0\tau_{10}}{1+\beta_0\tau_{20}^s} \sum_{j=2}^s n_j}]. \quad (3.5.1) \end{aligned}$$

3.6 Some Simple Properties

In the previous sections we developed a score-type statistic Z_T using historical control information to supplement the data from the control group in the current experiment. In this section, we briefly investigate the impact of using the historical data through computer simulation.

Consider first the score-type statistic without using the information from historical experiments. In this case, we use (3.2.1) to describe the moment structure of the data from the current experiment. In addition, only the moment equation (3.3.4) is used to estimate the unknown parameter τ_1 . After dropping the second term of the left hand side in estimating equation (3.3.3), and solving (3.3.3) and (3.3.8) under $H_0: \beta_1 = 0$, we obtain

$$\hat{\beta}_0 = \frac{\sum_{i=0}^k Y_{1i}}{\sum_{i=1}^k n_{1i}}. \quad (3.6.1)$$

We use (3.3.8) to obtain the estimated value of τ_1 . Therefore, (3.6.1) and (3.3.8) give the estimates of $\hat{\beta}_0$ and $\hat{\tau}_{10}$ of β_0 and τ_{10} respectively.

The score-type statistic without the historical control data is then

$$\begin{aligned} T' &= (1/(\sqrt{n_1})U_2(\hat{\beta}_0)) \\ &= \frac{1}{\sqrt{n_1}} \sum_{i=1}^k \frac{(Y_{1i} - n_{1i}\hat{\beta}_0)d_i}{\hat{\beta}_0(1 + \tau_1\hat{\beta}_0)} \end{aligned} \quad (3.6.2)$$

To find the variance of T' , we simply drop all terms involving the historical control data in (3.5.1). (Specifically we omit terms in $A_{22} - A_{21}A_{11}^{-1}A_{12}$). Then, the variance of T' can be written as

$$V(T') = \frac{1}{n_1} \frac{1}{(\hat{\beta}_0(1 + \tau_{10}\hat{\beta}_0))} \left\{ \sum_{i=0}^k n_{1i}d_i^2 - \frac{1}{n_1} \left(\sum_{i=0}^k n_{1i}d_i \right)^2 \right\}. \quad (3.6.3)$$

The standard simplified score statistic is now defined as

$$Z'_T = T' / \sqrt{V(T')}, \quad (3.6.4)$$

which is also asymptotically normally distributed, as $n_1 \rightarrow \infty$. Simulation is used to evaluate the type I and type II errors of both the score statistics Z_T and Z'_T given by (3.4.16) and (3.6.4) respectively. The random numbers are generated from mixed gamma and Poisson distributions using NAG subroutine. First, a random variable is generated from a gamma distribution. Then using this random variable as the mean parameter of a Poisson, a random variable is generated from the Poisson distribution. (The latter random variable actually follows a negative binomial distribution). Iterative methods are used to solve the three equations (3.3.7)-(3.3.9). For the statistic Z'_T developed in this section, the unknown parameters can be obtained from (3.6.1) and (3.3.8).

The type I error rates are estimated based on 1000 simulations with $\beta_1 = 0$ at a nominal significance level of 0.05. The results in Table 3.1 show that the type I errors close to or below the desired 5% for both Z_T and Z'_T .

The type II error rate, also based on 1000 simulations, was obtained for $\beta_1 = 4.25$. As expected, larger sample sizes yield greater power for both Z_T and Z'_T . (Table 3.1). However, Z_T does have better power than Z'_T which does not use historical control data information.

3.7 Summary

In this chapter, a score-type test for trend has been developed to test for increasing trend in mutation rate observed in the Ames assay with increasing dose. Provision was made for extra-Poisson variation within the current experiment that is characteristic of Ames assay data. Additional variation among the historical controls due to inter-study variability was also taken into account. This represents a refinement of a previous result by Leroux et al. (1994), who assumed that the historical and experimental data were subject to the same degree of overdispersion.

With this estimating equation approach to inference, only the first two moments of the data need be specified. The estimators of the regression parameters were obtained from classical estimating equation, and the estimators of the overdispersion parameters were estimated via companion moment equations. The test for trend was also obtained from one of the estimating equations, after replacing the parameters by their consistent estimators.

Under mild regularity conditions, the consistency of the estimated model parameters was established following a similar approach employed by Moore (1985, 1986), and Foutz (1977). The asymptotic normality distribution of the test for trend was obtained using the inverse mapping theorem and Taylor approximations. We also obtained a practical formula for computing the test for trend.

A simulation study was conducted to examine the impact of historical control data. The results indicate that the use of historical data can enhance the power of the test for trend significantly.

Table 3.1 Type I and Type II Error Rate of Test for Trend with and without Historical Controls

Sample Size n_{11} in Current Expt.	Number of Historical Controls	Error Rate
Type I Error ($\beta_0 = 0$)		
15	0	0.045
30	0	0.039
45	0	0.036
15	198	0.048
30	198	0.050
45	198	0.044
Type II Error ($\beta_1 = 4.25$)		
15	0	0.842
15	198	0.640
30	0	0.666
30	198	0.308
45	0	0.527
45	198	0.146

Chapter 4

Optimal Experimental Designs

4.1 Introduction

The information obtained from the Ames *Salmonella* assay can be enhanced through a judicious choice of the experimental design. The experimenter has considerable latitude in the selection of the number of dose groups, the actual dose levels, and the allocation of the available plates to each dose group. In this chapter, we develop optimal experimental designs for the Ames assay.

The theory of optimal experimental design has undergone vigorous development since about 1950, with early contributions by Elfving (1952, 1955, 1959), Chernoff (1953), de la Garza (1954, 1956), Guest (1958) and Kiefer (1958, 1959, 1960). Further developments were reported by Fedorov & Malynov (1972), Whittle (1973) and Kiefer (1974). Optimal designs in assay design and for low-dose extrapolation of carcinogen bioassay data were considered by Krewski et al. (1990), Krewski & Kovar (1982) and Krewski et al. (1984, 1986). The reader is referred to Portier (1991), and Kalish & Rosenberger (1978) for a review of optimal designs for carcinogen bioassay.

In section 4.2, we develop C, A, and D-optimal designs for the linear-exponential model. These optimal designs depend on the values of the model parameters because the model is nonlinear (linear-exponential). The optimal designs all involve only three dose levels, corresponding to the number of parameters in the linear-exponential model.

In section 4.3, we apply the optimal design results in section 4.2 to the collabora

tive trial data of the International Programme on Chemical Safety (IPCS). First, we obtain the C, A, D-optimal designs based on the estimates of the parameter values derived by Krewski et al. (1992). Then, we investigate the efficiency of selected 3-6 dose designs similar to those used in the IPCS study.

In section 4.4, we consider Bayesian approaches to optimal design, using a prior distribution for the values of the unknown model parameters. The Bayesian optimal designs involve only three points, even if the prior distribution of the unknown parameters is diffuse, providing little information about the values of the unknown parameters.

In section 4.5, the minimax criterion proposed by Sitter (1992) for binary dose response models is used to develop optimal designs for the Ames assay. Again, the optimal design involves only three points, even if the prior information on the model parameter is quite uncertain.

A summary of the main findings of the chapter is given in section 4.6.

4.2 C, A, and D-Optimal Designs

The variances of the estimated parameters in a dose response model depend upon the experimental design. An ideal experimental design would minimize the variances of all of the estimated parameters simultaneously. Unfortunately, such a design usually does not exist. We therefore investigate three frequently used optimal design criteria: D, A, and C-optimality.

Consider an experiment with K dose levels $x_1, x_2, \dots, x_K$, with n_i plates assigned to dose x_i ($i = 1, \dots, K$). We let $c_i = n_i/n$ denote the fraction of the total number $n = \sum n_i$ of plates assigned to the i th dose. Let $\mathbf{M}(\mathbf{x}, \mathbf{c})$ be the variance-covariance matrix of the estimated parameters $\hat{\beta}$, where $\mathbf{x} = (x_1, x_2, \dots, x_K)^T$ and $\mathbf{c} = (c_1, c_2, \dots, c_K)^T$. Under D -optimality, $\det \mathbf{M}(\mathbf{x}, \mathbf{c})$ is minimized. D -optimality was originally motivated by reference to the ellipsoidal confidence regions for the parameters of the linear model: a D -optimal design minimizes the content of this confidence region (i.e., the volume of the ellipsoid) (Burridge & Sebastiani (1994)). The A -optimal design minimizes $\text{tr} \mathbf{M}(\mathbf{x}, \mathbf{c})$. Equivalently, it minimizes the sum of

the variances of the parameter estimates. C -optimal designs are used to minimize the variance of a linear combination $\mathbf{l}^T \hat{\boldsymbol{\beta}}$ the model parameter estimates $\hat{\boldsymbol{\beta}}$. Thus, a C -optimal design seeks to minimize $\text{Var}(\mathbf{l}^T \hat{\boldsymbol{\beta}}) = \mathbf{l}^T \mathbf{M}(\mathbf{x}, \mathbf{c}) \mathbf{l}$.

4.2.1 C-Optimal Designs

Consider the linear-exponential dose response model

$$\mu(x) = EY(x) = (\beta_0 + \beta_1 x)e^{-\beta_2 x^2}, \quad (4.2.1)$$

with

$$\text{Var}(Y(x)) = \sigma^2[\mu(x)]^p. \quad (4.2.2)$$

Based on previous work by Snee & Irr (1984) and Krewski et al. (1992), we will take $p = 1.5$ in what follows.

The parameters $\boldsymbol{\beta} = (\beta_0, \beta_1, \beta_2)^T$ may be estimated by solving the estimating equations

$$\mathbf{D}^T \mathbf{V}^{-1}(\mathbf{Y} - \mathbf{EY}) = \mathbf{0}, \quad (4.2.3)$$

where $\mathbf{D} = (\partial\mu/\partial\beta_0, \partial\mu/\partial\beta_1, \partial\mu/\partial\beta_2)$. We wish to find the C -optimal design that minimizes $\text{Var}(\hat{\beta}_1) = \text{Var}(\mathbf{l}^T \hat{\boldsymbol{\beta}})$, where $\mathbf{l} = (0, 1, 0)^T$ and $\hat{\boldsymbol{\beta}} = (\hat{\beta}_0, \hat{\beta}_1, \hat{\beta}_2)^T$.

The model defined by (4.2.1) and (4.2.2) is a non-linear model. The optimal design, in general, depends on the unknown parameters. Chernoff (1953) showed that for large sample sizes, the (locally) optimal designs can be approximated by selecting a certain set of $r \leq k + (k-1) \cdots + (k-s+1)$ of the experiments available, and by repeating each of these r experiments in certain specified proportions. Here, s is the number of parameters involved in the linear combination $\mathbf{l}^T \boldsymbol{\beta}$ and k is the total number of parameters in the model. In our case, $s = 1$ and $k = 3$. Therefore, the maximum number of design points for the optimal design is $r \leq k = 3$. Because there are three parameters in the linear-exponential model, the number of design

points cannot be less than three. As a result, there must be exactly three design points in the C-optimal design.

Let $\theta = (\beta; \sigma)$. For $n_i \rightarrow \infty$ in such a way that $n_i/n \rightarrow c_i > 0$, the asymptotic variance-covariance matrix of $\hat{\beta}$ is

$$\text{Cov}(\hat{\beta}) = I(\theta)^{-1} = [D^T V^{-1} D]^{-1}, \quad (4.2.4)$$

where $I(\theta)$ is the analogous to the information matrix in likelihood theory, and $V = \text{Cov}(Y)$ (Liang and Zeger 1986; Theorem 2). It can be shown that

$$D^T V^{-1} D = \begin{bmatrix} C_{11} & C_{12} & C_{13} \\ C_{21} & C_{22} & C_{23} \\ C_{31} & C_{32} & C_{33} \end{bmatrix}, \quad (4.2.5)$$

where

$$C_{11} = n_0(\sigma^2 \mu^p)^{-1} e^{-2\beta_2 x_0^2} + n_1(\sigma^2 \mu^p)^{-1} e^{-2\beta_2 x_1^2} + n_2(\sigma^2 \mu^p)^{-1} e^{-2\beta_2 x_2^2},$$

$$C_{22} = n_0 x_0^2 e^{-2\beta_2 x_0^2} (\sigma^2 \mu^p)^{-1} + n_1 x_1^2 e^{-2\beta_2 x_1^2} (\sigma^2 \mu^p)^{-1} + n_2 x_2^2 e^{-2\beta_2 x_2^2} (\sigma^2 \mu^p)^{-1},$$

$$C_{33} = n_0 x_0^4 (\beta_0 + \beta_1 x_0)^2 e^{-2\beta_2 x_0^2} (\sigma^2 \mu^p)^{-1} + n_1 x_1^4 (\beta_0 + \beta_1 x_1)^2 e^{-2\beta_2 x_1^2} (\sigma^2 \mu^p)^{-1} \\ + n_2 x_2^4 (\beta_0 + \beta_1 x_2)^2 e^{-2\beta_2 x_2^2} (\sigma^2 \mu^p)^{-1},$$

$$C_{12} = C_{21} = n_0(\sigma^2 \mu^p)^{-1} x_0 e^{-2\beta_2 x_0^2} + n_1(\sigma^2 \mu^p)^{-1} x_1 e^{-2\beta_2 x_1^2} + n_2(\sigma^2 \mu^p)^{-1} x_2 e^{-2\beta_2 x_2^2},$$

$$C_{13} = C_{31} = -n_0(\sigma^2 \mu^p)^{-1} (\beta_0 + \beta_1 x_0) x_0^2 e^{-2\beta_2 x_0^2} - n_1(\sigma^2 \mu^p)^{-1} (\beta_0 + \beta_1 x_1) x_1^2 e^{-2\beta_2 x_1^2} \\ - n_2(\sigma^2 \mu^p)^{-1} (\beta_0 + \beta_1 x_2) x_2^2 e^{-2\beta_2 x_2^2},$$

and

$$C_{23} = C_{32} = -n_0(\sigma^2\mu^p)^{-1}x_0^3(\beta_0 + \beta_1x_0)e^{-2\beta_2x_0^2} - n_1(\sigma^2\mu^p)^{-1}x_1^3(\beta_0 + \beta_1x_1)e^{-2\beta_2x_1^2} \\ - n_2(\sigma^2\mu^p)^{-1}x_2^3(\beta_0 + \beta_1x_2)e^{-2\beta_2x_2^2}.$$

Consider the matrix $\mathbf{A}$ defined by

$$\mathbf{A} = \begin{bmatrix} e^{-\beta_2x_0^2} & e^{-\beta_2x_1^2} & e^{-\beta_2x_2^2} \\ x_0e^{-\beta_2x_0^2} & x_1e^{-\beta_2x_1^2} & x_2e^{-\beta_2x_2^2} \\ -(\beta_0 + \beta_1x_0)x_0^2e^{-\beta_2x_0^2} & -(\beta_0 + \beta_1x_1)x_1^2e^{-\beta_2x_1^2} & -(\beta_0 + \beta_1x_2)x_2^2e^{-\beta_2x_2^2} \end{bmatrix}. \quad (4.2.6)$$

It is easy to verify that

$$I(\theta) = D^T V^{-1} D = \mathbf{A} \mathbf{A}^T, \quad (4.2.7)$$

where

$$\mathbf{A} = \begin{bmatrix} n_{10}(\sigma^2\mu^p(x_0))^{-1} & 0 & 0 \\ 0 & n_{11}(\sigma^2\mu^p(x_1))^{-1} & 0 \\ 0 & 0 & n_{12}(\sigma^2\mu^p(x_2))^{-1} \end{bmatrix}. \quad (4.2.8)$$

Using (4.2.7), we have

$$I(\theta)^{-1} = (\mathbf{A} \mathbf{A}^T)^{-1} = (\mathbf{A}^{-1})^T \mathbf{A}^{-1}. \quad (4.2.9)$$

Letting $\mathbf{A} = [(a_{ij})]$ and $\mathbf{A}^{-1} = [(a^{ij})]$, we may write

$$I(\theta) = \mathbf{A} \mathbf{A}^T$$

$$= \begin{bmatrix} a_{11}n_{10}(\sigma^2\mu^p(x_0))^{-1} & a_{12}n_{11}(\sigma^2\mu^p(x_1))^{-1} & a_{13}n_{12}(\sigma^2\mu^p(x_2))^{-1} \\ a_{21}n_{10}(\sigma^2\mu^p(x_0))^{-1} & a_{22}n_{11}(\sigma^2\mu^p(x_1))^{-1} & a_{23}n_{12}(\sigma^2\mu^p(x_2))^{-1} \\ a_{31}n_{10}(\sigma^2\mu^p(x_0))^{-1} & a_{32}n_{11}(\sigma^2\mu^p(x_1))^{-1} & a_{33}n_{12}(\sigma^2\mu^p(x_2))^{-1} \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

$$= \left[\left(\sum_{i=1}^3 a_{ri} n_{i-1} (\mu^p(x_{i-1}) \sigma^2)^{-1} a_{si} \right) \right]. \quad (4.2.10)$$

Further,

$$\begin{aligned} I(\theta)^{-1} &= (\mathbf{A}^{-1})^T \mathbf{A}^{-1} \mathbf{A}^{-1} \\ &= \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \begin{bmatrix} \frac{(\sigma^2 \mu^p(x_0))}{n_{10}} & 0 & 0 \\ 0 & \frac{(\sigma^2 \mu^p(x_1))}{n_{11}} & 0 \\ 0 & 0 & \frac{(\sigma^2 \mu^p(x_2))}{n_{12}} \end{bmatrix} \begin{bmatrix} a^{11} & a^{12} & a^{13} \\ a^{21} & a^{22} & a^{23} \\ a^{31} & a^{32} & a^{33} \end{bmatrix} \\ &= \left[\left(\sum_{i=1}^3 a^{ir} \frac{\sigma^2 \mu^p(x_{i-1})}{n_{i-1}} a^{is} \right) \right]. \end{aligned} \quad (4.2.11)$$

We now have

$$\text{Var}(\hat{\beta}_1) = \mathbf{l}^T \mathbf{I}^{-1}(\theta) \mathbf{l} = \sum_{i=1}^3 a^{i2} \frac{[\sigma^2 \mu^p(x_{i-1})]}{n_{i-1}} a^{i2}, \quad (4.2.12)$$

where $\mathbf{l} = (0, 1, 0)^T$. Using the Cauchy inequality,

$$\begin{aligned} \text{Var}(\sqrt{n} \hat{\beta}_1) &= \sum_{i=1}^3 a^{i2} \frac{[\sigma^2 \mu^p(x_{i-1})]}{c_i} a^{i2} \\ &= \left(\sum_{i=1}^3 \frac{(a^{i2})^2 ([\sigma^2 \mu^p(x_{i-1})]^{1/2})^2}{(\sqrt{c_i})^2} \right) \left(\sum_{i=1}^3 (\sqrt{c_i})^2 \right) \\ &\geq \left\{ \sum_{i=1}^3 \frac{a^{i2} [\sigma^2 \mu^p(x_{i-1})]^{1/2}}{\sqrt{c_i}} \sqrt{c_i} \right\}^2 \\ &= \left\{ \sum_{i=1}^3 a^{i2} [\sigma^2 \mu^p(x_{i-1})]^{1/2} \right\}^2, \end{aligned} \quad (4.2.13)$$

where equality holds if and only if

$$c_i = \frac{a^{i2} \mu^{p/2}(x_{i-1})}{\sum_{i=1}^3 a^{i2} \mu^{p/2}(x_{i-1})} \equiv c_i^*. \quad (4.2.14)$$

Note that the optimal allocation c_i^* is independent of the parameter σ involved in the variance function in (4.2.2). We now illustrate the development of a C-optimal design for the Ames assay. Consider, for example, the linear-exponential

dose response model in (4.2.1) with parameter values $\beta_0 = 146$, $\beta_1 = 273\mu g^{-1}$, and $\beta_2 = 0.033$. These parameters correspond to those in a particular experiment with urban air particulate with TA100+S9 in the International Programme on Chemical Safety (Krewski et al., 1993).

Suppose that the allowable dose range is 0-6 μg , as in the original design. The C-optimal design is found using the following search routines.

- (1) Divide the interval $[0, 6]$ into M subintervals of equal length $h = 6/M$. Let $x_0 = k_1 * h$, $x_1 = k_2 * h$, and $x_3 = k_3 * h$, where k_1 , k_2 , and k_3 are non-negative integers.
- (2) Let x_0, x_1, x_2 assume all possible values $x_i = k_i * h$ ($i = 1, 2, 3$). We require $0 \leq k_1 < k_2 < k_3 \leq M$ when searching through all the possible values for x_0 , x_1 , and x_2 . It can be seen that $0 \leq k_1 \leq M - 2$, $1 \leq k_2 \leq M - 1$, and $2 \leq k_3 \leq M$.
- (3) At each step, the value of $Var(\hat{\beta}_1)$ is calculated as follows. Given $\mathbf{x} = (x_0, x_1, x_2)$, we first calculate the matrix $\mathbf{A}$, then use the NAG (1991) package F01AAF to find $\mathbf{A}^{-1}$. Then the right hand side of (4.2.13) is evaluated by replacing β_i by $\hat{\beta}_i$ ($i = 0, 1, 2$).
- (4) After searching through all of the possible values of x_0 , x_1 , and x_2 , we find the optimal values $\mathbf{x}^* = (x_0^*, x_1^*, x_2^*)$ that minimize $Var(\hat{\beta}_1)$.

As a second example, we also consider a linear-exponential model with $\beta_0 = 162$, $\beta_1 = 354\mu g^{-1}$, and $\beta_2 = 0.0639$. The parameters estimated are also obtained from a second experiment with the urban air particulate with TA100+S9 in the International Programme on Chemical Safety (Krewski et al., 1993). Table 4.1 shows the optimal designs for these two examples obtained using this procedure.

Note that in both cases, the lowest dose in the optimal design is zero (i.e., the control group) and the highest dose is equal to the maximum dose within the allowable dose range. The majority of the observations (over 71%) are taken at middle dose.

Table 4.1: Two Examples of C-Optimal Designs

Example	Number of Subintervals	$Var(\hat{\beta}_1)$	Optimal Doses			Optimal Allocations		
I^1	M	(μg^{-2})	x_0^*	x_1^*	x_2^*	c_0^*	c_1^*	c_2^*
	M=150	112.10	0.00	2.12	6.00	0.20	0.71	0.09
	M=90	112.12	0.00	2.13	6.00	0.19	0.71	0.09
II^2	M=90	136.7	0.00	1.93	6.00	0.19	0.72	0.09
	M=60	136.9	0.00	1.91	6.00	0.19	0.72	0.09

¹ Example I: $\beta_0 = 146$, $\beta_1 = 273$, $\beta_2 = 0.033$.

² Example II: $\beta_0 = 162$, $\beta_1 = 354$, $\beta_2 = 0.064$.

4.2.2 A-Optimal Designs

The A-optimal design seeks to minimize $Var(\hat{\beta}_0) + Var(\hat{\beta}_1) + Var(\hat{\beta}_2)$. Following Chernoff(1972), the number of optimal design points r is in the range $3 \leq r \leq k + (k - 1) + k - 3 + 1 = 6$. Therefore, the A-optimal design is a 3-6 point design.

In section 4.2.1, we obtained the variance-covariance matrix $I(\vartheta)^{-1}$ of $(\hat{\beta}_0, \hat{\beta}_1, \hat{\beta}_2)^T$. Therefore, we have

$$\begin{aligned}
 Var(\sqrt{n}\hat{\beta}_0) &+ Var(\sqrt{n}\hat{\beta}_1) + Var(\sqrt{n}\hat{\beta}_2) \\
 &= \sum_{i=1}^3 a^{i1} \frac{[\sigma^2 \mu^p(x_{i-1})]}{c_i} a^{i1} + \sum_{i=1}^3 a^{i2} \frac{[\sigma^2 \mu^p(x_{i-1})]}{c_i} a^{i2} \\
 &\quad + \sum_{i=1}^3 a^{i3} \frac{[\sigma^2 \mu^p(x_{i-1})]}{c_i} a^{i3} \\
 &= \sum_{i=1}^3 \frac{1}{(\sqrt{c_i})^2} \{ [\sum_{s=1}^3 (a^{is})^2 [\sigma^2 \mu^p(x_{i-1})]]^{1/2} \}^2 [\sum_{i=1}^3 (\sqrt{c_i})^2] \\
 &\geq \{ \sum_{i=1}^3 (\sum_{s=1}^3 (a^{is})^2 [\sigma^2 \mu^p(x_{i-1})])^{1/2} \}^2,
 \end{aligned}$$

where equality holds whenever

$$c_i = \frac{[\sum_{s=1}^3 (a^{is})^2 \mu^p(x_{i-1})]^{1/2}}{\sum_{i=1}^3 [\sum_{s=1}^3 (a^{is})^2 \mu^p(x_{i-1})]^{1/2}} \equiv c_i^*. \quad (4.2.15)$$

Numerical methods were used to find the A-optimal designs for the two examples

Table 4.2: Two Examples of A-Optimal Designs

Example	Number of Subintervals	$\text{Min } \sum_{i=0}^2 \text{Var}(\hat{\beta}_i)$	Optimal Doses			Optimal Allocation		
	M	μg^{-2}	x_0^*	x_1^*	x_2^*	c_0^*	c_1^*	c_2^*
I^1	M=60	137.1	0.00	2.00	6.00	0.35	0.58	0.07
	M=90	137.1	0.00	1.93	6.00	0.35	0.59	0.06
II^2	M=60	162.7	0.00	1.80	6.00	0.33	0.60	0.07
	M=90	162.6	0.00	1.80	6.00	0.33	0.60	0.07

¹ Example I: $\beta_0 = 146$, $\beta_2 = 273$, $\beta_2 = 0.033$.

² Example II: $\beta_0 = 162$, $\beta_1 = 354$, $\beta_2 = 0.0639$.

considered previously. Although 3-6 point designs were considered, it was found that the 4, 5, and 6 point designs all degenerated to a 3 point design. Consequently, we may use (4.2.15) to find A-optimal designs with three design points.

The A-optimal designs for our two examples are given in Table 4.2. As with the C-optimal designs, the lowest and highest doses in the A-optimal design correspond to zero and the maximum possible dose respectively; most of the observations are also assigned to the intermediate dose.

4.2.3 D-Optimal Designs

The general theory of D-optimality, via the Caratheodory theorem (Silvey, 1980), shows that there will be 3-6 design points in the D-optimal design when the expected response is a linear function of the unknown parameters. In the nonlinear case, a Taylor expansion can be used to expand the response function about the true values of the regression parameters. After dropping the higher order terms, approximate optimal designs can be obtained using the result for linear models.

To find the actual optimal design in the nonlinear case, the values of the regression parameter values must be known. These optimal designs are called locally optimal, because they apply only in a small neighbourhood of the true values of

the regression parameters. Unlike the linear case, the designs cannot be uniformly optimal for the entire range of the regression parameters.

Before investigating the linear-exponential model, we examine a quadratic dose response model which is linear in the model parameters. There are two reasons for investigating the quadratic model. First, the linear-exponential model can be well approximated by a quadratic model within the dose range of interest. Second, well-developed results due to Fedorov (1972) can be applied directly to the quadratic model. The results obtained for the quadratic model may provide some guidance on the linear-exponential model. Specifically, consider the quadratic form

$$\mu_i = E(Y_i|x_i) = \alpha_1 + \alpha_2 x_i + \alpha_3 x_i^2,$$

with

$$Var(Y_i|x_i) = \mu_i + \mu_i^2/c.$$

This will allow us to determine exactly the number of optimal design points theoretically.

Suppose $E(Y_i|x_i) = \sum_{t=1}^m \alpha_t x_i^{t-1}$ and $Var(Y_i|x_i) = \phi[\lambda(x_i)]^{-1}$, where $\lambda(x) = [P(x)]^{-1}$ and $P(x)$ is a positive polynomial of degree no higher than $2(m-1)$ defined on the domain $[a, b]$. It then follows from Theorem 2.3.2 of Fedorov (1972) that the D-optimal design is concentrated at m points.

In our case, $m = 3$, $\phi = 1$ and $P(x) = \mu_i(x) + \mu_i^2(x)/c$. Therefore, $P(x)$ is a polynomial of degree no higher than 4. Note that $2(m-1) = 2(3-1) = 4$ and $P(x) > 0$ on $[0, a]$, where $a > 0$ denotes the highest dose level in the dose response function. We conclude that the D-optimal design for the quadratic model involves only $m = 3$ points.

We now derive a formula to calculate the determinant of the "information" matrix $I(\theta)$. Let

$$D^T = \left[\frac{\partial \mu}{\partial \alpha_i} \right] = \begin{bmatrix} 1 & 1 & 1 \\ x_0 & x_1 & x_2 \\ x_0^2 & x_1^2 & x_2^2 \end{bmatrix}_{3 \times 3},$$

and

$$V^{-1} = \text{diag}([\sigma^2 \mu^p(x_{i-1})]^{-1}).$$

We then have

$$D^T V^{-1} D = \begin{bmatrix} Q_{11} & Q_{12} & Q_{13} \\ Q_{21} & Q_{22} & Q_{23} \\ Q_{31} & Q_{32} & Q_{33} \end{bmatrix},$$

where

$$Q_{11} = n_{10}[\sigma^2 \mu^p(x_0)]^{-1} + n_{11}[\sigma^2 \mu^p(x_1)]^{-1} + n_{12}[\sigma^2 \mu^p(x_2)]^{-1},$$

$$Q_{12} = Q_{21} = n_{10}x_0[\sigma^2 \mu^p(x_0)]^{-1} + n_{11}x_1[\sigma^2 \mu^p(x_1)]^{-1} + n_{12}x_2[\sigma^2 \mu^p(x_2)]^{-1},$$

$$Q_{13} = Q_{31} = n_{10}x_0^2[\sigma^2 \mu^p(x_0)]^{-1} + n_{11}x_1^2[\sigma^2 \mu^p(x_1)]^{-1} + n_{12}x_2^2[\sigma^2 \mu^p(x_2)]^{-1},$$

$$Q_{23} = Q_{32} = n_{10}x_0^3[\sigma^2 \mu^p(x_0)]^{-1} + n_{11}x_1^3[\sigma^2 \mu^p(x_1)]^{-1} + n_{12}x_2^3[\sigma^2 \mu^p(x_2)]^{-1},$$

and

$$Q_{33} = n_{10}x_0^4[\sigma^2 \mu^p(x_0)]^{-1} + n_{11}x_1^4[\sigma^2 \mu^p(x_1)]^{-1} + n_{12}x_2^4[\sigma^2 \mu^p(x_2)]^{-1}.$$

It is easy to verify that

$$I(\hat{\beta}) = D^T V^{-1} D = A \Lambda A^T,$$

where

$$A = \begin{bmatrix} 1 & 1 & 1 \\ x_0 & x_1 & x_2 \\ x_0^2 & x_1^2 & x_2^2 \end{bmatrix}$$

and

$$\Lambda = \begin{bmatrix} n_{10}[\sigma^2 \mu^p(x_0)]^{-1} & 0 & 0 \\ 0 & n_{11}[\sigma^2 \mu^p(x_1)]^{-1} & 0 \\ 0 & 0 & n_{12}[\sigma^2 \mu^p(x_2)]^{-1} \end{bmatrix}.$$

Thus,

$$\begin{aligned} \det I(\hat{\beta}) &= \det(D^T V^{-1} D) \\ &= \det(A \Lambda A^T) \\ &= (\det A)^2 [\sigma^2 \mu^p(x_0)]^{-1} [\sigma^2 \mu^p(x_1)]^{-1} [\sigma^2 \mu^p(x_2)]^{-1} n_0 n_1 n_2. \end{aligned}$$

Since $n_0 + n_1 + n_2 = n$ is constant, $\det \mathbf{I}(\hat{\beta})$ attains its maximum value when $n_0 = n_1 = n_2$, given the fixed design points x_0 , x_1 , and x_2 . Thus, the D-optimal design involves equal allocation.

To find the optimum dose levels, we need to minimize $\det \mathbf{I}(\hat{\beta})^{-1}$ or maximize $\det \mathbf{I}(\hat{\beta})$. This can be done using numerical methods to find maximum or minimum values of a function of three variables x_0 , x_1 , and x_2 .

We now consider 3 point D-optimal designs for the linear-exponential model. It follows from section 4.2.1 that

$$\det \mathbf{I}(\hat{\beta}) = \det(\mathbf{A} \mathbf{\Lambda} \mathbf{A}^T) = (\det \mathbf{A})^2 \det \mathbf{\Lambda},$$

where $\mathbf{A}$ and $\mathbf{\Lambda}$ are given in (4.2.6) and (4.2.8) respectively. Therefore,

$$\det \mathbf{I}(\hat{\beta}) = (\det \mathbf{A})^2 (\sigma^2 \mu^p(x_0))^{-1} (\sigma^2 \mu^p(x_1))^{-1} (\sigma^2 \mu^p(x_2))^{-1} n_0 n_1 n_2. \quad (4.2.17)$$

Since $n_0 + n_1 + n_2 = n$, the number of observations at each design point is $n_0 = n_1 = n_2 = n/3$.

Numerical methods can now be used to find optimal design points x_0^* , x_1^* , and x_2^* such that $\det \mathbf{I}(\hat{\beta})$ is maximized. D-optimal designs for both the quadratic dose response model $\mu(x) = \alpha_1 + \alpha_2 x + \alpha_3 x^2$ and the linear-exponential dose response model $\mu(x) = (\beta_0 + \beta_1 x) e^{-\beta_2 x^2}$ for our two examples are shown in Table 4.3.

The parameter α in the quadratic was estimated by fitting the model to the original data using GEEs as in (4.2.3). Note that the two models give identical D-optimal designs. Recall that the D-optimal designs for the quadratic model (which is linear in the parameters α_0 , α_1 and α_2) have exactly three design points (Fedorov, 1972). For the linear-exponential model, which is nonlinear in the parameters β_0 , β_1 and β_2 , no theoretical results on the number of points in the D-optimal design are available. However, our numerical results suggest that the D-optimal designs under the linear-exponential model must also have three design points. Thus, the D-optimal designs for the quadratic model may be used as a guide to studying D-optimal designs for the linear-exponential model.

Table 4.3: Two Examples of D-Optimal Designs

Example	Dose Response Model	Number of Subintervals M	Optimal Doses			Optimal Allocation		
			x_0^*	x_1^*	x_2^*	c_0^*	c_1^*	c_2^*
Ia^1	Quadratic	$M = 90$	0.0	2.2	6.0	0.333	0.333	0.333
Ib^2	Linear-exponential	$M = 90$	0.0	2.0	6.0	0.333	0.333	0.333
IIa^3	Quadratic	$M = 90$	0.0	2.86	6.0	0.333	0.333	0.333
IIb^4	Linear-exponential	$M = 90$	0.0	2.3	6.0	0.333	0.333	0.333

¹ Example Ia: $\alpha_1 = 145$, $\alpha_2 = 305$, $\alpha_3 = -40$.

² Example Ib: $\beta_0 = 146$, $\beta_1 = 273$, $\beta_2 = 0.033$.

³ Example IIa: $\alpha_1 = 169$, $\alpha_2 = 339$, $\alpha_3 = 55.3$.

⁴ Example IIb: $\beta_0 = 162$, $\beta_1 = 354$, $\beta_2 = 0.064$.

4.3 Suboptimal Designs

In the previous section, we derived 3-dose C, A, and D-optimal designs for the Ames assay. In practice, the application of such optimal designs is not straightforward for several reasons (Krewski & Goddard, 1990). First, construction of an optimal design requires knowledge of the parameters of the underlying stochastic model from which the data arise. Second, since optimal designs usually involve a relatively small number of critically placed doses, poor dose selection or the loss of an entire dose group may compromise the integrity of the original design. Third, the use of optimal designs with only three doses does not provide any degrees of freedom to test for lack of fit. Fourth, designs that are optimal for one purpose may not be optimal for another. Fifth, optimal designs may simply be unworkable because of operational constraints, such as limits on the range of doses that may be employed. Nonetheless, optimal designs do offer considerable insight into the development of practical experimental designs, and serve as a benchmark against which the efficiencies of suboptimal designs may be evaluated.

In this section, we consider suboptimal designs with more than three doses that

may be more suitable for use in practice. In particular, we consider suboptimal designs with 4-6 doses, patterned loosely after those used in a recent collaborative trial involving the Ames assay sponsored by the International Programme on Chemical Safety (IPCS) (Lewtas et al., 1992). Designs with both equal and unequal allocations are examined. In an attempt to develop recommendations for designs that might be used in practice, we consider the efficiency of a number of fixed suboptimal designs involving more than three doses under conditions observed in the IPCS collaborative trial. This analysis is useful in evaluating the efficiency of suboptimal designs under conditions that may be observed in practice.

4.3.1 Efficiency of Suboptimal Designs

Consider a design with $K > 3$ dose levels, say $\mathbf{x} = (x_0, x_1, \dots, x_{K-1})$. As in section 4.2, the unknown model parameter β is estimated by solving the estimating equation (4.2.3), and the variance of $\hat{\beta}$ is

$$\text{Var}(\hat{\beta}) = (\mathbf{D}^T \mathbf{V}^{-1} \mathbf{D})^{-1}, \quad (4.3.1)$$

where $\mathbf{D} = \partial \mu / \partial \beta$.

It can be shown that

$$\mathbf{D}^T \mathbf{V}^{-1} \mathbf{D} = \begin{bmatrix} C'_{11} & C'_{12} & C'_{13} \\ C'_{21} & C'_{22} & C'_{23} \\ C'_{31} & C'_{32} & C'_{33} \end{bmatrix}, \quad (4.3.2)$$

where

$$C'_{11} = \sum_{i=0}^{K-1} n_i (\sigma^2 \mu^p)^{-1} e^{-2\beta_2 x_i^2},$$

$$C'_{12} = C'_{21} = \sum_{i=0}^{K-1} n_i (\sigma^2 \mu^p)^{-1} e^{-2\beta_2 x_i^2} x_i,$$

$$C'_{13} = C'_{31} = - \sum_{i=0}^{K-1} n_i (\sigma^2 \mu^p)^{-1} e^{-2\beta_2 x_i^2} (\beta_0 + \beta_1 x_i) x_i^2,$$

$$C'_{22} = \sum_{i=0}^{K-1} n_i (\sigma^2 \mu^p)^{-1} e^{-2\beta_2 x_i^2} x_i^2,$$

$$C'_{23} = C'_{32} = - \sum_{i=0}^{K-1} n_i (\sigma^2 \mu^p)^{-1} e^{-2\beta_2 x_i^2} (\beta_0 + \beta_1 x_i) x_i^3,$$

and

$$C'_{33} = \sum_{i=0}^{K-1} n_i (\sigma^2 \mu^p)^{-1} e^{-2\beta_2 x_i^2} (\beta_0 + \beta_1 x_i)^2 x_i^4.$$

In contrast with section 4.2, the matrix $\mathbf{D}^T \mathbf{V}^{-1} \mathbf{D}$ cannot be written in the form $\mathbf{A} \mathbf{A} \mathbf{A}^T$, where $\mathbf{A}$ is a diagonal matrix, and the n_i ($i = 0, \dots, K-1$) only appear in $\mathbf{A}$. Therefore, the Cauchy inequality cannot be used to find the optimal allocations when the number of design points exceeds three. This will increase the number of variables involved in numerical searches to find the optimal dose levels and optimal allocation in designs with more than three dose groups.

We have demonstrated in section 4.2 that the optimal designs for C, A, and D-criteria are all 3-point designs. We have also shown that for the 3-point optimal designs, the C, A, and D-criteria can be achieved by minimizing the objective functions (4.2.13), (4.2.15) and (4.2.16) respectively. Let $C3$, $A3$ and $D3$ be the minimum values of these three criteria respectively. For an M -point experimental design, the variance-covariance matrix of the estimated parameters is given by (4.3.1). Let a_{ij} denote the (i, j) th element of the matrix (4.3.1). Further, let CM , AM and DM denote the minimum value of a_{22} , $a_{11} + a_{22} + a_{33}$, and $\det[a_{ij}]$ respectively. Then, the design efficiencies under the C, A, and D-criteria are defined as follows.

$$\text{C-design efficiency: } EC = \frac{C3}{CM}. \quad (4.3.3)$$

$$\text{A-design efficiency: } EA = \frac{A3}{AM}. \quad (4.3.4)$$

$$\text{D-design efficiency: } ED = \frac{D3}{DM}. \quad (4.3.5)$$

Using these criteria, the 3-point optimal designs are considered as benchmarks, against which the efficiencies of the M -point designs are evaluated. In the remainder of this section, efficiencies of the designs used in the IPCS study are investigated.

4.3.2 Designs in the IPCS Study

We now develop 3-dose optimal designs for a series of 1120 linear-exponential dose response models derived from a collaborative trial sponsored by the International Programme on Chemical Safety (Lewtas et al., 1992). Since the optimal designs depend on the unknown model parameters, we use the parameter estimates obtained in the collaborative trial to construct C, A, and D-optimal designs. A representative optimal design is then computed by averaging the dose levels and the allocations for the 1120 optimal designs obtained in this manner. These representative three dose designs can then be used in practice in the absence of knowledge of the values of the parameters in the underlying linear-exponential model. The efficiencies (4.3.3)-(4.3.5) of these representative designs relative to the optimal designs is investigated.

Twenty laboratories worldwide participated in a collaborative trial sponsored by the International Programme on Chemical Safety on the mutagenicity of complex mixtures as expressed in the *Salmonella*/microsome assay. The U.S. National Institute of Standards and Technology provided homogeneous reference samples of urban air and diesel particles and a coal tar solution to each participating laboratory, along with samples of benzo[a]pyrene and 1-nitropyrene which served as positive controls. Mutagenic potency was characterized by the slope of the initial linear component of the dose-response curve. Krewski et al. (1992) have obtained estimates of the regression parameters β_0 , β_1 , and β_2 for all 1120 individual experiments in the IPCS study. Since optimal designs for the linear-exponential depend on the unknown parameters β_0 , β_1 , and β_2 , we will develop optimal designs using the estimates $\hat{\beta}_0$, $\hat{\beta}_1$, and $\hat{\beta}_2$ of the true parameters.

The 3-dose C-optimal designs based on the results of the IPCS collaborative trial are summarized in table 4.4. In order to render the designs developed for different test agents more comparable, the highest dose has been scaled to unity throughout.

For each test agent, the average doses and the average allocations to those doses are reported for each strain of *Salmonella* (TA98 or TA100) and activation condition (+/- S9). Average optimal designs are also reported for each test agent and for all test agent combined.

The C-optimal designs are remarkably consistent across test agents, strain of *Salmonella* used, and activation conditions. Based on the overall average, a 3-dose design with doses in the ratio 0:0.4:1.0 (with the highest dose scaled to unity) and an allocation of 0.2, 0.7 and 0.1 to these three doses might be considered as somewhat representative of the individual three dose optimal designs.

D-optimal designs are given in Table 4.5. The D-optimal designs are highly consistent across test agents, strain of *Salmonella* used, and activation conditions. The overall average shows that a three dose design with dose spread in the ratio 0:0.4:1.0 and equal allocations of 0.33, 0.33 and 0.33 to three doses are representative of the individual three dose optimal designs.

A-optimal designs are given in Table 4.6. These designs are also quite consistent across test agents, strain of *Salmonella* used, and activation conditions. Overall, a three dose design with doses in the ratio 0:0.4:1.0 and allocations of 0.25, 0.65, and 0.1 to the three doses might be considered as somewhat representative of the individual three dose A-optimal designs.

The results in Tables 4.4-4.6 show that the optimal dose levels and the optimal allocations to these doses are quite comparable for the five chemical mixtures. Therefore, the average (weighted by number of experiments) of the dose levels and allocations are considered as the recommended empirical optimal designs. These empirical results show that the optimal designs are not highly sensitive to the values of the regression parameters.

The efficiencies of the recommended three dose designs and that of the original six dose design with equal allocation used in the IPCS study are given in Tables 4.7-4.9 for the C, D and A-criteria respectively. Whereas the original 6-dose design is, on average, only 57% as efficient as the C-optimal design, the representative 3-dose design enjoys an average relative efficiency of nearly 89% (Table 4.7). From Table 4.8, we see that the original 6-dose design is, on average, only 45% as efficient

as the D-optimal design, the representative 3-dose design has an average relative efficiency of over 97%. Table 4.9 shows that the original 6-dose design is, on average, only about 60% as efficient as the A-optimal design; however, the efficiency of the representative 3-dose design is over 87%.

These results indicate that the original 6-dose designs are somewhat inefficient relative to the optimal 3-dose designs. However, the representative 3-dose designs, which do not require knowledge of the unknown model parameters, enjoy relatively high efficiency.

Since the experimenters may be reluctant to employ a design with only three doses for reasons outlined in section 4.1, we now consider selected fixed designs with 4-6 doses. Specifically, we consider designs with 4-6 fixed doses, as well as the representative three dose designs discussed previously.

The designs with more than three doses were obtained by adding additional doses to the representative three dose designs. In addition to the unequal allocations shown above, designs with equal allocations will also be considered. The unequal allocations shown above were obtained by optimizing the allocation to the fixed doses separately for each of the 1120 experiments in the IPCS collaborative trial, and then averaging these allocations across the set of 1120 experiments. This optimal allocation was done under the constraint that no less than 10% of the total sample be allocated to any single dose. This constraint is imposed since the number of subjects might otherwise be allocated almost entirely to the three doses in the three dose representative design.

The efficiencies of designs with 3-6 fixed dose levels and equal allocation are shown in Table 4.10, Table 4.13 and Table 4.16 under the C-, D- and A-optimality criteria respectively. The results indicate that, in general, the efficiency of these fixed designs decreases as the number of dose levels increases. Nonetheless, the 6-dose design has an average efficiency of about 57% under all three criteria. Among the 3-6 dose fixed designs with equal allocation, the 4-dose designs have the highest average efficiencies (nearly 70%) under both the C and A-optimality criteria. Under D-optimality, the 3-dose designs enjoy the highest average efficiency (92.5%) (See Table 4.4-4.9 and Table 4.10, Table 4.13, and Table 4.16).

Optimal allocations for the fixed 3-6 dose designs with unequal allocation are shown in Table 4.11, Table 4.14, and Table 4.17. The optimal allocations for each test agent and the overall weighted average allocation under C, D and A-optimality are shown in Table 4.11, Table 4.14 and Table 4.17 respectively. The optimal allocations under the C and A-optimality criteria are similar. about 20% observations allocated to the zero dose level with about 10% allocated to the highest dose level. Under D-optimality, however, 30% of the subjects are assigned to the zero dose and the highest dose, with the remaining dose levels receive approximately equal allocations.

The results in Table 4.11, Table 4.14 and Table 4.17 indicate that unequal allocation leads to greater efficiency than equal allocation. Therefore, the efficiency of fixed dose designs based on these average optimal allocations shown in Table 4.11, Table 4.14, and Table 4.17 were computed. The results are presented in Table 4.12, Table 4.15, and Table 4.18 for the C, D and A-optimality respectively. These results show that the larger the number of fixed dose levels used, the lower the efficiency of the designs. Furthermore, optimal designs with unequal allocations enjoy an advantage in relative efficiency of about 10% compared to designs with equal allocations.

The efficiencies of these fixed 3-6 dose designs with either equal or unequal allocation are summarized in Table 4.19. These summary results reflect the following general findings noted previously.

1. The efficiency of the fixed designs considered here generally decreases as the number of dose levels increases.
2. The fixed designs considered here with equal allocation are notably less efficient than those with unequal allocation.
3. In the unequal allocation situation, the allocation ratios under the C and A-optimality are close to each other. In particular, 10% of the observations are at the highest dose level, and 20 – 30% are at the zero dose level. Under D-optimality, however, both the highest and the zero dose level have 30% of the observations.

Table 4.4 C - Optimal Designs for the Ames Assay

Test Agent	Strain +/- S9	Number of Expts.	Optimal Doses ^a			Max. Dose	Optimal Allocation		
			d_0	d_1	d_2		c_0	c_1	c_2
Coal Tar Solution	TA100+S9	40	0.000	0.408	1.000	5.0 μ g	0.249	0.649	0.101
	TA100-S9	32	0.000	0.489	1.000	10.0 μ g	0.353	0.522	0.125
	TA98+S9	40	0.000	0.387	1.000	5.0 μ g	0.219	0.689	0.092
	TA98-S9	38	0.000	0.461	1.000	10.0 μ g	0.317	0.566	0.117
	All ^b	150	0.000	0.433	1.000	-	0.280	0.612	0.108
Urban Air Particles	TA100+S9	79	0.000	0.465	1.000	2.0mg	0.311	0.575	0.115
	TA100-S9	80	0.000	0.465	1.000	2.0mg	0.314	0.570	0.116
	TA98+S9	80	0.000	0.320	1.000	8.0mg	0.106	0.827	0.067
	TA98-S9	80	0.000	0.286	1.000	10.0mg	0.078	0.861	0.062
	All ^b	319	0.000	0.383	1.000	-	0.202	0.708	0.090
Diesel Particles	TA100+S9	80	0.000	0.407	1.000	0.2mg	0.237	0.667	0.097
	TA100-S9	80	0.000	0.414	1.000	0.2mg	0.243	0.659	0.098
	TA98+S9	80	0.000	0.338	1.000	0.5mg	0.137	0.789	0.074
	TA98-S9	79	0.000	0.316	1.000	0.5mg	0.107	0.827	0.067
	All ^b	319	0.000	0.369	1.000	-	0.181	0.735	0.084
Benzopyrene	TA100+S9	79	0.000	0.406	1.000	2.0 μ g	0.238	0.664	0.098
	TA98+S9	79	0.000	0.379	1.000	2.0 μ g	0.198	0.714	0.087
	All ^b	158	0.000	0.393	1.000	-	0.218	0.689	0.092
1-Nitropyrene	TA100-S9	76	0.000	0.350	1.000	6.0 μ g	0.209	0.693	0.098
	TA98-S9	80	0.000	0.389	1.000	0.1 μ g	0.206	0.705	0.089
	All ^b	156	0.000	0.370	1.000	-	0.207	0.699	0.093
ALL	All ^b	1102	0.000	0.385	1.000	-	0.210	0.699	0.091

a: Highest dose scaled to unity.

b: Average weighted by the number of experiments.

Table 4.5 D - Optimal Designs for the Ames Assay

Test Agent	Strain +/- S9	Number of Expts.	Optimal Doses ^a			Max. Dose	Optimal Allocation		
			d_0	d_1	d_2		c_0	c_1	c_2
Coal Tar Solution	TA100+S9	40	0.000	0.415	1.000	5.0 μ g	0.333	0.333	0.333
	TA100-S9	32	0.000	0.490	1.000	10.0 μ g	0.333	0.333	0.333
	TA98+S9	40	0.000	0.399	1.000	5.0 μ g	0.333	0.333	0.333
	TA98-S9	38	0.000	0.464	1.000	10.0 μ g	0.333	0.333	0.333
	All ^b	150	0.000	0.439	1.000	-	0.333	0.333	0.333
Urban Air Particles	TA100+S9	79	0.000	0.468	1.000	2.0mg	0.333	0.333	0.333
	TA100-S9	80	0.000	0.467	1.000	2.0mg	0.333	0.333	0.333
	TA98+S9	80	0.000	0.354	1.000	8.0mg	0.333	0.333	0.333
	TA98-S9	80	0.000	0.327	1.000	10.0mg	0.333	0.333	0.333
	All ^b	319	0.000	0.404	1.000	-	0.333	0.333	0.333
Diesel Particles	TA100+S9	80	0.000	0.415	1.000	0.2mg	0.333	0.333	0.333
	TA100-S9	80	0.000	0.420	1.000	0.2mg	0.333	0.333	0.333
	TA98+S9	80	0.000	0.366	1.000	0.5mg	0.333	0.333	0.333
	TA98-S9	79	0.000	0.350	1.000	0.5mg	0.333	0.333	0.333
	All ^b	319	0.000	0.388	1.000	-	0.333	0.333	0.333
Benzopyrene	TA100+S9	79	0.000	0.415	1.000	2.0 μ g	0.333	0.333	0.333
	TA98+S9	79	0.000	0.393	1.000	2.0 μ g	0.333	0.333	0.333
	All ^b	158	0.000	0.404	1.000	-	0.333	0.333	0.333
1-Nitropyrene	TA100-S9	76	0.000	0.358	1.000	6.0 μ g	0.333	0.333	0.333
	TA98-S9	80	0.000	0.405	1.000	0.1 μ g	0.333	0.333	0.333
	All ^b	156	0.000	0.382	1.000	-	0.333	0.333	0.333
ALL	All ^b	1102	0.000	0.401	1.000	-	0.333	0.333	0.333

a: Highest dose scaled to unity.

b: Average weighted by the number of experiments.

Table 4.6 A - Optimal Designs for the Ames Assay

Test Agent	Strain +/- S9	Number of Expts.	Optimal Doses ^a			Max. Dose	Optimal Allocation		
			d_0	d_1	d_2		c_0	c_1	c_2
Coal Tar Solution	TA100+S9	40	0.000	0.381	1.000	5.0 μ g	0.407	0.523	0.070
	TA100-S9	32	0.000	0.438	1.000	10.0 μ g	0.676	0.272	0.052
	TA98+S9	40	0.000	0.362	1.000	5.0 μ g	0.360	0.574	0.066
	TA98-S9	38	0.000	0.412	1.000	10.0 μ g	0.638	0.312	0.050
	All ^b	150	0.000	0.396	1.000	-	0.510	0.430	0.060
Urban Air Particles	TA100+S9	79	0.000	0.455	1.000	2.0mg	0.364	0.535	0.100
	TA100-S9	80	0.000	0.453	1.000	2.0mg	0.368	0.530	0.102
	TA98+S9	80	0.000	0.300	1.000	8.0mg	0.230	0.720	0.049
	TA98-S9	80	0.000	0.270	1.000	10.0mg	0.194	0.758	0.048
	All ^b	319	0.000	0.369	1.000	-	0.289	0.636	0.075
Diesel Particles	TA100+S9	80	0.000	0.408	1.000	0.2mg	0.237	0.666	0.097
	TA100-S9	80	0.000	0.414	1.000	0.2mg	0.243	0.659	0.098
	TA98+S9	80	0.000	0.339	1.000	0.5mg	0.139	0.788	0.074
	TA98-S9	80	0.000	0.316	1.000	0.5mg	0.108	0.826	0.067
	All ^b	319	0.000	0.369	1.000	-	0.182	0.734	0.084
Benzopyrene	TA100+S9	79	0.000	0.396	1.000	2.0 μ g	0.280	0.632	0.088
	TA98+S9	79	0.000	0.379	1.000	2.0 μ g	0.198	0.714	0.087
	All ^b	158	0.000	0.388	1.000	-	0.239	0.673	0.088
1-Nitropyrene	TA100-S9	76	0.000	0.320	1.000	6.0 μ g	0.360	0.572	0.068
	TA98-S9	80	0.000	0.390	1.000	0.1 μ g	0.204	0.702	0.094
	All ^b	156	0.000	0.356	1.000	-	0.280	0.639	0.082
ALL	All ^b	1102	0.000	0.374	1.000	-	0.280	0.642	0.078

a: Highest dose scaled to unity.

b: Average Weighted by the number of experiments.

**Table 4.7 Efficiency of Original 6 - Dose Design
and Recommended 3 - Dose Design Under C-Optimality
Relative to C-Optimal 3 - Dose Design**

Test Agent	Strain +/- S9	Number of Expts.	Efficiency of Original 6-Dose Design ^a			Efficiency of Recommended 3-Dose Design ^c		
			Min	Avg	Max	Min	Avg	Max
Coal Tar Solution	TA100+S9	40	0.555	0.569	0.600	0.720	0.951	0.999
	TA100-S9	32	0.550	0.551	0.554	0.700	0.745	0.882
	TA98+S9	40	0.550	0.579	0.599	0.884	0.967	1.000
	TA98-S9	38	0.549	0.557	0.595	0.700	0.823	0.997
	All	150	0.549 ^b	0.565 ^d	0.600 ^b	0.700 ^b	0.879 ^d	1.000 ^b
Urban Air Particles	TA100+S9	79	0.551	0.558	0.581	0.709	0.835	0.996
	TA100-S9	80	0.551	0.556	0.578	0.709	0.832	0.999
	TA98+S9	80	0.546	0.580	0.601	0.756	0.876	0.999
	TA98-S9	80	0.505	0.562	0.601	0.608	0.796	0.981
	All	319	0.546 ^b	0.564 ^d	0.601 ^b	0.608 ^b	0.835 ^d	0.999 ^b
Diesel Particles	TA100+S9	80	0.547	0.576	0.601	0.703	0.934	1.000
	TA100-S9	80	0.550	0.574	0.597	0.752	0.954	0.999
	TA98+S9	80	0.550	0.586	0.600	0.760	0.897	0.999
	TA98-S9	79	0.535	0.578	0.600	0.735	0.858	0.993
	All	319	0.535 ^b	0.579 ^d	0.601 ^b	0.703 ^b	0.911 ^d	1.000 ^b
Benzopyrene	TA100+S9	79	0.550	0.574	0.600	0.779	0.944	0.999
	TA98+S9	79	0.550	0.583	0.600	0.717	0.955	0.999
	All	158	0.550 ^b	0.579 ^d	0.600 ^b	0.748 ^b	0.949 ^d	0.999 ^b
1-Nitropyrene	TA100-S9	76	0.472	0.570	0.601	0.041	0.849	0.997
	TA98-S9	80	0.539	0.575	0.601	0.673	0.930	0.999
	All	156	0.472 ^b	0.573 ^d	0.601 ^b	0.041 ^b	0.890 ^d	0.999 ^b
ALL	All	1102	0.472 ^b	0.572 ^d	0.601 ^b	0.365 ^b	0.887 ^d	1.000 ^b

a: Equal allocation with dose levels in the ratios 1: 2: 4: 6: 8.

b: Minimum or maximum column values.

c: Dose levels: $d_0 = 0, d_1 = 0.4, d_2 = 1.0$; Allocation: $c_0 = 0.2, c_1 = 0.7, c_2 = 0.1$.

d: Average weighted by the number of experiments.

**Table 4.8 Efficiency of Original 6 - Dose Design
and Recommended 3 - Dose Design Under D-Optimality
Relative to D-Optimal 3 - Dose Design**

Test Agent	Strain +/- S9	Number of Expts.	Efficiency of Original 6-Dose Design ^a			Efficiency of Recommended 3-Dose Design ^c		
			Min	Avg	Max	Min	Avg	Max
Coal Tar Solution	TA100+S9	40	0.438	0.456	0.486	0.924	0.991	0.999
	TA100-S9	32	0.485	0.488	0.489	0.921	0.937	0.982
	TA98+S9	40	0.438	0.449	0.469	0.978	0.994	0.999
	TA98-S9	38	0.439	0.475	0.489	0.921	0.959	1.000
	All	150	0.438 ^b	0.466 ^d	0.489 ^b	0.921 ^b	0.972 ^d	1.000 ^b
Urban Air Particles	TA100+S9	79	0.446	0.470	0.487	0.921	0.961	0.999
	TA100-S9	80	0.448	0.472	0.488	0.921	0.962	0.999
	TA98+S9	80	0.437	0.440	0.458	0.957	0.985	0.999
	TA98-S9	80	0.438	0.451	0.518	0.867	0.964	0.998
	All	319	0.437 ^b	0.458 ^d	0.518 ^b	0.916 ^b	0.968 ^d	0.999 ^b
Diesel Particles	TA100+S9	80	0.438	0.450	0.488	0.921	0.985	1.000
	TA100-S9	80	0.440	0.450	0.485	0.935	0.991	0.999
	TA98+S9	80	0.438	0.440	0.479	0.940	0.985	0.999
	TA98-S9	79	0.438	0.440	0.471	0.950	0.981	1.000
	All	319	0.438 ^b	0.445 ^d	0.486 ^b	0.935 ^b	0.986 ^d	1.000 ^b
Benzopyrene	TA100+S9	79	0.438	0.453	0.477	0.945	0.987	0.999
	TA98+S9	79	0.438	0.447	0.486	0.923	0.990	0.999
	All	158	0.438	0.450 ^d	0.486 ^b	0.923 ^b	0.988 ^d	0.999 ^b
1-Nitropyrene	TA100-S9	76	0.440	0.470	0.595	0.131	0.925	0.999
	TA98-S9	80	0.438	0.449	0.486	0.901	0.989	1.000
	All	156	0.438 ^b	0.459 ^d	0.595 ^b	0.131 ^b	0.958 ^d	1.000 ^b
ALL	All	1102	0.437 ^b	0.454 ^d	0.595 ^b	0.131 ^b	0.975 ^d	1.000 ^b

a: Equal allocation with dose levels in the ratios 1: 2: 4: 6: 8.

b: Minimum or maximum column values.

c: Recommended design: $d_0 = 0$, $d_1 = 0.4$, $d_2 = 1.0$, $c_0 = 0.333$, $c_1 = 0.333$, $c_2 = 0.333$.

d: Average weighted by the number of experiments.

**Table 4.9 Efficiency of Original 6 - Dose Design
and Recommended 3 - Dose Design Under A- Optimality
Relative to A- Optimal 3 - Dose Design**

Test Agent	Strain +/- S9	Number of Expts.	Efficiency of Original 6-Dose Design ^a			Efficiency of Recommended 3-Dose Design ^c		
			Min	Avg	Max	Min	Avg	Max
Coal Tar Solution	TA100+S9	40	0.537	0.576	0.643	0.691	0.901	0.981
	TA100-S9	32	0.448	0.448	0.454	0.536	0.550	0.617
	TA98+S9	40	0.546	0.596	0.637	0.809	0.938	0.993
	TA98-S9	38	0.447	0.459	0.566	0.536	0.606	0.890
	All	150	0.447 ^b	0.524 ^d	0.643 ^b	0.536 ^b	0.761 ^d	0.981 ^b
Urban Air Particles	TA100+S9	79	0.563	0.569	0.594	0.780	0.880	0.998
	TA100-S9	80	0.563	0.568	0.593	0.780	0.878	0.998
	TA98+S9	80	0.502	0.636	0.661	0.728	0.908	0.978
	TA98-S9	80	0.553	0.629	0.665	0.643	0.854	0.969
	All	319	0.502 ^b	0.601 ^d	0.665 ^b	0.643 ^b	0.880 ^d	0.998 ^b
Diesel Particles	TA100+S9	80	0.547	0.576	0.601	0.788	0.935	0.987
	TA100-S9	80	0.550	0.574	0.597	0.827	0.959	0.987
	TA98+S9	80	0.535	0.576	0.600	0.712	0.856	0.975
	TA98-S9	79	0.535	0.579	0.600	0.687	0.811	0.988
	All	319	0.535 ^b	0.576 ^d	0.601 ^b	0.687 ^b	0.890 ^d	0.988 ^b
Benzopyrene	TA100+S9	79	0.565	0.587	0.615	0.836	0.954	0.998
	TA98+S9	79	0.564	0.597	0.616	0.740	0.948	0.998
	All	158	0.564 ^b	0.592 ^d	0.616 ^b	0.740 ^b	0.951 ^d	0.998 ^b
1-Nitropyrene	TA100-S9	76	0.493	0.584	0.638	0.043	0.827	0.988
	TA98-S9	80	0.539	0.578	0.634	0.638	0.918	0.985
	All	156	0.493 ^b	0.581 ^d	0.638 ^b	0.043 ^b	0.874 ^d	0.988 ^b
ALL	All	1102	0.447 ^b	0.579 ^d	0.665 ^b	0.043 ^b	0.876 ^d	0.998 ^b

a: Equal allocation with dose levels in the ratios 1: 2: 4: 6: 8

b: Minimum or maximum group value.

c: Recommended design: $d_0 = 0, d_1 = 0.4, d_2 = 1.0$; $c_0 = 0.3, c_1 = 0.6, c_2 = 0.1$.

d: Average weighted by the number of experiments.

Table 4.10 Efficiency of Fixed 3-6 Dose Designs with Equal Allocation (C- Optimality)

Test Agent	Strain +/- S9	Number of Expts.	K=3			K=4			K=5			K=6			
			0.0	0.4	1.0 ^a	Min	Avg	Max	0.0	0.2	0.4	0.7	1.0	Min	Avg
Coal Tar Solution	TA100+S9	40	0.475	0.644	0.810	0.678	0.733	0.761	0.607	0.656	0.672	0.550	0.569	0.600	
	TA100-S9	32	0.751	0.809	0.824	0.675	0.689	0.731	0.644	0.651	0.664	0.550	0.550	0.551	
	TA98+S9	40	0.445	0.588	0.750	0.669	0.723	0.751	0.580	0.645	0.673	0.550	0.579	0.599	
	TA98-S9	38	0.521	0.758	0.824	0.675	0.708	0.756	0.633	0.656	0.669	0.549	0.557	0.595	
	All ^{b,c}	150	0.445	0.699	0.824	0.675	0.713	0.761	0.580	0.652	0.673	0.550	0.563	0.600	
Urban Air Particles	TA100+S9	79	0.555	0.753	0.814	0.675	0.708	0.747	0.640	0.661	0.673	0.551	0.558	0.581	
	TA100-S9	80	0.585	0.760	0.815	0.676	0.710	0.754	0.648	0.661	0.671	0.551	0.556	0.578	
	TA98+S9	80	0.335	0.418	0.703	0.551	0.627	0.731	0.474	0.550	0.671	0.546	0.580	0.601	
	TA98-S9	80	0.270	0.362	0.507	0.501	0.582	0.701	0.410	0.503	0.628	0.505	0.562	0.601	
	All ^{b,c}	319	0.270	0.573	0.815	0.501	0.656	0.754	0.410	0.594	0.673	0.505	0.564	0.601	
Diesel Particles	TA100+S9	80	0.423	0.623	0.817	0.656	0.717	0.747	0.563	0.648	0.673	0.547	0.576	0.601	
	TA100-S9	80	0.416	0.638	0.813	0.669	0.728	0.742	0.557	0.659	0.673	0.550	0.574	0.597	
	TA98+S9	80	0.337	0.462	0.795	0.559	0.651	0.734	0.477	0.578	0.673	0.550	0.586	0.600	
	TA98-S9	79	0.325	0.415	0.768	0.535	0.620	0.739	0.460	0.545	0.673	0.535	0.578	0.600	
	All ^{b,c}	319	0.325	0.534	0.817	0.535	0.679	0.747	0.460	0.607	0.673	0.535	0.578	0.601	
Benzopyrene	TA100+S9	79	0.435	0.624	0.790	0.654	0.723	0.743	0.573	0.651	0.672	0.550	0.574	0.600	
	TA98+S9	79	0.345	0.555	0.811	0.579	0.707	0.742	0.489	0.632	0.673	0.550	0.583	0.600	
	All ^{b,c}	158	0.345	0.589	0.811	0.579	0.715	0.743	0.489	0.642	0.673	0.550	0.578	0.600	
1-Nitropyrene	TA100-S9	76	0.301	0.511	0.794	0.294	0.688	0.773	0.286	0.595	0.672	0.172	0.570	0.601	
	TA98-S9	80	0.300	0.579	0.803	0.552	0.698	0.740	0.449	0.631	0.673	0.539	0.575	0.601	
	All ^{b,c}	156	0.300	0.545	0.803	0.294	0.693	0.773	0.286	0.604	0.673	0.472	0.572	0.601	
All	All ^{b,c}	1102	0.270	0.588	0.824	0.294	0.691	0.773	0.286	0.620	0.673	0.472	0.571	0.601	

a: Allocation: $c_0, c_1, \dots, c_k$.

b: Minimum or maximum of the group(test agents) value.

c: Average weighted by the number of experiments.

Table 4.11 Optimal Allocation for Fixed 3-6 Dose Designs (C- Optimality)

Test Agent	Strain +/- S9	Number of Expts.	Optimal Allocation for Design with K=3-6 Doses			
			K=3 $c_0 \ c_1 \ c_3$	K=4 $c_0 \ c_1 \ c_2 \ c_3$	K=5 $c_0 \ c_1 \ c_2 \ c_3 \ c_4$	K=6 $c_0 \ c_1 \ c_2 \ c_3 \ c_4 \ c_5$
Coal Tar Solution	TA100+S9	40	0.249 0.649 0.101	0.235 0.173 0.480 0.112	0.218 0.168 0.413 0.100 0.102	0.265 0.100 0.140 0.355 0.100 0.100
	TA100-S9	32	0.353 0.522 0.125	0.359 0.100 0.441 0.100	0.300 0.100 0.400 0.100 0.100	0.261 0.100 0.105 0.334 0.100 0.100
	TA98+S9	40	0.219 0.698 0.092	0.220 0.260 0.418 0.102	0.203 0.248 0.350 0.100 0.100	0.198 0.100 0.196 0.305 0.100 0.100
	TA98-S9	38	0.317 0.546 0.117	0.308 0.111 0.474 0.108	0.279 0.111 0.408 0.100 0.103	0.281 0.100 0.105 0.334 0.100 0.100
	All	150	0.280 0.612 0.108 ^a	0.280 0.161 0.453 0.105	0.250 0.156 0.393 0.100 0.101 ^c	0.231 0.100 0.137 0.332 0.100 0.100 ^b
Urban Air Particles	TA100+S9	79	0.311 0.575 0.115	0.300 0.103 0.499 0.100	0.270 0.103 0.419 0.100 0.104	0.244 0.100 0.101 0.354 0.100 0.100
	TA100-S9	80	0.314 0.570 0.116	0.299 0.103 0.499 0.100	0.285 0.101 0.408 0.100 0.107	0.250 0.100 0.100 0.350 0.100 0.100
	TA98+S9	80	0.106 0.827 0.067	0.129 0.626 0.145 0.100	0.121 0.550 0.129 0.100 0.100	0.121 0.100 0.450 0.129 0.100 0.100
	TA98-S9	80	0.078 0.861 0.062	0.105 0.690 0.105 0.100	0.104 0.595 0.101 0.100 0.100	0.104 0.100 0.495 0.101 0.100 0.100
	All	319	0.202 0.708 0.090 ^b	0.206 0.360 0.312 0.100 ^b	0.195 0.337 0.263 0.100 0.103 ^b	0.180 0.100 0.286 0.234 0.100 0.100 ^b
Diesel Particles	TA100+S9	80	0.237 0.667 0.097	0.231 0.236 0.433 0.100	0.225 0.225 0.350 0.100 0.100	0.215 0.100 0.180 0.305 0.100 0.100
	TA100-S9	80	0.243 0.659 0.098	0.224 0.148 0.528 0.100	0.210 0.148 0.443 0.100 0.100	0.208 0.100 0.126 0.366 0.100 0.100
	TA98+S9	80	0.137 0.769 0.074	0.155 0.563 0.183 0.100	0.143 0.506 0.153 0.100 0.100	0.135 0.100 0.410 0.155 0.100 0.100
	TA98-S9	79	0.107 0.827 0.067	0.119 0.637 0.144 0.100	0.115 0.551 0.134 0.100 0.100	0.114 0.100 0.459 0.127 0.100 0.100
	All	319	0.181 0.735 0.084 ^b	0.182 0.366 0.322 0.100 ^b	0.173 0.357 0.270 0.100 0.108 ^b	0.168 0.100 0.294 0.238 0.100 0.100 ^b
Benzopyrene	TA100+S9	79	0.220 0.694 0.098	0.220 0.227 0.435 0.100	0.220 0.164 0.393 0.100 0.130	0.208 0.100 0.173 0.310 0.100 0.100
	TA98+S9	79	0.196 0.714 0.087	0.200 0.303 0.397 0.100	0.184 0.290 0.327 0.100 0.100	0.181 0.100 0.234 0.285 0.100 0.100
	All	158	0.218 ^b 0.690 ^b 0.092 ^b	0.219 0.265 0.416 0.100 ^b	0.208 0.222 0.355 0.100 0.113 ^b	0.195 0.100 0.204 0.302 0.100 0.100 ^b
1-Nitropyrene	TA100-S9	76	0.200 0.693 0.098	0.216 0.454 0.235 0.100	0.203 0.405 0.180 0.100 0.100	0.183 0.111 0.325 0.182 0.100 0.100
	TA98-S9	80	0.206 0.705 0.099	0.196 0.261 0.443 0.100	0.185 0.238 0.378 0.100 0.100	0.183 0.100 0.205 0.313 0.100 0.100
	All	156	0.207 ^b 0.690 ^b 0.093 ^b	0.208 0.357 0.334 0.100 ^b	0.194 0.321 0.283 0.101 0.100 ^b	0.183 0.105 0.265 0.248 0.100 0.100 ^b
ALL	All	1102	0.210 ^b 0.690 ^b 0.091 ^b	0.219 0.311 0.367 0.102 ^b	0.203 0.278 0.312 0.100 0.104 ^b	0.191 0.101 0.237 0.271 0.100 0.100 ^b

a: Allocation: $c_0, c_1, \dots, c_5$.

b: Minimum or maximum of the group(test agents) value.

c: Average weighted by the number of experiments.

2 of/de 2

PM-1 3½"x4" PHOTOGRAPHIC MICROCOPY TARGET
NBS 1010a ANSI/ISO #2 EQUIVALENT

PRECISIONSM RESOLUTION TARGETS

Table 4.12 Efficiency of Fixed 3-6 Dose Designs with Unequal Allocation (C-optimality)

Test Agent	Strain +/- S9	Number of Expts.	K=3			K=4			K=5			K=6				
			2:	7:	1	2:	3:	4:	1	2:	3:	1:	2:	3:	1:	1
Coal Tar Solution	TA100+S9	40	0.719	0.951	0.999	0.701	0.824	0.859		0.661	0.750	0.781	0.651	0.690	0.700	
	TA100-S9	32	0.700	0.745	0.882	0.687	0.712	0.781		0.610	0.662	0.730	0.640	0.651	0.680	
	TA98+S9	40	0.884	0.966	1.000	0.782	0.837	0.863		0.711	0.762	0.780	0.620	0.681	0.700	
	TA98-S9	38	0.700	0.823	0.997	0.687	0.753	0.865		0.640	0.691	0.781	0.641	0.671	0.693	
	All ^{b,c}	150	0.700	0.871	1.000	0.687	0.781	0.865		0.610	0.716	0.781	0.620	0.673	0.700	
Urban Air Particles	TA100+S9	79	0.709	0.835	0.996	0.694	0.762	0.855		0.650	0.702	0.770	0.644	0.676	0.700	
	TA100-S9	80	0.709	0.832	0.999	0.691	0.759	0.855		0.651	0.699	0.771	0.610	0.674	0.700	
	TA98+S9	80	0.756	0.876	0.999	0.682	0.779	0.865		0.673	0.760	0.781	0.607	0.672	0.701	
	TA98-S9	80	0.608	0.796	0.981	0.571	0.716	0.856		0.511	0.650	0.771	0.420	0.531	0.661	
	All ^{b,c}	319	0.608	0.834	0.999	0.571	0.754	0.865		0.511	0.703	0.781	0.420	0.638	0.700	
Diesel Particles	TA100+S9	80	0.703	0.934	1.000	0.690	0.821	0.866		0.651	0.747	0.781	0.590	0.680	0.700	
	TA100-S9	80	0.752	0.954	0.999	0.716	0.739	0.831		0.661	0.751	0.780	0.581	0.690	0.700	
	TA98+S9	80	0.760	0.879	0.999	0.685	0.796	0.860		0.620	0.720	0.781	0.510	0.611	0.700	
	TA98-S9	79	0.735	0.858	0.993	0.664	0.766	0.855		0.601	0.692	0.780	0.490	0.580	0.700	
	All ^{b,c}	319	0.703	0.910	0.999	0.664	0.780	0.866		0.601	0.727	0.781	0.490	0.610	0.700	
Benzopyrene	TA100+S9	79	0.773	0.914	0.999	0.734	0.824	0.864		0.680	0.751	0.780	0.600	0.681	0.700	
	TA98+S9	79	0.717	0.955	0.999	0.696	0.699	0.866		0.641	0.750	0.782	0.521	0.671	0.700	
	All ^{b,c}	158	0.717	0.919	0.999	0.734	0.761	0.866		0.641	0.750	0.782	0.521	0.676	0.700	
1-Nitropyrene	TA100-S9	76	0.041	0.849	0.997	0.315	0.763	0.862		0.360	0.701	0.781	0.250	0.621	0.690	
	TA98-S9	80	0.673	0.929	0.999	0.622	0.815	0.864		0.580	0.741	0.780	0.461	0.660	0.700	
	All ^{b,c}	156	0.041	0.889	0.999	0.315	0.789	0.864		0.360	0.721	0.781	0.250	0.610	0.700	
All	All ^{b,c}	1102	0.041	0.894	1.000	0.315	0.773	0.866		0.360	0.723	0.762	0.250	0.653	0.700	

a: Allocation: $c_0, c_1, \dots, c_k$.

b: Minimum or maximum of the group(test agents) value.

c: Average weighted by the number of experiments.

Table 4.13 Efficiency of Fixed 3-6 Dose Designs with Equal Allocation (D-Optimality)

Test Agent	Strain +/- S9	Number of Expts.	K=3			K=4			K=5			K=6			
			0.0	0.4	1.0 ^a	Min	Avg	Max	0.0	0.2	0.4	0.7	1.0	Min	Avg
Coal Tar Solution	TA100+S9	40	0.883	0.946	1.000	0.723	0.764	0.780	0.561	0.572	0.588	0.550	0.569	0.600	
	TA100-S9	32	0.981	0.998	1.000	0.722	0.731	0.753	0.588	0.590	0.591	0.550	0.550	0.554	
	TA98+S9	40	0.854	0.932	0.981	0.753	0.769	0.779	0.562	0.568	0.578	0.550	0.579	0.599	
	TA98-S9	38	0.919	0.982	1.000	0.722	0.743	0.778	0.565	0.582	0.591	0.549	0.557	0.595	
	All ^{b,c}	150	0.854	0.964	1.000	0.722	0.751	0.780	0.561	0.578	0.591	0.550	0.563	0.600	
Urban Air Particles	TA100+S9	79	0.906	0.987	1.000	0.722	0.742	0.774	0.567	0.579	0.589	0.551	0.558	0.581	
	TA100-S9	80	0.920	0.987	1.000	0.723	0.742	0.772	0.568	0.579	0.589	0.551	0.556	0.578	
	TA98+S9	80	0.817	0.889	0.976	0.756	0.781	0.785	0.558	0.560	0.573	0.516	0.580	0.601	
	TA98-S9	80	0.661	0.841	0.923	0.749	0.776	0.785	0.557	0.561	0.588	0.505	0.562	0.601	
	All ^{b,c}	319	0.661	0.926	1.000	0.722	0.760	0.785	0.557	0.570	0.589	0.505	0.564	0.601	
Diesel Particles	TA100+S9	80	0.854	0.947	1.000	0.723	0.763	0.780	0.562	0.570	0.590	0.547	0.576	0.601	
	TA100-S9	80	0.805	0.954	1.000	0.728	0.763	0.778	0.561	0.570	0.588	0.550	0.574	0.597	
	TA98+S9	80	0.821	0.898	1.000	0.730	0.777	0.785	0.558	0.560	0.583	0.550	0.586	0.600	
	TA98-S9	79	0.802	0.880	0.994	0.741	0.779	0.785	0.557	0.562	0.579	0.535	0.578	0.600	
	All ^{b,c}	319	0.802	0.919	1.000	0.723	0.770	0.785	0.557	0.566	0.590	0.535	0.578	0.601	
Benzopyrene	TA100+S9	79	0.825	0.944	1.000	0.733	0.763	0.782	0.563	0.570	0.582	0.550	0.574	0.600	
	TA98+S9	79	0.804	0.924	1.000	0.723	0.770	0.780	0.559	0.567	0.588	0.550	0.583	0.600	
	All ^{b,c}	158	0.804	0.934	1.000	0.723	0.766	0.782	0.559	0.568	0.588	0.550	0.578	0.600	
1-Nitropyrene	TA100-S9	76	0.070	0.824	0.998	0.365	0.746	0.780	0.385	0.570	0.657	0.472	0.570	0.601	
	TA98-S9	80	0.718	0.938	1.000	0.726	0.768	0.785	0.558	0.568	0.586	0.539	0.575	0.601	
	All ^{b,c}	156	0.070	0.881	0.999	0.365	0.753	0.785	0.385	0.569	0.657	0.472	0.572	0.601	
All	All ^{b,c}	1102	0.070	0.925	1.000	0.365	0.760	0.785	0.385	0.570	0.657	0.472	0.571	0.601	

a: Allocation: $c_0, c_1, \dots, c_k$.

b: Minimum or maximum of the group(test agents) value.

c: Average weighted by the number of experiments.

Table 4.14 Optimal Allocation for Fixed 3-6 Dose Designs (D-Optimality)

Test Agent	Strain +/- S9	Number of Expts.	Optimal Allocation for Design with K=3-6 Doses			
			K=3 c ₀ c ₁ c ₂	K=4 c ₀ c ₁ c ₂ c ₃	K=5 c ₀ c ₁ c ₂ c ₃ c ₄	K=6 c ₀ c ₁ c ₂ c ₃ c ₄ c ₅
Coal Tar Solution	TA100+S9	40	0.333 0.333 0.333	0.300 0.110 0.290 0.300	0.300 0.110 0.190 0.100 0.300	0.290 0.100 0.100 0.110 0.100 0.300
	TA100-S9	32	0.333 0.333 0.333	0.300 0.100 0.300 0.300	0.300 0.100 0.200 0.100 0.300	0.203 0.100 0.100 0.197 0.100 0.300
	TA98+S9	40	0.333 0.333 0.333	0.300 0.140 0.260 0.300	0.300 0.125 0.175 0.100 0.300	0.300 0.100 0.100 0.100 0.100 0.300
	TA98-S9	38	0.333 0.333 0.333	0.300 0.100 0.300 0.300	0.300 0.100 0.200 0.100 0.300	0.232 0.100 0.100 0.168 0.100 0.300
	All	150	0.333 0.333 0.333 ^a	0.300 0.112 0.288 0.300	0.300 0.109 0.191 0.100 0.300 ^a	0.256 0.100 0.100 0.144 0.100 0.300 ^a
Urban Air Particles	TA100+S9	79	0.333 0.333 0.333	0.300 0.100 0.300 0.300	0.300 0.101 0.199 0.100 0.300	0.242 0.100 0.100 0.158 0.100 0.300
	TA100-S9	80	0.333 0.333 0.333	0.300 0.100 0.300 0.300	0.300 0.100 0.200 0.100 0.300	0.300 0.100 0.100 0.100 0.100 0.300
	TA98+S9	80	0.333 0.333 0.333	0.300 0.256 0.144 0.300	0.300 0.187 0.114 0.100 0.300	0.300 0.100 0.100 0.100 0.100 0.300
	TA98-S9	80	0.333 0.333 0.333	0.300 0.295 0.105 0.300	0.300 0.147 0.154 0.100 0.300	0.300 0.100 0.100 0.100 0.100 0.300
	All	319	0.333 0.333 0.333 ^a	0.300 0.187 0.212 0.300 ^a	0.195 0.337 0.263 0.100 0.103 ^a	0.286 0.100 0.100 0.115 0.100 0.100 ^a
Diesel Particles	TA100+S9	80	0.333 0.333 0.333	0.300 0.124 0.276 0.300	0.300 0.125 0.175 0.100 0.300	0.281 0.100 0.100 0.119 0.100 0.300
	TA100-S9	80	0.333 0.333 0.333	0.300 0.103 0.298 0.300	0.300 0.105 0.195 0.100 0.300	0.290 0.100 0.100 0.110 0.100 0.300
	TA98+S9	80	0.333 0.333 0.333	0.300 0.235 0.165 0.300	0.300 0.179 0.121 0.100 0.300	0.294 0.100 0.100 0.106 0.100 0.300
	TA98-S9	79	0.333 0.333 0.333	0.300 0.277 0.123 0.300	0.300 0.191 0.108 0.100 0.300	0.300 0.100 0.100 0.100 0.100 0.300
	All	319	0.333 0.333 0.333 ^a	0.300 0.184 0.215 0.300 ^a	0.300 0.150 0.150 0.100 0.300 ^a	0.291 0.100 0.100 0.109 0.100 0.300 ^a
Benzopyrene	TA100+S9	79	0.333 0.333 0.333	0.300 0.127 0.273 0.300	0.300 0.122 0.178 0.100 0.300	0.289 0.100 0.100 0.111 0.100 0.300
	TA98+S9	79	0.333 0.333 0.333	0.300 0.149 0.251 0.300	0.300 0.132 0.168 0.100 0.300	0.297 0.100 0.100 0.103 0.100 0.300
	All	158	0.333 ^b 0.333 ^b 0.333 ^a	0.300 0.138 0.262 0.300 ^a	0.300 0.127 0.173 0.100 0.300 ^a	0.293 0.100 0.100 0.107 0.100 0.300 ^a
1-Nitropyrene	TA100-S9	76	0.333 0.333 0.333	0.300 0.229 0.179 0.300	0.300 0.184 0.126 0.106 0.283	0.291 0.103 0.109 0.109 0.100 0.300
	TA98-S9	80	0.333 0.333 0.333	0.300 0.149 0.251 0.300	0.300 0.126 0.174 0.100 0.300	0.294 0.100 0.100 0.106 0.100 0.300
	All	156	0.333 ^b 0.333 ^b 0.333 ^a	0.300 0.189 0.215 0.300 ^a	0.300 0.155 0.150 0.103 0.292 ^a	0.293 0.101 0.104 0.102 0.100 0.300 ^a
ALL	All	1102	0.333 ^b 0.333 ^b 0.333 ^a	0.300 0.163 0.238 0.300 ^a	0.300 0.137 0.163 0.100 0.298 ^a	0.284 0.101 0.100 0.115 0.100 0.300 ^a

a: Allocation: $c_0, c_1, \dots, c_k$

b: Minimum or maximum of the group(test agents) value.

c: Average weighted by the number of experiments.

Table 4.15 Efficiency of Fixed 3-6 Dose Designs with Unqual Allocation (D-Optimality)

Test Agent	Strain +/- S9	Number of Expts.	Efficiency of Designs with K=3-6 Doses (Allocation Ratio)															
			K=3			K=4			K=5			K=6						
			1/3:	1/3:	1/3:	3:	1:	3:	3	3:	1:	2:	1:	3	Min	Avg	Max	
			Min	Avg	Max	Min	Avg	Max	Min	Avg	Max	Min	Avg	Max	Min	Avg	Max	
Coal Tar Solution	TA100+S9	40	0.924	0.991	0.999	0.882	0.900	0.912		0.790	0.810	0.830	0.681	0.700	0.730			
	TA100-S9	32	0.921	0.937	0.982	0.902	0.906	0.911		0.821	0.830	0.830	0.680	0.690	0.695			
	TA98+S9	40	0.978	0.994	1.000	0.882	0.882	0.911		0.790	0.810	0.821	0.680	0.710	0.730			
	TA98-S9	38	0.921	0.959	0.999	0.890	0.906	0.911		0.800	0.820	0.825	0.680	0.690	0.720			
	All ^{bc}	150	0.921	0.970	0.999	0.882	0.898	0.912		0.790	0.817	0.830	0.690	0.697	0.730			
Urban Air Particles	TA100+S9	79	0.921	0.961	1.000	0.886	0.908	0.913		0.796	0.826	0.833	0.685	0.689	0.711			
	TA100-S9	80	0.921	0.962	0.999	0.890	0.908	0.912		0.799	0.825	0.833	0.681	0.688	0.708			
	TA98+S9	80	0.957	0.985	0.999	0.894	0.907	0.921		0.760	0.790	0.822	0.690	0.740	0.750			
	TA98-S9	80	0.867	0.964	0.981	0.895	0.915	0.922		0.700	0.771	0.800	0.720	0.751	0.760			
	All ^{bc}	319	0.867	0.968	1.000	0.886	0.909	0.922		0.700	0.803	0.833	0.681	0.716	0.760			
Diesel Particles	TA100+S9	80	0.921	0.985	1.000	0.888	0.902	0.912		0.771	0.810	0.833	0.680	0.710	0.730			
	TA100-S9	80	0.935	0.991	0.999	0.887	0.905	0.912		0.755	0.810	0.830	0.681	0.700	0.730			
	TA98+S9	80	0.940	0.985	0.999	0.892	0.904	0.920		0.761	0.800	0.831	0.690	0.730	0.760			
	TA98-S9	79	0.950	0.981	0.993	0.892	0.910	0.922		0.750	0.790	0.832	0.690	0.741	0.760			
	All ^{bc}	319	0.921	0.985	1.000	0.887	0.905	0.922		0.750	0.802	0.833	0.680	0.720	0.760			
Benzopyrene	TA100+S9	79	0.945	0.987	0.999	0.886	0.902	0.912		0.761	0.811	0.833	0.690	0.700	0.740			
	TA98+S9	79	0.923	0.990	0.999	0.885	0.903	0.918		0.755	0.800	0.832	0.690	0.720	0.750			
	All ^{bc}	158	0.923	0.988	0.999	0.885	0.902	0.918		0.755	0.805	0.833	0.690	0.710	0.710			
1-Nitropyrene	TA100-S9	76	0.131	0.925	0.999	0.788	0.876	0.915		0.275	0.762	0.831	0.390	0.700	0.770			
	TA98-S9	80	0.901	0.989	0.999	0.891	0.907	0.919		0.722	0.810	0.833	0.680	0.710	0.760			
	All ^{bc}	156	0.131	0.957	0.999	0.788	0.891	0.919		0.275	0.786	0.833	0.390	0.705	0.770			
ALL	All ^{bc}	1102	0.131	0.973	1.000	0.788	0.901	0.919		0.275	0.802	0.833	0.390	0.709	0.760			

a: Allocation: $c_0, c_1, \dots, c_k$.

b: Minimum or maximum of the group(test agents) value.

c: Average weighted by the number of experiments.

Table 4.16 Efficiency of Fixed 3-6 Dose Designs with Equal Allocation (A-optimality)

Test Agent	Strain +/- S9	Number of Expts.	Efficiency of Design with K=3-6 Doses (Dose Spacing)																						
			K=3			K=4			K=5			K=6													
			0.0	0.4	1.0 ^a	Min	Avg	Max	0.0	0.2	0.4	0.7	1.0	Min	Avg	Max	0.0	0.1	0.2	0.4	0.7	1.0	Min	Avg	Max
Coal Tar Solution	TA100+S9	40	0.491	0.621	0.699	0.629	0.739	0.777	0.558	0.637	0.667	0.537	0.576	0.643											
	TA100-S9	32	0.365	0.372	0.404	0.510	0.519	0.558	0.435	0.439	0.466	0.447	0.448	0.454											
	TA98+S9	40	0.555	0.650	0.699	0.693	0.752	0.777	0.597	0.649	0.677	0.546	0.596	0.637											
	TA98-S9	38	0.365	0.402	0.585	0.510	0.553	0.735	0.435	0.465	0.621	0.447	0.459	0.566											
	All ^{b,c}	150	0.365	0.511	0.699	0.510	0.640	0.777	0.435	0.547	0.677	0.447	0.519	0.643											
Urban Air Particles	TA100+S9	79	0.594	0.648	0.717	0.679	0.718	0.762	0.637	0.659	0.681	0.563	0.569	0.594											
	TA100-S9	80	0.594	0.647	0.718	0.679	0.718	0.770	0.637	0.657	0.677	0.563	0.568	0.593											
	TA98+S9	80	0.477	0.651	0.687	0.611	0.712	0.774	0.525	0.618	0.669	0.502	0.636	0.661											
	TA98-S9	80	0.516	0.627	0.680	0.555	0.679	0.771	0.452	0.581	0.664	0.553	0.629	0.665											
	All ^{b,c}	319	0.477	0.613	0.718	0.555	0.706	0.762	0.452	0.628	0.681	0.502	0.600	0.665											
Diesel Particles	TA100+S9	80	0.617	0.692	0.718	0.656	0.717	0.747	0.563	0.648	0.672	0.547	0.576	0.601											
	TA100-S9	80	0.638	0.705	0.719	0.669	0.728	0.742	0.557	0.659	0.673	0.550	0.574	0.597											
	TA98+S9	80	0.541	0.637	0.713	0.559	0.651	0.734	0.477	0.578	0.673	0.535	0.576	0.600											
	TA98-S9	79	0.519	0.606	0.718	0.535	0.620	0.739	0.460	0.545	0.673	0.535	0.579	0.600											
	All ^{b,c}	319	0.519	0.660	0.719	0.535	0.679	0.747	0.460	0.607	0.673	0.535	0.576	0.601											
Benzopyrene	TA100+S9	79	0.623	0.691	0.718	0.672	0.739	0.761	0.586	0.659	0.681	0.565	0.587	0.615											
	TA98+S9	79	0.564	0.690	0.718	0.589	0.726	0.761	0.497	0.644	0.680	0.564	0.597	0.616											
	All ^{b,c}	158	0.564	0.690	0.718	0.589	0.732	0.761	0.497	0.651	0.680	0.564	0.592	0.616											
1-Nitropyrene	TA100-S9	76	0.236	0.604	0.691	0.312	0.714	0.775	0.304	0.601	0.673	0.493	0.584	0.638											
	TA98-S9	80	0.517	0.683	0.771	0.552	0.703	0.784	0.449	0.635	0.704	0.539	0.578	0.634											
	All ^{b,c}	156	0.236	0.643	0.771	0.312	0.708	0.784	0.304	0.618	0.704	0.493	0.581	0.638											
All	All ^{b,c}	1102	0.236	0.629	0.771	0.312	0.693	0.784	0.304	0.609	0.704	0.447	0.573	0.665											

a: Allocation: $c_0, c_1, \dots, c_k$.

b: Minimum or maximum of the group(test agents) value.

c: Average weighted by the number of experiments.

Table 4.17 Optimal Allocation for Fixed 3-6 Dose Designs (A- Optimality)

Optimal Allocation for Design with K=3-6 Doses						
Test Agent	Strain +/- S9	Number of Expts.	K=3	K=4	K=5	K=6
			$c_0 \ c_1 \ c_2$	$c_0 \ c_1 \ c_2 \ c_3$	$c_0 \ c_1 \ c_2 \ c_3 \ c_4$	$c_0 \ c_1 \ c_2 \ c_3 \ c_4 \ c_5$
Coal Tar Solution	TA100+S9	40	0.407 0.523 0.070	0.373 0.205 0.323 0.100	0.345 0.198 0.258 0.100 0.100	0.313 0.100 0.150 0.238 0.100 0.100
	TA100-S9	32	0.676 0.272 0.052	0.600 0.100 0.200 0.100	0.588 0.100 0.113 0.100 0.100	0.488 0.100 0.100 0.100 0.100 0.112
	TA98+S9	40	0.360 0.574 0.066	0.330 0.270 0.300 0.100	0.320 0.260 0.220 0.100 0.100	0.275 0.100 0.198 0.228 0.100 0.100
	TA98-S9	38	0.638 0.312 0.050	0.579 0.121 0.200 0.100	0.542 0.118 0.139 0.100 0.100	0.471 0.100 0.108 0.111 0.100 0.111
	All	150	0.510 0.430 0.060 ^a	0.470 0.174 0.255 0.100	0.449 0.168 0.182 0.100 0.100 ^a	0.387 0.100 0.139 0.169 0.100 0.100 ^a
Urban Air Particles	TA100+S9	79	0.364 0.535 0.100	0.346 0.102 0.451 0.100	0.324 0.103 0.373 0.100 0.100	0.282 0.100 0.101 0.316 0.100 0.100
	TA100-S9	80	0.368 0.530 0.102	0.351 0.108 0.438 0.100	0.308 0.102 0.390 0.100 0.100	0.286 0.100 0.100 0.314 0.100 0.100
	TA98+S9	80	0.230 0.720 0.049	0.226 0.558 0.115 0.100	0.203 0.493 0.104 0.100 0.100	0.199 0.100 0.394 0.108 0.100 0.100
	TA98-S9	80	0.194 0.758 0.048	0.183 0.618 0.100 0.100	0.169 0.531 0.100 0.100 0.100	0.163 0.100 0.438 0.100 0.100 0.100
	All	319	0.289 0.636 0.075 ^a	0.277 0.346 0.276 0.100 ^a	0.251 0.307 0.242 0.100 0.100 ^a	0.233 0.100 0.258 0.210 0.100 0.100 ^a
Diesel Particles	TA100+S9	80	0.237 0.667 0.097	0.231 0.236 0.433 0.100	0.225 0.225 0.350 0.100 0.100	0.215 0.100 0.180 0.305 0.100 0.100
	TA100-S9	80	0.243 0.659 0.098	0.224 0.148 0.528 0.100	0.210 0.148 0.443 0.100 0.100	0.208 0.100 0.126 0.366 0.100 0.100
	TA98+S9	80	0.137 0.789 0.074	0.155 0.563 0.183 0.100	0.145 0.505 0.150 0.100 0.100	0.135 0.100 0.410 0.155 0.100 0.100
	TA98-S9	79	0.107 0.827 0.067	0.119 0.637 0.144 0.100	0.145 0.505 0.150 0.100 0.100	0.114 0.100 0.459 0.127 0.100 0.100
	All	319	0.181 0.735 0.084 ^a	0.182 0.396 0.322 0.100 ^a	0.181 0.346 0.273 0.100 0.100 ^a	0.168 0.100 0.294 0.238 0.100 0.100 ^a
Benzopyrene	TA100+S9	79	0.280 0.632 0.088	0.267 0.228 0.408 0.100	0.244 0.215 0.341 0.100 0.100	0.225 0.100 0.173 0.301 0.100 0.100
	TA98+S9	79	0.198 0.714 0.087	0.222 0.300 0.378 0.100	0.208 0.284 0.309 0.100 0.100	0.191 0.100 0.227 0.282 0.100 0.100
	All	158	0.239 ^a 0.673 ^a 0.088 ^a	0.244 0.264 0.393 0.100 ^a	0.226 0.249 0.325 0.100 0.100 ^a	0.208 0.100 0.200 0.292 0.100 0.100 ^a
1-Nitropyrene	TA100-S9	76	0.360 0.572 0.068	0.333 0.407 0.155 0.105	0.304 0.351 0.145 0.100 0.100	0.288 0.111 0.275 0.126 0.100 0.100
	TA98-S9	80	0.204 0.702 0.094	0.196 0.261 0.439 0.104	0.184 0.240 0.374 0.100 0.100	0.183 0.100 0.205 0.312 0.100 0.100
	All	156	0.280 ^a 0.630 ^a 0.082 ^a	0.264 0.334 0.297 0.104 ^a	0.260 0.278 0.261 0.100 0.100 ^a	0.236 0.105 0.240 0.219 0.100 0.100 ^a
ALL	All	1102	0.280 ^a 0.642 ^a 0.078 ^a	0.287 0.302 0.308 0.103 ^a	0.203 0.278 0.312 0.100 0.104 ^a	0.246 0.101 0.226 0.226 0.100 0.100 ^a

a: Allocation: $c_0, c_1, \dots, c_k$.

b: Minimum or maximum of the group (test agents) value.

c: Average weighted by the number of experiments.

Table 4.18 Efficiency of Fixed 3-6 Dose Designs with Unequal Allocation (A-optimality)

Test Agent	Strain +/- S9	Number of Expts.	Efficiency of Designs with K=3-6 Doses(Allocation Ratio)											
			K=3			K=4			K=5			K=6		
			3:	6:	1	3:	3:	1	2:	3:	3:	1:	2:	1:
			Min	Avg	Max	Min	Avg	Max	Min	Avg	Max	Min	Avg	Max
Coal Tar Solution	TA100+S9	40	0.691	0.904	0.981	0.582	0.738	0.871	0.570	0.701	0.780	0.600	0.690	0.750
	TA100-S9	32	0.536	0.551	0.617	0.445	0.452	0.468	0.441	0.451	0.480	0.480	0.490	0.511
	TA98+S9	40	0.809	0.938	0.993	0.652	0.782	0.862	0.620	0.730	0.790	0.640	0.710	0.750
	TA98-S9	38	0.536	0.606	0.890	0.446	0.468	0.711	0.440	0.481	0.680	0.481	0.510	0.670
	All ^c	150	0.536	0.749	0.981	0.445	0.614	0.871	0.440	0.591	0.780	0.480	0.600	0.750
Urban Air Particles	TA100+S9	79	0.779	0.880	0.981	0.677	0.746	0.855	0.641	0.696	0.781	0.656	0.685	0.727
	TA100-S9	80	0.779	0.878	0.999	0.676	0.742	0.855	0.640	0.692	0.780	0.655	0.683	0.724
	TA98+S9	80	0.728	0.908	0.999	0.572	0.820	0.863	0.560	0.750	0.790	0.570	0.720	0.760
	TA98-S9	80	0.643	0.854	0.981	0.628	0.789	0.859	0.590	0.730	0.791	0.570	0.700	0.760
	All ^c	319	0.643	0.834	0.999	0.572	0.774	0.863	0.560	0.717	0.791	0.570	0.697	0.760
Diesel Particles	TA100+S9	80	0.788	0.934	0.987	0.690	0.821	0.866	0.651	0.747	0.781	0.650	0.700	0.720
	TA100-S9	80	0.827	0.954	0.987	0.716	0.739	0.866	0.661	0.751	0.780	0.650	0.710	0.720
	TA98+S9	80	0.712	0.879	0.975	0.685	0.796	0.866	0.620	0.720	0.781	0.600	0.680	0.720
	TA98-S9	79	0.687	0.858	0.988	0.664	0.766	0.855	0.601	0.692	0.780	0.580	0.650	0.710
	All ^c	319	0.687	0.890	0.988	0.664	0.780	0.866	0.601	0.727	0.781	0.580	0.685	0.720
Benzopyrene	TA100+S9	79	0.779	0.944	0.999	0.713	0.820	0.871	0.670	0.751	0.790	0.670	0.710	0.730
	TA98+S9	79	0.717	0.955	0.999	0.681	0.836	0.872	0.640	0.760	0.790	0.621	0.710	0.740
	All ^c	158	0.717	0.949	0.999	0.681	0.828	0.872	0.640	0.756	0.790	0.620	0.710	0.740
1-Nitropyrene	TA100-S9	76	0.043	0.849	0.988	0.332	0.718	0.854	0.380	0.691	0.790	0.420	0.670	0.750
	TA98-S9	80	0.638	0.929	0.985	0.622	0.816	0.864	0.580	0.740	0.781	0.561	0.690	0.730
	All ^c	156	0.043	0.889	0.988	0.332	0.767	0.864	0.380	0.715	0.790	0.420	0.680	0.750
All	All ^c	1102	0.043	0.862	0.999	0.332	0.747	0.872	0.380	0.701	0.791	0.420	0.674	0.760

a: Allocation: $c_0, c_1, \dots, c_k$.

b: Minimum or maximum of the group(test agents) value.

c: Average weighted by the number of experiments.

Table 4.19 Efficiency of Fixed 3-6 Dose Designs

Optimality Criterion	Number of Doses	Dose Spacing	Allocation Ratios	Efficiency		
				Min	Avg	Max
C - Optimality	3	0.0 0.4 1.0	1: 1: 1	0.270	0.588	0.824
	4	0.0 0.2 0.4 1.0	1: 1: 1: 1	0.294	0.691	0.773
	5	0.0 0.2 0.4 0.7 1.0	1: 1: 1: 1: 1	0.286	0.620	0.673
	6	0.0 0.1 0.2 0.4 0.7 1.0	1: 1: 1: 1: 1: 1	0.472	0.571	0.601
D - Optimality	3	0.0 0.4 1.0	1: 1: 1	0.070	0.925	1.000
	4	0.0 0.2 0.4 1.0	1: 1: 1: 1	0.365	0.760	0.785
	5	0.0 0.2 0.4 0.7 1.0	1: 1: 1: 1: 1	0.385	0.570	0.657
	6	0.0 0.1 0.2 0.4 0.7 1.0	1: 1: 1: 1: 1: 1	0.472	0.571	0.601
A - Optimality	3	0.0 0.4 1.0	1: 1: 1	0.236	0.629	0.771
	4	0.0 0.2 0.4 1.0	1: 1: 1: 1	0.312	0.693	0.784
	5	0.0 0.2 0.4 0.7 1.0	1: 1: 1: 1: 1	0.304	0.609	0.704
	6	0.0 0.1 0.2 0.4 0.7 1.0	1: 1: 1: 1: 1: 1	0.477	0.573	0.665
C - Optimality	3	0.0 0.4 1.0	2: 7: 1	0.041	0.894	1.000
	4	0.0 0.2 0.4 1.0	2: 3: 4: 1	0.315	0.773	0.866
	5	0.0 0.2 0.4 0.7 1.0	2: 3: 3: 1: 1	0.360	0.723	0.762
	6	0.0 0.1 0.2 0.4 0.7 1.0	2: 1: 2: 3: 1: 1	0.250	0.653	0.700
D - Optimality	3	0.0 0.4 1.0	3: 3: 3	0.131	0.973	1.000
	4	0.0 0.2 0.4 1.0	3: 1: 3: 3	0.788	0.901	0.919
	5	0.0 0.2 0.4 0.7 1.0	3: 1: 2: 1: 3	0.275	0.802	0.833
	6	0.0 0.1 0.2 0.4 0.7 1.0	3: 1: 1: 1: 1: 3	0.390	0.709	0.760
A - Optimality	3	0.0 0.4 1.0	3: 6: 1	0.043	0.862	0.999
	4	0.0 0.2 0.4 1.0	3: 3: 3: 1	0.332	0.747	0.872
	5	0.0 0.2 0.4 0.7 1.0	2: 3: 3: 1: 1	0.380	0.701	0.791
	6	0.0 0.1 0.2 0.4 0.7 1.0	3: 1: 2: 2: 1: 1	0.420	0.674	0.760

4.4 Bayesian Optimal Designs

4.4.1 Bayesian D-Optimal Designs

As shown in section 4.3, the “information” matrix is of the form

$$I(\boldsymbol{\beta}, \mathbf{c}) = \mathbf{D}^T \mathbf{V}^{-1} \mathbf{D},$$

so that

$$\text{Cov}(\hat{\boldsymbol{\beta}}) = (\mathbf{D}^T \mathbf{V}^{-1} \mathbf{D})^{-1} = I(\boldsymbol{\beta}, \mathbf{c}).$$

Recall that since

$$\mu(\boldsymbol{\beta}) = (\beta_0 + \beta_1 x) e^{-\beta_2 x^2}$$

is not a linear function of $\boldsymbol{\beta}$, the D-optimal design will in general depend on the values of β_0 , β_1 , and β_2 .

In practice, β_0 , β_1 , and β_2 are usually unknown. Consequently, estimates $\hat{\boldsymbol{\beta}}$ of $\boldsymbol{\beta}$ were used to obtain the optimal designs given in section 4.3. To allow for uncertainty in the parameter values, we now adopt a Bayesian point of view and introduce a prior distribution for the unknown parameter.

Specifically, we consider $\boldsymbol{\beta}$ as a random vector having a certain prior distribution. Following Chaloner & Larntz (1989), the D-optimal criterion in this Bayesian context is defined as

$$\phi(\mathbf{c}) = E_{\boldsymbol{\beta}} \log \det I(\boldsymbol{\beta}, \mathbf{c}). \quad (4.4.1)$$

Here, $E_{\boldsymbol{\beta}}$ denotes the expectation with respect to random variable $\boldsymbol{\beta}$. $E_{\boldsymbol{\beta}} \log \det I(\boldsymbol{\beta}, \mathbf{c})$ is a concave function of $\mathbf{c}$, satisfying

$$E_{\boldsymbol{\beta}} \log \det I(\boldsymbol{\beta}, \alpha \mathbf{c}_1 + (1 - \alpha) \mathbf{c}_2) \geq \alpha E_{\boldsymbol{\beta}} \log \det I(\boldsymbol{\beta}, \mathbf{c}_1) + (1 - \alpha) E_{\boldsymbol{\beta}} \log \det I(\boldsymbol{\beta}, \mathbf{c}_2).$$

A concave function has a unique maximum value. Although there are at most six design points in the optimal design for the three parameter linear-exponential dose response model in the non-Bayesian designs discussed previously (Chernoff,

1953), there is no *a priori* upper bound of the number of design points (dose levels) in Bayesian optimal designs.

Finding the Bayesian optimal design is a problem of numerical optimization. Specifically, we need to maximize (4.4.1) as a function of the design points $\mathbf{x}$ and the allocations $\mathbf{c}$. In a 3-dose design, for example, we need to find the optimal allocation $\mathbf{c} = (c_1, c_2, c_3)$ once the design points $\mathbf{x} = (x_1, x_2, x_3)$ are fixed, where $0 \leq c_i \leq 1$ with $\sum_{i=1}^3 c_i = 1$, such that $\phi(\mathbf{c})$ is maximized.

The first example in section 4.2.3 will be used to demonstrate the way in which Bayesian optimal designs are developed. Since the exact number of design points for the Bayesian optimal design is unknown, we confine ourselves to finding Bayesian optimal designs with 3-5 doses based on numerical search methods. Consider first a 5-dose design in which the range of doses $[0, 6]$ is divided into 24 subintervals. For simplicity, we only allow x_i to assume values equal to the upper endpoint of these subintervals. For each set of fixed values of x_i ($i = 1, 2, \dots, 5$), the maximized value of $\phi(\mathbf{c})$ and the corresponding values of c_i ($i = 1, 2, \dots, 5$) can be obtained by using the simplex method implemented using the Nelder-Mead algorithm (Nelder, 1965; O'Neill, 1971; NAG, 1992). The optimum choice of $\mathbf{x} = (x_1, x_2, \dots, x_5)$ which maximizes $\phi(\mathbf{c})$ can then be found.

The simplex method is used to find the minimum value of the objective function with respect to the allocations $c_1, c_2, \dots, c_k$. In general, the simplex method used has no restriction on the domain of the variables in these functions. However, for our objective function $\phi(\mathbf{c})$, the variables $c_1, c_2, \dots, c_k$ must satisfy $0 \leq c_i \leq 1$ with $\sum_{i=1}^k c_i = 1$. Using the transformation $c_i = t_i^2 / \sum_{i=1}^k t_i^2$ ($i = 1, \dots, k$), we have $\phi(\mathbf{c}) = \phi^*(\mathbf{t})$. As a result, $\phi^*(\mathbf{t})$ is a function defined on R^k . The simplex method can be applied to the function $\phi^*(\mathbf{t})$. The uniqueness of the maximum value of the function $\phi(\mathbf{c})$ is guaranteed because $\phi(\mathbf{c})$ is a concave function.

In example 1b in section 4.2.3, the estimated values of the model parameters were $\beta_0 = 146$, $\beta_1 = 273\mu g^{-1}$, and $\beta_2 = 0.033$. In practice, we do not know the exact value of β_1 before the experiment is conducted, although we may have some prior knowledge about β_1 . We will use the empirical distribution of the estimated values of β_1 , based on the data for 1-nitropyrene using TA100-S9 as the prior distribution

Table 4.20: An Example of a Bayesian D-optimal Design ^a

No. of Dose levels	Optimal Design Points	Optimal Allocation
3	0.0 2.4 6.0	0.33 0.33 0.33
4	0.0 2.4 1.0 6.0	0.33 0.33 0.00 0.33
5	0.0 0.5 1.8 2.4 6.0	0.33 0.00 0.00 0.33 0.33

^a $\beta_0 = 146$, $\beta_2 = 0.033$, β_1 has prior distribution (4.4.2).

of β_1 given below.

$$P\{\beta_1 = (k - 1) * 100\} = \begin{cases} 0.174 & k = 1, 2, 3, 4 \\ 0.092 & k = 5, 6 \\ 0.0244 & k = 7, 8, 9, 10, 11. \end{cases} \quad (4.4.2)$$

The true values of β_0 and β_2 are assumed to be 143 and 0.033 respectively. The Bayesian D-optimal design are given in Table 4.4.

These results show that only three design points are involved in the Bayesian optimal design. This result is not the same as those obtained by Chaloner & Larntz (1989), who developed Bayesian optimal design for binary data. In particular, they found that as the uncertainty about the unknown parameters increases, the number of design points required to attain the Bayesian optimal design increases for the scale parameter μ in the logistic dose response model, although not for the shape parameter β . Our results for count data agree with the results for the shape parameter in their binary data model. Specifically, our results indicate that even if uncertainty in the values of the unknown regression parameters β is introduced using a prior distribution, a three dose design is sufficient to maximize the Bayesian D-optimality criterion $\phi(c)$.

4.4.2 Bayesian C-Optimal Designs

Recall that the non-Bayesian C-optimality criterion is defined by

$$Var(\hat{\beta}_1) = \sigma^2(0, 1, 0)(\mathbf{D}^T \mathbf{V}^{-1} \mathbf{D})^{-1}(0, 1, 0)^T.$$

Table 4.21: An Example of a Bayesian C-optimal Design^a

No. of dose levels	Optimal Design Points	Optimal Allocation
3	0.0 1.78 6.0	0.160 0.780 0.059
4	0.0 1.78 5.7 6.0	0.163 0.777 0.00 0.067
5	0.0 0.9 1.78 5.7 6.0	0.160 0.00 0.777 0.078

^a $\beta_0 = 146$, $\beta_2 = 0.033$, β_1 has prior distribution (4.4.2).

Let a_{ij} denote the (i, j) element of matrix $\mathbf{D}^T \mathbf{V}^{-1} \mathbf{D}$. Minimizing $\text{Var}(\hat{\beta}_1)$ is equivalent to minimizing the $(2, 2)$ element of the matrix $(\mathbf{D}^T \mathbf{V}^{-1} \mathbf{D})^{-1}$, say a^{22} . Using the adjunct matrix, a^{22} can be calculated as follows:

$$a^{22} = \frac{1}{\sigma^2} \text{Var}(\hat{\beta}_1) = \frac{a_{11}a_{33} - a_{13}a_{31}}{\det \mathbf{D}^T \mathbf{V}^{-1} \mathbf{D}}.$$

Consequently, it is not necessary to calculate the inverse matrix of $\mathbf{D}^T \mathbf{V}^{-1} \mathbf{D}$ in order to obtain the $(2, 2)$ element of $[\mathbf{D}^T \mathbf{V}^{-1} \mathbf{D}]^{-1}$.

The Bayesian C-optimality criteria is

$$\phi_c(\mathbf{c}) = E_{\beta_1} \text{Var}(\hat{\beta}_1).$$

We now find the Bayesian C-optimal design for our example. Let $\beta_0 = 146$ and $\beta_2 = 0.033$, and consider the prior distribution for β_1 given in (4.4.2). Partitioning the dose domain $[0, 6]$ into 20 equal subintervals and using the same numerical methods as in section 4.4.1, we find the Bayesian C-optimal design given in Table 4.5. As with Bayesian D-optimal designs in section 4.4.1, the Bayesian C-optimal designs have only three dose levels.

4.4.3 Bayesian A-Optimal Designs

Recall that the A-optimal designs minimize $\phi_a = \text{Var}(\hat{\beta}_0) + \text{Var}(\hat{\beta}_1) + \text{Var}(\hat{\beta}_2)$. Let a^{ij} denote the (i, j) element of $(\mathbf{D}^T \mathbf{V}^{-1} \mathbf{D})^{-1}$. Using the adjunct matrix, ϕ_a can be calculated as follows:

$$\phi_a = a^{11} + a^{22} + a^{33} = \frac{(a_{22}a_{33} - a_{23}a_{32}) + (a_{11}a_{33} - a_{13}a_{31}) + (a_{11}a_{22} - a_{21}a_{12})}{\det \mathbf{D}^T \mathbf{V}^{-1} \mathbf{D}}.$$

Table 4.22: An Example of a Bayesian A-optimal Design^a

No. of Dose Levels	Optimal Design Points	Optimal Allocation
3	0.0, 0.19, 6.0	0.33, 0.59, 0.07
4	0.0, 0.19, 5.4, 6.0	0.33, 0.59, 0.0, 0.07
5	0.0, 0.8, 0.19, 5.4, 6.0	0.33, 0.0, 0.589, 0.0, 0.07

^a $\beta_0 = 146$, $\beta_2 = 0.033$. β_1 has the prior distribution (4.4.2).

The Bayesian A-optimal design minimizes $E_{\beta_1} \{ \text{Var}(\hat{\beta}_0) + \text{Var}(\hat{\beta}_1) + \text{Var}(\hat{\beta}_2) \}$. The Bayesian A-optimal designs for our example are given in Table 4.6. As with Bayesian C-optimal designs, the Bayesian A-optimal designs also have only three dose levels.

It is of interest to compare the Bayesian optimal designs with the non-Bayesian optimal designs. Suppose the true values of the three regression parameters are $\beta_0 = 146$, $\beta_1 = 273$ and $\beta_2 = 0.033$. Our numerical results have shown that the D-optimal design has three design points, 0.0, 2.0 and 6.0. The corresponding allocations are 0.33, 0.33 and 0.33. In the Bayesian D-optimal design, the three design points are 0.0, 2.5 and 6.0. The use of prior distribution thus shifts the design point x_2 from 2.0 to 2.5. Similar results can also be observed in the Bayesian C and A-optimal designs. Based on this example, introducing prior uncertainty results in a shift of the design points, but does not appear to increase the number of design points as in the case of binary data.

4.5 Optimal Design under Minimax Criteria

For nonlinear models, the optimal designs depend on the unknown model parameters. Therefore, initial estimates or guesses of the values of these unknown parameters are necessary. Bayesian optimal designs provide one way to address uncertainty in the model parameters by using a prior distribution for the parameter values. In this section, another approach to accommodate uncertainty based on the minimax criteria proposed by Sitter (1992) is discussed. Although Sitter (1992) proposed the

use of minimax criteria for binary data, we consider the application of his methods to count data.

Let $\beta^{(0)}$ be an initial estimate of β . Let D denote the set of all the possible designs, i.e., $(x, c) \in D$, and let P be a subset of the parameter space Ω . The region P characterizes the desired region of robustness of the design to poor initial parameter estimates and thus reflects the experimenter's uncertainty about $\beta^{(0)}$ as initial estimates.

The minimax design c^* is chosen to satisfy

$$R(c^*) = \text{Min}_{c \in D} \text{Max}_{\beta \in P} R((x, c), \beta),$$

where $R((x, c), \beta) = \det I^{-1}(x, c, \beta)$ under the D-optimality. Under C-optimality,

$$R((x, c), \beta) = (0, 1, 0) I^{-1}(x, c, \beta) (0, 1, 0)^T = \text{Var}(\hat{\beta}_1).$$

Numerical methods are needed to find the minimax optimal design. However, the optimization problem is more complicated than previously because $\text{Max}_{\beta \in P} R(c, \beta)$ may not be a concave function of c . For this reason, the initial estimates used in the simplex method may affect the maximum value if the criterion function has more than one local maximum. One solution is to evenly distribute a large number of initial parameter values across the domain of c . For a given initial value, one locally maximum design can be obtained by the simplex method. The simplex method is then repeated using another initial estimate value. In this way, we can obtain as many local maximum values (some of which may be identical) as the number of initial estimates. We then choose the design giving the overall maximum value.

In our example, we calculate the maximum values directly. The main idea behind the algorithm used is to discretize the optimality function as well as the design (x, c) . Consider first a 4-dose design with design points x_1, x_2, x_3 , and x_4 and the corresponding allocations c_1, c_2, c_3 , and c_4 , with $0 \leq c_i \leq 1$ and $\sum c_i = 1$. Equivalently, c_1 can be assumed to satisfy the conditions $0 \leq c_1 \leq 1$, $0 \leq c_2 \leq 1 - c_1$, $0 \leq c_3 \leq 1 - c_1 - c_2$, and $0 \leq c_4 \leq 1 - c_1 - c_2 - c_3$. Partitioning $[0, 1]$ into m_s subintervals with equal length of $h = 1/m_s$, we then have $0 \leq kw_1 \leq m_s$ with $c_1 = kw_1 * h$, $0 \leq kw_2 \leq m_s - kw_1$ with $c_2 = kw_2 * h$, $0 \leq kw_3 \leq m_s - kw_1 - kw_2$ with $c_3 = kw_3 * h$ and $c_4 = 1 - c_1 - c_2 - c_3$.

The numerical search can be done under these restrictions. Given fixed design points x_i^* ($i = 1, \dots, M$), the optimal allocations c_i^* ($i = 1, \dots, M$) can be obtained numerically. We can repeat the same numerical search using another set of fixed design points x_i ($i = 1, \dots, M$) and continue until all of the possible sets of fixed design points are evaluated. Finally the overall optimal minimax design can be found by comparing the values of the locally optimal values.

Given the region P , the value of $\text{Max}_{\beta \in P} R(\mathbf{c}, \beta)$ can be found first. In order to find $\min_{\mathbf{c}} \text{Max}_{\beta \in P} R(\mathbf{c}, \beta)$, the numerical search goes through all the possible x_i and for each fixed set of $\{x_i\}$, the optimal allocation c_i are found.

In the D-optimal design,

$$R(\mathbf{c}, \beta) = \det \mathbf{I}^{-1}(\mathbf{c}, \beta)$$

Under C-optimality,

$$R(\mathbf{c}, \beta) = I^{(22)}(\mathbf{c}, \beta),$$

where $I^{(22)}(\mathbf{c}, \beta)$ is the (2,2) element of matrix $\mathbf{I}^{-1}(\mathbf{c}, \beta)$.

When implementing the algorithm, one technical difficulty is that for certain designs $(\mathbf{x}, \mathbf{c})$, $\det \mathbf{I}(\mathbf{c}, \beta) = 0$. Our objective is to find $(\mathbf{x}, \mathbf{c})$ such that $\text{Max}_{\beta \in P} R(\mathbf{c}, \beta)$ attains its minimum with respect to $(\mathbf{x}, \mathbf{c})$. When certain designs, $(\mathbf{x}, \mathbf{c})$ render $R(\mathbf{c}, \beta)$ very large or infinite, this design will not satisfy the minimax criterion. Therefore, the numerical search skips those $\mathbf{c}(\mathbf{x})$ for which $\det \mathbf{I}(\mathbf{x}, \mathbf{c}, \beta) = 0$.

In the case of C-optimality, $R(\mathbf{c}, \beta)$ is the (2,2) element of $[\mathbf{I}(\mathbf{c}, \beta)]^{-1}$. For A-optimality, $R(\mathbf{c}, \beta)$ is the sum of (1,1), (2,2) and (3,3) elements of $[\mathbf{I}(\mathbf{c}, \beta)]^{-1}$. We will use the same example and prior distribution (4.2.2) given in section 4.1.1. We partition the dose range $[0,6]$ into 24 subintervals with equal length. We also partition the interval $[0,1]$ into 30 equal subintervals. Let $h_x = \frac{6.0}{24}$ and $h_c = \frac{1}{30}$. Then the numerical search can be conducted by using $x_i = k_i * h_x$, and $c_i = kw_i * h_c$, where kw_i and k_i are positive integers and kw_i satisfy condition (4.5.2).

Suppose P only consists of a single point

$$\beta^{(0)} = (\beta_0, \beta_1, \beta_2)^T = (146, 273, 0.033)^T.$$

Table 4.23: An Example of a Minimax Optimal Design ^a

Design Criterion	Optimal Design Point	Optimal Allocation
D	0.00 0.25 2.25 6.00	0.33 0.00 0.33 0.33
C	0.00 0.70 2.10 6.00	0.21 0.00 0.69 0.10
A	0.00 0.90 2.11 6.00	0.27 0.00 0.65 0.08

^a $\beta_0 = 146$, $\beta_2 = 0.033$, and β_1 has the prior distribution (4.4.2).

Then maximizing

$$R(\mathbf{c}, \beta^{(0)}) = \det \mathbf{I}^{-1}(\mathbf{c}, \beta^{(0)})$$

yields the optimal design points $\mathbf{x} = (0.00, 0.25, 2.25, 6.0)^T$ and allocation $\mathbf{c} = (0.33, 0.00, 0.33, 0.33)^T$. The same result is obtained via the simplex method.

The following Table 4.7 gives the minimax optimal designs for the same example and prior distribution of β_1 considered in section 4.4.1. Under the minimax criterion, the 4-point designs degenerate to 3-point designs. Similar results were obtained for the M-point design ($M > 3$) under the Bayesian criterion considered in section 4.3.

4.6 Summary

In this chapter, optimal designs for the Ames assay have been developed under the linear-exponential dose response model. Although the primary emphasis is on the development of C, A, and D-optimal designs, consideration was also given to Bayesian optimal designs and designs based on minimax criteria. Since the classical C, A, and D-optimal designs for the linear-exponential model considered here are nonlinear in the model parameters, these designs are dependent on the parameter values.

In an attempt to develop C, A and D-optimal designs of practical relevance, optimal designs were developed for 1120 linear-exponential models derived from a recent collaborative trial with the Ames assay sponsored by the International Programme on Chemical Safety. These designs all involved three dose levels, corresponding to

the number of parameters in the linear-exponential model. Whereas the C and D-optimal designs involve an unbalanced allocation of observations to doses, the A-optimal designs call for equal allocation.

Representative optimal designs were obtained by averaging the C, A, and D-optimal designs across the 1120 different experiments included in the IPCS study. These fixed 3-dose optimal designs generally enjoyed high efficiency relative to the individual 3-dose optimal designs (and were much more efficient than the original 6-dose designs used in the IPCS study), and could be considered for use in practice in the absence of knowledge of the model parameters

Fixed suboptimal designs with 4 - 6 doses obtained by augmenting the representative 3-dose designs were also considered. Additional design points are useful in defining the shape of the dose-response curve, testing for lack of fit, and protecting against the loss of data at one or more doses. The fixed designs with equal allocation were notably less efficient than the fixed designs with unequal allocation. In general, the efficiency of these fixed 3-6 dose designs decreased with the number of design points.

In comparison with this in-depth analysis of C, A, and D-optimal designs for the Ames assay, Bayesian and minimax optimal designs were treated only briefly. The purpose of both the Bayesian and minimax designs is to allow for uncertainty in the values of the model parameters. (In the previous analysis, such uncertainty is implicitly allowed for through the development of representative designs obtained by averaging the optimal designs over a variety of conditions encountered in practice.) In the Bayesian case, uncertainty is expressed by means of a prior distribution on the model parameters; in the minimax case, the maximum value of the objective function is minimized within a specified region of the parameter space. In contrast with previous results for binary data, the Bayesian and minimax optimal designs for count data arising in the Ames assay did not involve more than three design points in the limited number of examples considered in this chapter.

Chapter 5

Rao-Scott Transformation

Counts of mutant colonies obtained from the Ames *Salmonella*/microsome assay are subject to extra-Poisson variation. Consequently, direct methods of analysis of data from the Ames assay need to provide for such overdispersion. Rao & Scott (1992) proposed a simple data transformation that can be used to eliminate overdispersion. The Rao-Scott transformation is based on the concept of a design effect in survey sampling, which relates the variance of the observations under cluster sampling to the variance under simple random sampling. Recently, Scott & Rao (1994) extended the Rao-Scott transformation method to count data. After applying the transformation, standard methods of analysis designed for Poisson data can be used.

In this chapter, the Rao-Scott transformation is applied to develop simplified methods of (i) testing for trend, (ii) fitting the linear-exponential model, and (iii) assessing goodness-of-fit. In section 5.1, the Rao-Scott transformation is used to modify the tests for trend in mutagenic response with increasing dose discussed in chapter 3. Previously Fung et al. (1994) used the Rao-Scott transformation in developing simplified tests for trend with overdispersed binomial data. The transformation is applied in fitting the linear-exponential model in section 5.2. This application is similar in spirit to work done by Krewski & Zhu (1994) in fitting dose response models to developmental toxicity data subject to extra-multinomial variation. One important advantage of modeling the transformed data is that the need to use quadratic estimating equations to estimate the overdispersion parameter is eliminated. In section 5.3, the application of goodness-of-fit tests on both

the original and transformed scales is discussed. Again, testing for goodness of fit after transformation avoids the need to allow for overdispersion in constructing an appropriate test statistic.

5.1 Test for Trend

5.1.1 Test Statistics

Depending on how the design effect reflecting intraclass correlation is estimated, different versions of the Rao-Scott transformation are possible (cf. Fung et al., 1994). Here, we consider two versions of the transformation for use in the Ames assay. In order to avoid complete specification of the distribution of the data, we will work only with the first two moments. The first version is essentially an *ad hoc* method designed to ensure that the mean and variance of the transformed data are equal. The second version is the method given by Scott & Rao (1994). Following the application of either transformation, generalized score tests based on generalized estimating equations are developed under the assumption that the mean and variance of the data are equal.

Method 1

Let μ_i and V_i denote the mean and variance of the observation Y_{ij} respectively. The design effect is then defined by $D_i = V_i/\mu_i$. Applying the transformation

$$\hat{Y}_{ij} = \frac{Y_{ij}}{D_i}, \quad (5.1.1)$$

we have

$$\left. \begin{aligned} E\hat{Y}_{ij} &= \mu_i/D_i \\ \text{Var}\hat{Y}_{ij} &= V_i/[D_i]^2 = \mu_i/D_i \end{aligned} \right\}. \quad (5.1.2)$$

Hence, $E\hat{Y}_{ij} = \text{Var}\hat{Y}_{ij}$.

Based on the transformed data $\hat{Y}_{ij}$ in (5.1.1), the score-type statistic T in (3.6.2) can be written as

$$T = \sum_{i=1}^K (\hat{Y}_i - n_i \hat{\mu}^*) d_i \quad (5.1.3)$$

under the null hypothesis $\beta_i = 0$, where $\hat{Y}_i = \sum_{j=1}^{n_i} Y_{ij} / D_i = Y_i / D_i$, $\hat{\mu}^* = \sum_{i=1}^K \hat{Y}_i / \sum_{i=1}^K n_i$. The variance of T is estimated by

$$\hat{V}(T) = \hat{\mu}^* \left\{ \sum_{i=1}^K n_i d_i^2 - n_T^{-1} \left(\sum_{i=1}^K n_i d_i \right)^2 \right\}, \quad (5.1.4)$$

where $n_T = \sum_{i=1}^K n_i$ (also see Tarone (1982)).

Under the null hypothesis $\beta_i = 0$, consistent estimates of μ_i and V_i are given by

$$\hat{\mu}_i = \bar{Y} = \frac{\sum_{i=1}^K Y_i}{\sum_{i=1}^K n_i} \quad (5.1.5)$$

and

$$\hat{V}_i = \frac{\sum_{i=1}^K \sum_{j=1}^{n_i} (Y_{ij} - \bar{Y})^2}{\sum_{i=1}^K n_i} \quad (5.1.6)$$

respectively. Therefore, a consistent estimate $\hat{D}_i$ of D_i is

$$\hat{D}_i = \frac{\hat{V}_i}{\hat{\mu}} = \frac{\hat{V}_i}{\bar{Y}}. \quad (5.1.7)$$

Note that since we are working under the null hypothesis, $\hat{D}_i$ is the same for all $i = 1, \dots, K$.

The trend statistic based on the transformed data is given by

$$Z = \frac{T}{[\hat{V}(T)]^{1/2}}. \quad (5.1.8)$$

This statistic is a special case of (3.6.2) with $\tau_1 = 0$, and is asymptotically standard normal as $\min\{n_i\} \rightarrow \infty$.

The test statistic given in (5.1.8) is of exactly the same form as that developed in (3.6.2), if the mean and variance satisfy the relationship $Variance = Mean(1 + \phi Mean)$. (This can be verified by noticing that in (3.6.2), $\hat{D}_i = 1 + \hat{\mu}_i \hat{\phi}$). The advantage of the trend test given in (5.1.8) based on the transformed data is that it does not require specification of the relationship between the mean and variance of the data beyond that implied by the Poisson distribution. Since the transformed data are not subject to extra-Poisson variation, it is also not necessary to estimate the value of the overdispersion parameter ϕ .

Next, we consider another method of test for trend.

Method 2

Let $Y_i = \sum_{j=1}^{n_i} Y_{ij}$ and $\dot{Y}_i = Y_i / \hat{D}_i^*$, where $\hat{D}_i^*$ is the estimated design effect given by Rao & Scott (1994), and defined by

$$\hat{D}_i^* = \frac{n_i}{\bar{Y}_i} \sum_{j=1}^{n_i} (Y_{ij} - \frac{Y_i}{n_i})^2 / (n_i - 1). \quad (5.1.9)$$

The moments of the transformed data are given by

$$\mu_i^* = E\dot{Y}_i \approx \text{Var}\dot{Y}_i \approx \mu_i \dot{t}_i, \quad (5.1.10)$$

where $\dot{t}_i = n_i / \hat{D}_i^*$.

Under the null hypothesis $\beta_1 = 0$, ($\mu_i = \beta_0$) Applying the estimating equation (3.3.6) to the $\dot{Y}_i$, we obtain the score-type statistic

$$T' = \sum_{i=1}^K (\dot{Y}_i - \hat{\beta}_{0i}) d_i \quad (5.1.11)$$

with

$$\text{Var}T' = \sum_{i=1}^K \hat{\beta}_{0i} d_i^2 - \frac{(\sum_{i=1}^K \hat{\beta}_{0i} d_i)^2}{\sum_{i=1}^K \hat{\beta}_{0i}}, \quad (5.1.12)$$

where $\hat{\beta}_{0i} = \hat{\beta}_0 n_i / \hat{D}_i^*$. The standardized statistic

$$\frac{T'}{\sqrt{\text{Var}T'}} \quad (5.1.13)$$

is asymptotically standard normal as K and $\min\{n_i\} \rightarrow \infty$.

The estimates $\hat{\beta}_{0i}$ of β_{0i} can be obtained by solving the equation

$$\sum_{i=1}^K \frac{\dot{Y}_i - \hat{\beta}_{0i}}{\hat{\beta}_{0i}} = 0. \quad (5.1.14)$$

In fact, only $\hat{\beta}_0$ needs to be estimated from (5.1.14), since $\hat{\beta}_{0i} = \hat{\beta}_0 n_i / \hat{D}_i^*$. Under the null hypothesis $\beta_1 = 0$, we can use the pooled design effect

$$\hat{D}^* = \frac{n_T}{Y} \sum_{i=1}^K \sum_{j=1}^{n_i} (Y_{ij} - \frac{Y}{n_T})^2 / (n_T - 1),$$

where $Y = \sum_{i=1}^K Y_i$ and $n_T = \sum_{i=1}^K n_i$. If $n_i = n_{i'}$, $\hat{D}_i^* = \hat{D}_{i'}^* = \hat{D}^*$ for $i \neq i'$, we have $\hat{\beta}_{0i} = \sum_{i=1}^K \dot{Y}_i / K$. More generally, $\hat{\beta}_0$ can be solved by $\hat{\beta}_0 = [\sum_{i=1}^K (\dot{Y}_i / \dot{n}_i)] / K$, where $\dot{n}_i / \hat{D}_i^*$

5.1.2 Simulation Results

A limited simulation study was conducted to evaluate the type I and type II errors of the statistics in (5.1.8) and (5.1.13). The random variables Y_{ij} are generated by the following two steps.

- (1) Generate $\lambda_{ij} \sim \text{Gamma}(g, h_i)$ with $gh_i = \beta_0 + \beta_1 d_i$. Then, for each λ_{ij} , $E\lambda_{ij} = gh_i$ and $\text{Var}\lambda_{ij} = gh_i(1 + g^{-1}gh_i) = gh_i(1 + h_i)$.
- (2) For each random variable λ_{ij} generated in (1), generate $Y_{ij} \sim \text{Poisson}(\lambda_{ij})$. It is easy to verify that

$$EY_{ij} = gh_i = \beta_0 + \beta_1 d_i$$

and

$$\text{Var}(Y_{ij}) = gh_i(1 + g^{-1}gh_i) = gh_i(1 + h_i).$$

In the simulation, we took $g = 6.5$ and $\beta_0 = 8.19$.

The type I errors of tests for trend based on methods 1 and 2 are shown in Table 5.1 based on a nominal significance level of 5%. The type II errors are calculated under the alternative hypothesis with $\beta_1 = 25$. Two sample sizes are considered: $n_i = 9$ with $K = 5$ and $\sum_{i=1}^K n_i = 45$, and $n_i = 50$ with $K = 5$ and $\sum_{i=1}^K n_i = 250$. The results in Table 5.1 are based on 1000 simulations.

Although the type I error rates for method 1 is lower than that for method II, both methods have an inflated type I error rate when $n_i = 9$. With $n_i = 50$, however, method 1 produces a type I error rate close to the rate of 5%. Despite its inflated type I error rate, method 2 does not give greater power than method 1.

5.2 Parameter Estimation

In this section, we consider the problem of estimating the regression parameters in the linear-exponential model using estimating equation methods. In the presence of overdispersion, an additional estimation equation is needed to estimate the overdispersion parameter. To apply these methods, we must specify the relationship between the mean and variance of the data. Specincation of the mean-variance

Table 5.1: Type I and Type II Errors of Tests for Trend at a Significance Level of 5%

Trend Test	Type I Error $n_i = 9, K = 5$	Type I Error $n_i = 50, K = 5$	Type II Error $n_i = 9, K = 5$	Type II Error $n_i = 50, K = 5$
<i>Method - 1</i>	0.075	0.04	0.54	0.001
<i>Method - 2</i>	0.20	0.075	0.59	0.001

relationship and estimation of the overdispersion parameter is not required if the Rao-Scott transformation is used.

5.2.1 Parameter Estimation without Using the Rao-Scott Transformation

As discussed in chapter 3, we suppose that the observations $\{Y_{ij}\}$ are generated as follows. Consider random variates $\lambda_{ij} \sim \text{Gamma}(g, h_i)$ such that $E\lambda_{ij} = gh_i$ and $\text{Var}\lambda_{ij} = gh_i^2$. Conditional on λ_{ij} , we suppose that $Y_{ij}|\lambda_{ij} \sim \text{Poisson}(\lambda_{ij})$. Unconditionally,

$$EY_{ij} = gh_i = \mu_i = (\beta_0 + \beta_1 d_i) \exp(-\beta_2 d_i^2) \quad (5.2.1)$$

and

$$V(\mu_i, \phi) = \text{Var}Y_{ij} = \mu_i(1 + \mu_i\phi), \quad (5.2.2)$$

where $\phi = g^{-1}$.

Consider now the quasi-information matrix $\mathbf{B}$, with

$$\mathbf{B}_{st} = \sum_{i=1}^K \frac{n_i}{V(\mu_i, \phi)} \left(\frac{\partial \mu_i}{\partial \beta_s} \right) \left(\frac{\partial \mu_i}{\partial \beta_t} \right). \quad (5.2.3)$$

The Newton-Raphson procedure for estimating $\beta = (\beta_0, \beta_1, \beta_2)^T$ can be written as

$$\hat{\beta}_s^{\text{new}} = \hat{\beta}_s^{\text{old}} + \sum_{t=0}^2 (\mathbf{B}^{-1})_{st} \sum_{i=1}^K [(Y_i - n_i \mu_i) / V(\mu_i, \phi)] \frac{\partial \mu_i}{\partial \beta_s} \quad (5.2.4)$$

($s = 0, 1, 2$). The moment equation for the overdispersion parameter is

$$\sum_{j=1}^{n_i} \frac{(Y_{ij} - \mu_i)^2}{\mu_i(1 + \mu_i\phi)} = n_i, \quad (5.2.5)$$

which leads to

$$\hat{\phi}_i = \mu_i^{-1} [\mu_i^{-1} n_i^{-1} \sum_{j=1}^{n_i} (Y_{ij} - \mu_i)^2 - 1]. \quad (5.2.6)$$

Averaging over the K dose levels, we have

$$\hat{\phi} = \sum_{i=1}^K \hat{\phi}_i / K. \quad (5.2.7)$$

As shown by Moore (1986), the estimators $\hat{\beta}$ and $\hat{\phi}$ are consistent as $\min\{n_i\} \rightarrow \infty$. The simulation results based on (5.2.4) and (5.2.7) will be given in section 5.2.3.

5.2.2 Parameter Estimation Using the Rao-Scott Transformation

In this section, we apply the Rao-Scott transformation to the observations Y_{ij} , and then estimate the model parameters based on the transformed data. Specifically, consider the Rao-Scott transformation based on method 2 in section 5.1.1. As shown in section 5.1.1, the mean and variance of the transformed data $\hat{Y}_i$ are approximately equal.

Using (3.3.1), the Newton-Raphson equations can be written as

$$\hat{\beta}_s^{new} = \hat{\beta}_s^{old} + \sum_{t=0}^2 (\mathbf{B}^*)_{st}^{-1} \sum_{i=1}^K [(\hat{Y}_i - \mu_i^*) / \mu_i^*] \frac{\partial \mu_i^*}{\partial \beta_s}, \quad (5.2.8)$$

where the matrix $\mathbf{B}_{st}^*$ is defined by

$$\mathbf{B}_{st}^* = \sum_{i=1}^K \frac{1}{\mu_i^*} \left(\frac{\partial \mu_i^*}{\partial \beta_s} \right) \left(\frac{\partial \mu_i^*}{\partial \beta_t} \right), \quad (5.2.9)$$

The estimated regression parameters $\hat{\beta}_0$, $\hat{\beta}_1$ and $\hat{\beta}_2$ are completely determined by (5.2.8), since there is no overdispersion parameter to be estimated.

Table 5.2: Estimates of the Parameter in the Linear-Exponential Model Before and After Applying the Rao-Scott Transformation^a

Model Fitting	Sample Size [*]	$\hat{\beta}_0 \pm SD$	$\hat{\beta}_1 \pm SD$	$\hat{\beta}_2 \pm SD$
Before Rao-Scott Transformation	$n_i = 5$	9.49 ± 1.05	20.06 ± 5.43	0.11 ± 0.22
	$n_i = 10$	7.1 ± 0.60	24.8 ± 3.45	0.17 ± 0.13
	$n_i = 50$	7.37 ± 0.27	21.8 ± 1.50	0.06 ± 0.06
After Rao-Scott Transformation	$n_i = 5$	10.35 ± 0.77	22.25 ± 4.20	0.17 ± 0.11
	$n_i = 10$	7.09 ± 0.58	25.01 ± 3.31	0.17 ± 0.12
	$n_i = 50$	7.38 ± 0.25	21.6 ± 1.44	0.07 ± 0.05

^{*} $K = 5$.

^a Based on 1000 simulations with $\beta_0 = 8.09$, $\beta_1 = 20$, and $\beta_2 = 0.2$.

5.2.3 Simulation Results

A brief simulation study was conducted using the parameter estimation methods given in both section 5.2.1 and section 5.2.2. The true parameter values used in the simulation studies are $\beta_0 = 8.19$, $\beta_1 = 20$, and $\beta_2 = 0.2$. The total number of dose levels is set to $K = 5$, with sample sizes of $n_i = 5, 10$ or 50 . The average value of the parameter estimates based on 1,000 simulations are shown in Table 5.2, along with the standard derivation (SD) of these estimates. These results suggest that the regression parameter estimates obtained using the Rao-Scott transformation are similar to those obtained using the original data.

We now examine these two sets of estimates in more detail. In particular, we consider the relative difference

$$\delta_i = \frac{\hat{\beta}_i^{(B)} - \hat{\beta}_i^{(A)}}{\hat{\beta}_i^{(A)}} \times 100\% \quad (5.2.10)$$

of the estimated values of β_i before (B) and after (A) transformation, as well as the relative standard error

$$\Delta_i = \frac{SE(\hat{\beta}_i^{(B)}) - SE(\hat{\beta}_i^{(A)})}{SE(\hat{\beta}_i^{(A)})} \times 100\% \quad (5.2.11)$$

The average values of these two measures indicate the extent to which the parameter estimates and their estimated standard errors differ before and after transformation. The results in Table 5.3 indicate that the two sets of parameter estimates are in good agreement. However, the standard errors of the estimates appear smaller following transformation.

We also consider the relative bias defined by

$$RB = \frac{\hat{\beta}_i - \beta_i}{\beta_i} \times 100\%$$

and the relative standard deviation

$$RSD = \left(\frac{V(\hat{\beta}_i)}{\beta_i^2} \right)^{0.5} \times 100\%$$

These two measures may be combined in the form of the relative mean square error

$$RMSE = [RB^2 + RSD^2]^{1/2}.$$

The results given in Table 5.4 demonstrate that relative bias, relative standard deviation and relative mean squared error are comparable before and after transformation even if the sample size n , is as small as 10. However, the relative bias and relative standard deviation are larger when the true value of β_2 is small.

5.3 Goodness-of-Fit Test

In the previous chapter, we used the linear-exponential model to describe the mean response in relation to dose. We now develop goodness-of-fit tests to examine whether or not the mean response is of the linear-exponential form. Two goodness-of-fit methods are proposed, based on the original data and the transformed data. It is shown that evaluating goodness-of-fit after applying the Rao-Scott transformation has several advantages compared to assessing goodness-of-fit based on the original data.

Table 5.3: Comparison of Parameter Estimates Before and After Applying the Rao Scott Transformation^a

β_2	Sample Size (n_i)	Avg. δ_i			Avg. Δ_i		
		β_0	β_1	β_2	β_0	β_1	β_2
0.2	10	1.1e-2	1.5e-3	-0.46	0.61	0.61	0.62
	50	4.2e-4	3.2e-3	-5.0e-2	0.65	0.65	0.64
0.5	10	-2.3e-3	2.8e-2	-1.5e-2	0.57	0.61	0.61
	50	-1.5e-4	8.9e-3	1.2e-2	0.63	0.64	0.66
1.0	10	-1.4e-3	3.0e-2	5.7e-2	0.59	0.59	0.60
	50	-9.0e-5	-1.8e-4	3.5e-4	0.64	0.65	0.65

^a Based on 50 simulations with $\beta_0 = 8.19$, $\beta_1 = 20$.

5.3.1 Goodness-of-Fit Tests without Using the Rao-Scott Transformation

Consider the linear-exponential dose response model

$$\left. \begin{aligned} \mu_i &= EY_{ij} = (\beta_0 + \beta_1 d_i) \exp(-\beta_2 d_i^2) \\ V(\mu_i, \phi) &= \text{Var}(Y_{ij}) = \mu_i(1 + \mu_i \phi) \end{aligned} \right\}. \quad (5.3.1)$$

Recall that the parameters $\beta_0, \beta_1, \beta_2$ and ϕ are estimated simultaneously using the estimating equations

$$\sum_{i=1}^K \frac{Y_i - n_i \mu_i}{V(\mu_i, \phi)} \frac{\partial \mu_i}{\partial \beta_s} = 0 \quad (5.3.2)$$

($s = 0, 1, 2$) and the moment equation

$$\sum_{i=1}^K \left[\frac{(Y_i - n_i \mu_i)^2}{n_i V(\mu_i, \phi)} - 1 \right] = 0 \quad (5.3.3)$$

or

$$\sum_{i=1}^K \sum_{j=1}^{n_i} \left[\frac{(Y_{ij} - \mu_i)^2}{\mu_i(1 + \mu_i \phi)} - 1 \right] = 0. \quad (5.3.3')$$

Table 5.4: Relative Bias (RB), Relative Standard Deviation (RSD), and Relative Mean Squared Error (RMSE) of Regression Parameters Estimation Before and After the Rao-Scott Transformation^a

Ex.	β_2	S^3 n_i	RB			RSD			RMSE		
			β_0	β_1	β_2	β_0	β_1	β_2	β_0	β_1	β_2
I^1	0.2	10	5.1e-3	-7.1e-2	-0.52	3.0e-2	7.5e-5	3.0	3.0e-2	0.15	3.3
		50	2.3e-4	-2.1e-2	-7.4e-2	6.6e-3	1.7e-5	0.58	6.6e-3	3.2e-2	0.59
	0.5	10	1.1e-2	-1.6e-2	-0.1	3.8e-2	9.6e-5	0.50	3.8e-2	0.19	0.51
		50	-4.4e-3	1.9e-2	4.6e-2	9.9e-3	2.5e-5	0.1	9.8e-3	3.6e-2	0.10
	1.0	10	3.2e-3	-2.2e-2	-4.1e-2	2.4e-2	6.1e-5	0.01	2.4e-2	0.11	0.10
		50	-2.4e-2	3.2e-2	2.4e-2	6.2e-3	1.5e-5	3.1e-2	6.8e-3	3.9e-2	3.2e-2
II^2	0.2	10	2.0e-2	2.4e-2	0.17	3.2e-2	8.0e-5	1.11	3.2e-2	0.14	1.12
		50	3.6e-3	3.3e-2	0.16	5.9e-3	2.3e-5	0.55	9.2e-3	3.7e-2	0.58
	0.5	10	-1.6e-2	-2.2e-2	-1.8e-3	2.5e-2	6.3e-5	0.32	2.5e-2	9.5e-2	0.32
		50	-1.5e-2	1.4e-3	4.9e-3	5.4e-3	1.3e-5	8.1e-2	5.6e-3	2.5e-2	8.1e-2
	1.0	10	7.0e-3	2.3e-2	-3.6e-2	3.1e-2	7.8e-5	0.11	3.1e-2	0.15	0.11
		50	-5.1e-3	-1.5e-2	-1.8e-2	7.1e-3	1.8e-5	2.8e-2	7.1e-3	3.8e-2	2.8e-2

^a: Based on 50 simulation with $\beta_0 = 8.19$ and $\beta_1 = 20.0$.

¹: Before the Rao-Scott Transformation.

²: After the Rao-Scott Transformation.

³: Sample Size.

where $Y_i = \sum_{j=1}^{n_i} Y_{ij}$. The objective of the goodness-of-fit test is to examine whether or not the mean μ_i is of the linear-exponential form (5.3.1).

The development of the goodness-of-fit test is based on quasi-likelihood theory (see e.g. Dobson, 1990). Before the discussion of the goodness-of-fit test, we first briefly review some of the relevant concepts of the quasi-likelihood function.

The quasi-likelihood function $l(\mu, Y)$ for a random variable Y with mean $E(Y) = \mu$ satisfies the partial differential equation (Wedderburn, 1974)

$$\frac{\partial l(\mu, Y)}{\partial \mu} = V^{-1}(\mu, \phi)(Y - \mu).$$

Following Leroux and Krewski (1993), consider a quasi-likelihood function with K distinct parameters (excluding any overdispersion parameters). Such a saturated or maximal model uses a different parameter to represent the mean of the observations at different dose levels. Let $\mu^* = (\mu_1^*, \dots, \mu_K^*)^T$ denote the parameters in the maximal model and let $Y = (Y_1, \dots, Y_K)^T$ denote the observation at each dose. The deviance is defined by

$$D(Y, \mu) = 2[l(\hat{\mu}^*, Y) - l(\hat{\mu}, Y)] = 2 \sum_{i=1}^K [l(\hat{\mu}_i^*, Y_i) - l(\hat{\mu}_i, Y_i)], \quad (5.3.4)$$

where $l(\cdot, \cdot)$ is the logquasi-likelihood of the data Y_i . $\hat{\mu}_i^*$ denotes the quasi-likelihood of the mean under the maximal model, and $\hat{\mu}_i$ estimated mean response, under the linear-exponential model. We consider $l(\mu^*, Y)$ as the maximum logquasi-likelihood achievable for an exact fit in which fitted values equal the data (McCullagh & Nelder, 1983).

The maximal or saturated model has K parameters μ_i^* whereas the model of interest has three parameters, β_0 , β_1 , and β_2 , excluding the overdispersion parameter ϕ . Because the maximal model can be regarded as providing a complete description of the data, $l(\hat{\mu}_i^*, Y_i) - l(\hat{\mu}_i, Y_i)$, the difference between the log quasi-likelihood function for the maximal model and the linear-exponential model of interest, provides a measure of goodness-of-fit for the model of interest. If the model $l(\mu_i, Y_i)$ describes the data well, then the value of $l(\hat{\mu}_i, Y_i)$ should be close to that of $l(\hat{\mu}_i^*, Y_i)$.

It is easy to verify that $\hat{\mu}_i^* = \bar{Y}_i = Y_i/n_i$. Therefore, Leroux & Krewski (1993)

proposed the goodness-of-fit test

$$\begin{aligned} \hat{GF} &= 2 \sum_{i=1}^K [l(\bar{Y}_i, \phi; Y_i) - l(\hat{\mu}_i, \phi; Y_i)] \\ &= 2 \sum_{i=1}^K [Y_i \ln \frac{\bar{Y}_i}{\hat{\mu}_i} - (Y_i + n_i \phi^{-1}) \ln \frac{1 + \bar{Y}_i \phi}{1 + \hat{\mu}_i \phi}] \end{aligned} \quad (5.3.5)$$

Since K is fixed, the asymptotic distribution of $\hat{GF}$ is chi-squared with $K - 3$ degrees of freedom as $\min\{n_i\} \rightarrow \infty$. If β_0 , β_1 , and β_2 are known, then $\hat{GF} \sim \chi_K^2$.

For this method to be usable, we require that the observations Y_{ij} satisfy the moment conditions

$$\left. \begin{aligned} EY_{ij} &= \mu_i^*, \\ \text{Var}(Y_{ij}) &= \mu_i^* (1 + \mu_i^* \phi) \end{aligned} \right\}. \quad (5.3.6)$$

If this mean-variance relationship does not hold, or the estimate of ϕ is inconsistent, then $\hat{GF}$ in (5.3.5) may not follow a chi-squared distribution even if $\mu_i = EY_{ij}$ is of the linear-exponential form.

In order to address these concerns we will use the Rao-Scott transformation to develop a goodness-of-fit test in section 5.3.2. After applying this transformation the mean-variance relationship (5.3.1) is no longer a requirement, nor is estimation of the overdispersion parameter ϕ .

5.3.2 Goodness-of-Fit Test Using the Rao-Scott Transformation

Let V_i denote the variance of the observation Y_{ij} at dose level d_i . After applying the Rao-Scott transformation, $\hat{Y}_{ij} = Y_{ij}/D_i$, we have

$$\hat{\mu}_i = E\hat{Y}_{ij} = \frac{\mu_i}{D_i}$$

and

$$\text{Var}(\hat{Y}_{ij}) = \frac{V_i}{D_i^2} = \frac{D_i \mu_i}{D_i^2} = \frac{\mu_i}{D_i}.$$

Therefore, for the transformed data, $\hat{Y}_{ij}$, we have $E\hat{Y}_{ij} = \text{Var}(\hat{Y}_{ij})$.

The quasi-likelihood function for the transformed data is

$$\begin{aligned}\frac{\partial l(\tilde{\mu}, \hat{Y})}{\partial \tilde{\mu}} &= V^{-1}(\tilde{\mu})(\hat{Y} - \tilde{\mu}) \\ &= \frac{1}{\tilde{\mu}}(\hat{Y} - \tilde{\mu}) = \frac{\hat{Y}}{\tilde{\mu}} - 1.\end{aligned}\quad (5.3.7)$$

with

$$l(\tilde{\mu}, \hat{Y}) = \hat{Y} \ln \tilde{\mu} - \tilde{\mu} + C(\hat{Y}). \quad (5.3.8)$$

Let μ_i^* denote the mean of the observations at the dose level d_i under a maximal model with K parameters. Solving the quasi-likelihood function

$$\frac{\partial l(\mu_i^*, \hat{Y}_i)}{\partial \mu_i^*} = V^{-1}(\mu_i^*)(\hat{Y}_i - n_i \mu_i^*) = 0$$

yields

$$\hat{\mu}_i^* = \frac{\sum_{j=1}^{n_i} \hat{Y}_{ij}}{n_i} = \frac{\hat{Y}_i}{n_i}. \quad (5.3.9)$$

Hence,

$$l(\hat{\mu}_i^*, \hat{Y}_i) = \hat{Y}_i \ln \frac{\hat{Y}_i}{n_i} - \hat{Y}_i + C(\hat{Y}_i). \quad (5.3.10)$$

The quasi-likelihood ratio test statistic, GF^* , can be written as

$$GF^* = 2 \sum_{i=1}^K [l(\hat{\mu}_i^*, \hat{Y}_i) - l(\tilde{\mu}_i, \hat{Y}_i)]. \quad (5.3.11)$$

Substituting (5.3.8), (5.3.9) and (5.3.10) into (5.3.11) yields

$$GF^* = 2 \sum_{i=1}^K [\hat{Y}_i \ln \frac{\hat{Y}_i/n_i}{\tilde{\mu}_i} - (\frac{\hat{Y}_i}{n_i} - \tilde{\mu}_i) n_i]. \quad (5.3.12)$$

We can also write GF^* in terms of original data Y_i as follows:

$$\begin{aligned}GF^* &= 2 \sum_{i=1}^K \left\{ \frac{Y_i}{D_i} \ln \frac{\bar{Y}_i}{\mu_i} - n_i \frac{\bar{Y}_i - \mu_i}{D_i} \right\} \\ &= 2 \sum_{i=1}^K \frac{1}{D_i} \{ Y_i \ln \frac{\bar{Y}_i}{\mu_i} - n_i (\bar{Y}_i - \mu_i) \}.\end{aligned}\quad (5.3.13)$$

where $\mu_i = (\beta_0 + \beta_1 d_i) \exp(-\beta_2 d_i^2)$. Given the values of β_0 , β_1 and β_2 , the asymptotic distribution of the test statistic GF^* is chi-squared with K degrees of freedom (χ_K^2), as $\min\{n_i\} \rightarrow \infty$ (see section 5.3.4), provided the linear-exponential model holds.

Following Scott & Rao (1994), we will use the estimated design effect

$$\hat{D}_i = \frac{\sum_{j=1}^{n_i} (Y_{ij} - \bar{Y}_i)^2 / (n_i - 1)}{\bar{Y}_i}. \quad (5.3.14)$$

Since the numerator and the denominator in (5.3.14) are consistent estimates of V_i and μ_i respectively, $\hat{D}_i$ is also a consistent estimate of D_i . Replacing D_i by $\hat{D}_i$ in (5.3.13) gives

$$\hat{GF}^* = 2 \sum_{i=1}^K \frac{1}{\hat{D}_i} [Y_i \ln \frac{\bar{Y}_i}{\mu_i} - n_i (\bar{Y}_i - \mu_i)]. \quad (5.3.15)$$

It follows from Slutsky's theorem that $\hat{GF}^*$ and GF^* have the same asymptotic distribution. We therefore propose to use $\hat{GF}^*$ as goodness-of-fit test.

There are two advantages of (5.3.15) as compared to (5.3.5). First, in (5.3.15) the relationship between the mean and variance of the observations need not be specified. However, we must assume that $Variance = Mean(1 + \phi Mean)$ in (5.3.5). Recall that the objective of the goodness-of-fit test is to examine whether or not the mean response is of the linear-exponential form. Violation of the mean-variance relation $Variance = Mean(1 + \phi Mean)$ when using (5.3.5) may affect the value of the test statistic $\hat{GF}$ in (5.3.5): a large value of $\hat{GF}$ may not be due to lack-of-fit of the linear-exponential dose response model, but rather to incorrect specification of the the mean-variance relationship. Second, (5.3.15) does not involve the overdispersion parameter ϕ . However, one possible advantage of using (5.3.5) is that if the data actually satisfy the assumed mean-variance relationship at all dose levels, then we may get a better estimate of ϕ by using all of the data from all of the dose groups.

5.3.3 Simulation Results

In this section, we present some simulation results of the sensitivity study of the goodness-of-fit tests given in (5.3.15) and (5.3.5). Recall that the objective of the goodness-of-fit test is to examine whether or not the mean of observations is of the linear-exponential form with respect to dose. A desirable goodness-of-fit test should be sensitive to departures from the linear-exponential form. We will consider the following cases separately.

Case 1.

The observations are generated by mixing gamma and Poisson variates. Consider the random variable $\lambda_{ij} \sim G(g, h_i)$ with $E\lambda_{ij} = gh_i$ and $Var(\lambda_{ij}) = gh_i^2$. We further assume that $Y_{ij}|\lambda_{ij} \sim Poisson(\lambda_{ij})$. Unconditionally

$$\left. \begin{aligned} \mu_i &= EY_{ij} = gh_i, \\ VarY_{ij} &= gh_i[1 + g^{-1}(gh_i)] = \mu_i(1 + \mu_i\phi). \end{aligned} \right\} \quad (5.3.16)$$

Since

$$\mu_i = (\beta_0 + \beta_1 d_i) \exp(-\beta_2 d_i^2), \quad (5.3.17)$$

h_i can be written as

$$h_i = (\beta_0 + \beta_1 d_i) \exp(-\beta_2 d_i^2) / g. \quad (5.3.18)$$

In the simulation study, we calculate the type I and type II errors based on 1000 repeated simulations. The results are summarized in Table 5.5 with GOF(B) and GOF(A) denoting the goodness-of-fit statistics without and with the Rao-Scott transformation respectively.

After the data are generated, we use the estimating equation (5.3.2) and moment equation (5.3.3) or (5.3.3') to estimate the parameters. Then, GOF(B) and GOF(A) were calculated, based on the estimated values of the parameters, by (5.3.5) and (5.3.15) respectively. The results are as follows.

GOF(A) gives a more reasonable type I error than GOF(B). One possible explanation for this is that when calculating GOF(B), the value of ϕ needs to be estimated. If the estimate $\hat{\phi}$ of ϕ based on (5.3.3) is inaccurate, the value of GOF(B) will be affected. To verify this, we also used (5.3.3') to estimate the parameter ϕ , which gives a much better estimate. The resulting type I error of GOF(B) reduces to 5.09%.

Case 2.

In this example, the random variables are not generated from the linear-exponential form as in the previous example. Instead, we use the model

$$EY_{ij} = \mu_i = \alpha_0 + \alpha_1 d_i^\theta.$$

However, in applying the data, we still incorrectly assume that the linear-exponential model (5.3.17) holds. Then the parameters β_0 , β_1 , β_2 and ϕ are estimated using the estimating equation (5.3.2). The GOF(B) is not sensitive to the violation of the linear-exponential form (5.3.17). In contrast, GOF(A) is much more powerful.

Case 3:

The random variables are generated with mean

$$\mu_i = \alpha_0 + \alpha_1(1 + d_i)^\phi.$$

As in Case 2, we still assume the data Y_{ij} come from a linear-exponential mean form (5.3.17).

In Case 3, we find that GOF(A) has type I error close to the nominal level of 5%, and has smaller type II error than GOF(B) when the linear-exponential model does not hold. Therefore, GOF(A) seems sensitive to the violation of the the linear-exponential model. In contrast, GOF(B) is insensitive. Generally, an inflated value of ϕ reduces the magnitude of GOF(B) significantly. Note that in Case 3b, the type II error of GOF(B) is 99.3% while the type II error of GOF(A) is only 30.5%. This is mainly due to the poor estimate of ϕ because of the moment equation (5.3.3). In Case 4, we will use the moment equation (5.3.3') to investigate the same example and examine the change of the type II error.

In Case 4-6 of Table 5.5, we continue to examine if the the violation of the linear-exponential model will result in inflated GOF(B) or GOF(A). In particular, we use (5.3.3') instead of (5.3.3) to estimate the overdispersion parameter ϕ . If the data are not generated from a linear-exponential model, we may obtain an inaccurate estimate of ϕ even if using (5.3.3'). This inaccurate estimate of ϕ may affect the value of GOF(B), which can generate bigger type II error than that of GOF(A). However, it is actually more difficult to verify this using type II error. In our example when the data are generated not from the linear-exponential model the GOF(B) and GOF(A) values are well beyond the 5% critical value of chi-squared distribution that both GOF(B) and GOF(A) suppose to follow under the linear-exponential model. Therefore, it is not easy to detect the differences by using the type II errors. Besides, the values of GOF(A) and GOF(B) are quite close, although

the values of $GOF(A)$ are generally slightly larger than that of $GOF(B)$. Thus, the examples in Cases 4-6 are carefully selected in order to compare the performance of $GOF(B)$ and $GOF(A)$.

Case 4:

In Cases 1-3, we used (5.3.3) to estimate the overdispersion parameter ϕ . Since (5.3.3) cannot yield good estimates of ϕ when K is small, the inflated type I error rate may be due to the poor estimates of ϕ . To verify this, in Case 4, we used (5.3.3') to estimate the parameter ϕ . We immediately find that the type I errors of $GOF(B)$ given in Case 1 reduced from 10.2% to 5.13%. This implies that the $GOF(B)$ is very sensitive to the estimate of ϕ .

Case 5:

In this case, we generated the data using the mean $\alpha_0 + \alpha_1 d_i^{0.63}$, and the variance $Var = mean(1 + \phi mean)$. Table 5.5 shows that the type II errors at 5% significant level are 41.6% and 40.0% for $GOF(B)$ and $GOF(A)$ respectively. Thus, $GOF(B)$ and $GOF(A)$ have very close type II errors. Comparing the type II errors to those in Case 5, we can see that the major effect to the difference of type II error between $GOF(B)$ and $GOF(A)$ is the mean-variance relation.

Case 6:

This is an example in which the data Y_{ij} are generated with $EY_{ij} = \alpha_0 + \alpha_1 d_i^{0.63}$, and $Var(Y_{ij}) = (1 + h)\mu_i$, where h is a constant. Then we incorrectly assumed the data come from a linear-exponential model with

$$VarY_{ij} = \mu_i(1 + \phi\mu_i).$$

The parameters β_i and ϕ were estimated using the estimating equation (5.3.2) and the moment equation (5.3.3'). Since the assumed mean-variance relation is different from that the data were generated, a poor estimate $\hat{\phi}$ of ϕ likely enlarges $GOF(B)$. The bigger type II error of $GOF(B)$ reported in Table 5.5 is likely due to the incorrectly assumed mean-variance relation. The violation of the linear-exponential model seems contributing little to the larger type II error of $GOF(B)$.

From the limited cases in the above discussion, we find that the large type I or type II error of $GOF(B)$ is primarily due to the poor estimates of ϕ , either di-

rectly (small sample size, inefficient method (5.3.3)) or indirectly (the data generated from non-linear-exponential model, incorrect assumed relation between mean and variance). Based on our limited simulation results, we conclude that GOF(A) is a better goodness-of-fit test than GOF(B) in terms of both type I and II error.

5.3.4 The Asymptotic Distribution of the Proposed Goodness-of-Fit Test

In this section, we show that the test GF^* given in (5.3.15) asymptotically follows a χ_K^2 distribution, as $\min\{n_i\} \rightarrow \infty$.

Note that the observations Y_i and $Y_{i'}$ are independent for $i \neq i'$. Thus we need only show that

$$GF_i^* = \frac{2}{D_i} [Y_i \ln \frac{\bar{Y}_i}{\mu_i} - n_i(\bar{Y}_i - \mu_i)] \sim \chi_1^2$$

asymptotically. Applying a Taylor expansion to $\ln \bar{Y}_i$ about μ_i in

$$GF_i^* = \frac{2}{D_i} [Y_i(\ln \bar{Y}_i - \ln \mu_i) - n_i(\bar{Y}_i - \mu_i)]$$

we obtain

$$\begin{aligned} GF_i^* &= \frac{2}{D_i} \left\{ Y_i \left[\frac{(\bar{Y}_i - \mu_i)}{\mu_i} - \frac{(\bar{Y}_i - \mu_i)^2}{2(\mu_i^{(\theta)})^2} \right] - n_i(\bar{Y}_i - \mu_i) \right\} \\ &= \frac{2}{D_i} \left\{ (\bar{Y}_i - \mu_i) \frac{(Y_i - n_i \mu_i)}{\mu_i} - \frac{Y_i}{2(\mu_i^{(\theta)})^2} (\bar{Y}_i - \mu_i)^2 \right\} \\ &= \frac{2}{D_i} \left\{ \frac{n_i(\bar{Y}_i - \mu_i)^2}{\mu_i} - n_i(\bar{Y}_i - \mu_i)^2 \frac{\bar{Y}_i}{2(\mu_i^{(\theta)})^2} \right\} \\ &= 2 \frac{(\bar{Y}_i - \mu_i)^2 \mu_i}{V_i/n_i} \left\{ \frac{1}{\mu_i} - \frac{\bar{Y}_i}{2(\mu_i^{(\theta)})^2} \right\} \\ &= 2 \frac{(\bar{Y}_i - \mu_i)^2}{V_i/n_i} \left\{ 1 - \mu_i \frac{\bar{Y}_i}{2(\mu_i^{(\theta)})^2} \right\}, \end{aligned}$$

where $\mu_i^{(\theta)}$ is between $\bar{Y}_i$ and μ_i . Since $\bar{Y}_i$ is consistent estimate of μ_i ,

$$\mu_i \frac{\bar{Y}_i}{2(\mu_i^{(\theta)})^2} \xrightarrow{p} \frac{1}{2} \quad (5.3.19)$$

Table 5.5: Type I and Type II Error Rates (%) of Goodness-of-Fit Tests Before and After Applying the Rao-Scott Transformation at a Nominal Significance Level of 5%

Case ^a	Sample Size K=5	Before Transformation		After Transformation	
		Type I ^b Error	Type II Error	Type I ^b Error	Type II Error
1. $EY_{ij} = (\beta_0 + \beta_1 d_i) \exp(-\beta_2 d_i^2)$ $Var(Y_{ij}) = \mu_i(1 + \phi\mu_i)$, $\phi = 0.154$ $\beta_0 = 8.19$, $\beta_1 = 20.0$, $\beta_2 = 0.30$.	$n_i = 200$	10.2	-	5.10	-
2. $\mu_i = EY_{ij} = \alpha_0 + \alpha_1 d_i^\theta$ $V(Y_{ij}) = \mu_i(1 + \phi\mu_i)$, $\phi = 0.154$ 2a $\alpha_0 = 8.19$, $\alpha_1 = 20$, $\theta = 1$ 2b $\alpha_0 = 8.19$, $\alpha_1 = 20$, $\theta = 0.6$	$n_i = 200$	8.4 ^c -	- 97.5	5.0 ^c -	- 40.5
3. $\mu_i = \alpha_0 + \alpha_1(1 + d_i)^\theta$ $Var(Y_{ij}) = \mu_i(1 + \phi\mu_i)$, $\phi = 0.154$. 3a $\alpha_0 = 8.19$, $\alpha_1 = 15$, $\theta = 1$ 3b $\alpha_0 = 8.19$, $\alpha_1 = 15$, $\theta = 2.0$	$n_i = 200$	3.0 ^c -	- 99.3	5.8 ^c -	- 30.5
4. $\mu_i = (\beta_0 + \beta_1 d_i) \exp(\beta_2 d_i^2)$ $V(Y_{ij}) = \mu_i(1 + \phi\mu_i)$, $\phi = 0.154$ $\beta_0 = 8.19$, $\beta_1 = 20.0$, $\beta_2 = 0.30$	$n_i = 100$	- 10.2	-	- 5.13	-
5. $\mu_i = EY_{ij} = \alpha_0 + \alpha_1 d_i^{0.63}$ $Var(Y_{ij}) = \mu_i(1 + \phi\mu_i)$ $\alpha_0 = 8.19$, $\alpha_1 = 50$, $\phi = 0.154$	$n_i = 100$	-	41.6	-	40.0
6. $\mu_i = EY_{ij} = \alpha_0 + \alpha_1 d_i^{0.63}$ $Var(Y_{ij}) = (1 + h)\mu_i$ $\alpha_0 = 8.19$, $\alpha_1 = 50$, $h = 1.26$	$n_i = 100$	-	7.1	-	2.5

^a ϕ estimated by (5.3.3) in cases 1-3 and by (5.3.3') in cases 4-6.^b Type I error not shown since the underlying model is not of the linear-exponential form.^c Note that since this model is a special case of the linear-exponential model, only type I error is considered.

as $n_i \rightarrow \infty$. By the Central Limit Theorem,

$$\frac{(\bar{Y}_i - \mu_i)^2}{V_i/n_i} \sim \chi_1^2 \quad (5.3.20)$$

as $n_i \rightarrow \infty$. Combining the results of (5.3.19) and (5.3.20) and applying Slutsky's theorem, we have $GF_i^* \sim \chi_1^2$ as $n_i \rightarrow \infty$. Finally, since $GF^* = \sum_{i=1}^K GF_i^*$, with GF_i^* is independent of $GF_{i'}^*$ for $i \neq i'$, we have $GF^* \sim \chi_K^2$ as $\min\{n_i\} \rightarrow \infty$.

5.4 Summary and Conclusions

In this chapter, we have explored the use of Rao-Scott transformation in the analysis of data from the Ames *Salmonella*/microsome assay. This simple data transformation, introduced by Rao & Scott (1992, 1994), is based on the concept of a design effect widely used in survey sampling, and can be used to eliminate overdispersion induced by cluster sampling. Specifically, application of the Rao-Scott transformation to Ames assay data eliminates extra-Poisson variation, permitting the use of standard methods of statistical analysis based on the Poisson distribution.

Three separate analysis issues were considered: (i) testing for trend in mutation rates, (ii) fitting the linear-exponential model to Ames assay data, and (iii) evaluating goodness-of-fit of the linear-exponential model. In each case, methods of analysis based on both the original and transformed data were compared. Tests for trend using the Rao-Scott transformation considered here are similar in spirit to those previously used by Fung et al. (1994) for overdispersed binomial data arising in studies of developmental toxicity. Based on the limited simulation studies conducted in this chapter, the Rao-Scott transformation appears to provide trend tests which are both valid, at least for moderately large sample sizes, and sensitive. The use of the Rao-Scott transformation is similar to previous applications of this technique in modeling dose-response relationships in developmental toxicity experiments by Krewski & Zhu (1994). Here, similar estimates of the regression parameters in the linear-exponential model for the Ames assay were obtained based on either the original or transformed data. Leroux & Krewski (1993) proposed a goodness-of-fit test based on the deviance function. Application of the Rao-Scott transformation

prior to assessing goodness-of-fit appears to be advantageous in terms of both type I and type II error.

The advantages of the Rao-Scott transformation in the present context are several fold. First, the transformed data are not subject to extra-Poisson variation, eliminating the need to characterize overdispersion during the analysis of the transformed data. Second, analysis of the transformed data may be done using simpler methods designed for use with Poisson data. And third, analyses on the transformed scale may lead to enhanced statistical properties, as appears to be the case in testing for goodness-of-fit.

Chapter 6

Summary and Further Research

This thesis considers statistical modeling and analysis of the Ames assay. First, we consider the modeling of the Ames assay based on fundamental biological principle governing the Ames assay. The proposed model can reduce to both the Margolin et al. (1981) model and Krewski et al. (1993) model when comparable assumptions are made. We recommend the linear-exponential dose-response model given by Krewski et al. (1993), and use this model in the subsequent developments.

Secondly, the problem of testing for trend with historical controls is studied. A method is proposed allowing for extra-Poisson variation both between and within laboratories. Limited simulation studies demonstrated that the use of historical control data enhances the power of the test for trend.

Third, we developed optimal designs for the Ames assay. Three optimal criteria are studied under the linear-exponential model. Sub-optimal designs are proposed after analyzing the data sets in the International Programme on Chemical Safety. The use of Bayesian and minimax optimality criteria is also illustrated.

Finally, the Rao-Scott transformation is used to study the test for trend, parameter estimation, and goodness-of-fit. The goodness-of-fit test is proposed to examine whether or not a dose response function is of a certain form. With the Rao-Scott transformation, no assumptions about the nature of the mean-variance relationship of the data are necessary. Limited simulation studies showed that the goodness-of-fit has the better operating characteristics than that based on the original data.

With respect to future research, a more intensive simulation study to examine

the performance of the test for trend in chapter 3 would be of interest. Further simulation work is also needed to compare the different methods proposed in chapter 5 for testing for trend, parameter estimation, and goodness-of-fit.

Another research prospect is related to the International Programme on Chemical Safety (IPCS) study. Investigating the inter-laboratory and intra-laboratory source of variation for the mutagenic potency in Ames assay is the primary objective of the IPCS study. Krewski et al. (1992) have studied the sources of variation by applying classical variance component methodology. However, in their method, the unknown regression parameters are estimated first, and then the analysis of variance is used to decompose the variance of the mutagenic potency estimate into within and between laboratory components. In contrast to their methods, two different methods should be considered to investigate the same issue.

First, it is natural to consider a Bayesian analysis of the IPCS data, given the hierarchical structure of the problem. The hierarchical Bayesian approach allows us to simultaneously estimate the mutagenic potency parameters and the variability of both between and within laboratories. The advantage of using the hierarchical model is the simplicity of organizing the concepts. In addition, it enables unified estimation of the parameters based on combined information from all twenty laboratories participated in the IPCS study and prior knowledge. The sources of variation can be estimated by obtaining the marginal posterior means for both intra- and inter-laboratory variabilities. Gibbs sampling algorithm will play a major role in this computation. The results obtained from the hierarchical model can also be compared to those obtained from the variance component methodology. Secondly mixed-effects or random-effects modeling can be applied to the same problem. In this context, the regression parameters can be considered as random variables, rather than unknown constants. The mixed-effects model can be solved by applying the generalized estimating equation methods. Although the random-effect model has been considered by Burnett et al. (1992), finding efficient statistical computing method to solve the multi-level random-effect model and compare the results with those of variance component and hierarchical Bayesian requires further study.

Bibliography

- [1] Abdelbasit, K.M. and Plackett, R.L. (1983). Experimental design for binary data. *Journal of American Statistical Association* 78, 90-98.
- [2] Ames, B.N. (1971). The detection of chemical mutagens with enteric bacteria. In: *Chemical mutagens: principles and methods for their detection*. Vol. 1. (A. Hollaender, ed). Plenum Press, New York, pp. 267-282.
- [3] Ames, B.N., Durston, W.E., Yamasaki, E. and Lee, F.D. (1973). Carcinogens are mutagens-a simple test system combining liver homogenates for activation and bacteria for detection. *Proceedings of the National Academy of Sciences* 70, 2281-2285.
- [4] Ames, B.N., McCann, J. and Yamasaki, E. (1975). Methods for detecting carcinogens and mutagens with the *Salmonella*/mammalian-microsome mutagenicity test. *Mutation Research* 31, 347-363.
- [5] Armitage, P. (1955). Tests for linear trends in proportions and frequencies. *Biometrics* 11, 375-386.
- [6] Bernstein, L., Kaldor, J., McCann, J., and Pike, M.C. (1982). An empirical approach to the statistical analysis of mutagenesis data from the *Salmonella* test. *Mutation Research* 97, 267-281.
- [7] Bickis, M. and Krewski, D. (1985). Statistical design and analysis of the long-term carcinogenicity bioassay. In: D.B. Clayson, D. Krewski and I. Munro (eds), *Toxicological Risk Assessment. Vol. 1: Biological and Statistical Criteria*. CRC Press, Boca Raton, Fl., pp. 125-147.

- [8] Billingsley, P. (1986). *Probability and Measure* (2nd Edition). John Wiley & Sons. New York.
- [9] Breslow, N.E. (1984). Extra-Poisson variation in log-linear models. *Applied Statistics* 33, 38-44.
- [10] Breslow, N. (1990). Tests of hypotheses in overdispersed Poisson regression and other quasi likelihood models. *Journal of the American Statistical Association* 85, 565-571.
- [11] Broekhoven, L.H. and Nestmann, E.R. (1991). Statistical analysis of the *Salmonella* mutagenicity assay. In *Statistics in Toxicology* (Krewski, D. and Franklin, C. eds.), Gordon and Breach, New York, pp. 205-264.
- [12] Burnett, R.T., Shedden, J., and Krewski, D. (1992). Nonlinear regression models for correlated count data. *Environmentrics* 3, 211-222.
- [13] Burrige, J. and Sebastiani, P. (1994). D-optimal designs for generalised linear models with variance proportional to the square of the mean. *Biometrika* 81, 295-304.
- [14] Chaloner, K. (1984). Optimal Bayesian experimental design for linear models. *Annals of Statistics* 12, 283-300.
- [15] Chaloner, K. and Larntz, K. (1989). Optimal Bayesian design applied to logistic regression experiments. *Journal of Statistical Planning and Inference* 21, 191-208.
- [16] Chernoff, H. (1953). Locally optimal designs for estimating parameters. *Annals of Mathematical Statistics* 24, 586-602.
- [17] Cochran, W.G. (1954). Some methods for strengthening the common chi-squared tests. *Biometrics* 10, 417-451.
- [18] Consul, P.C. (1989). *Generalized Poisson Distributions*, Marcel Dekker. New York.

- [19] Cox, D.R. (1983). Some remarks on over-dispersion. *Biometrika* 70, 269-274.
- [20] Crowder, M. (1987). On linear and quadratic estimating functions. *Biometrika* 74, 591-597.
- [21] de la GARZA, A. (1954). Spacing of information in polynomial regression. *Annals of Mathematical Statistics* 25, 123-130.
- [22] de la GARZA, A. (1956). Quadratic extrapolation and a related test of hypothesis. *Journal of American Statistical Association* 27, 644-649.
- [23] Dean, C., and Lawless, J.F. (1989a). Testing for overdispersion in Poisson regression models. *Journal of American Statistical Association* 84, 467-472.
- [24] Dean, C., and Lawless, J.F. (1989b). Comments on "An extension of quasilielihood estimation", by Godambe and Thompson. *Journal of Statistical Planning and Inference* 22, 155-158.
- [25] Dobson, A.J. (1990). *An Introduction to Generalized Linear Models*, Chapman and Hall, London.
- [26] Elfving, G. (1952). Optimum allocation in linear regression theory. *Annals of Mathematical Statistics* 23, 255-262.
- [27] Elfving, G. (1955). Geometric allocation theory. In: *Skand. Aktuarietidskr.* 37, 170-190.
- [28] Elfving, G. (1959). Design of linear experiments. *Cramer Festschrift Volume*. Wiley, New York, pp. 58-74.
- [29] Fedorov, V.V. (1972). *Theory of Optimal Experiments*. (Translated from the Russian.) Academic Press, New York.
- [30] Fedorov, V.V. and Malyutov, M.B. (1972). Optimal design in regression experiments. *Math. Operationsforsch. Statist.* 14, 237-324.

- [31] Ford, I., Torsney, B., and Wu, C.F.J. (1992) The use of a canonical form in the construction of locally optimal designs for nonlinear problems. *Journal of the Royal Statistical Society. Series B* 54, 569-583.
- [32] Foutz, R.V. (1977). On the unique consistent solution to the likelihood equations. *Journal of the American Statistical Association* 72, 147-148.
- [33] Fung, K.Y., Krewski, D., Rao, J.N.K. and Scott, A.J. (1994). Test for trend in developmental toxicity experiments with correlated binary data. *Risk Analysis* 14, 621-630.
- [34] Guest, P.G. (1958). The spacing of observations in polynomial regression. *Annals of Mathematical Statistics* 29, 294-299.
- [35] Hoel, P.G. and Jennrich, R.I. (1979). Optimal designs for dose response experiments in cancer research. *Biometrika* 66, 307-316.
- [36] Hoel, D.G. and Yanagawa, T. (1986). Incorporating historical controls in testing for a trend in proportions. *Journal of the American Statistical Association* 81, 1095-1099.
- [37] Kalish, L.A. and Rosenberger, J.L. (1978). Optimal designs for the estimation of the logistic function. Technical Reports and Preprint No. 33, The Pennsylvania State University.
- [38] Kiefer, J. (1958). On the nonrandomized optimality and randomized non-optimality of symmetrical designs. *Annals of Mathematical Statistics* 29, 675-699.
- [39] Kiefer, J. (1959). Optimum experimental designs. *Journal of the Royal Statistical Society Series* 21, 272-319.
- [40] Kiefer, J. and Wolfowitz, J. (1960). The equivalence of two extremum problems. *Canadian Journal of Mathematics* 12, 363-366.
- [41] Kiefer, J. (1974). General equivalence theory for optimum designs. (approximate theory). *Annals of Statistics* 2, 849-879.

- [42] Kikuchi, Y. & Yanagawa, T. (1988). Incorporating historical information in testing for trend in Poisson means. *Annals of the Institute of Statistical Mathematics* 40, 367-379.
- [43] Kilbey, B.J., Legator, M., Nichols, W. and Ramel, C. (1984). *Handbook of Mutagenicity Test Procedures*. (2nd edition) Elsevier, Amsterdam.
- [44] Knuiman, M.W., Laird, N.M. and Louis, T.A. (1987). Inter-laboratory variability in Ames assay results. *Mutation Research* 180, 171-182.
- [45] Krewski, D. and Franklin, C.A. (eds.) (1991). *Statistics in Toxicology*. Gordon and Breach, New York.
- [46] Krewski, D., Bickis, M., Kovar, J. and Arnold, D.L. (1986). Optimal experimental designs for low dose extrapolation I. The case for zero background. *Utilitas Mathematica* 29, 245-262.
- [47] Krewski, D., Colin, D. and Dewanji, A. (1990). Statistical considerations in the designs of toxicological experiments with chemical mixtures. In: *Progress in Predictive Toxicology* (Clayson, D.B., Munro, I.C., Shubik, P. and Swenberg, J.A., editors) 1990 Elsevier Science Publishers B.V. (Biomedical Division).
- [48] Krewski, D. and Goddard, M. (1990). Principles of bioassay design. *Drug Information Journal* 24, 384-394.
- [49] Krewski, D. and Kovar, J. (1982). Low dose extrapolation under single parameter dose response models. *Communications in Statistics B* 11, 27-46.
- [50] Krewski, D., Kovar, J., and Bickis, M., (1984). Optimal experimental designs for low dose extrapolation II. The case of non-zero background. In: *Topics in Applied Statistics*, Concordia University, Montreal. pp. 167-191.

- [51] Krewski, D., Leroux, B.G., Creason, J. and Claxton, L. (1992). Sources of variation in the mutagenic potency of complex chemical mixtures based on the *Salmonella*/microsome assay. *Mutation Research* 276, p.33-59.
- [52] Krewski, D., Leroux, B.G., Bleuer, S.R. and Broekhoven, L.H. (1993). Modeling the Ames *Salmonella*/microsome assay. *Biometrics* 49, 499-510.
- [53] Krewski, D. and Rao, J.N.K. (1981). Inference from stratified samples: properties of the linearization, jackknife and balanced repeated replication methods. *Annals of Statistics* 9, 1010-1019.
- [54] Krewski, D., Smythe, R.T., Fung, K. and Burnett, R.T. (1991). Conditional and unconditional tests with historical controls. *Canadian Journal of Statistics* 19, 407-423.
- [55] Krewski, D. and Van Ryzin, J. (1981). Dose response models for quantal response toxicity data. In *Statistics and Related Topics* (M. Csorgo, D. Dawson, J.N.K. Rao and E. Saleh, eds), North-Holland, Amsterdam, pp. 201-231.
- [56] Krewski, D. and Zhu, Y. (1994). Applications of multinomial dose-response models in developmental toxicity risk assessment. *Risk Analysis* 14, 595-609.
- [57] Krewski, D. and Zhu, Y. (1995). A simple data transformation for estimating benchmark doses in developmental toxicity experiments. *Risk Analysis*. In press.
- [58] Lawless, J.F. (1987). Regression methods for Poisson process data. *Journal of the American Statistical Association* 82, 808-815.
- [59] Lawless, J.D. (1987). Negative binomial and mixed Poisson regression. *The Canadian Journal of Statistics* 15, 209-225.

- [60] Leroux, B.G., Fung, K.Y., Krewski, D. and Prentice, R.L. (1994). The use of historical control data in testing for trend in counts. *Statistica Sinica* 4, 581-601.
- [61] Leroux, B.G. and Krewski, D. (1993). Components of variation in mutagenic potency values based on the Ames *Salmonella* test. *Canadian Journal of Statistics* 21, 448-459.
- [62] Leroux, B.G. and Krewski, D. (1993). AMESFIT: A microcomputer program for fitting linear-exponential dose-response models in the Ames *Salmonella* assay. *Environmental and Molecular Mutagenesis* 22, 78-84.
- [63] Lewtas, J., Claxton, L.D., Rosenkranz, H.S., Schuetzle, D., Shelby, M., Matsushita, M., Wurgler, F.E., Zimmerman, F.K., Lofroth, G., May, W.E., Krewski, D., Matsushima, T. & Becking, G.C. (1992). Design and implementation of a collaborative study on the mutagenicity of complex mixtures in *Salmonella typhimurium*. *Mutation Research* 276, 3-6.
- [64] Liang, K-Y. and Zeger, S.L. (1986). Longitudinal data analysis using generalized linear models. *Biometrika* 73, 13-22.
- [65] Margolin, B.H., Kaplan, N. and Zeiger, E. (1981). Statistical analysis of the Ames *Salmonella*/microsome test. *Proceedings of the National Academy of Sciences* 78, 3779-3783.
- [66] Margolin, B.H., Kim, B.S. and Risko, K.J. (1989). The Ames *Salmonella*/Microsome Mutagenicity Assay: Issues of Inference and Validation. *Journal of the American Statistical Association* 84, 651-661.
- [67] Maron, D.M. and Ames, B.N. (1983). Revised methods for the *Salmonella* mutagenicity test. *Mutation Research* 113, 173-215.
- [68] McGregor, D., Prentice, R.D., McConville, M., Lee Y.J. and Caspary, W.J. (1984). Reduced mutant yield at high doses in the

- Salmonella*/activation assay: the cause is not always toxicity. *Environmental Mutagenesis* 6, 545-557.
- [69] McCullagh, P. (1983). Quasi-likelihood functions. *Annals of Statistics* 11, 59-67.
- [70] McCullagh, P. and Nelder, J.A. (1989). *Generalized Linear Models* (2nd edition). Chapman and Hall, London.
- [71] Moore, D.F. (1985). Method of moment estimation for overdispersed counts and proportions. Ph.D thesis, University of Washington, Seattle.
- [72] Moore, D.F. (1986). Asymptotic properties of moment estimators for overdispersed counts and proportions. *Biometrika* 73, 583-588.
- [73] Moore, D.F. (1987). Modelling the extraneous variation in the presence of extra-binomial variation. *Applied Statistics* 36, 8-14.
- [74] Moore, D.F. and Tsiatis, A. (1991). Robust estimation of the variance in moment methods for extra-binomial and extra-Poisson variation. *Biometrics* 47, 383-401.
- [75] Most, D.E. (1993). An empirical comparison of statistical models for *Salmonella* mutagenicity test data. M.Sc. thesis, Department of Biostatistics, University of North Carolina, Chapel Hill.
- [76] Myers, L.E., Sexton, N.H., Southerland, L.I. & Wolff, T.J. (1981). Regression analysis of Ames test data. *Environmental Mutagenesis* 3, 575-586.
- [77] NAG (Numerical Algorithms Group) (1993). NAG. Oxford.
- [78] Nelder, J.A. and Mead, R. (1965). A simplex method for function minimization. *Computer Journal* 7, 308-313.
- [79] O'Neill, R. (1971). Function minimization using a simplex procedure. Algorithm AS 47. *Applied Statistics* 20, 338-345.

- [80] Portier, C.J. (1991). Design of two-year carcinogenicity experiments: dose allocation, animal allocation and sacrifice times. In: *Statistical in Toxicology*. (Krewski, D. and Franklin, C. eds.). Gordon and Breech, New York. pp. 457-469.
- [81] Prentice, R.L., Smythe, R.T., Krewski, D. and Mason, M. (1992). On the use of historical control data to estimate dose response trends in quantal bioassay. *Biometrics* 48, 459-478.
- [82] Pukelsheim, F. (1993). *Optimal Design of Experiments*. John Wiley & Sons, New York.
- [83] Rao, J.N.K. and Scott, A.J. (1992). A simple method for the analysis of clustered binary data. *Biometrics* 48, 577-585.
- [84] Scott, A.J. and Rao, J.N.K. (1994). A simple method for the analysis of count data. (Submitted).
- [85] Silvey, S.D. (1980). *Optimal Design*. Chapman and Hall, New York.
- [86] Sitter, R.R. (1992). Robust designs for binary data *Biometrics* 48, 1145-1155.
- [87] Sitter, R.R. and Torsney, B. (1994). Optimal designs for binary response experiment with two design variables. *Statistica Sinica*, In press.
- [88] Sitter, R.R. and Wu, C.F.J. (1993a). Optimal designs in binary response experiments: Fieller, D, and A criteria. *Scandinavian Journal of Statistics* 20, 329-342.
- [89] Sitter, R.R. and Wu, C.F.J. (1993b). On the accuracy of Fieller intervals for binary response data. *Journal of the American Statistical Association* 88, 1021-1025.
- [90] Simpson, D.G. and Margolin, B.H. (1986). Recursive nonparametric testing for dose-response relationships subject to downturns at high doses. *Biometrika* 73, 589-596.

- [91] Simpson, D.G. and Margolin, B.H. (1990). Nonparametric testing for dose-response curves subject to downturns: asymptotic power consideration. *Annals of Statistics* 18, 373-390.
- [92] Snee, R.D. and Irr, J.D. (1984). A procedure for the statistical evaluation of Ames *Salmonella* assay results. Comparison of results among 4 laboratories. *Mutation Research* 128, 115-125.
- [93] Stead, A.G., Hasselblad, B., Creason, J.P. and Claxton, L. (1981). Modeling the Ames test. *Mutation Research* 85, 13-17.
- [94] Smythe, R.T., Krewski, D., and Murdoch, D. (1986). The use of historical controls in modelling dose-response relationships in carcinogenesis. *Statistics and Probability Letters* 4, 87-93.
- [95] Tarone, R.E. (1982). The use of historical control information in testing for a trend in Poisson means. *Biometrics* 38, 457-462.
- [96] Tsutakawa, R.K. (1972). Design of an experiment for bioassay. *Journal of American Statistical Association* 67, 584-590.
- [97] Waleh, N.S., Rapport, S.J. and Mortelmans, K. (1982). Development of a toxicity test to be coupled to the Ames *Salmonella* assay and the method of construction of the required strains. *Mutation Research* 97, 247-256.
- [98] Wedderburn, R.W.M. (1974). Quasi-likelihood functions, generalized linear models and the Gauss-Newton method. *Biometrika* 61, 439-447.
- [99] Whittle, P. (1973). Some general points in the theory of optimal experimental design. *Journal of the Royal Statistical Society, Series B* 35, 123-130.
- [100] Williams, D.A. (1982). Extra-binomial variation in logistic linear models. *Applied Statistics* 31, 144-148.

- [101] Wu, C.F.J. (1988). Optimal design for percentile estimation of a quantal response curve. In: *Optimal Design and Analysis of Experiments* (Dodge, Y., Fedorov, V., and Wynn, H.P. eds.) Elsevier Science Publishers B. V., Amsterdam, pp. 213-223.

END

1 4-1 1-9 5

FIN