1

MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-
STANDARD REFERENCE MATERIAL 1010a
(ANSI and ISO TEST CHART No. 2)

National Library
of Canada

Bibliothèque nationale
du Canada

Canadian Theses Service Service des thèses canadiennes

Ottawa, Canada
K1A 0N4

NOTICE

The quality of this microform is heavily dependent upon the quality of the original thesis submitted for microfilming. Every effort has been made to ensure the highest quality of reproduction possible.

If pages are missing, contact the university which granted the degree.

Some pages may have indistinct print especially if the original pages were typed with a poor typewriter ribbon or if the university sent us an inferior photocopy.

Reproduction in full or in part of this microform is governed by the Canadian Copyright Act, R.S.C. 1970, c. C-30, and subsequent amendments.

AVIS

La qualité de cette microforme dépend grandement de la qualité de la thèse soumise au microfilmage. Nous avons tout fait pour assurer une qualité supérieure de reproduction.

S'il manque des pages, veuillez communiquer avec l'université qui a conféré le grade.

La qualité d'impression de certaines pages peut laisser à désirer, surtout si les pages originales ont été dactylographiées à l'aide d'un ruban usé ou si l'université nous a fait parvenir une photocopie de qualité inférieure.

La reproduction, même partielle, de cette microforme est soumise à la Loi canadienne sur le droit d'auteur, SRC 1970, c. C-30, et ses amendements subséquents.

**APPROACHES TO THE SYNTHESIS
AND STEREOCHEMICAL ANALYSIS
OF AZA CROWN ETHERS**

by

Denis Joseph Landry

A Thesis submitted to the Faculty of
Graduate Studies in partial fulfilment
of the requirements for the degree of
Master Science

Department of Chemistry
Carleton University
Ottawa, Ontario

© September, 1989.

N_2O_4 MODELS

Many macrocyclic polyethers have attracted attention because of their ability to form stable complexes. In order to fulfill the stereochemical requirement of some metals, many crown systems have been prepared with ring sizes from 14 to 32.²²

However, two of the most widely investigated systems are dibenzo 18-crown-6 and dicyclohexyl 18-crown-6. These products have different properties which result in different binding abilities.

Figure 9 : Stability constants ($\text{Log } k_1$) for dibenzo and dicyclohexyl 18-C-6²³

	Na^+	K^+	Cs^+	
	4.36	5.00	3.55
c-s-c	4.08	6.01	4.61	
c-a-c	3.68	5.38	3.49	
t-s-t	2.99	4.14	3.00
t-a-t	2.52	3.26	2.27	

The difference in selectivity shown in Figure 9 reflects both the electrostatic and steric effects of the annelated ring. The strength of the C-O dipoles are affected by the nature of the substituent. Rigidity and flexibility of a ligand skeleton can dictate the cavity size of the ligand. Ligands of the flexible type show a plateau selectivity because they can flex to accommodate a host cation. This is not the case for smaller more rigid systems showing a peak selectivity for the cation that best matches its cavity size.²²

The undersigned recommend to the
Faculty of Graduate Studies acceptance of the thesis
**"Approaches to the Synthesis and
Stereochemical Analysis of Aza Crown Ehters"**
Submitted by Denis J. Landry
in partial fulfilment of the requirements for the
degree of Master of Science.

Dr. G.W. Buchanan
Supervisor

Chairman
Department of Chemistry

ABSTRACT

The synthesis of some configurational isomers of dicyclohexyl tetraaza-14-crown-4 ethers has been attempted. In all cases, the major isolated product is the 7 membered ring analogue, resulting from a retro Michael type reaction in solution.

A second series of target molecules was the corresponding dicyclohexyl diazatetraoxa-18-crown-6. Although the final hydrogenation leading to these systems was not successful, a number of interesting stereochemical features of tosylated and benzylated precursors are found. Specifically, large ^{13}C nmr chemical shift differences were noted between 9 and 18 membered analogues. From ^1H - ^1H nuclear Overhauser effects, conformational differences in the macrocyclic protons of these materials are deemed responsible for the chemical shift trends.

^{13}C spin lattice relaxation times have been employed to study the relative mobilities of the free and complexed 9 vs 18 membered ring materials.

ACKNOWLEDGEMENTS

I would like to thank the chemistry department for the use of the computer facilities in obtaining this document. The financial support and guidance of Dr. G. Buchanan throughout my studies has been appreciated.

In attempting to thank all the other people who have played in a part in this story, I can only faithfully start from a period that begins long before my arrival at Carleton. I was born into a family of 6 children. Both my parents believed that life was essentially two things; family and work. It was my oldest brother that was to sow the seed. As a child my admiration for Peter was to have a profound effect on my life. In an attempt to be just like my big brother, I quickly adopted his interest in both athletics and science.

Throughout the difficult periods of my life, I have often needed the moral fiber basic to my family (thanks mom and dad). I emerged from high school relatively intact with only a few cuts and bruises. David Wildsmith had given me my love of music and the Martial arts, Ronny Dunn had given me a taste for something wild and I..... gave myself another chance.

My stay at Carleton has been lined with many periods of joy and pain (sunshine and rain). Throughout these times, there have been some special people who were there to see the good and the bad in me, yet their lasting friendship did not falter and I was able to endure. Some of these people are Bernie (thanks for those 2" thick roast beef sandwiches)Gareau, Ken(I really can hear someone talking to me)Stripelis, Paul (boy did we have some rough times)Krushinske, Paul(let's get the F___ out of here)Parisien, Eddy boy (look at those calves)Dowdall, Lee (we are not here to talk, well actually we are here to talk) Collier, and Gerry (I don't really want to be Prime Minister)Crotty - Thanks guys.

TABLE OF CONTENTS

TITLE	PAGE
ABSTRACT	i
ACKNOWLEDGEMENTS	ii
TABLE OF CONTENTS	iii
LIST OF TABLES	v
LIST OF FIGURES	vi
Historical	I
Introduction	5
Theory and Background	
A. Macrocyclic effect	14
B. Complexation	16
C. ^{13}C N.M.R.	
i. General	19
ii. Magnetization vector - M_z	20
iii. Relaxation times - T_1	22
iv. The nuclear Overhauser effect (n.O.e.)	23
 Results and Discussion	
I. Attempts of the synthesis of trans dicyclohexyl tetraaza 14-crown-4	25
Attempts at the stabilization of 1	28
Mechanistic description of retro Michael addition	29
Attempted template synthesis of trans dicyclohexyl tetraaza 14-crown-4	31
Attempts at the synthesis of cis dicyclohexyl tetraaza 14-crown -4	36
II. Attempted synthesis of dicyclohexyl diaminotetraoxa 18-crown-6	38

Results and Discussion cont'd**III. NMR Studies****Conformational analysis**

a) Low temperature 40

b) Characterization of conformational differences between 9 and 18 membered ring tosylates 41

c) Complexation properties of the seven membered trans fused ring 56

Mobility studies

a) Complexation properties of the tosylated and benzylated dibenzo diaminotetraoxa 18-C-6 58

Conclusion 60

Experimental 61

References 75

Appendices 82

LIST OF TABLES

TABLE	TITLE	PAGE
1	^{13}C nmr chemical shifts for 8 in CDCl_3	44
2	^{13}C nmr chemical shifts for 2 in CDCl_3	45
3	^{13}C nmr chemical shifts for 14 in CDCl_3	49
4	Theoretical ABCD spin system parameters for simulated spectrum in Figure 56	53
5	Elemental analysis for $\text{C}_{18}\text{H}_{36}\text{N}_4\text{NiCl}_2$	55
6	^{13}C nmr chemical shifts for 3 during the titration with $\text{NiCl}_2\cdot\text{H}_2\text{O}$	56

LIST OF FIGURES

FIGURE	TITLE	PAGE
1	Complexes of some natural products	1
2	Structure of some early models	2
3	Cyclam	3
4	Mixed donor crowns	3
5	N ₄ and N ₂ O ₄ models	5
6	Conformations of cis-anti-cis dicyclohexyl tetraoxa 14-crown-4	6
7	Conformation of cis-syn-cis dicyclohexyl tetraoxa 14-crown-4	6
8	Target molecules	7
9	Stability constants for dibenzo and dicyclohexyl 18-C-6	8
10	Selectivity characteristics of some crown ethers	9
11	Stability constants for some dibenzo crown ethers	9
12	Decomplexation energy for dibenzo and dicyclohexyl 18-C-6	10
13	Stability constants for dibenzo 18-crown-6 in various solvents	11
14	Conformation of the major isomers of cis-syn-cis dicyclohexyl 18-C-6	11
15	Conformation of the minor isomer of cis-anti-cis dicyclohexyl 18-C-6 in solution	12
16	Target molecules for the N ₂ O ₄ system	12
17	Benzylated and tosylated 18 membered precursors	13
18	Stability constants of some K ⁺ and Ag ⁺ complexes	14
19	Cu ²⁺ complex of diaza tetraoxa 18-crown-6	15
20	Stability constants of some Ni ²⁺ complexes	15

LIST OF FIGURES CONT'D

FIGURE	TITLE	PAGE
21	Stepwise substitution of oxygen donor atoms in 18-crown-6 ethers	16
22	Relative stability of various membered ethyleneoxy chelates	17
23	Cartesian coordinate description of a perturbed magnetization vector	19
24	Cartesian coordinate description of the relaxation along the Z axis	21
25	Cartesian coordinate description of the relaxation in the XY plane	21
26	Cartesian coordinate description of the inversion recovery method for determination of T_1	22
27	Proposed synthetic scheme for the synthesis of dicyclohexyl tetraoxa 14-crown-4	24
28	Intramolecular reaction product from Figure 27	25
29	Proposed synthetic scheme for the synthesis of dicyclohexyl tetraaza 14-crown-4	26
30	Possible cleavage pattern of 3	27
31	Structure of the intramolecular reaction product 2	28
32	Proposed scheme for the synthesis of 1,2 bis[3methyl ester-N-tosylamine] cyclohexane	28
33	Proposed scheme for the synthesis of 1,2 bis [3 methyl ester-N-tosylamine]	29
34	Proposed mechanism for retro Michael addition product	30
35	Methyl acrylate	30
36	Mechanism of a template reaction	31

LIST OF FIGURES CONT'D

FIGURE	TITLE	PAGE
37	Proposed scheme for the synthesis of dicyclohexyl tetraaza 14-crown-4 using metal template	32
38	Conformation of a cis coordinated Ni^{2+} Complex of 1	33
39	IR of 1 in methanol- titration with nickel	34
40	UV of $\text{NiCl}_2 \cdot \text{H}_2\text{O}$ in methanol	35
41	Synthetic scheme for the synthesis of 1 and 6	36
42	Proposed synthetic reaction scheme for the synthesis of cis dicyclohexyl tetraaza 14-crown-4 ether	37
43	Synthetic scheme for the synthesis of dibenzo-diaazatetraoxa 18-crown-6 ether	38
44	Synthetic scheme for the synthesis of dibenzo-diazatetraoxa 18-crown-6 ether	39
45	N tosyl-substituted crown ether aza analogues	40
46	The ^{13}C nmr spectrum of 8 in CDCl_3	42
47	The ^{13}C nmr spectrum of 9 in CDCl_3	43
48	Carbon numbering scheme for 8 , 9 and 14 analogues	44
49	Proton numbering scheme for 8 , 9 and 14 analogues	45
50	Homonuclear 2-D shift correlated (H, H NOESY) spectrum in CDCl_3 with the 1-D spectrum	46
51	Homonuclear 2-D shift correlated (H, H NOESY)spectrum in CDCl_3 with the 1-D spectrum	47
52	MM calculations of dihedral angles for 8 and 9 using TRIBL	48

LIST OF FIGURES CONT'D

FIGURE	TITLE	PAGE
53	MM calculations of interatomic distances for the protons of 8 and 9 as determined by the MM program TRIBL	50
54	The ^{13}C nmr spectrum of 14 in CDCl_3	51
55	Homonuclear 2-D shift correlated (^1H , ^{13}C HETCOR) spectrum of 14 in CDCl_3 with the 1-D spectrum above.	52
56	Homonuclear 2-D shift correlated (^1H , ^1H NOESY) spectrum of 14 in CDCl_3 with the 1-D spectrum .	53
57	Experimental and theoretical spectrum of the $\text{CH}_2\text{-O}$ and $\text{CH}_2\text{-N}$ fragments in 9	55
58	Structure of the cyclic amide and its reduced product.	56
59	Plot of the observed ^{13}C nmr chemical shifts upon titration of 3 with $\text{NiCl}_2\cdot\text{H}_2\text{O}$ in methanol	57
60	Correlation time vs relaxation time graph	58

ACKNOWLEDGEMENTS

I would like to thank the chemistry department for the use of the computer facilities in obtaining this document. The financial support and guidance of Dr. Buchanan throughout my studies has been appreciated.

HISTORICAL

I. Coordination compounds, or more commonly called complexes, play a very important role in our lives. Perhaps the most important of these naturally occurring complexes are the Mg complex of chlorophyll and the Fe complex of hemoglobin.

Figure 1: Complexes of some natural products

Each system is affected by and receives much of its properties from the presence of the metal. Because of the structural complexity of these natural products, model compounds were prepared to facilitate the characterization of the complex in the natural system. From the beginning, a systematic approach was taken to elucidate the role of the metal and ligand. A system was considered an appropriate model if it emulated or contained analogous features of the natural product. Subsequent alteration of the model and investigation of the resulting changes imparted (physical, kinetic, thermodynamic) were studied. Often, construction of a ligand to approximate the environment about a certain metal was used as an alternate approach.

Complexes of porphyrin, corrins and phthalocyanines have been investigated as model systems because of their relation to important natural products shown in Figure 1. The major similarity is the structure of the donor sites held in a very rigid planar conformation.

Figure 2: Structure of some early N_4 models

The use of phthalocyanine complexes has now found applications in paints, fabrics, plastics and dyestuffs. Recently, there has been interest in developing their use in electrochemical devices such as organic catalysts, electrocatalysts, photocatalysts and fuel cells.²

It was recognized that the successful application of a model would be the result of a basic requirement of four donor nitrogen atoms more or less rigidly confined to a plane leaving two axial sites available for interaction with the substrate. From this point, a systematic approach could be taken to determine the effects of substituents on the

properties of the ligand. One of the early models used for this purpose was the cyclic compound "CYCLAM" originally synthesized by Van Alphen.³

Figure 3: Cyclam

Since its introduction, this system has led to an understanding of the phenomenon of enhanced stability called the macrocyclic effect as well as hole size-best fit concepts.⁴ A large amount of work has been done on the aza crown systems and several reviews and books have been written.^{5,22}

II.

In 1964, the mode of action of the antibiotic, valinomycin was found to be related to its ability to induce the transport of K^+ across the lipophilic membrane of mitochondria.⁶ At this time, no appropriate model system was available to study the process. However, in 1967, a number of cyclic polyethers or "crown compounds" were synthesized by Pederson.⁷ These compounds were able to strongly coordinate to alkali and alkaline earth metals. The possibility of applying these crown systems to the full range of guest metals which had not been previously investigated resulted in a broadening of the field of Host-Guest Chemistry.

Shortly after the initial report by Pederson, different potential donor sites were introduced into the cyclic structure to provide new chemical environments as well as the ability to functionalize the ring.⁸

Figure 4: Mixed donor crowns

Since their discovery, the macrocyclic crown systems have evolved far beyond their use as biomimetic models. They are currently used in many different applications including preconcentrating metals in water⁹, solvent extraction in nuclear technology¹⁰, ion transport¹¹ and reagent modifiers in organic chemistry.¹²

INTRODUCTION

The two systems that will be discussed are the N_4 and N_2O_4 crowns.

Figure 5: N_4 and N_2O_4 models

N_4 MODELS

Cyclic N_4 systems are not only important in the disproportionation of superoxides but also as oxygen carriers.^{13 a,b,c} They have also been found useful as potential catalysts for insertion of oxygen into organic substrates.^{14 a,b}

The amine analogues of crown ethers have been developed in several ways in order to obtain ligands with new properties. Stability, selectivity and solubility were addressed by introducing functionalized side chains attached to the macrocyclic ring or the nitrogen sites.^{15 a,b} In other cases, the host metal is accommodated by the functional group of the side chain as well as the crown itself.¹⁶ Redox potentials of host metal are greatly affected by pendant functional groups.¹⁷

It has been observed that other changes to the cyclic structure that do not have a direct influence on the binding of the metal can also affect the behavior of the macrocyclic compound. Much work has been done in the crown ether systems to study the effect of small conformational changes to the ring structure by the introduction of cyclohexyl and benzyl moieties.¹⁸ These components have altered the physical properties of their non-substituted counterparts. It is observed that the aliphatic substrate is the more flexible of the two and more closely echoes the binding property of the unsubstituted ring^{19 a,b,c}

Previous work with dicyclohexyl 14-crown-4 has shown the existence of two unequally populated conformations for the cis-anti-cis isomer in solution.²⁰

Figure 6: Conformations of cis-anti-cis dicyclohexyl 14-crown-4

Interconversion between these two conformations is a result of one cyclohexane ring inverting along with reorganization of the 14 member ring. For the cis-syn-cis isomer, there exists a single degenerate conformational process attributed to ring inversion of both cyclohexane rings. This inversion is also accompanied by a reorganization of the 14 membered ring geometry.

Figure 7: Conformation of cis-syn-cis dicyclohexyl 14-crown-4

In order to study the influence of nitrogen vs oxygen on the stereochemical dynamics in solution, the synthesis of several dicyclohexyl tetraaza 14-crown-4 ethers depicted in Figure 8 has been attempted.

Figure 8: Target molecules

Another component that has been used to impose change on the macrocyclic ring is the presence of an amide as a donor group. These dioxo systems are of interest because they possess both the novel ligand properties of the saturated tetraamine and oligopeptides.²¹ Thus, the amide and cyclohexyl substituents can be used to alter the ring flexibility in the cyclam skeleton. The resulting conformation equilibria in solution can be studied to observe these structural effects on the dynamic stereochemical behavior.

N₂O₄ MODELS

Many macrocyclic polyethers have attracted attention because of their ability to form stable complexes. In order to fulfill the stereochemical requirement of some metals, many crown systems have been prepared with ring sizes from 14 to 32.²²

However, two of the most widely investigated systems are dibenzo 18-crown-6 and dicyclohexyl 18-crown-6. These products have different properties which result in different binding abilities.

Figure 9 : Stability constants (Log k_1) for dibenzo and dicyclohexyl 18-C-6²³

	Na ⁺	K ⁺	Cs ⁺	
	4.36	5.00	3.55
c-s-c	4.08	6.01	4.61	
c-a-c	3.68	5.38	3.49

t-s-t	2.99	4.14	3.00	
t-a-t	2.52	3.26	2.27	

The difference in selectivity shown in Figure 9 reflects both the electrostatic and steric effects of the annelated ring. The strength of the C-O dipoles are affected by the nature of the substituent. Rigidity and flexibility of a ligand skeleton can dictate the cavity size of the ligand. Ligands of the flexible type show a plateau selectivity because they can flex to accommodate a host cation. This is not the case for smaller more rigid systems showing a peak selectivity for the cation that best matches its cavity size.²²

Figure 10 : Selectivity characteristics of some crown ethers

Figure 11 : Stability constants for some dibenzo crowns.

Benzene and cyclohexane rings are able to alter the complex constants themselves as well as the selectivity of the ligand. This can be seen by considering decomplexation energy for Na^{+} .

Figure 12: Decomplexation energy of Na^+ for dibenzo and dicyclohexyl 18-C-6

The barrier to removal of Na^+ seems to be a conformational change. The smaller activation energy of decomplexation of the dicyclohexyl material is attributed to the greater flexibility of the ligand.²⁴

Conformational changes to the ring structure have a profound effect on the stability of the complex. Figure 10 indicates that in all three cases, the stabilities are affected by the stereochemistry of the substituents on the 18-crown-6. In the crystal structure of the trans-syn-trans isomer, four of the oxygens point towards the center of the molecular cavity.²⁵ The trans-anti-trans isomer only has one of the oxygen donor groups pointing towards the center. The cis isomers both have four oxygen atoms pointing into the center of their molecular cavities. Figure 10 clearly indicates the cis isomers form the more stable complexes. The reason for the difference is rooted in the known flexibility of the cis disubstituted cyclohexane compared to its trans isomer. The cis isomers are more prepared to flex in order to direct the remaining two oxygen donor atoms towards the center of the macrocyclic cavity.

The stability and selectivity of the ligands are also determined by the interaction of the cation with the solvent.^{26 a,b} The stability constants of both metals in Figure 13 indicates that the degree of solvation of the metal is a function of the solvent. When methanol is used as the solvent, the metal is less solvated and consequently the ligand binding ability is enhanced.

Figure 13: Stability constants for dibenzo 18-C-6 in various solvents²⁵

	Log K_s	
	H ₂ O	MeOH
Na ⁺	1.21	4.08
K ⁺	2.02	6.01

The varied properties of these isomers have generated interest in structural studies of these ligands in solution.

Many of the conformational studies of these systems have been carried out with the cis isomers. The greater flexibility in these products provide some very interesting conformational changes. The cis-syn-cis and cis-anti-cis isomers of the dicyclohexyl 18-crown-6 contain different conformations in solution. The major conformational isomer is one having the characteristic all gauche O-CH₂-CH₂-O network in the macrocyclic ring.^{27 a,b}

Figure 14: Conformation of the major isomer of cis-syn-cis dicyclohexyl 18-C-6 in solution

The minor isomer is suggested to be one that has been deformed in the central macrocyclic region of the ring in the macrocycle.

Figure 15: Conformation of the minor isomer of cis-anti-cis dicyclohexyl 18-C-6 in solution

It is known that in solution torsional angle differences can exert pronounced influences on ^{13}C chemical shifts. In every instance carbons experiencing gauche interactions are shielded by approximately 5 ppm. relative to their trans counterparts.²⁸ This fact has been used extensively to identify the presence of various conformational isomers in solution. The bond angles for sp^3 hybridized carbon and oxygen are qualitatively similar and it should be reasonable to assume angular dependence of ^{13}C chemical shifts in the O-C-C-O and C-C-O-C network. However, it is unknown whether we can make this assumption with nitrogen. Thus, the dihedral angle dependence of ^{13}C chemical shifts may be different for O-C-C-N and C-C-N-C. To further study the process in solution and attempt to elucidate the structure of the nitrogen analogues of the dicyclohexyl 18-C-6, the synthesis of the cis isomers of dicyclohexyl diaza 18-C-6 was attempted.

Figure 16: Target molecules for the N_2O_4 system

Product I

cis-s-cis
cis-a-cis

In the course of our attempted synthesis of the cis isomers of the product I (Figure 16), we obtained some benzylated and tosylated aza analogues that exhibited a number of interesting stereochemical features. Large ^{13}C nmr chemical shift differences between g

and **9** were found to be the result of conformational differences in the macrocyclic portions of the molecule.

Figure 17 : Benzylated and tosylated precursors

Since molecular flexibility of crown ethers has led to difficulties in freezing out particular conformations in solution, conformational studies are rare.²⁹ However, Kleinpeter has used N-tosyl substituted crown ether aza analogues to study ring inversion processes in solution.³⁰ Using the tosyl substituents, he has been able to impose enough destabilization to increase the energy of transition resulting in separate nmr signals being observed for the preferred conformer and its less populated counterpart. Hence, the tosylated and benzylated 18 membered crowns can be used to study the inversion processes occurring in these compounds.

THEORY AND BACKGROUND

I. MACROCYCLIC EFFECT

The term macrocyclic effect was introduced by Cabbiness and Margerum in 1969 to account for the greater thermodynamic stability of complexes containing macrocyclic ligands when compared with those of non macrocyclic ligands of similar structure.³¹ Early work has ascribed the macrocyclic effect exclusively to either enthalpy³² or entropy^{33 a,b} terms. Recently it has been shown that ΔS is always positive whereas ΔH varies in magnitude with the matching of the size of the metal ion and the hole size of the ligand.^{34 a,b} When ether oxygens are replaced by nitrogen or sulfur, the stabilizing influence of cyclization of the ligand disappears.

Figure 18 : Stability constants of some K^+ and Ag^+ complexes³⁵

In Figure 18, the increase in the stability of the open chain ligand to the cyclized ether system indicates the magnitude of the macrocyclic effect. However, there is very little difference in the stability constants for the cyclized diamine going from 7.8 in the open chain ligand to 7.9 in the cyclic ligand. This effect has been attributed to the fact that in oxygen crown systems, the potassium metal is bound to all the donor atoms. However, when an oxygen is replaced, the metal is sometimes bound to only a few of the donor atoms. This is the case for the copper complex of diazatetraoxa 18-crown-6.

Copper(II) is coordinated to two nitrogen atoms and to only two of the four oxygen atoms.

Figure 19 : Cu^{2+} complex of diaza 18-C-6³⁶

However, with a system such as cyclen, all four donor sites are involved with the binding of the metal ions. The stability constant difference reflects the macrocyclic effect.

Figure 20 : Stability constants of some Ni^{2+} complexes²²

Log k = 15.3

Log k = 22.2

II. COMPLEXATION

Two factors affecting complexation are the type of cation and the donor atoms in the ligand. Substitution of nitrogen or sulfur for oxygen in 18-crown-6 has a profound influence on the complexation of metals. Stepwise substitution of nitrogen or sulfur in the crown skeleton has led to conclusions about their stabilizing/destabilizing influences on complexation.²³ In Figure 21, the substitution of oxygen for nitrogen reduces the affinity of the ligand for the alkali and alkaline earth metals while enhancing the stability of some of the transition metals that are of comparable size.

Figure 21 : Stepwise substitution of oxygen donor atoms in 18-crown-6.

K_s				Solvent
K^+	6.10	3.90	2.04	MeOH
Ag^+	1.60	3.30	7.80	H ₂ O

In macrocyclic chemistry, it is generally accepted that the cavity at the center of the ligand produces the enhanced complex stability when there is a match between the size of the cavity and the metal ion being complexed. Molecular mechanic (MM) calculations have shown a steady increase in the hole size from 12-ane-N₄ to 16-ane-N₄.³⁷ The flexibility of the ligand is also an important factor in the stability of the complex. Flexibility can be introduced by increasing size of some or all of the the chelates in the ligand.

The structure of the groups connecting the donor atoms has been the subject of much research.³⁸ This work has led to the observation that complexation is often affected by the structure of the binding chelate.

N₄

The ethylenoxy group plays a very important role in complexation. The intervening CH₂-CH₂ fragment and the coordinating metal are more stable than the 6 membered and the four membered ones.^{39 a,b}

Figure 22 : Relative stability of various membered ethylenoxy chelates

The effect of increasing the chelate size on the metal ion selectivity of the ligand is also found in tetraaza systems.⁴⁰ The high Ligand Field strength found in some complexes of N₄ donor ligands has been used to postulate that the N₄ macrocyclic ligand is rigid enough to compress a metal that is larger than what a size-match fit would dictate. Increasing the chelate size results in an increase in ligand flexibility. Metal ion compression leads to an out of plane mode of coordination in an attempt to lessen compressive forces.⁴⁰ The out of plane mode of coordination decreases the size-match selectivity. Since selectivity has been a major objective in the design of these host systems, the factors involved in complexation of particular metals has always been of interest.

The stability of the small metal complexes sometimes level off or completely reverse after a number of 5 membered chelates are changed to 6 membered chelates. Results show that this reversal occurs after two increases going from 12-ane-N₄ to 14-ane-N₄.⁴⁰ A quantitative assessment of the metal ion size relationship has also been made by Bush and co-workers using Co(II) and Ni(II) couples. A minimum strain energy was found for the 14 membered ring.⁴¹

Thus in order to achieve both stable and selective ligands, the structure must contain the proper cavity diameter as well as a degree of rigidity that will not allow the ligand to distort in order to accomodate the metal.

The ligand must also satisfy the geometric requirements of the metal. Every asymmetry of charge distribution in the crown also disturbs the complexation of the spherical metal ion.⁴² The arrangement of the donor atoms is important to the binding ability of the crown. Every deformation of the inner charge shell which is not in keeping with the geometry of the guest, reduces the binding ability of the ligand and the stability of

the complex. This is the reason Ni(II) complexes are so stable with 14-ane-4 (Cyclam). Cyclam incorporates the transition metal in a very flexible conformation which accommodates the geometry requirement of the Ni perfectly to permit the formation of especially strong metal-nitrogen interactions.⁴³ It has always been recognized that these ligands complex strongly with those metals whose ionic radius matches the radius of the cavity. When cyclam incorporates the Ni (II) in its cavity, it adopts the characteristics of a square planar diamagnetic complex of the ligand without distortion of the central cavity.⁴⁴

With the understanding that the 14 membered ring is an ideal ligand for the complexation of nickel, the addition of the cyclohexyl rings gives us an excellent opportunity to impose extra rigidity in the two 5 membered chelates. The presence of the amide donor group in the two remaining 6 membered chelates is an attempt to increase the rigidity of the macrocyclic ring. The increase in the overall rigidity, should in effect, reduce the ability of the ligand to distort its conformation in order to fit the metal. This factor is important because it is at this point that selectivity can be achieved. If the compression forces the metal induces are severe enough to induce a conformational change in order to fit the metal, the ligand will not bind. However, if the metal fits the cavity without imposing a change in the ring it can be selectively bound. Thus, both factors could be used to enhance the binding ability as well as the selectivity of the trans substituted tetraaza 14-crown-4 ethers. Since the rigidity of the trans disubstituted pattern will not allow the interconversion of the isomers, the stereochemical dynamics in the dicyclohexyl tetraaza 14-crown-4 ethers would be limited to macrocyclic ring conformational changes. As the more flexible substituent, the cis isomer should, however, provide enhanced stereochemical dynamics in solution giving rise to interesting conformational changes in both the macrocyclic ring and the cyclohexyl substituent.

N₂O₄

The binding of alkali and alkaline earth metal ions to macrocyclic crown ethers is electrostatic in nature. Ligand basic groups try to orientate themselves as evenly as possible over a sphere of positive charge found on the metal. Thus, the requirements for these ligands are not as specific as with the N₄ systems. Wide variations in coordination number and geometry are thus possible.⁴⁴ All that is necessary is that the ligand provide an electronic environment to replace all or part of the metal's solvation shell. When some oxygen donor groups are replaced by nitrogen in the crown skeleton there is a resulting change in the binding ability of these crowns. The binding of a metal is now a result of

electrostatic and covalent bonds. This offers an expansion of the variety of metals capable of being complexed by crown ethers alone.

III. ¹³C Nuclear Magnetic Resonance^{45,46}

In the presence of a magnetic field H_0 , a nucleus may undergo magnetic resonance if it possesses nuclear spin. Nuclei with spin quantum number $I \neq 0$ interact with magnetic fields due to their magnetic moments.

$$\mu = \gamma \hbar I$$

The gyromagnetic ratio, γ , characterizes each nucleus. When exposed to a static magnetic field, H_0 , a spinning nucleus behaves like a gyroscope in a gravitational field.

The frequency of precession about H_0 is known as the Larmor frequency. The Larmor precession is accelerated by increased H_0 .

$$\nu_0 = \frac{\gamma}{2\pi} H_0$$

However, it is more realistic to study the influence of magnetic fields on an assembly of a large number of identical nuclei than on one spinning nucleus. With the proper frequency ν_1 , the magnetic moment vectors μ representing the nuclear magnetization are forced to precess in phase. The resulting magnetization vector is no longer parallel to H_0 and is now composed of three components along the x, y, and z axes.

$$\mathbf{M} = M_x \mathbf{i} + M_y \mathbf{j} + M_z \mathbf{k}$$

Figure 23: Cartesian coordinate description of a perturbed magnetization vector

Magnetization vector- M_z

M_z is the magnetization vector along the z-axis. Relaxation along this axis is called spin lattice relaxation with a time T_1 . Relaxation occurs by the loss of energy from the excited nuclear spins to the surrounding molecular lattice.

^{13}C relaxation times cover a range from a few milliseconds to several hundred seconds. There are a number of mechanisms that contribute to spin-lattice relaxation. The most common of these are

dipole-dipole	T_{1DD}
spin-rotation	T_{1SR}
quadrupolar	T_{1Q}
scalar	T_{1SC}
chemical shift anisotropy	T_{1CSA}

Each of these combine to give an overall spin-lattice relaxation time T_1

$$\frac{1}{T_1} = \frac{1}{T_{1DD}} + \frac{1}{T_{1SR}} + \frac{1}{T_{1Q}} + \frac{1}{T_{1SC}} + \frac{1}{T_{1CSA}}$$

Carbons bearing a second nucleus possessing a magnetic spin relax more efficiently than nonsubstituted carbons. For carbons bearing directly bonded protons, T_{1DD} often constitutes the only relaxation mechanism available. The relaxation rate (R) is defined as the reciprocal of the relaxation time. For most non-viscous solutions, R is equated as

$$R_{1DD} = \frac{1}{T_{1DD}} = \left(\gamma_H^2 \gamma_C^2 \frac{\hbar}{2\pi} \right)^2 \tau_c \sum \frac{1}{r_{C-H}^6}$$

R_{1DD} is directly proportional to the correlation time τ_c and inversely proportional to the internuclear distance r^6 . Hence, protons that are not directly attached to the carbon will not

contribute to the relaxation. Relaxation can also occur from T_{1SR} . This mechanism is, however, accompanied by a decrease in the effectiveness of the dipole-dipole relaxation mechanism which results in a longer T_{1DD} . The physical observation of carbons having longer relaxation times may result in the low intensity absorption bands found in the ^{13}C nmr.

Figure 24: Cartesian coordinate description of the relaxation along the Z axis

M_x, M_y

M_x , and M_y are the transverse magnetization components in the xy plane. Relaxation in this plane is expressed in terms of spin-spin relaxation with a time T_2 . Relaxation occurs by the interchange of energy from nuclei spins. The result of this is a fanning out in the xy plane. The net result is that the positive and negative signals cancel each other, leading to no detectable signal along the x or y axis.

Figure 25: Cartesian coordinate description of the relaxation along the XY plane

Relaxation times - T_1 Measurement

The spin lattice relaxation times (T_1) are usually the parameter of choice when attempting to extract information from the molecular motion of the atoms of the molecule. This is because T_c is the time required for the C-H vector to rotate through one radian. A common method used for the determination of spin lattice relaxation times is known as the inversion recovery method. A 180 degree pulse is applied to invert the magnetization to the -z axis.

Figure 26: Cartesian coordinate description of the Inversion Recovery method for determination of T_1

Immediately after the pulse, M_z will begin to relax back along the z axis towards the equilibrium value M_z . This process is expressed as:

$$M_z = M_z (1 - 2\exp(-t/T_1))$$

When M_z passes through the zero point, T_1 is calculated in the following manner;

$$0 = M_z (1 - 2\exp(-t/\Gamma_1))$$

$$T_1 = \frac{t_0}{2.303 \log 2} = \frac{t_0}{0.693}$$

t_0 = time when $M_z = 0$

n.O.e.

The enhancement of peak intensity during double irradiation experiments is known as the nuclear Overhauser effect (n.O.e.). The effect can be attributed to a cross relaxation between spatially proximate nuclei. When no other relaxation mechanism is operative other than dipole-dipole, n.O.e. enhancement will reach maximum enhancement value. For cross relaxation between protons, this maximum is 50%, whereas for ^1H - ^{13}C cross relaxation, the value is approximately 300%. The utility of n.O.e. conformational studies comes from the dependence of n.O.e. on nuclear distances as shown in the r^{-6} term in equation .

$$R_{1DD} = \frac{1}{T_{1DD}} = \left(\gamma_{\text{H}}^2 \gamma_{\text{H}}^2 \frac{\hbar}{2\pi} \right)^2 \gamma_c^2 \sum \frac{1}{r_{\text{H-H}}^6}$$

RESULTS AND DISCUSSION

I Attempted synthesis of dicyclohexyl tetraaza 14-crown-4

In our laboratory, previous attempts at the synthesis of configurational isomers of dicyclohexyl tetraaza-14-crown-4 using tosylated starting materials failed to give the desired 14 membered ring.⁴⁷ The synthetic scheme that was followed is shown in Figure 27.

Figure 27 : Proposed synthetic scheme for synthesis of dicyclohexyl tetraaza 14-crown-4

Scheme I

Scheme II

Scheme III

The final cyclization of the ditosylate in scheme III (Figure 27) was found to proceed via the intramolecular pathway to produce a seven membered ring as depicted in Figure 28.

Figure 28: Intramolecular reaction product from Scheme shown in Figure 27

In our attempts at the total synthesis of the isomers of the dicyclohexyl tetraaza 14-crown-4, we have focused on two different approaches:

- method 1: The addition of amines to α,β unsaturated carboxylic esters followed by cyclization by aminolysis of the ester
- method 2: Metal assisted cyclizations using nickel.

METHOD 1

Attempted total synthesis of dicyclohexyl tetraaza 14-crown-4

Several review articles have appeared that deal primarily with the synthesis of complexes of synthetic macrocyclic ligands.^{48 a,b,c} Many of these reactions have involved complexation between a presynthesized ligand and a metal in solution. This has also been our method of choice for the synthesis of dicyclohexyl tetraaza 14 crown-4 ethers. This approach has been extensively used for the synthesis of complexes containing synthetic macrocycles including cyclic polyethers (coronands) and macrobicyclic ligands (cryptands). In order to reduce side reactions such as polymerization, the final ring closure is usually carried out at high dilutions. The advantages of this method are found in its generality. In the final cyclization step, all isomeric products are theoretically possible (see Figure 8).

One potential disadvantage to this method is that the synthesis often results in low yields of the desired product. The second method involves the synthesis of the ligand in the presence of the metal ion. There have been many reported methods that have used the metal to enhance intermolecular reactions to improve yields of larger cyclic structures.^{49 a,b,c} It is known that the metal ion plays an important role in directing the steric course of the cyclization.⁵⁰ This is a potential disadvantage in that some isomer may not be formed upon cyclization due to unfavorable interaction of the intermediates.

The addition of linear polyamines to α,β unsaturated carboxylic esters involves a Michael addition followed by intramolecular lactam formation in one step.^{15,45} The addition of a racemic mixture of trans 1,2 diaminocyclohexane and methyl acrylate in

MeOH gives qualitative yields of **1** as an oil (see scheme in Figure 29). The ^{13}C nmr of **1** shows 7 resonances with chemical shifts: 172.47(C=O), 60.59(C^{*}H-N), 50.81(C^{*}H₃-O), 41.49 (C^{*}H₂-N), 34.40 (C^{*}H₂-C=O), 30.78 (CH₂-C^{*}H-N) and 24.39 (C^{*}H₂-CH₂-CH). The mass spectrum contains the parent ion at 287 m/e. The IR shows the ester peak at 1736 cm⁻¹. Aminolysis with **1** using a second equivalent of the trans 1,2 diaminocyclohexane yielded a cyclic amide product. The proposed scheme is found in Figure 29.

Figure 29 : Proposed synthetic scheme for the synthesis of dicyclohexyl-tetraaza 14-crown-4

The reaction progress was monitored by IR. The formation of the amide peak at 1645 cm⁻¹ was followed by the disappearance of the ester peak at 1736 cm⁻¹. The scheme in Figure 29 was designed to minimize the intramolecular reaction by forming the diester before the final cyclization step. Initial synthesis of **1** also fixes the location of the amide donor sites on the macrocyclic ring.

Aminolysis of **1** with trans 1,2 diaminocyclohexane yielded a reaction product with a MS peak at 168 m/e. The ^{13}C nmr showed 9 peaks with chemical shifts 177.43 (C=O), 61.15, 57.15 (C^{*}H-N), 43.11 (C^{*}H₂-C=O), 40.63 (C^{*}H₂-N), 32.70, 31.27 (C^{*}H₂-CH-N), 23.91, 23.71 (C^{*}H₂-CH₂-CH-N). Diborane reduction of **2** gives **3** as a colorless oil. The reaction was also monitored by IR for the disappearance of the amide

peak at 1645 cm^{-1} . The ^{13}C NMR spectrum of **3** showed 5 resonances with chemical shifts at 67.2 ($\text{C}^*\text{H-N}$), 46.59 ($\text{C}^*\text{H}_2\text{-N}$), 33.45 ($\text{C}^*\text{H}_2\text{-CH-N}$), 31.38 ($\text{C}^*\text{H}_2\text{-CH}_2\text{-N}$), and 25.16 ($\text{C}^*\text{H}_2\text{-CH}_2\text{-CH-N}$). The mass spectrum of this material gave a peak at 154 m/e (E.I.)

The MS data tend to indicate that the **3** is not fully cyclized 14 membered ring. In view of the fact that the **1** was isolated and purified before the aminolysis step, the result of the mass spectrum could lead to the conclusion that the parent ion was not obtained because of the cleavage pattern of the target molecule. Mass spectral data for some polyamines have been reported to generate structural units one half of the original structures⁵¹

Figure 30 : Possible cleavage pattern of **3**

In order to clarify the ambiguous nature of the results for the molecular weight of **2**, an alternate method for molecular weight determination had to be employed. The measurement of the colligative properties of the unknown are well documented methods for molecular weight determination.⁵² Freezing point depression data reveal that the molecular weight of the present product is approximately 145 g/mol. (see Appendix 1,1-i, and 1-ii for freezing point data and calculations)

The experimental setup for freezing point depression was evaluated by comparing the experimental molecular weight determination with a known standard. The standard deviations for the three experiments were well within acceptable limits. The relative error for the experimental determination of the molecular weight of benzophenone was 5%.

The experimental value of the unknown was found to be within 15 % of the expected value of the 7 membered ring. However, it is obvious that the molecular weight of **2** is far from the value of the 14 membered product (340 g/mol). Using the freezing point depression and mass spectral data, **2** has been assigned as the intramolecular 7 membered ring product.

Figure 31: Structure of intramolecular reaction product.

Since **1** had been isolated and purified before the cyclization step, the formation of the intramolecular product indicates an unstability of **1** in solution. It was later observed that when **1** was allowed to remain in solution for several days at room temperature, it converted to the same product as depicted in Figure 31.

Attempts at the Stabilization of 1

Since the target molecule was the 14 membered ring, cyclization of **1** could not occur until it was stabilized. Attempts to stabilize **1** followed two different approaches. The first involved chemical modification and the second involved complexation with a metal chloride. The second method will be discussed in a later section (**METHOD II**)

In an attempt to stabilize **1**, large tosyl groups were used to reduce the basic strength of the nitrogen as well as introduce some steric hinderance. Aminolysis of **1** with various other nucleophiles was also attempted. Since the tosyl group is large and bulky, it is possible that if the nitrogen atoms were tosylated the approach of the carbonyl group might be inhibited and the resulting formation of the 14 membered diamide would be possible.

Figure 32 : Proposed scheme for the synthesis 1,2 bis [3 methyl ester-N-tosylamine] cyclohexane

Although many 2^o amines have been tosylated in pyridine⁵³, the nitrogens in **1** are now too hindered to induce a nucleophilic attack on the TsCl.

A second approach to the synthesis of **1** involved tosylating the starting material before the Micheal addition.

Figure 33 : Scheme for the synthesis of 1,2 bis
[3 methyl ester-N-tosylamine] cyclohexane

The tosylation of the *trans* 1,2 diaminocyclohexane gives a 70 % yield of **4** as white plates. The ^{13}C nmr of **4** shows 8 resonances with chemical shifts : 144.06 ($\text{C}^*\text{H-SO}_2$), 137.58 ($\text{C}^*\text{H-CH}_3$), 130.12 ($\text{C}^*\text{H-CH-CH}_3$), 127.59 ($\text{C}^*\text{H-CH-SO}_2$), 57.03 ($\text{C}^*\text{H-N}$), 33.56 ($\text{C}^*\text{H}_2\text{-CH-N}$) and 24.60 ($\text{C}^*\text{H}_2\text{-CH}_2\text{-CH-N}$). When **4** is refluxed with methyl acrylate in solution, only starting material is recovered. The inductive effects of the tosyl group decreases the basicity of the nitrogen atoms and **4** is no longer a strong enough base to induce a Micheal type reaction on the methyl acrylate. Typically, sulfonamides have pKa's of -1.5 .

The second group of experiments dealt with changing the nucleophile in the aminolysis reaction. If steric consideration of *trans* 1,2 diaminocyclohexane inhibits nucleophilic attack on the carbonyl of the ester group in the methyl acrylate, a reduction in the size of the nucleophile could possibly induce an intermolecular cyclization reaction. However, the use of 1,2 ethylenediamine and ethylamine were also found to be ineffective in the formation of the 14 membered ring.

Mechanistic description of the intramolecular reaction product.

Under the present synthetic conditions, the only product isolated is the intramolecular reaction product. The formation of this product is possibly due to a retro type Michael reaction as shown in Figure 34.

METHOD 2*Attempted Template synthesis of the dicyclohexyl tetraaza 14-crown-4*

By isolating intermediates in the formation of the macrocyclic complexes, Curtis and co-workers have been able to deduce some mechanistic features of reactions involving a metal template.^{54 a,b,c}

Figure 36 : Mechanism of a template reaction

Curtis has proposed that in the condensation reactions, the initial product is the isopropylideneimine species which undergoes an Aldol type reaction between the imino group and an acetone molecule to yield a β amino ketone. At this point, the molecule is either stabilized by coordinating with the keto group or undergoes an imine formation with the available amine.

Using scheme I shown in Figure 36 as a model, our synthetic approach involved the complexation of **1** with nickel followed by aminolysis with trans 1,2 diamino cyclohexane as shown in Figure 37. The preference for nickel(II) in complexation reactions reflects, in part, the ability of the metal to act as a metal template for ligand forming reaction, the lack of interfering redox reaction, and a well understood ligand field description of the various coordination geometries of the nickel (II).⁵⁵

Figure 37 : Proposed scheme for the synthesis of tetraaza dicyclohexyl tetraaza 14-crown-4 ether using metal template

The Ni(II) complexes of many polyamine ligands in solution produce a mixture of violet, octahedral paramagnetic $\text{Ni.L}(\text{H}_2\text{O})_2^{2+}$ and yellow square planar diamagnetic Ni.L_2^{2+} complexes.^{56a,b} Similarly, when Ni^{2+} is added to **1** in methanol, the solution immediately turns violet. This is indicative of the cis coordination of the nickel.

Figure 38: Proposed conformation of the cis coordinated Ni^{2+} complex of **1**

Evidence for the complexation of **1** is found in the titration of the ligand with $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ in methanol. Figure 39 indicates that the titration of the ligand with the metal results in a carbonyl shift from 1739.5 cm^{-1} to 1680.0 cm^{-1} in the IR. As **1** is titrated, the disappearance of the band at 1738 cm^{-1} is followed by the formation of a band at 1680 cm^{-1} . When an equivalent amount of the metal is added, the band at 1738 cm^{-1} disappears. This can be rationalized by considering the electron withdrawing effect of the Ni on electron density of the carbonyl. As the Ni is coordinated, electron density is removed from the carbonyl into the d_{z^2} orbital of the nickel making the carbonyl bond vibrate at a higher frequency.

More evidence for the complexation of **1** is also found in the visible spectrum in Figure 40. When $\text{NiCl}_2 \cdot \text{H}_2\text{O}$ is dissolved in solution, the electronic transition that occur are from ${}^3\text{T}_{2g} - {}^3\text{A}_{2g}$ ($V_1 = 1100\text{ cm}^{-1}$), ${}^3\text{T}_{1g} - {}^3\text{A}_{2g}$ ($V_2 = 730\text{ cm}^{-1}$), and ${}^3\text{T}_{1g} - {}^3\text{A}_{2g}$ ($V_3 = 400\text{ cm}^{-1}$).⁵⁷ The replacement of the water ligands by the donor sites in **1** should cause an increase in the Dq of the complex. This increase is anticipated because amines are found higher in the spectrochemical series than water. The result of this replacement would be a hypsochromic shift of the observed bands. The visible spectrum shows the expected shifts towards higher energy. The V_1 and V_2 transitions have shifted to 900 cm^{-1} and 590 cm^{-1} . The V_3 has trailed off into the UV. The electronic spectrum of **5** is typical of other cis isomerized complexes of nickel.⁵⁸

Figure 39 : Infra-red spectra of the titration of **1** with $\text{NiCl}_2 \cdot \text{H}_2\text{O}$

in methanol (salt plates)

a) ligand with 0.0 equiv Ni b) ligand with 0.25 equiv Ni
c) ligand with 0.5 equiv Ni d) ligand with 1.0 equiv Ni

Figure 40 : The U.V. spectrum of a) $\text{NiCl}_2 \cdot \text{H}_2\text{O}$ in methanol
 b) $\text{NiCl}_2 \cdot \text{H}_2\text{O}$ with 1.0 equiv of **1** in methanol

When the **1** is reacted with an equivalent amount of $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ in methanol, the resulting complex forms the cis coordinated product as shown in Figure 38. The resulting addition of trans 1,2 diaminocyclohexane to this product gives a low yield of a product that contains an $M+1$ peak at 337 m/z (C.I.). This peak had not been observed in any of the earlier attempts in method 1. Since the product requires further purification ^{13}C nmr could not be used at this point. However, the mass spectral data is encouraging and the template method seems to have given the required conditions for cyclization of **1**.

Attempts at the synthesis of cis dicyclohexyl tetraaza 14-crown-4

When a mixture of cis and trans 1,2 diaminocyclohexane was reacted with methyl acrylate, two products were obtained.

Figure 41 : Synthetic scheme for the synthesis of cis and trans diester

Both **1** and **6** were obtained in high yields. We have discussed earlier the properties of **1** in solution. When a mixture of **1** and **6** in methanol was allowed to reflux for 48 hours, the only product isolated was the intramolecular product **2** and **6**. The chemical inertness of **6** prompted us to attempt its cyclization under different conditions. The synthetic approach is shown in Figure 42.

Figure 42: Proposed synthetic reaction scheme for the synthesis of cis dicyclohexyl tetraaza 14-crown-4 ether

Isolation of the racemic cis 1,2 diaminocyclohexane from a mixture of cis and trans

isomers was accomplished using the method described by Santo and Kidani⁵⁹. When the cis 1,2 diaminocyclohexane was refluxed with methyl acrylate in methanol, quantitative yields of **6** were obtained. The ¹³C NMR of **6** shows 7 resonances with chemical shift 173.30 (C=O), 56.52 (CH₂-N), 51.45 (CH₃-O), 42.56 (CH₂-N), 35.0 (C^{*}H₂-C=O), 28.00 (C^{*}H₂-CH-N) and 22.30 (C^{*}H₂-CH₂-CH). The mass spectrum contains the M+1 peak at 287 m/z (C.I.). The IR shows a peak at 1740 cm⁻¹ indicative of the ester functional group. The solvents used for the attempted cyclization of the cis diester were methanol, dimethoxy ethane and THF. After several days refluxing, only starting material was isolated.

It is interesting to note the difference in stability of **6** compared to **1**. In solution **1** converts to the seven membered ring through an intramolecular reaction pathway (Figure 34). The product formed contains a strain free cyclohexyl ring trans fused to a quasi chair form seven membered ring. The intramolecular product of **6** is not observed. The stability of **6** could be perhaps due the strain involved in the formation of two cis fused rings.

II Attempted synthesis of dicyclohexyl diazatetraoxa

18-crown-6

Two synthetic approaches were taken for the synthesis of the dicyclohexyl diaminotetraoxa 18-crown-6.

METHOD I

The synthesis of **10** was initially attempted using the method of Hogberg.⁶⁰ The proposed scheme is found in figure 43.

Figure 43 : Synthetic scheme for the synthesis of Dibenzo-diazatraoxa-18-crown-6

Products **8** and **9** were isolated and purified. The yields were similar to those reported. However, Hogberg reported a 45% detosylation yield of **8** using glacial acetic acid saturated with HBr. In our hands, this method did not produce the detosylated product. Other methods attempted Na metal in HMPT⁶¹, and LiAlH₄ in THF⁶². The latter method was successful, however, the yield was too low for practical synthetic purposes.

METHOD II

The second synthetic approach was outlined by Pederson.⁶³

Figure 44 : Synthetic scheme for the synthesis of dibenzo-diazatetraoxa 18-crown-6

All products were isolated and purified by reported methods. However, Pederson reports hydrogenation of **14** at 50 °C and 100 psi for 6 hours debenzylated the 18 membered product. Our attempts involved hydrogenation of this product 70 °C and 800 psi. for 20 hours. Only starting material was recovered.

Catalytic transfer hydrogenation had been useful for removal of benzyl and benzyloxycarbonyl protecting groups in peptide synthesis.⁶⁴ Using Palladium black as the catalyst, **15** was obtained in quantitative yields. The ¹³C nmr of **15** showed 5 resonances with chemical shifts 148.09(C*H=C-O), 120.60(CH=C*H-C-O), 111.14

(C*H=CH-C-O), 67.46(C*H₂-N), 49.01(CH₂-O). The mass spectrum gives a peak at 358 m/e (MW=358 g/mol). This product was identical to a sample of **10** isolated earlier.

The hydrogenation of the dibenzo 18-crown-6 using ruthenium has been successful in obtaining the cis-syn-cis and cis-anti-cis isomers.⁶⁵ Another catalyst that can be used is rhodium. This catalyst is recommended because they cause the least amount of hydrogenolysis of ether linkages during reduction.⁶⁶ It is, however, affected by the presence of strong nitrogen bases, as in the unsuccessful reduction of aryl amines.⁶⁷ Benzyl amines have been hydrogenated using ruthenium and rhodium catalysts.⁶⁸ Unfortunately, all attempts at hydrogenation of the N₂O₄ analogue using these catalyst did not provide the cis isomer and only starting material was recovered.

NMR STUDIES

Conformational studies using low temperature ¹³C nmr

The ¹³C nmr spectra of **8** and **14** are temperature independent down to 193 K. This indicates that there must be rapid interconversion between conformational isomers on the nmr timescale. In **8** and **14**, the tosyl substituent are four atoms removed from the adjacent aromatic rings, and hence little steric interaction occurs between them.

Relative compounds studied by Kleinpeter possessed steric hinderance between the tosyl moiety and the more proximate neighbouring aromatic functions, which apparently raised the barrier to conformational interconversion.³⁰

Figure 45 : N-tosyl substituted aza crown ethers

Hence, spectra for separate isomers are observed at low temperature via ¹³C nmr.

Characterization of conformational differences between 9 and 18 membered ring tosylates

The ^{13}C nmr of **8** and **2** are shown in Figure 46 and 47. Assignments of resonances for individual carbons are listed in Table 1 and 2. (see Figure 48 for numbering scheme)

NOV	FREQ	PPM	INTEGRITY
51	7429.7	147.482	59.457
52	7223.2	143.576	10.145
53	6859.1	134.339	24.071
54	6531.9	129.827	121.935
55	6374.4	127.182	114.127
56	6090.8	121.052	119.583
57	5632.2	111.932	113.278
58	5213.9	77.795	31.120
59	3904.1	77.483	195.490
60	3851.6	77.283	36.416
61	3481.0	72.143	38.519
62	3274.2	77.009	167.879
63	3059.3	74.711	25.643
64	3051.2	74.540	32.212
65	2831.9	74.264	173.114
66	2404.4	47.478	113.469
67	2446.0	49.435	101.114
68	1581.8	21.504	46.212
69	35.4	0.504	39.247
70	14.1	0.328	41.431
71	-0.1	-0.003	597.245

Figure 46. The ^{13}C NMR spectrum of 2.

Figure 47: The ^{13}C NMR spectrum of 9

Figure 48 : Carbon numbering scheme for **8**, **9** and **14** Table 1: ¹³C nmr chemical shifts for **8** in CDCl₃ (numbering scheme in Figure 48)

CARBON #	CHEMICAL SHIFT	CARBON #	CHEMICAL SHIFT
1	121.05	6	143.58
2	111.95	7	127.14
3	147.68	8	129.84
4	67.67	9	136.58
5	48.64	10	21.50

CARBON #	CHEMICAL SHIFT	CARBON #	CHEMICAL SHIFT
1	124.36	6	143.46
2	122.87	7	127.10
3	151.25	8	129.77
4	74.00	9	135.80
5	51.19	10	21.48

With the exception of the tosyl carbons, all resonances for **8** are shielded upfield relative to those of **2**. The most pronounced differences are, however, at C₄ (6 ppm) and the ortho carbon C₂ (11 ppm). These chemical shift differences are most likely due to the stereochemical influences. Because of the difficulties in obtaining crystalline products of **8** and **2**, X-ray studies could not be used to determine the exact conformation. However, the ¹H-¹H nuclear Overhauser effect (n.O.e.) is a well known technique for determining internuclear distances in solution.⁶⁹ Hence, a proton homonuclear shift correlated 2-D nmr (H, H NOESY) experiment was performed and the spectrum is shown in Figure 50 for **8** and Figure 51 for **2**. The numbering scheme for the protons in **8**, **2** and **14** are found in Figure 49.

Figure 49 : The numbering scheme for the protons in the benzylated and tosylated precursors.

Figure 50: Homonuclear 2-D shift correlated (^1H , ^1H NOESY) spectrum of **8** in CDCl_3 with the 1-D spectrum above. The corresponding assignment is also shown

Figure S1: Homonuclear 2-D shift correlated (^1H , ^1H NOESY) spectrum of **2** in CDCl_3 with the 1-D spectrum above. The corresponding assignment is also shown.

As a result of these experiments, it was possible to determine which protons were interacting by cross relaxation. Using the numbering scheme found in Figure 48, a strong n.O.e. was observed between H5 and H6 in **8**. Cross relaxation of H5 and H6 must involve a particular conformation of the carbons bearing these protons. Using molecular models we can see that the protons are close when the CH₂-O carbon approaches the plane of the aromatic rings. As the ring carbon approaches the plane of the ring, the dihedral angle between the ortho carbon and the CH₂-O carbon approaches zero. In support of such a geometry, X-ray data on the related dibenzo 18-crown-6 has indicated that the CH₂-O carbons are found in the plane of the aromatic ring.⁶⁹

In contrast, the ring constraints of **2** force the C₄ carbons out of the plane of the aromatic ring in order to adopt an extended chair conformation as shown by molecular models. In TRIBL⁷⁰, MMX force field calculation are used to calculate geometric, spectroscopic and thermodynamic properties.^{72a,b} The close agreement between the experimental and MM calculated methods reinforce the predictive power of molecular mechanic calculations. Figure 52 illustrates the dihedral angles between the ortho carbon C₂ and the C₄ carbon in **8** and **2** as determined by TRIBL.

Figure 52 : MM calculations for dihedral angle between the CH₂-O and the ortho carbons in **8** and **2** (numbering scheme in Figure 48)

The smaller dihedral angles for the ortho carbon and the CH₂-O carbons in **8** results in a greater degree of π -steric interaction for these carbons compared to **2**. The increase in the number of contours for the H5 and H6 protons in the 2-D proton NOESY spectrum of **8** compared to **2**, indicates a greater degree of cross relaxation. Since the protons H5 and H6 are closer to each other in **8** than in **2**, the closer association of these protons in **8** is accompanied by the observed upfield shift of C₂ and C₄ resonances relative to **2**. Since n.O.e. is distance dependent, the proximity of H5 and H6 could be compared in **8** and **2**. The number of contours for the n.O.e. between H5 and H6 are found to be greater in the 2-D nmr of **8**. In addition the interatomic distances given by TRIBL show the closer association of H5 and H6 in **8** than in **2**. The calculated distances are found in Figure 53.

The average interatomic distance for H5 and H6 in **8** is 2.4 Å. The distance for these protons in **2** is found to be 2.8 Å. Because of the conformational change in **2**, this gauche interaction is reduced by an increase in the torsional angle between the carbons of concern. The negligible n.O.e. of H5 and H6 in **2** mirrors the decrease in the gauche interaction by noting the decrease in the proximity of the H5 and H6 protons from 2.4 Å in **8** to 2.8 Å in **2**.

The assignment of the ¹³C nmr spectrum of **14** in Figure 54, was accomplished with the aid of a heteronuclear 2-D shift correlated (HETCOR) experiment shown in Figure 55. Individual chemical shifts are presented in Table 3.

Table 3: ¹³C nmr chemical shifts for **14** in CDCl₃ (numbering scheme in Figure 48)

CARBON #	CHEMICAL SHIFT	CARBON #	CHEMICAL SHIFT
1	120.49	6	58.57
2	111.79	7	139.62
3	148.39	8	127.00
4	66.94	9	128.62
5	53.53	10	128.29

With the exception of the benzyl group, the ¹³C nmr shifts in **14** are remarkably similar to those of **8**. In addition, the ¹H-¹H n.O.e. results for **14** shown in Figure 56, are nearly identical to **8** suggesting similar crown ether conformations in **8** and **14**.

Figure 53: Interatomic distances for the protons of **8** and **2** as determined by the MMX program TRIBL.

Figure 54 The ^{13}C NMR spectrum of **14**.

Figure 55: Homonuclear 2-D shift correlated (^1H , ^{13}C HETCOR) spectrum of **14** in CDCl_3 with the 1-D spectra included.

Figure S6: Homonuclear 2-D shift correlated (H,H NOESY) spectrum of **14** in CDCl₃ with the 1-D spectrum above. The corresponding assignment is also shown

The ^1H nmr spectra offer some additional insight into the dynamics of **8** and **9**. In **8**, the $\text{CH}_2\text{-O}$ and $\text{CH}_2\text{-N}$ protons are simple triplets with area ratios of 1:2:1. An average coupling constant of approximately 6 Hz is observed. This suggests some rotational averaging of local conformations in the $\text{O-CH}_2\text{-CH}_2\text{-N}$ portion of the molecule.

By contrast in **9**, a more complex pattern is observed. If **9** is essentially locked into an extended chair conformation, then these 4 protons represent an ABCD spin system, rather than an A_2B_2 as is represented in **8**. To verify this, the ^1H nmr spectrum of **9** was simulated using the LAME⁷³ program, with chemical shifts and coupling constants shown in Table 4.

Table 4: Theoretical ABCD spin system parameters for simulated spectrum in Figure 57

<u>CHEMICAL SHIFT</u>		<u>COUPLING CONSTANT</u>	
SYSTEM	C.S. (Hz)	SYSTEM	COUPLING J (Hz)
<hr/>			
A	1317	J_{AB}	-12.00
B	1313	J_{AC}	3.00
C	1040	J_{AD}	6.00
D	1043	J_{BC}	6.00
		J_{BD}	3.00

The good agreement between experimental and calculated spectra indicates that the postulate of a "locked " extended chair conformation in the 9 membered ring of **9** is reasonable. Figure 56 shows the experimental and calculated spectra.

Figure 57 : Experimental and simulated spectra for the $\text{CH}_2\text{-O}$ and $\text{CH}_2\text{-N}$ in **9**.

Complexation properties of the 7 membered trans 1,2 diamine cyclohexane and its oxo precursor.

Since the presence of the oxo group would inevitably alter the flexibility of the seven membered ring, it is anticipated that the above materials would also have different properties. Titration of **2** and **3** with a metal capable of binding with the nitrogen donor groups, gives some insight into the effect of the amide on the property of the aza crown ether.

Figure 58: Structure of cyclic amide and its reduced product

When $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ is added to a methanol solution of **3**, the resulting yellow solution is consistent with the formation of a square planar complex. The complex was isolated and the elemental analysis was consistent for a 2:1 complex. The results of the elemental analysis are listed in table 1.

Table 5: Elemental analysis results for the $\text{C}_{18}\text{H}_{36}\text{N}_4\text{NiCl}_2$

Atom	Calculated	Experimental
% C	49.35	52.68
% N	12.79	13.33
% H	6.28	9.37
% Cl	16.18	17.33

Since the sample was found to be extremely hygroscopic, the complex was dried for several hours at reduced pressures before elemental analysis. The determination of the nickel was complicated and results were inconclusive. However, the close agreement with the chloride contents lends some support to the hypothesis that a 2:1 complex was isolated. From Figure 58, we can also see that there is a plateau levelling at approximately 0.5

equivalents of Ni where any further addition of the metal causes no further changes in chemical shifts. This stoichiometry supports the 2:1 complex proposed earlier. Considering the reduced size of the cavity, the seven membered ring in **3** would be unable to accommodate the metal in a 1:1 complex, as in the metal cyclam complex. As a result the metal is bound by two molecule in a sandwich type compound. Sandwich compounds have been well documented.³⁸

Table 6 : ^{13}C nmr chemical shifts for **3** during the titration with $\text{NiCl}_2 \cdot \text{H}_2\text{O}$

CARBON POSITION	CHEMICAL SHIFT DUE TO EQUIVALENTS Ni^{a}				
	0.00	0.12	0.24	0.36	0.48
C_1	67.4	66.8	66.1	65.7	65.2
C_2	46.8	46.5	46.2	46.0	46.0
C_3	30.7	30.0	29.5	28.7	28.4
C_4	33.9	33.4	33.2	33.0	33.0
C_5	26.1	25.9	25.6	25.5	25.4

a : All the chemical shifts are reported from a methanol solution of **3**

Figure 59: Plot of the observed ^{13}C nmr chemical shift upon titration of **3** with $\text{NiCl}_2 \cdot \text{H}_2\text{O}$ in methanol

The persistence of the green color during the titration of **2** with a methanol solution of $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ indicates the absence of any complexation by the ligand.

Complexation properties of the tosylated and benzylated dibenzo 18-crown-6

In our work on the dibenzo diazatetraoxa 18-crown-6 ethers, the tosylated and benzylated precursors contained some relevant information that could be extracted using relaxation studies. The three products of concern are **8**, **9** and **14**.

The relationship between relaxation time, T_1 , and correlation time is given in Figure 59.⁷⁵

Figure 59 : Correlation time vs relaxation time graph

The correlation time is defined as the average time for a molecule to rotate through one radian. Since large molecules will tumble relatively slowly in solution because of their size and the reordering of the solvent cage, Figure 59 indicates that these molecules will relax more efficiently than small rapidly tumbling molecules. Table 2 indicates that this is indeed the case for product **8** and **9**. The T_1 's for **9** are all higher than those of **8**. If the tumbling of the molecule is isotropic, the T_1 is found to be inversely proportional to the number of directly attached protons. The observed T_1 's for the aromatic C-H and the methine C-H₂ follows the expected 2:1 ratio. The T_1 for C-H, CH₂ are 2.06 ± 0.05 and 1.07 ± 0.04 respectively.

From Table 1 in appendix II, the absence of any change in the relaxation times of

the carbons in **8**, **9** and **14** indicates no complexation with KNCS. Because of the extended chair conformation of **9**, molecular models show that the donor sites are not in favorable positions for complexation. However, earlier results with the n.O.e. have shown that the 18 membered systems are found in a more flexible planar conformation. The more flexible 18 membered systems provide a more favorable conformation for complexation because the central cavity region allows for greater interaction with a guest metal. Since complexes have been isolated with the dibenzo diaminotetraoxa 18-crown-6, it is assumed that the substituents on the macrocyclic rings are a major factor in the overall inability of **8** and **14** to form complexes with potassium.

CONCLUSIONS

The initial attempts at the synthesis of dicyclohexyl tetraaza-14-crown-4 ethers involved cyclization by aminolysis of **1**. However even at high dilution, the dominance of the intramolecular reaction resulted in the formation of the trans fused 7 membered ring. Attempts at inducing the intermolecular reaction by using smaller nucleophiles in the final cyclization step did not provide the desired 14 membered ring product. The stability of the **1** was achieved by complexation with nickel in methanol. The complex was characterized using ultra-violet and infra-red spectroscopy. The cyclization of the complex with 1,2-diaminocyclohexane has provided some hope for the eventual synthesis of the target molecule. Although the product could not be isolated at this time, M.S. data indicates the presence of the 14 membered ring.

On the other hand, **6** is stable in solution. Although the cyclization of this product could not be achieved with trans-1,2-diaminocyclohexane, it does present the required stability for studies using various nucleophiles for cyclization.

The presence of the amide functional group in the macrocyclic ring provides some interesting results when we compare **2** and **3**. There is a distinct difference in the binding abilities of the ligands with nickel. The ligand containing the amide in the macrocyclic ring structure, does not form a complex with nickel. However, **2** was found to give a 2:1 complex with the metal.

The initial scheme used to synthesize the dibenzo diaza-18-crown-6 provided the tosylated precursor in a 30% yield. Many attempts at detosylating the product failed to give the target molecule. The use of LiAlH_4 was found to be effective for detosylation of the 18 membered ring. However, the yields were too low for practical purposes. The second scheme provided a 10% yield of the benzylated precursor. Using Palladium black, the product was debenzylated in a quantitative yield. However, all attempts at the hydrogenation of the dibenzo diaza-18-crown-4 could were unsuccessful.

The precursors to the dibenzo diaza-18-crown-4 have provided some interesting results about their conformations of these systems in solution nmr. Conformational differences between the 9 and 18 membered product give rise to notable chemical shift differences. The differences are a result of the stereochemical interactions of the shielding nuclei.

EXPERIMENTAL

1.0 GENERAL :

Synthesis:

All reagent grade solvents were doubly distilled over KOH.

Other solvents were obtained as absolute (having less than 0.01 % water)

THF and DMF were dried over Na metal.

All starting materials were obtained from freshly opened bottles or distilled at reduced pressures. Solids were recrystallized from appropriate solvents.

Purification:

All purification of synthesized products involved the use of Silica gel -Grade 923 (mesh 100-200) activated at 120 °C for 48 hours.

Instrumentation:

^{13}C (50.3 MHz) NMR spectra : Varian XL-200 - All NMR shifts are listed in ppm

^1H (300 MHz) NMR spectra : Varian EM-360A- All NMR shifts are listed in ppm.

Mass Spectrometry : VG 7070E

Infra red spectra : Perkin-Elmer 1600 series FTIR

The visible spectra : Perkin-Elmer Lambda 4B UV/VIS

Freezing Point Depression:

The apparatus used for freezing point depression included a test tube containing a thermometer and a wire stirrer supported by a cork. Once the solids were dissolved, readings were taken starting at 190.0 °C. In order to avoid any cooling due to air convection, the test tube was immediately placed in a larger test tube supported by another cork sleeve. As the solids were cooling the solution was continuously stirred to ensure consistent cooling.

Nomenclature:

The nomenclature used in naming the following cyclohexane derivatives and crown ethers is used throughout the literature.

2.0 ATTEMPTED SYNTHESIS OF TRANS DICYCLOHEXANO-TETRAAZA-14-C-4

2.1 1,2 Bis[3-methyl ester-amine] cyclohexane (**1**)

A mixture of ($\pm$) trans-1,2 diaminocyclohexane (3.30 g, 29.0 mmol) and methyl acrylate (4.97, 58.0 mmol) was refluxed in absolute methanol (260 ml) under N_2 for 48 hours. The solvent was removed by evaporation under reduced pressure and the resulting oil was purified by column chromatography using silica gel (100-200 mesh). The product was eluted with methanol (2% triethylamine). The product eluted with an R_f value of 0.22 on a TLC, using methanol as the mobile phase. The resulting product was obtained as an oil

Yield: 80%

IR (methanol) 1740 cm^{-1}

MS M^+ 287 m/z

^{13}C NMR

2.2 Attempted Cyclization of **1** using ($\pm$) trans, 1,2-cyclohexanediamine (**2**)

A mixture of **1** (1.00 g, 3.5 mmol) and ($\pm$) trans 1,2-diaminocyclohexane (0.40 g, 3.5 mmol) was refluxed for 48 hours in absolute methanol under N_2 . The reaction was monitored by observing the disappearance of the ester peak at $1740\ cm^{-1}$ in the IR. After removal of the methanol under reduced pressure, an orange solid was obtained. This material was purified by column chromatography using silica gel (100-200 mesh) and MeOH (2% trimethylamine). The final product was obtained by recrystallization in ethyl acetate to yield white needles.

Yield 80%

Melting point: 146-147 $^{\circ}C$

IR: (KBr) amide 1664 cm⁻¹

MS: 168 m/z

¹³C NMR **3.0 Cyclohexyl diaza 7-crown-2**

The racemic mixture of **2** (0.40g, 1.2 mmol) was dissolved in dry THF (2 mL) under argon and cooled in an ice bath. The reducing agent BH₃-THF (13.5 mL) was added and the resulting mixture stirred for 5 minutes. Slow addition of the diborane was necessary to reduce effervescence. The reaction mixture was allowed to warm to room temperature and then was refluxed at 95 °C for 48 hours. Hydrolysis was completed by adding H₂O (2mL) and 6N HCl (4mL). The mixture was refluxed for an additional 10 minutes. After cooling to room temperature, the mixture was extracted with ether (50mL). The aqueous layers were combined and heated with 25% NaOH and then extracted with CH₂Cl₂(50mL). The organic phase was dried over Na₂SO₄ and the solvent removed to yield an oil.

Yield: 44%

MS: 154 m/z

IR (methanol) no amide peak at 1664 cm⁻¹¹³C NMR:

4.0 Attempted Cyclization of **1** using various nucleophiles

4.1. Attempted Cyclization of **1** using 1,2 diethylamine

A mixture of (**1**) (0.80 g, 2.8 mmol) and 1,2 diethylamine (0.16 g, 2.7 mmol) was refluxed in absolute methanol(200 ml) for 48 hours under N₂. The reaction was monitored by observing the disappearance of the ester peak at 1740 cm⁻¹. After removal of the solvent by rotary evaporation, the solid was recrystallized in ethyl acetate to yield a white solid

Yield: 85%

Melting Point: 146-148 °C

IR (methanol) amide 1664 cm⁻¹

MS: 168 m/z

¹³C NMR: same as in 2.2

4.2 Attempted cyclization **1** using ethylamine

A mixture of **1** (1.10 g, 3.8 mmol) and ethylamine (0.17 g, 3.8 mmol) was refluxed in absolute methanol (600 ml) for 48 hours. Using the same work up as in 2.1, a white solid was obtained.

Yield: 60%

Melting Point: 146-147 °C

IR (methanol) amide 1664 cm⁻¹

MS: 168 m/z

¹³C NMR: same as 2.2

4.3. Attempted cyclization of **1** using tosylated intermediates.

4.3.1 1,2 bistosylated cyclohexane diamine (**4**)

The (±) trans 1,2 diaminocyclohexane(3.00 g, 26.3 mmol) was dissolved in CH₂Cl₂(200 ml) and cooled to 3 °C on an ice bath. Pyridine(4.16 g, 50.0 mmol) was then added to the flask. While stirring the reaction mixture, three equivalents of p-toluenesulfonyl chloride(15.5 g, 78.8 mol) was slowly added being careful to maintain temperature below 5 °C. The yellow slurry was allowed to sit for 2 hours at room temperature. Water was added to remove the pyridinium hydrochloride formed. The excess pyridine was removed by adding an additional aliquot of 1 M HCl (100 ml). The CH₂Cl₂ phases were collected and dried over MgSO₄. The solvent was removed by

rotary evaporation to yield the crude material. The product was recrystallized from CH_2Cl_2 /Hexane mixture to yield white plates.

Yield: 74%

Melting Point: 146-147 °C

^{13}C NMR:

4.3.2 Attempted synthesis of 1,2-bis[3-methyl ester-N-tosylamino] cyclohexane

Method 1

4.3.2a

The bistosylate **1** (0.85 g, 2.0 mmol) was dissolved in a solution of absolute methanol (300 ml) and dichloromethane (20 ml). The CH_2Cl_2 was added to cause dissolution of the methanol insoluble tosyl material. The methyl acrylate (0.17 g, 2.0 mmol) was then added in one portion. The solution was allowed to reflux for 76 hours. Evaporation of the solvents yielded a thick semi solid. The solid was dissolved in methanol to yield the starting material methyl acrylate.

The methanol insoluble product was recrystallized in CH_2Cl_2 /Hexane mixture to yield starting tosylate material as white plates.

Method 11

1 (0.69 g, 2.41 mmol) was dissolved in pyridine (40 ml) and cooled to 0 °C. The p-toluenesulfonyl chloride (3.8 g, 19.3 mmol) was added to the solution making sure to maintain a temperature below 5 °C. The dark orange solution was placed in the refrigerator at 5 °C for 76 hours. No pyridinium hydrochloride needles were formed. The reaction was not worked up any further.

5.0 Attempted Cyclization of 1 using a Nickel Template

5.1 trans tetraaza cyclohexyl 14-crown-4

1 (0.74 g, 2.6 mmol) was dissolved in absolute methanol (400 ml) and the $\text{NiCl}_2 \cdot \text{H}_2\text{O}$ (0.61 g, 2.6 mmol) was added over a 30 minute period. Upon the first addition of the nickel, the solution turned violet. The solution was allowed to sit for one day undisturbed. The ($\pm$) trans 1,2 trans diaminocyclohexane (0.30 g, 2.6 mmol) was added in one portion and the solution refluxed for 4 days. The solution was cooled and NaCN (0.49 g, 10.2 mmol) was added to the violet solution. Immediately the nickel cyanide complex was formed turning the solution yellow. The solution was refluxed for 2 hours. The reaction mixture was evaporated at reduced pressures to yield a yellow solid. CHCl_3 (20 ml) was added to the flask and transferred to a fritted funnel. The solid was washed with two more portion of CHCl_3 (20 ml). The organic extract were dried over NaSO_4 and evaporated to yield a yellow oil. The oil was purified by column chromatography to yield an oil. The crude product was repeatedly purified using silica gel columns and the resulting crude product isolated gave an $R_f=0.37$ (TLC-eluting with MeOH). The product contained other impurities.

Yield : 2%

M.S.: $M+1$ 337 m/e (C.I.)

6.0. Attempted Synthesis of cis tetraaza cyclohexyl-14-crown-4

6.1 Isolation of cis 1,2-diaminocyclohexane

The cis and trans 1,2 diaminocyclohexane (125.0 g, 1.09 mol) was dissolved in MeOH (800 mL) and added to a solution of $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ (130.0 g, 0.55 mol) in MeOH (2220 ml). The mixture was stirred at room temperature for 3 hours. The yellow solid was filtered and washed with MeOH until washings were colorless.

Yield: 21%

The yellow solid was then dissolved in excess H_2SO_4 . Once dissolved, the green solution was treated with MeOH to yield the white sulfate salt.

Yield: 90% Melting point: $>300^\circ\text{C}$

A large excess of 25% NaOH solution was added to an aqueous solution of the sulfate salt. The aqueous layer was then extracted with CHCl_3 . The CHCl_3 was then dried over K_2CO_3 . The CHCl_3 was then evaporated to yield the cis diamine as a clear colorless oil.

Derivative: cis diamine.HCl

Melting pt.: 304-305 °C (Lit 304-309 °C)

^{13}C of cis diamine	CH-N	52.3
	$\text{C}^*\text{H}_2\text{-CH-N}$	31.4
	$\text{C}^*\text{H}_2\text{-CH}_2\text{-CH-N}$	28.6

6.2 Synthesis of cis 1,2 Bis(3 methyl ester-amine) cyclohexane (6)

Methyl acrylate (4.43g, 0.051 mol) was added to a solution of cis 1,2 diaminocyclohexane (2.83g, 0.025 mol) in absolute MeOH (400 mL) and refluxed for 3 days under Argon. The solvent was removed and the resulting oil was purified by column chromatography using silica gel (100-200 mesh) and MeOH(2% tetramethylamine) as the eluting solvent. The resulting product was colorless oil.

Yield: 60 %

IR (methanol) 1736 cm^{-1} MS: $\text{M}+1$ @ 287 m/z (C.I.) ^{13}C NMR: 6.3 Attempted cyclization of 6 in Methanol

A mixture of **6** (0.75 g, 2.6 mmol) was dissolved in absolute methanol (500 ml). The ($\pm$) trans 1,2 cyclohexanediamine (0.30 g, 2.6 mmol) was added and the solution was refluxed for seven days. The methanol was removed in vacuo to yield starting materials.

IR (methanol) : 1730 cm^{-1} ^{13}C NMR : same as in 5.2

6.4 Attempted cyclization of 6 in THF

6 (0.95 g, 3.3 mmol) was dissolved in dry THF (30 ml). The (±) trans 1,2 diaminocyclohexane (0.38 g, 3.3 mol) was added in one portion. After refluxing for six days, the reaction was worked up as in 5.3 to yield starting materials.

IR (methanol) 1730 cm^{-1}

^{13}C NMR :same as in 5.2

6.5 At, mpted cyclization of 6 in Glyme

6 (0.50 g, 1.7 mmol) was dissolved in dimethyl ether (40 ml) along with (±) trans 1,2 diaminocyclohexane (0.19 g, 1.7 mmol). The solution was refluxed for six days. The solution was then evaporated to yield starting materials.

IR : 1730 cm^{-1}

^{13}C NMR : same as in 5.2

7.0 Attempted synthesis of the cis isomers of 5,6,14,15-Dicyclohexyl-4,7,13,16-tetraoxa-1,10-diazacyclooctadeca

METHOD A

7.1 1,2 bis[2-(p-toluenesulfonyloxy)ethoxy]amine (7)

A solution of diethanolamine (7.063 g, 67.2 mmol) in pyridine (106 ml) was cooled to $3\text{ }^{\circ}\text{C}$. The pyridine was taken from a freshly opened bottle. The toluenesulfonyl chloride (43.46 g, 0.23 mol) was then added in small portions, ensuring that the temperature never increased above $5\text{ }^{\circ}\text{C}$. After the solution was complete, the flask was stored in the refrigerator until no more crystals of the pyridinium hydrochloride formed (24 hours). The entire mixture was then poured into an ice-water mixture (570 g). The water was decanted and the solid slurry was placed in a separatory funnel and extracted with three portions of water (100 ml) followed by three portions of diethyl ether (100 ml). The ether was then washed with two portions of 1 M HCl (150 ml). The organic phase was then treated with K_2CO_3 (5 g) and Na_2CO_3 (5 g) and then evaporated to yield a dark orange viscous oil. The oil was recrystallized in a 60:40 ethyl acetate/hexane mixture to yield a colorless solid.

IR: no OH or NH bands

Melting Point : 78-80 °C (Lit. val. - 80 °C)⁶¹¹³C NMR

7.2 N,N'-ditosyl-5,6,14,15-dibenzo-4,7,13,16-tetraoxa-1,10-diaza-cyclooctadeca-5-14-diene

Catechol(6.4 g, 58.1 mmol)was dissolved in dry DMF (90 ml). To this solution, K₂CO₃(4.0 g, 29.0 mmol) was added to obtain the salt. **2** (18.0 g, 29.0 mmol) was then dissolved in DMF(40 ml) and slowly added to the salt of catechol. After 24 hours, another addition of K₂CO₃(4.0 g, 29.0 mmol) along with the final equimolar amount of **2** (18.0 g, 29.0 mmol) was completed. The reaction was allowed to reflux for another 48 hours. The reaction was stopped and the solvent removed under reduced pressure to yield a black tar. This solid was extracted with 10% NaOH and the CH₂Cl₂ phase was then dried over MgSO₄. The insoluble material was then washed in MeOH to yield a light brown solid. This solid was recrystallized in CH₂Cl₂/Acetone to give a colorless solid

Yield: 20 %

Melting Pt: 218-219 °C (LIT- 215-216 °C)⁶¹

MS: M+ @ 358 m/z

^{13}C NMR: **8** ^1H NMR: **8**

2.39 (6 H, singlet,)	6.78 (8H, split AA'BB'system)
3.72 (8 H, triplet, $J_{AB}=6.2$ Hz)	7.66 (4H, multiplet)
4.10 (8 H, triplet, $J_{AB}=6.2$ Hz)	7.69 (4H, multiplet)

The methanol soluble product was recrystallized in heptane to yield a white solid

Yield : 30 %

Melting Point: 172-173 °C (Lit- 174-175 °C)⁶¹

 ^{13}C NMR: **9** ^1H NMR: **9**

2.39 (3 H, singlet)	6.96 (4H, singlet)
3.48 (4 H, AA'BB' multiplet)	7.25 (2 H, multiplet of a doublet)
4.39 (4 H, AA'BB' multiplet)	7.62 (2 H, multiplet of a doublet)

7.3 5,6,4,15-Dibenzo-4,7,13,16-tetraoxa-1,10-diazacyclooctadeca-5,14-diene (10)

7.3.1 Method I:

A mixture of **8** (1.09 g, 0.15 mmol), phenol (0.5 g, 5.2 mmol) and glacial acetic acid saturated with HBr (4 ml) was stirred for 48 hours at 80 °C. The solution was concentrated under reduced pressure and then diluted with water (10 ml). The solution was extracted with CH₂Cl₂, and the aqueous phase was evaporated to dryness. No solids were obtained. The reaction product was not treated further.

7.3.2 Method II

To a three necked quickfit 25 mL flask, Hexamethyl phosphotriamide (HMPT) (2 mL) was added along with Na metal (0.046g, 0.02 mol). This solution was cooled to 0 °C. The solution went a deep blue color. After all the Na metal had dissolved, **8** (1.0 g, 0.15 mmol) was dissolved in HMPT (2 mL) and added to the reaction mixture. The reaction was allowed to stir for 5 hours at 3 °C. The reaction mixture was distilled at approximately 3 torr to remove most of the solvents. The undistilled material was extracted with several portions of diethyl ether (20 mL). The ether was dried over MgSO₄ and evaporated to yield no solid. The reaction mixture was then acidified with 1 M HCl (1 mL) and extracted with several portions of diethyl ether (20 mL). The aqueous phase was filtered and treated with a 10 % aqueous NaOH (20 mL). The solution was extracted with CH₂Cl₂ (40 mL) and evaporated to yield a white solid.

Yield: 20 %

Melting Point: 220-221 °C

¹H NMR : same as in 6.2 for **8**

7.3.3 Method III

8 (1.03 g, 0.15 mmol) was dissolved in dry THF (100 mL) along with LiAlH₄ (1.40 g, 60 equiv) under argon. After cooling, the mixture was diluted with THF (100 mL) and quenched with a 5% LiOH (20 mL). Filtration and evaporation provided a black tar. This residue was refluxed in methanol (100 mL) for 24 hours to yield a solid white material. The solid was recrystallized in toluene to yield a white solid. The solid was found to be air sensitive and was kept under argon at all times.

Yield: 2%

Melting Point: 174-175 °C

^{13}C NMR :

METHOD B

7.5 1,2 Bis (2-hydroxyethoxy)benzene (11)

2-Chloroethanol (37.7g, 0.54 mol) was added slowly to a solution of pyrocatechol (20.5g, 0.18 mol) and sodium hydroxide (18.8 g, 0.54 mol) in 95% ethanol (181 mL). The mixture was stirred and refluxed for 48 hours. The reaction mixture was evaporated to give a thick black oil. The product was washed with several portions of 10% NaOH until the aqueous phase was colorless. The CH_2Cl_2 phase was then dried over Na_2SO_4 and evaporated to give a light brown solid. The solid was recrystallized in 60:40 EtOAc/Hexane.

Yield: 45%

Melting point: 79-80 °C (Lit 80 °C)⁶⁵

7.6 1,2 Bis[2-(p-toluenesulfonyloxy)-ethoxy]benzene (12)

A round bottom flask, equipped with a stirrer, condenser and thermometer was charged with dry pyridine (109 mL) and 1,2 bis[2-(hydroxyethoxy)]benzene (19.6g, 0.099 mol). The mixture was cooled to 3 °C and p-toluenesulfonyl chloride (40.2g, 0.21 mol) was added at such a rate that the temperature did not exceed 15 °C. Agitation was continued for 3 hours. A chilled solution of 1 M HCl (90 mL) in water was added dropwise to the flask. Solid ditosylate was filtered off as a white solid and air dried.

Yield: 86%

Melting point: 94 °C (Lit. 95-96 °C)⁶⁵

7.7 1,2-bis[2-(N-benzylamino)ethoxy]benzene (13)

Benzylamine (60.3g, 0.56 mol) was added to a solution of **12** (28.2 g, 0.055 mol) in xylene (560 mL). The resulting mixture was agitated at reflux for 16 hours, and then cooled to precipitate the white fibrous byproduct which was removed by filtration. Concentration of the filtrate under vacuum gave an orange oil. The diamine was used without purification in the next step.

7.8 N,N'-Dibenzyl-5,6,14,15-Dibenzo-4,5,13,16-tetraoxa-1,10-diazacyclooctadeca-5,14-diene (14)

The crude **13** (28.0g,), xylene (700 mL), **12** (28.2g, 0.055 mol) and n-propylamine (14.9g, 0.10 mol) were mixed and refluxed for 24 hours. The reaction mixture was cooled and the solution was decanted from the white fibrous byproduct and concentrated under vacuum to yield an orange oil. Boiling methanol was used to extract the impurities leaving undissolved the white solid.

Yield: 10% Melting point: 175-176 °C (Lit 176 °C)⁶⁵

¹³C NMR- see Table 3 (page 49)

79 5,6,4,15-dibenzo-4,7,13,16-teraoxa-1,10-diazacyclooctadeca-5,14-diene (15)

7.9.1 Attempted debenzoylation -Method I

14 (0.40g, 0.78 mmol) was dissolved in dry THF (30 mL). The catalyst, palladium/carbon (0.10 g) was added and the mixture hydrogenated at 50 °C and 800 psi for 20 hours. Solids were removed by filtration. The THF was then evaporated at reduced pressure to yield starting material.

Yield: 80% **Melting point:** 175-176 °C

^{13}C NMR : same as 7.8

7.9.2 Method II

14 (0.16g, 0.31 mmol) was added to a solution of methanol (12 mL) along with the catalyst palladium black (0.16g). Formic acid 4,4% (0.8 mL) was added to cause dissolution of the dibenzyl product. The solution was stirred at room temperature over argon for 3 days. The catalyst was removed by filtration and the filtrate evaporated to yield a white methanol soluble product.

Yield: 95% Melting Point 175-176 °C (Lit 175-176 °C)⁶⁵

^{13}C NMR : same as in 6.3.4

9.0 Attempted synthesis of 5,6,14,15-Dicyclohexyl-4,7,13,16-tetraoxa-1,10-diazaacyclooctadeca

9.1 Method 1

15 (0.10 g, 0.28 mmol) was dissolved in butanol (40 mL) and the ruthenium catalyst (0.100g) was added to the solution. The mixture was hydrogenated at 60 °C for 20 hours at 1000 psi. Filtration of the solids and evaporation yielded starting material.

Yield: 95%

Melting point: 175-176 °C

¹H NMR: same as in 6.2 for **8**

9.2 Method 2

15 (0.14g, 0.40 mmol) was dissolved in butanol (30 mL). The rhodium catalyst (0.020g) was added to the solution and hydrogenated at 65 °C and 1000 psi for 20 hours. Filtration of the solids and evaporation yielded starting material.

Yield: 98%

Melting Point 175-176 °C

¹H NMR: same as in 6.2 for **8**

REFERENCES

1. Berezin, B.D., Coordination Compounds of Porphyrin and Phthalocyanines, John Wiley & Sons, New York, 1983.
2. Lever, A.B.P, Hempstead, M.R., Leznoff, C.C., Liu, W., Melnik, M., Nevin, W.A., Seymor, P., Pure and Applied Chem., 58(11), 1467-1476, (1986).
3. Alphen, J.V., Rec Trans. Chem. Phys Res, 55, 835, (1936).
4. Hancock, R., Pure & Applied Chem, 58(11), 1445-1452, (1986).
5. Krakowiak, K., Bradshaw, J., Zamecka-Krakowiak. D.; Chem Rev., 89, 929, (1989)
6. Vogtle, F. (Editor), Painter, G.K., Pressman, B.C., Host Guest Complex Chemistry, Springer Verlag, p.33, (1982).
7. Pederson, C.J., J. Am. Chem. Soc., 89, 7017, (1967).
8. Arnold, K.A., Echegoyan, L., Frencezek, F.R., Gandour, R.D., Gatto, V.J., White, B.D., Gokel, G.W., J. Am. Chem, Soc., 109, 3716-3721, (1987).
9. Zolotov, Y.A., Malofeeva, G.I., Petrukhim, D.M., Timberbaev, A.R., Pure & Applied Chem., vol 59(4), 497-509, (1987).
10. Navratil, J.D, Pure and Applied Chem, 58(6), 885-888, (1986).
11. Izatt, R.M., Lindh, G.C., Bruening, R.L., Bradshaw, J.S., Lamb, J.D., Christensen, J.J., Pure and Applied Chem, 58(11), 1453-1450, (1986).
12. Gokel, G.W., Durst, H.D.; Synthesis, 168, (1976)

- 13 a) Peacock S.C., Cram, D.J., J. Chem. Soc., Chem Commun, 282, (1976).
- b) Kimura, E., Saiconalca, A., Namoto, M.; Biochim. Biophys Acta, 67, 172-179, (1981).
- c) Kimura, E., Pure and Applied Chem, 58(11), 1461-1466, (1986).
- 14 a) Curtis, W.D., Laidler, D.A., Stoddart, J.F., Jones, G.H., J. Chem. Soc., Chem Commun, 833-835. (1975).
- b) Nishinaga, A., Tomita, H., Nishizawa, L., Matsuura, T., J. Chem Soc., Dalton Trans, 1504, (1981).
- 15 a) Curtis W.D., King, R.M., Stoddart, J.F., Jones, G.H., J.C.S. Chem. Commun., 284, (1976).
- b) Kodem, T.A., Topics Curr. Chem, 121, 157, (1984).
16. Lotz, J., Kodem, T.A., J. Chem. Soc. Chem. Commun., 15, (1977).
- 17 Lotz, J.J., Lodem, T.A., Inorg, Chem, 25, 404, (1986).
18. Buchanan, G.W., Kirby, R.A., Charland, J.P., J. Am. Chem Soc., 110, 2477. (1988).
- 19 a) Kodama, M., Kimura, E.; J. Chem. Soc., Dalton Trans, 116. (1976).
- b) Kodama, M., Kimura, E., J. Am. Chem Soc., 91, 6540. (1969).
- c) Kodama, M., Kimura, E., J. Chem. Soc., Dalton Trans, 1720, (1976).
20. Buchanan, G.W., Kirby, R.A., Bourque, K., Morat, C., Can. J. Chem., 66, 2204-2209. (1988).

21. Kodama, M, Kimura, E., J. Chem. Soc., Dalton Trans, 694, (1981).
22. Melson, G.A. (Ed.), Coordination Chemistry of Macrocyclic Compounds , Plenum Press, New York, 1979
23. Frensdorff, H.K., J. Am. Chem. Soc., 93(3), 600-605, (1971)
24. Patai, S (Ed.), The chemistry of ethers, crown ethers, hydroxyl groups and their sulfur analogues, Wiley and Sons, New York, 1988
25. Burden, I.J.; Coxon, A.C.; Stoddart, J.F.; Wheatly, C.M.; J. Chem. Soc.; Perkin Trans I, 220,1977
- 26 a) Knochel, A.; Oehler, J.; Rudolph, G.; Simwell,V., Tetrahedron, 33, 119, (1977)
b) Izatt, R.M.; Eatough, D.J.; Christensen, D.; Struct. Bonding, 16, 161, (1973)
- 27 a) Buchanan, G.W., Bourque, K., Bovencamp, J.W., Rodrique, A., Can J Chem, 63, 2747, (1985).
b) Buchanan, G.W., Kirby, R.A., Tetrahedron Letters, 28(14), 1507-1510. (1987).
28. Buchanan, G.W., Khan, M.Z., Ripmeester, J.A., Bovencamp,J.W., Rodrique, A., Can J. Chem, 65, 2564, (1987).
29. Kleinpeter, E.; Gabler, M.; Schroth, W.; Magnetic Resonance in Chemistry, 26, 387-391, (1988)
- 30 Kleinpeter, E., Gabler, M., Schroth, W., Magnetic Resonance, 26, 380, (1988).
31. Cabbiness, D.K., Margerum, D.W., J. Am. Chem. Soc, 91, 6540, (1969).

32. Hinz, F., Margerum, D.W., Inorg. Chem., 13, 2911, (1974).
- 33 a) Kodama, M., Kimura, E.; J. Chem. Soc. Chem. Commun., 326, (1975).
- 34 a). Anichini, A., Fabrizzi, L., Paolettia, P., J. Chem. Soc. Chem. Commun. , 244, (1977).
- b) Gallori, E., Martini, E., Micheloni, M., Paoletti, P., J. Chem. Soc. Dalton Trans I, 1722, (1980).
- 35 Frensdorff, H.K., J. Am. Chem. Soc., 93, 600, (1971)
36. Herceg, P. M., Weiss, R., Acta Cryst,29, 542, (1973).
37. Hancock, R.D., Dobson, S.M., Evers, A., Wade, P W., Ngwenya, M.P., Boeyens, J.C.A., Wainwright, K.P., J. Am. Chem. Soc., 110, 2788-2794, (1988).
- 38 Pederson, C.J., J. Am. Chem. Soc, 92, 391, (1970).
- 39 Dunitz,J.D., Hemmerich, P., Jorgensen, C.K., Reiner, D. (Editors); Structure and Bonding, 16, (1973).
- b) Lehn, M.; J. Chem. Soc., Perkin Trans I, 767, (1977).
40. Hancock, R.D., Pure & Applied Chem, 58(11), 1445-1452, (1986).
41. Martin, L.Y., Dehayes, L.J., Zompa, L.J., Buisch. D.H., J. Am. Chem. Soc, 96, 4046, (1974).
42. Coxon, A.C., Laidler, D.A., Pettman, R.B., Stoddart, J.F., J. Am. Chem. Soc, 100, 8260, (1978).
43. Martin, Y.L., Dettayes, J., Zompa, L.Z., Busch, D.H., J. Am. Chem. Soc.,96, 4046 , (1974).

44. Daley, N.K., Synthetic Multidentate Macrocyclic Compounds, Academic Press, New York, (1978)
45. Derome, A.: Modern NMR techniques for Chemistry Research Pergamon Press, New York, (1978)
46. Abraham, R.J., Loftus, P.; Proton and Carbon-13 NMR Spectroscopy, John Wiley and sons, New York. 1978
47. Kodama T.; Carleton University 498 thesis, 1987
- 48 a). Dietrich, B.; Journal of Chemical Education, 62(11), 954-964, (1985).

b) Gokel, G.W., Dishong, D.M., Schultz, R.A., Gatto. V.J.; Synthesis, 997-1012, (1982).

c) Gradshaw, J.S., Stott, P.E., Tetrahedron, 36, 461-510, (1980).
- 49 a). Hodges, K.D., Barefield, E.K., Wagner, F., Inorg, Chem., 15(6), 1370-1377, (1976).

b) Barefield, E.K., Wagner, F.I, Herlinger, A.W., Dahl, A., R., Inorg Synthesis, 16, 220, (1976).

c) Bosnich, B., Poon, C.K., Tobe, M.L., Inorg Chemistry, 4(8), 1102-1108, (1965).
50. Jolly, W. (Editor), Linday, L.F., Busch, D.H., Preperative Inorganic Reactions, 6 , Interscience, New York, (1971).
51. Beynor, J.H., Suanders, R.A., Williams, A.E.; The Mass Spectra of Organic Molecules, Elsevier Publishing Co., Amsterdam. p. 289, (1968).
52. Atkins, P.W.; Physical Chemistry, Freeman and Company, San Fransisco, 1982

53. Remers, W.A., Roth, R.H., Gibs, G.J., Weiss, M.J.; J. Org. Chem 36, 1232, (1971)
- 54 a). Curtis, N.F.; J. Chem. Soc. Dalton Trans., 1357, (1972).
- b) Curtis, N.F.; J. Chem. Soc. Dalton, Trans., 863, (1973).
- c) Curtis, N.F., Cook, D.F.; J. Chem. Soc. Dalton Trans., 1076, (1973).
55. Busch, D.H., Farmery, K., Goedken, V., Katovic, V., Melnyk, A.C., Sperati, C.R., Tokel, N.; Adv. Chem. Ser., 100, 44, (1971)
- 56 a) Billo, E.J.; Inorg. Chem., 20, 4019-4021. (1981).
- b) Hay, R.W., Pujari, M.P.; J. Chem. Soc. Dalton Trans, 1483-1489, (1973).
- 57 Potterfield, W.W., Inorganic Chemistry, Addison Wesley Publishing, 1984
58. Hay, R.w., Pujari, M.; J. Chem. Soc., Dalton Trans, 1485, (1986)
59. Saito, R., Kadani, Y.; Chem. Letters, 123-126, (1976).
60. Hogberg, Cram, D.J.; J. Org. Chem., 40(1), 151, (1975).
61. Cuvigney, Larcheveque, J. Organomet. Chem, 64, 315, (1974)
62. Pietraszkiewicz, M., Jurczak, J.; Tetrahedron, 40(15), 2967-2970, (1984).
63. Pederson, C.J., Bromels, M.H., U.S. Patent 3847949 (1974)
Dupont, C.A. 82, 73049 (1975)
64. ElAmin, B., Anantharamaiah, G., Royer, G., Means G.; J. Org. Chem., 44(19), 3442, (1979)
65. Buchanan, G.W., Bourque, K., Bovencamp, J.W., Rodrigue, A., Bannard, R.A., Tetrahedron Letters, 3963, (1984)

66. Freifelder, M., J. Org. Chem., 26, 1835, (1961)
67. Smith, H.A., Thompson, R.G., Advances in Catalysis, 9, 727, (1957)
68. Engelhard Industries of Canada Limited, Aurora, Ont, Canada
69. Noggle, J.H., Schirmer, R.E.; The Nuclear Overhauser Effect, Academic Press, New York, 1971
70. Busch, T., J. Chem. Soc. B, 1440, (1971)
71. Tribl is a complete molecular modeling software system. Originally written by Pensak, D., Wendoloski, J., Wasserman, Z. from Du Pont Experimental Station, the program has been modified by Herman, L., Roberson, W., and ApSimon, J. W. from Carleton University to run primarily on a TEKTRONIX 4014 graphics terminal.
- 72 a) Bovill, M.J., Chadwick, D.J., Sutherland I.O.; J. Chem. Soc., Perkin II, 1529, (1980)

b) Gue, R., Jacobson, S.H., Pizer, R.; J. Am. Chem. Soc., 100, 1150, (1985)
73. LAME is a fortran program that has been written for the XL-series to simulate spectra for spin-1/2 nuclei.
74. Levy, G., Nelson, G., Carbon-13 Nuclear Magnetic Resonance for Chemists, Wiley-Interscience, New York, 1972

APPENDIX 1

FREEZING POINT DEPRESSION

Table 1: Temperature data for three experiments using the solvent Camphor. The resulting average is given along with the standard deviation.

TIME (MINUTES)	TEMPERATURE (± 0.2 °C)			Average
	I	II	III	
0	190.0	190.0	190.0	190.0 ± 0.0
	182.0	182.0	181.5	181.8 ± 0.2
1	181.0	181.0	181.0	181.0 ± 0.0
	180.5	180.0	180.3	180.3 ± 0.2
2	180.0	179.5	180.0	179.8 ± 0.3
	180.0	179.0	179.5	179.5 ± 0.5
3	179.5	179.0	179.0	179.2 ± 0.3
	178.5	178.0	178.5	178.3 ± 0.3
4	177.0	177.0	177.5	177.2 ± 0.3
	174.5	175.0	174.0	174.5 ± 0.5
5	171.0	171.0	171.5	171.2 ± 0.3
	166.5	166.5	166.0	166.3 ± 0.5
6	162.0	163.0	162.5	162.5 ± 0.5
	158.0	159.0	158.5	158.5 ± 0.5
7	154.0	155.0	155.0	155.7 ± 0.5
	149.0	150.0	149.5	149.5 ± 0.2
8	145.5	146.0	146.0	145.8 ± 0.3
	142.0	142.0	142.5	142.2 ± 0.3
9	138.0	138.5	138.0	138.2 ± 0.3
	134.5	134.5	134.0	134.3 ± 0.3
10	131.5	132.0	131.0	131.5 ± 0.5

APPENDIX 1

Table 2: Temperature data for the freezing point depression of 1.00 g of **2** in 21.50g CAMPHOR. The resulting average of all three runs is given along with the standard deviation

TIME (MINUTES)	TEMPERATURE (± 0.2 °C)			AVERAGE
	I	II	III	
0	190.0	190.0	190.0	190.0 ± 0.0
1	169.2	169.0	169.0	169.1 ± 0.1
2	162.0	162.1	161.9	162.0 ± 0.1
3	158.0	158.0	158.1	158.0 ± 0.1
4	153.5	153.6	153.5	153.5 ± 0.1
5	147.5	147.5	147.6	147.5 ± 0.1
6	141.0	141.1	141.0	141.0 ± 0.1
7	133.5	133.6	133.5	133.5 ± 0.1
8	127.5	127.6	127.5	127.5 ± 0.1
9	121.5	121.5	121.5	121.5 ± 0.0
10	115.5	115.7	115.4	115.5 ± 0.2

APPENDIX 1

Table 3: Temperature data for freezing point depression of 2.10 g of **2** in 21.20 g CAMPHOR. The resulting average of all three runs is given along with the standard deviation.

TIME (MINUTES)	TEMPERATURE ($\pm 0.2^{\circ}\text{C}$)			
	<u>I</u>	<u>II</u>	<u>III</u>	AVERAGE
0	190.0	190.0	190.0	190.0 ± 0.0
1	153.5	153.2	153.5	153.5 ± 0.2
2	150.6	150.5	150.5	150.5 ± 0.1
3	149.1	149.0	149.1	149.1 ± 0.1
4	145.0	145.0	145.1	145.0 ± 0.1
5	139.1	139.0	139.0	139.0 ± 0.1
6	133.5	133.6	133.5	133.5 ± 0.1
7	128.7	128.4	128.4	128.5 ± 0.2
8	123.0	123.0	123.1	123.0 ± 0.1
9	117.1	117.1	117.0	117.1 ± 0.1
10	113.1	113.0	113.0	113.0 ± 0.1

2

OF/DE

2

MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS
STANDARD REFERENCE MATERIAL 1010a
(ANSI and ISO TEST CHART No. 2)

APPENDIX 1

Table 4: Temperature data for the freezing point depression of the 2.05 g Benzophenone in 20.00g CAMPHOR.
The resulting average of the three runs is given along with the standard deviation.

TIME (MINUTES)	TEMPERATURE ($\pm 0.2^{\circ}\text{C}$)			
	I	II	III	AVERAGE
0	190.0	190.0	190.0	190.0 ± 0.0
1	160.0	160.0	160.1	160.0 ± 0.1
2	156.6	156.6	156.4	165.5 ± 0.1
3	154.5	154.5	154.5	154.5 ± 0.1
4	152.6	152.4	152.5	152.5 ± 0.1
5	149.5	149.5	149.4	149.5 ± 0.1
6	145.0	145.0	145.1	145.0 ± 0.1
7	140.2	139.8	139.4	139.8 ± 0.4
8	139.5	139.3	139.6	139.5 ± 0.2
9	129.2	129.0	129.0	129.1 ± 0.1
10	123.5	123.5	123.4	123.5 ± 0.1

APPENDIX I

Graph #1 of the data in tables 1-4

APPENDIX 1

CALCULATIONS:

$$\Delta T = K_f m$$

$$M.W. = \frac{(g \text{ solute})(K_f)}{(kg \text{ solvent})(\Delta T)}$$

m = molality of solution

K_f = freezing point constant for the solvent

ΔT = freezing point lowering

a) BENZOPHENONE

$$M.W. = \frac{(2.17 \pm 0.01 \text{ g}) (37.7^\circ \text{C kg solvent/ moles solute})}{((2.12 \pm 0.01) \times 10^{-2} \text{ kg solvent})(20.0^\circ \text{C})}$$

$$= 193 \pm 1 \text{ g/mol} \quad (\text{actual value } 182.42 \text{ g/mol})$$

$$\text{relative error} = \frac{193 - 182.34}{182.42} \times 100\% = 5\%$$

b) UNKNOWN 1

$$M.W. = \frac{(1.00 \pm 0.01 \text{ g}) (37.7^\circ \text{C kg solvent/ moles solute})}{((2.15 \pm 0.01) \times 10^{-2} \text{ kg solvent}) (12.5^\circ \text{C})}$$

$$= 141 \pm 1 \text{ g/mol}$$

UNKNOWN 2

$$M.W. = \frac{(2.10 \pm 0.01 \text{ g}) (37.7^\circ \text{C kg solvent/ moles solute})}{((2.12 \pm 0.01) \times 10^{-2} \text{ kg solvent}) (25.0^\circ \text{C})}$$

$$= 149 \pm 1 \text{ g/mol}$$

$$\text{Average} = \frac{141 + 149}{2} = 145 \pm 1 \text{ g/mol}$$

APPENDIX II

Table 1 : ^{13}C Spin Lattice Relaxation Times (T_1)(s) for the
9 and 18 membered crown ethers. The T_1 for the free
ligands dissolved in CDCl_3 with 1.0 equiv of KNCS
(for numbering scheme see Figure 48)

Carbon Position	<u>9</u> ^a	<u>9</u> +KNCS	<u>8</u> ^a	<u>8</u> +KNCS	<u>14</u> ^a
C1	1.97±0.06	1.97±0.14	0.57±0.01	0.55±0.55	0.61±0.02
C2	2.06±0.05	1.93±0.31	0.53±0.02	0.050±0.02	0.54±0.01
C3	31.46±1.25		12.95±1.74	7.59± 1.42	10.86±2.99
C4	1.07±0.04	1.22±0.05	0.34±0.02	0.033±0.01	0.39±0.01
C5	1.00±0.03	1.11±0.07	0.31±0.02	0.32±0.01	0.36±0.01
C6	27.43±0.73		8.85±1.36	5.49±0.79	0.56±0.05
C7	2.15±0.05	1.90±0.22	0.96±0.03	0.97±0.03	8.39±0.91
C8	2.30±0.11	1.30±0.14	0.97±0.06	1.00±0.04	1.01±0.04
C9					1.01±0.04
C10	2.84±0.12	2.45±0.41	2.48±0.16	2.04±0.27	0.73±0.03

^a 0.1 M solution in CDCl_3

END

2002/9 1

FIN